

# *Weather and climate extreme events in a changing climate*

Book or Report Section

Accepted Version

Seneviratne, S. I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Luca, A. D., Ghosh, S., Iskandar, I., Kossin, J., Lewis, S., Otto, F., Pinto, I., Satoh, M., Vicente-Serrano, S. M., Wehner, M., Zhou, B. and Allan, R. ORCID: <https://orcid.org/0000-0003-0264-9447> (2021) Weather and climate extreme events in a changing climate. In: Masson-Delmotte, V. P., Zhai, A., Pirani, S. L. and Connors, C. (eds.) Climate Change 2021: The Physical Science Basis: Working Group I contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK, pp. 1513-1766. doi: <https://doi.org/10.1017/9781009157896.013> Available at <https://centaur.reading.ac.uk/101846/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/9781009157896.013>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

Caption: Table A.11.2. Synthesis table summarising assessments presented in Tables 11.4-11.21 for hot extremes (HOT EXT.), heavy precipitation (HEAVY PRECIP.), agriculture and ecological droughts (AGR./ECOL. DROUGHT), and hydrological droughts (HYDR. DROUGHT). It shows the direction of change and level of confidence in the observed trends (column OBS.), human contribution to observed trends (ATTR.), and projected changes at 1.5°C, 2°C and 4°C of global warming for each AR6 region. Projections are shown for two different baseline periods, 1850-1900 (pre-industrial) and 1995-2014 (modern or recent past)(see section 1.4.1 for more details). Direction of change is represented by an upward arrow (increase) and a downward arrow (decrease). Level of confidence is reported for LOW: *low*, MED.: *medium*, HIGH: *high*; levels of likelihood (only in cases of *high confidence*) include: L: *likely*, VL: *very likely*, EL: *extremely likely*, VC: *virtual certain*. See section 11.9, Tables 11.4-11.21 for details. Dark orange shading highlights *high confidence* (also including *likely*, *very likely*, *extremely likely* and *virtually certain* changes) increases in hot temperature extremes, agricultural and ecological drought, or hydrological droughts. Light orange indicates *medium confidence* increases in these extremes, and blue shadings indicate decreases in these extremes. *High confidence* increases in heavy precipitation are highlighted in dark blue, while *medium confidence* increases are highlighted in light blue. No assessment for changes in drought with respect to the 1995-2014 baseline is provided, which is why the respective cells are empty.

Sub-Region		OBS.	ATTR.	BASELINE: PRE-INDUSTRIAL			BASELINE: 1995-2014		
				1.5°C	2°C	4°C	1.5°C	2°C	4°C
Mediterranean (same region as for Europe) MED	HOT EXT.	↑ V. L.	↑ L.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ HIGH	↑ HIGH	LOW	↑ MED.	↑ HIGH
	AGR./ECOL. DROUGHT	↑ MED.	↑ MED.	↑ MED.	↑ HIGH	↑ V. L.			
	HYDR. DROUGHT	↑ HIGH	↑ MED.	↑ MED.	↑ HIGH	↑ V. L.			
Sahara SAH	HOT EXT.	↑ L.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
West-Africa WAF	HOT EXT.	↑ L.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	↑ MED.	LOW	LOW	LOW	LOW			
N.Eastern-Africa NEAF	HOT EXT.	↑ MED.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	↓ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↓ MED.			
Central-Africa CAF	HOT EXT.	LOW	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
S.Eastern-Africa SEAF	HOT EXT.	↑ MED.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
W.Southern-Africa WSAF	HOT EXT.	↑ L.	↑ L.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	LOW	↑ MED.	↑ L.	LOW	LOW	↑ HIGH
	AGR./ECOL. DROUGHT	↑ MED.	LOW	↑ MED.	↑ HIGH	↑ L.			
	HYDR. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ MED.			
E.Southern-Africa ESAF	HOT EXT.	↑ L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ HIGH	↑ L.	↑ E. L.	↑ MED.	↑ HIGH	↑ V. L.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	↑ MED.	↑ MED.	↑ HIGH			
	HYDR. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ MED.			
Madagascar MDG	HOT EXT.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ HIGH	↑ L.	↑ E. L.	MED.	↑ HIGH	↑ V. L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ HIGH			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ MED.			
Russian Arctic RAR	HOT EXT.	↑ V. L.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
Arabian-Peninsula ARP	HOT EXT.	↑ V. L.	↑ MED.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ HIGH	↑ V. L.	LOW	↑ MED.	↑ L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
W.C.Asia WCA	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	LOW	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ MED.			
W.Siberia WSB	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ HIGH	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
E.Siberia ESB	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
Russian-Far-East RFE	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
E.Asia EAS	HOT EXT.	↑ V. L.	↑ L.	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ L.	↑ V. L.	↑ V. C.	↑ HIGH	↑ L.	↑ E. L.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	LOW	↑ MED.			
	HYDR. DROUGHT	↑ MED.	LOW	LOW	LOW	LOW			
E.C.Asia ECA	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
Tibetan-Plateau TIB	HOT EXT.	↑ V. L.	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
S.Asia SAS	HOT EXT.	↑ HIGH	↑ HIGH	↑ V. L.	↑ E. L.	↑ V. C.	↑ L.	↑ V. L.	↑ V. C.
	HEAVY PRECIP.	↑ HIGH	LOW	↑ HIGH	↑ L.	↑ E. L.	↑ MED.	↑ HIGH	↑ V. L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	↓ MED.			

	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
S.E.Asia SEA	HOT EXT.	↑ HIGH	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ HIGH	↑ L.	↑ E.L.	↑ MED.	↑ HIGH	↑ V.L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
N.Australia NAU	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HOT EXT.	↑ HIGH	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ MED.	↑ HIGH	↑ V.L.	LOW	↑ MED.	↑ L
C.Australia CAU	AGR./ECOL. DROUGHT	↓ MED.	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HOT EXT.	↑ L.	↑ L.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
E.Australia EAU	HEAVY PRECIP.	LOW	LOW	LOW	↑ MED.	↑ L.	LOW	LOW	↑ HIGH
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ HIGH			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
S.Australia SAU	HOT EXT.	↑ L.	↑ L.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	LOW	↑ MED.	↑ L.	LOW	LOW	↑ HIGH
	AGR./ECOL. DROUGHT	↑ MED.	LOW	↑ MED.	↑ MED.	↑ HIGH			
New-Zealand NZ	HYDR. DROUGHT	↑ MED.	LOW	LOW	↑ MED.	↑ MED.			
	HOT EXT.	↑ L.	LOW	↑ L.	↑ V.L.	↑ V.C.	↑ HIGH	↑ L	↑ E.L.
	HEAVY PRECIP.	LOW	LOW	LOW	↑ MED.	L.	LOW	LOW	HIGH
S.Central- America SCA	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ MED.			
	HOT EXT.	↑ MED.	↑ MED.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
Caribbean CAR	HEAVY PRECIP.	LOW	LOW	LOW	LOW	LOW	LOW	LOW	LOW
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
N.W.South- America NWS	HOT EXT.	↑ L.	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	LOW	LOW	LOW	LOW	LOW	LOW
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
N.South- America NSA	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HOT EXT.	↑ L.	↑ MED.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ MED.	↑ MED.	LOW	↑ MED.	↑ MED.
South-American- Monsoon SAM	AGR./ECOL. DROUGHT	LOW	LOW	↑ MED.	↑ HIGH	↑ HIGH			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ HIGH			
	HOT EXT.	↑ L.	↑ MED.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
N.E.South- America NES	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ MED.	↑ MED.	LOW	↑ MED.	↑ MED.
	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	↑ MED.	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
S.W.South- America SWS	HOT EXT.	↑ L.	↑ MED.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	LOW	LOW	LOW	LOW	LOW	LOW
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ HIGH			
S.E.South- America SES	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ HIGH			
	HOT EXT.	↑ HIGH	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	↑ HIGH	LOW	↑ MED.	↑ HIGH	↑ L.	LOW	↑ MED.	↑ L
S.South- America SSA	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	↓ MED.	LOW	LOW	LOW	LOW			
	HOT EXT.	LOW	LOW	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
Greenland/Iceland GIC	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ HIGH	↑ V.L.	LOW	↑ MED.	↑ L
	AGR./ECOL. DROUGHT	LOW	LOW	↑ MED.	↑ HIGH	↑ HIGH			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ HIGH			
Mediterranean (same region as for Africa)	HOT EXT.	↑ V.L.	↑ L.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ HIGH	↑ HIGH	LOW	↑ MED.	↑ HIGH
	AGR./ECOL. DROUGHT	↑ MED.	↑ MED.	↑ MED.	↑ HIGH	↑ V.L.			
West&Central- Europe WCE	HYDR. DROUGHT	↑ HIGH	↑ MED.	↑ MED.	↑ HIGH	↑ V.L.			
	HOT EXT.	↑ V.L.	↑ L.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	↑ MED.	LOW	↑ HIGH	↑ L.	↑ E.L.	↑ MED.	↑ HIGH	↑ V.L.
E.Europe EEU	AGR./ECOL. DROUGHT	↑ MED.	LOW	LOW	↑ MED.	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	↑ MED.			
	HOT EXT.	↑ V.L.	↑ L.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
N.Europe NEU	HEAVY PRECIP.	↑ HIGH	↑ HIGH	↑ HIGH	↑ L.	↑ E.L.	↑ MED.	↑ HIGH	↑ V.L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HYDR. DROUGHT	↓ MED.	LOW	LOW	LOW	↑ MED.			
N.Central- America NCA	HOT EXT.	↑ L.	↑ MED.	↑ L.	↑ V.L.	↑ V.C.	↑ HIGH	↑ L	↑ E.L.
	HEAVY PRECIP.	LOW	LOW	↑ HIGH	↑ L.	↑ E.L.	↑ MED.	↑ HIGH	↑ V.L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ L.			
W.North- America WNA	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HOT EXT.	↑ L.	↑ MED.	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	↑ MED.	↑ HIGH	↑ V.L.	LOW	↑ MED.	↑ L
C.North- America CNA	AGR./ECOL. DROUGHT	↑ MED.	↑ MED.	LOW	↑ MED.	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	↑ MED.	↑ MED.			
	HOT EXT.	LOW	LOW	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
E.North- America ENA	HEAVY PRECIP.	↑ HIGH	↑ MED.	↑ HIGH	↑ L.	↑ E.L.	↑ MED.	↑ HIGH	↑ V.L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	↑ MED.			
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
N.E.North- America NEN	HOT EXT.	↑ V.L.	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V.L.	↑ V.C.	↑ HIGH	↑ L	↑ E.L.
	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW			
N.W.North- America	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW			
	HOT EXT.	↑ V.L.	↑ HIGH	↑ V.L.	↑ E.L.	↑ V.C.	↑ L	↑ V.L.	↑ V.C.
	HEAVY PRECIP.	LOW	LOW	↑ L.	↑ V.L.	↑ V.C.	↑ HIGH	↑ L.	↑ E.L.

NWN	AGR./ECOL. DROUGHT	LOW	LOW	LOW	LOW	LOW				
	HYDR. DROUGHT	LOW	LOW	LOW	LOW	LOW				

## Chapter 11: Weather and climate extreme events in a changing climate

### Coordinating Lead Authors:

Sonia I. Seneviratne (Switzerland), Xuebin Zhang (Canada)

### Lead Authors:

Muhammad Adnan (Pakistan), Wafae Badi (Morocco), Claudine Dereczynski (Brazil), Alejandro Di Luca (Australia, Canada/Argentina), Subimal Ghosh (India), Iskhag Iskandar (Indonesia), James Kossin (United States of America), Sophie Lewis (Australia), Friederike Otto (United Kingdom/Germany), Izidine Pinto (South Africa/Mozambique), Masaki Satoh (Japan), Sergio M. Vicente-Serrano (Spain), Michael Wehner (United States of America), Botao Zhou (China)

### Contributing Authors:

Mathias Hauser (Switzerland), Megan Kirchmeier-Young (Canada/United States of America), Lisa V. Alexander (Australia), Richard P. Allan (United Kingdom), Mansour Almazroui (Saudi Arabia), Lincoln M. Alvez (Brazil), Margot Bador (Australia, France/France), Rondrotiana Barimalala (South Africa/Madagascar), Richard A. Betts (United Kingdom), Suzana J. Camargo (United States of America/Brazil, USA), Pep G. Canadell (Australia), Erika Coppola (Italy), Markus G. Donat (Spain/Germany, Australia), Hervé Douville (France), Robert J. H. Dunn (United Kingdom/Germany, United Kingdom), Erich Fischer (Switzerland), Hayley J. Fowler (United Kingdom), Nathan P. Gillett (Canada), Peter Greve (Austria/Germany), Michael Grose (Australia), Lukas Gudmundsson (Switzerland/Germany, Iceland), Jose Manuel Gutierrez (Spain), Lofti Halimi (Algeria), Zhenyu Han (China), Kevin Hennessy (Australia), Richard G. Jones (United Kingdom), Yeon-Hee Kim (Republic of Korea), Thomas Knutson (United States of America), June-Yi Lee (Republic of Korea), Chao Li (China), Georges-Noel T. Longandjo (South Africa/Democratic Republic of Congo), Kathleen L. McInnes (Australia), Tim R. McVicar (Australia), Malte Meinshausen (Australia/Germany), Seung-Ki Min (Republic of Korea), Ryan S. Padron Flasher (Switzerland/Ecuador, United States of America), Christina M. Patricola (United States of America), Roshanka Ranasinghe (The Netherlands/Sri Lanka, Australia), Johan Reynolds (The Netherlands/Belgium), Joeri Rogelj (United Kingdom/Belgium), Alex C. Ruane (United States of America), Daniel Ruiz Carrascal (United States of America/Colombia), Bjørn H. Samset (Norway), Jonathan Spinoni (Italy), Qiaohong Sun (Canada/China), Ying Sun (China), Mouhamadou Bamba Sylla (Rwanda/Senegal), Claudia Tebaldi (United States of America), Laurent Terray (France), Wim Thiery (Belgium), Jessica Tierney (United States of America), Maarten K. van Aalst (The Netherlands), Bart van den Hurk (The Netherlands), Robert Vautard (France), Wen Wang (China), Seth Westra (Australia), Jakob Zscheischler (Germany)

### Review Editors:

Johnny Chan (China), Asgeir Sorteberg (Norway), Carolina Vera (Argentina)

### Chapter Scientists:

Mathias Hauser (Switzerland), Megan Kirchmeier-Young (Canada/ United States of America), Hui Wan (Canada)

This Chapter should be cited as:

Seneviratne, S. I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F. Otto, I. Pinto, M. Satoh, S. M. Vicente-Serrano, M. Wehner, B. Zhou, 2021, Weather and Climate Extreme Events in a Changing Climate. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.

**Date:** August 2021

**This document is subject to copy-editing, corrigenda and trickle backs.**

1 Table Of Content

2

3 **Executive Summary ..... 6**

4

5 **11.1 Framing .....11**

6 11.1.1 Introduction to the chapter ..... 11

7 11.1.2 What are extreme events and how are their changes studied? ..... 11

8 11.1.3 Types of extremes assessed in this chapter ..... 12

9 11.1.4 Effects of greenhouse gas and other external forcings on extremes ..... 13

10

11 **BOX 11.1: Thermodynamic and dynamic changes in extremes across scales.....15**

12

13 11.1.5 Effects of large-scale circulation on changes in extremes ..... 17

14 11.1.6 Effects of regional-scale processes and forcings and feedbacks on changes in extremes..... 18

15 11.1.7 Global-scale synthesis..... 19

16

17 **BOX 11.2: Low-likelihood high-impact changes in extremes .....24**

18

19 **11.2 Data and Methods .....26**

20 11.2.1 Definition of extremes ..... 26

21 11.2.2 Data..... 27

22

23 **BOX 11.3: Extremes in paleoclimate archives compared to instrumental records.....28**

24

25 11.2.3 Attribution of extremes ..... 31

26 11.2.4 Projecting changes in extremes as a function of global warming levels..... 32

27

28 **Cross-Chapter Box 11.1: Translating between regional information at global warming levels vs**

29 **scenarios for end users .....34**

30

31 **11.3 Temperature extremes .....38**

32 11.3.1 Mechanisms and drivers ..... 38

33 11.3.2 Observed trends ..... 40

34 11.3.3 Model evaluation ..... 43

35 11.3.4 Detection and attribution, event attribution..... 45

36 11.3.5 Projections..... 47

37

38 **11.4 Heavy precipitation .....51**

39 11.4.1 Mechanisms and drivers ..... 51

40 11.4.2 Observed Trends..... 52

41 11.4.3 Model evaluation ..... 56

42 11.4.4 Detection and attribution, event attribution..... 57

43 11.4.5 Projections..... 59

1  
2 **11.5 Floods.....63**  
3 11.5.1 Mechanisms and drivers ..... 64  
4 11.5.2 Observed trends ..... 65  
5 11.5.3 Model evaluation ..... 65  
6 11.5.4 Attribution..... 66  
7 11.5.5 Future projections ..... 67  
8  
9 **11.6 Droughts .....68**  
10 11.6.1 Mechanisms and drivers ..... 68  
11 11.6.1.1 Precipitation deficits ..... 69  
12 11.6.1.2 Atmospheric evaporative demand ..... 69  
13 11.6.1.3 Soil moisture deficits ..... 70  
14 11.6.1.4 Hydrological deficits..... 71  
15 11.6.1.5 Atmospheric-based drought indices..... 71  
16 11.6.1.6 Relation of assessed variables and metrics for changes in different drought types ..... 71  
17 11.6.2 Observed trends ..... 72  
18 11.6.2.1 Precipitation deficits ..... 72  
19 11.6.2.2 Atmospheric evaporative demand ..... 72  
20 11.6.2.3 Soil moisture deficits ..... 73  
21 11.6.2.4 Hydrological deficits..... 74  
22 11.6.2.5 Atmospheric-based drought indices..... 74  
23 11.6.2.6 Synthesis for different drought types ..... 75  
24 11.6.3 Model evaluation ..... 75  
25 11.6.3.1 Precipitation deficits ..... 75  
26 11.6.3.2 Atmospheric evaporative demand ..... 75  
27 11.6.3.3 Soil moisture deficits ..... 76  
28 11.6.3.4 Hydrological deficits..... 77  
29 11.6.3.5 Atmospheric-based drought indices..... 77  
30 11.6.3.6 Synthesis for different drought types ..... 77  
31 11.6.4 Detection and attribution, event attribution..... 78  
32 11.6.4.1 Precipitation deficits ..... 78  
33 11.6.4.2 Soil moisture deficits ..... 79  
34 11.6.4.3 Hydrological deficits..... 79  
35 11.6.4.4 Atmospheric-based drought indices..... 80  
36 11.6.4.5 Synthesis for different drought types ..... 80  
37 11.6.5 Projections..... 81  
38 11.6.5.1 Precipitation deficits ..... 81  
39 11.6.5.2 Atmospheric evaporative demand ..... 82  
40 11.6.5.3 Soil moisture deficits ..... 83



1	11.6.5.4	Hydrological deficits.....	84
2	11.6.5.5	Atmospheric-based drought indices.....	85
3	11.6.5.6	Synthesis for different drought types.....	86
4			
5	<b>11.7</b>	<b>Extreme storms .....</b>	<b>87</b>
6	11.7.1	Tropical cyclones.....	88
7	11.7.1.1	Mechanisms and drivers.....	88
8	11.7.1.2	Observed trends .....	88
9	11.7.1.3	Model evaluation .....	90
10	11.7.1.4	Detection and attribution, event attribution.....	92
11	11.7.1.5	Projections.....	94
12	11.7.2	Extratropical storms.....	97
13	11.7.2.1	Observed trends .....	97
14	11.7.2.2	Model evaluation .....	98
15	11.7.2.3	Detection and attribution, event attribution.....	98
16	11.7.2.4	Projections.....	98
17	11.7.3	Severe convective storms.....	99
18	11.7.3.1	Mechanisms and drivers.....	100
19	11.7.3.2	Observed trends .....	101
20	11.7.3.3	Model evaluation .....	102
21	11.7.3.4	Detection and attribution, event attribution.....	103
22	11.7.3.5	Projections.....	103
23	11.7.4	Extreme winds.....	104
24			
25	<b>11.8</b>	<b>Compound events .....</b>	<b>106</b>
26	11.8.1	Overview.....	106
27	11.8.2	Concurrent extremes in coastal and estuarine regions.....	107
28	11.8.3	Concurrent droughts and heat waves.....	108
29			
30	<b>BOX 11.4: Case study: Global-scale concurrent climate anomalies at the example of the 2015-2016</b>		
31	<b>extreme El Niño and the 2018 boreal spring/summer extremes .....</b>		<b>109</b>
32			
33	<b>11.9</b>	<b>Regional information on extremes.....</b>	<b>113</b>
34	11.9.1	Overview.....	113
35	11.9.2	Temperature extremes.....	114
36	11.9.3	Heavy precipitation.....	114
37	11.9.4	Droughts.....	115
38			
39	<b>Frequently Asked Questions.....</b>		<b>117</b>
40	FAQ 11.1:	How do changes in climate extremes compare with changes in climate averages?.....	117
41	FAQ 11.2:	Will unprecedented extremes occur as a result of human-induced climate change? .....	119

1      FAQ 11.3:    Did climate change cause that recent extreme event in my country?..... 120

2

3      **Large tables** .....**122**

4

5      **Acknowledgements** .....**233**

6

7      **References** .....**234**

8

9      **Appendix 11.A** .....**314**

10

11     **Figures** .....**316**

12

13

## 1 Executive Summary

2  
3 **This chapter assesses changes in weather and climate extremes on regional and global scales, including**  
4 **observed changes and their attribution, as well as projected changes.** The extremes considered include  
5 temperature extremes, heavy precipitation and pluvial floods, river floods, droughts, storms (including  
6 tropical cyclones), as well as compound events (multivariate and concurrent extremes). Changes in marine  
7 extremes are addressed in Chapter 9 and Cross-Chapter Box 9.1. Assessments of past changes and their  
8 drivers are from 1950 onward, unless indicated otherwise. Projections for changes in extremes are presented  
9 for different levels of global warming, supplemented with information for the conversion to emission  
10 scenario-based projections (Cross-Chapter Box 11.1; Chapter 4, Table 4.2). Since AR5, there have been  
11 important new developments and knowledge advances on changes in weather and climate extremes, in  
12 particular regarding human influence on individual extreme events, on changes in droughts, tropical  
13 cyclones, and compound events, and on projections at different global warming levels (1.5°C–4°C). These,  
14 together with new evidence at regional scales, provide a stronger basis and more regional information for the  
15 AR6 assessment on weather and climate extremes.

16  
17 **It is an established fact that human-induced greenhouse gas emissions have led to an increased**  
18 **frequency and/or intensity of some weather and climate extremes since pre-industrial time, in**  
19 **particular for temperature extremes.** Evidence of observed changes in extremes and their attribution to  
20 human influence (including greenhouse gas and aerosol emissions and land-use changes) has strengthened  
21 since AR5, in particular for extreme precipitation, droughts, tropical cyclones and compound extremes  
22 (including dry/hot events and fire weather). Some recent hot extreme events would have been *extremely*  
23 *unlikely* to occur without human influence on the climate system. {11.2, 11.3, 11.4, 11.6, 11.7, 11.8}

24  
25 **Regional changes in the intensity and frequency of climate extremes generally scale with global**  
26 **warming. New evidence strengthens the conclusion from SR1.5 that even relatively small incremental**  
27 **increases in global warming (+0.5°C) cause statistically significant changes in extremes on the global**  
28 **scale and for large regions (*high confidence*). In particular, this is the case for temperature extremes**  
29 **(*very likely*), the intensification of heavy precipitation (*high confidence*) including that associated with**  
30 **tropical cyclones (*medium confidence*), and the worsening of droughts in some regions (*high***  
31 ***confidence*).** The occurrence of extreme events unprecedented in the observed record will increase with  
32 increasing global warming, even at 1.5°C of global warming. Projected percentage changes in frequency are  
33 higher for the rarer extreme events (*high confidence*). {11.1, 11.2, 11.3, 11.4, 11.6, 11.9, CC-Box 11.1}

## 34 Methods and Data for Extremes

35  
36  
37 **Since AR5, the confidence about past and future changes in weather and climate extremes has**  
38 **increased due to better physical understanding of processes, an increasing proportion of the scientific**  
39 **literature combining different lines of evidence, and improved accessibility to different types of climate**  
40 **models (*high confidence*). There have been improvements in some observation-based datasets,**  
41 **including reanalysis data (*high confidence*). Climate models can reproduce the sign of changes in**  
42 **temperature extremes observed globally and in most regions, although the magnitude of the trends**  
43 **may differ (*high confidence*).** Models are able to capture the large-scale spatial distribution of precipitation  
44 extremes over land (*high confidence*). The intensity and frequency of extreme precipitation simulated by  
45 Coupled Model Intercomparison Project Phase 6 (CMIP6) models are similar to those simulated by CMIP5  
46 models (*high confidence*). Higher horizontal model resolution improves the spatial representation of some  
47 extreme events (e.g., heavy precipitation events), in particular in regions with highly varying topography  
48 (*high confidence*). {11.2, 11.3, 11.4}

## 49 Temperature Extremes

50  
51  
52 **The frequency and intensity of hot extremes have increased and those of cold extremes have decreased**  
53 **on the global scale since 1950 (*virtually certain*). This also applies at regional scale, with more than**

1 **80% of AR6 regions<sup>1</sup> showing similar changes assessed to be at least *likely*.** In a few regions, *limited*  
 2 *evidence* (data or literature) prevents the reliable estimation of trends. {11.3, 11.9}

3  
 4 **Human-induced greenhouse gas forcing is the main driver of the observed changes in hot and cold**  
 5 **extremes on the global scale (*virtually certain*) and on most continents (*very likely*).** The effect of  
 6 enhanced greenhouse gas concentrations on extreme temperatures is moderated or amplified at the regional  
 7 scale by regional processes such as soil moisture or snow/ice-albedo feedbacks, by regional forcing from  
 8 land use and land-cover changes, or aerosol concentrations, and decadal and multidecadal natural variability.  
 9 Changes in anthropogenic aerosol concentrations have *likely* affected trends in hot extremes in some regions.  
 10 Irrigation and crop expansion have attenuated increases in summer hot extremes in some regions, such as the  
 11 U.S. Midwest (*medium confidence*). Urbanization has *likely* exacerbated changes in temperature extremes in  
 12 cities, in particular for night-time extremes. {11.1, 11.2, 11.3}

13  
 14 **The frequency and intensity of hot extremes will continue to increase and those of cold extremes will**  
 15 **continue to decrease, at both global and continental scales and in nearly all inhabited regions<sup>1</sup> with**  
 16 **increasing global warming levels.** This will be the case even if global warming is stabilized at 1.5°C.  
 17 Relative to present-day conditions, changes in the intensity of extremes would be at least double at 2°C, and  
 18 quadruple at 3°C of global warming, compared to changes at 1.5°C of global warming. The number of hot  
 19 days and hot nights and the length, frequency, and/or intensity of warm spells or heat waves will increase  
 20 over most land areas (*virtually certain*). In most regions, future changes in the intensity of temperature  
 21 extremes will *very likely* be proportional to changes in global warming, and up to 2–3 times larger (*high*  
 22 *confidence*). The highest increase of temperature of hottest days is projected in some mid-latitude and semi-  
 23 arid regions, at about 1.5 time to twice the rate of global warming (*high confidence*). The highest increase of  
 24 temperature of coldest days is projected in Arctic regions, at about three times the rate of global warming  
 25 (*high confidence*). The frequency of hot temperature extreme events will *very likely* increase non-linearly  
 26 with increasing global warming, with larger percentage increases for rarer events. {11.2, 11.3, 11.9; Table  
 27 11.1; Figure 11.3}

### 28 **Heavy Precipitation and Pluvial Floods**

29  
 30  
 31 **The frequency and intensity of heavy precipitation events have *likely* increased at the global scale over**  
 32 **a majority of land regions with good observational coverage. Heavy precipitation has *likely* increased**  
 33 **on the continental scale over three continents: North America, Europe, and Asia.** Regional increases in  
 34 the frequency and/or intensity of heavy precipitation have been observed with at least *medium confidence* for  
 35 nearly half of AR6 regions, including WSAF, ESAF, WSB, SAS, ESB, REF, WCA, ECA, TIB, EAS, SEA,  
 36 NAU, NEU, EEU, GIC, WCE, SES, CNA, and ENA. {11.4, 11.9}

37  
 38 **Human influence, in particular greenhouse gas emissions, is *likely* the main driver of the observed**  
 39 **global scale intensification of heavy precipitation in land regions.** It is *likely* that human-induced climate  
 40 change has contributed to the observed intensification of heavy precipitation at the continental scale in North  
 41 America, Europe and Asia. Evidence of a human influence on heavy precipitation has emerged in some  
 42 regions. {11.4, 11.9, Table 11.1}

43  
 44 **Heavy precipitation will generally become more frequent and more intense with additional global**  
 45 **warming. At global warming levels of 4°C relative to the pre-industrial, very rare (e.g., 1 in 10 or more**

---

<sup>1</sup> See Figure 1.18 in Chapter 1 for definition of AR6 regions. Acronyms for inhabited regions: ARP: Arabian Peninsula; CAF: C. Africa; CAR: Caribbean; CAU: C. Australia; CNA: C. North America; EAS: E. Asia; EAU: E. Australia; ECA: E. Central Asia; EEU: E. Europe; ENA: E. North America; ESAF: E. Southern Africa; ESB: E. Siberia; GIC: Greenland/Iceland; MDG: Madagascar; MED: Mediterranean; NAU: N. Australia; NCA: N. Central America; NEAF: N.E. Africa; NEN: N.E. North America; NES: N.E. South America; NEU: N. Europe; NSA: N. South America; NWN: N.W. North America; NWS: N.W. South America; NZ: New Zealand; RAR: Russian Arctic; RFE: Russian Far East; SAH: Sahara; SAM: South American Monsoon; SAS: South Asia; SAU: Southern Australia; SCA: S. Central America; SEAF: S.E. Africa; SES: S.E. South America; SSA: S. South America; SWS: S.W. South America; TIB: Tibetan Plateau; WAF: Western Africa; WCA: W. Central Asia; WCE: Western & Central Europe; WNA: W. North America; WSAF: W. Southern Africa; WSB: W. Siberia.

1 **years) heavy precipitation events would become more frequent and more intense than in the recent**  
2 **past, on the global scale (*virtually certain*) and in all continents and AR6 regions. The increase in**  
3 **frequency and intensity is *extremely likely* for most continents and *very likely* for most AR6 regions.** At  
4 the global scale, the intensification of heavy precipitation will follow the rate of increase in the maximum  
5 amount of moisture that the atmosphere can hold as it warms (*high confidence*), of about 7% per 1°C of  
6 global warming. The increase in the frequency of heavy precipitation events will accelerate with more  
7 warming and will be higher for rarer events (*high confidence*), with a *likely* doubling and tripling in the  
8 frequency of 10-year and 50-year events, respectively, compared to the recent past at 4°C of global warming.  
9 Increases in the intensity of extreme precipitation at regional scales will vary, depending on the amount of  
10 regional warming, changes in atmospheric circulation and storm dynamics (*high confidence*). {11.4, Box  
11 11.1}

12  
13 **The projected increase in the intensity of extreme precipitation translates to an increase in the**  
14 **frequency and magnitude of pluvial floods – surface water and flash floods – (*high confidence*), as**  
15 **pluvial flooding results from precipitation intensity exceeding the capacity of natural and artificial**  
16 **drainage systems. {11.4}**

### 17 **River Floods**

18  
19  
20 **Significant trends in peak streamflow have been observed in some regions over the past decades (*high***  
21 ***confidence*).** This includes increases in RAR, NSA, and parts of SES, NEU, ENA and  
22 decreases in NES, SAU, and parts of MED and EAS). The seasonality of river floods has changed in cold  
23 regions where snow-melt is involved, with an earlier occurrence of peak streamflow (*high confidence*).  
24 {11.5}

25  
26 **Global hydrological models project a larger fraction of land areas to be affected by an increase in**  
27 **river floods than by a decrease in river floods (*medium confidence*).** River floods are projected to become  
28 more frequent and intense in some AR6 regions (RAR, SEA, SAS, NWS) (*high confidence*) and less  
29 frequent and intense in others (WCE, EEU, MED) (*high confidence*). Regional changes in river floods are  
30 more uncertain than changes in pluvial floods because complex hydrological processes and forcings,  
31 including land cover change and human water management, are involved. {11.5}

### 32 **Droughts**

33  
34  
35 **Different drought types exist, and they are associated with different impacts and respond differently to**  
36 **increasing greenhouse gas concentrations.** Precipitation deficits and changes in evapotranspiration (ET)  
37 govern net water availability. A lack of sufficient soil moisture, sometimes amplified by increased  
38 atmospheric evaporative demand (AED), results in agricultural and ecological drought. Lack of runoff and  
39 surface water result in hydrological drought. {11.6}

40  
41 **Human-induced climate change has contributed to decreases in water availability during the dry**  
42 **season over a predominant fraction of the land area due to evapotranspiration increases (*medium***  
43 ***confidence*).** Increases in evapotranspiration have been driven by AED increases induced by increased  
44 temperature, decreased relative humidity and increased net radiation (*high confidence*). Trends in  
45 precipitation are not a main driver in affecting global-scale trends in drought (*medium confidence*), but have  
46 induced drying trends in a few AR6 regions (NES: *high confidence*; WAF, CAF, ESAF, SAM, SWS, SSA,  
47 SAS: *medium confidence*). Increasing trends in agricultural and ecological droughts have been observed on  
48 all continents (WAF, CAF, WSAF, ESAF, WCA, ECA, EAS, SAU, MED, WCE, WNA, NES: *medium*  
49 *confidence*), but decreases only in one AR6 region (NAU: *medium confidence*). Increasing trends in  
50 hydrological droughts have been observed in a few AR6 regions (MED: *high confidence*; WAF, EAS, SAU:  
51 *medium confidence*). Regional-scale attribution shows that human-induced climate change has contributed to  
52 increased agricultural and ecological droughts (MED, WNA), and increased hydrological drought (MED) in  
53 some regions (*medium confidence*). {11.6, 11.9}

54  
55 **The land area affected by increasing drought frequency and severity expands with increasing global**

1 **warming (*high confidence*)**. Several regions will be affected by more severe agricultural and ecological  
2 droughts even if global warming is stabilized in a range of 1.5°C–2°C of global warming (*high confidence*),  
3 including WCE, MED, EAU, SAU, SCA, NSA, SAM, SWS, SSA, NCA, CAN, WSAF, ESAF and MDG  
4 (*medium confidence*). At 4°C of global warming, about 50% of all inhabited AR6 regions would be affected  
5 (WCE, MED, CAU, EAU, SAU, WCA, EAS, SCA, CAR, NSA, NES, SAM, SWS, SSA, NCA, CAN, ENA,  
6 WNA, WSAF, ESAF, MDG; *medium confidence* or higher), and only two regions (NEAF, SAS) would  
7 experience decreases in agricultural and ecological drought (*medium confidence*). There is *high confidence*  
8 that the projected increases in agricultural and ecological droughts are strongly affected by ET increases  
9 associated with enhanced AED. Several regions are projected to be more strongly affected by hydrological  
10 droughts with increasing global warming (at 4°C of global warming: NEU, WCE, EEU, MED, SAU, WCA,  
11 SCA, NSA, SAM, SWS, SSA, WNA, WSAF, ESAF, MDG; *medium confidence* or higher). There is *low*  
12 *confidence* that effects of enhanced atmospheric CO<sub>2</sub> concentrations on plant water-use efficiency alleviate  
13 extreme agricultural and ecological droughts in conditions characterized by limited soil moisture  
14 and enhanced AED. There is also *low confidence* that these effects will substantially reduce global plant  
15 transpiration and the severity of hydrological droughts. There is *high confidence* that the land carbon sink  
16 will become less efficient due to soil moisture limitations and associated drought conditions in some regions  
17 in higher-emission scenarios, in particular under global warming levels above 4°C. {11.6, 11.9, CC-Box 5.1}

### 18 19 **Extreme Storms, Including Tropical Cyclones (TCs)**

20  
21 **The average and maximum rain rates associated with TCs, extratropical cyclones and atmospheric**  
22 **rivers across the globe, and severe convective storms in some regions, increase in a warming world**  
23 **(*high confidence*)**. Available event attribution studies of observed strong TCs provide *medium confidence*  
24 for a human contribution to extreme TC rainfall. Peak TC rain rates increase with local warming at least at  
25 the rate of mean water vapour increase over oceans (about 7% per 1°C of warming) and in some cases  
26 exceeding this rate due to increased low-level moisture convergence caused by increases in TC wind  
27 intensity (*medium confidence*). {11.7, 11.4, Box 11.1}

28  
29 **It is *likely* that the global proportion of major TC (Category 3–5) intensities over the past four decades**  
30 **has increased**. The average location where TCs reach their peak wind intensity has *very likely* migrated  
31 poleward in the western North Pacific Ocean since the 1940s, and TC translation speed has *likely* slowed  
32 over the conterminous USA since 1900. Evidence of similar trends in other regions is not robust. The global  
33 frequency of TC rapid intensification events has *likely* increased over the past four decades. None of these  
34 changes can be explained by natural variability alone (*medium confidence*).

35  
36 **The proportion of intense TCs, average peak TC wind speeds, and peak wind speeds of the most**  
37 **intense TCs will increase on the global scale with increasing global warming (*high confidence*)**. The  
38 total global frequency of TC formation will decrease or remain unchanged with increasing global warming  
39 (*medium confidence*). {11.7.1}

40  
41 **There is *low confidence* in past changes of maximum wind speeds and other measures of dynamical**  
42 **intensity of extratropical cyclones. Future wind speed changes are expected to be small, although**  
43 **poleward shifts in the storm tracks could lead to substantial changes in extreme wind speeds in some**  
44 **regions (*medium confidence*)**. There is *low confidence* in past trends in characteristics of severe convective  
45 storms, such as hail and severe winds, beyond an increase in precipitation rates. The frequency of springtime  
46 severe convective storms is projected to increase in the USA, leading to a lengthening of the severe  
47 convective storm season (*medium confidence*); evidence in other regions is limited. {11.7.2, 11.7.3}.

### 48 49 **Compound Events, Including Dry/Hot events, Fire Weather, Compound Flooding, and Concurrent** 50 **Extremes**

51  
52 **The probability of compound events has *likely* increased in the past due to human-induced climate**  
53 **change and will *likely* continue to increase with further global warming**. Concurrent heat waves and  
54 droughts have become more frequent and this trend will continue with higher global warming (*high*  
55 *confidence*). Fire weather conditions (compound hot, dry and windy events) have become more probable in

1 some regions (*medium confidence*) and there is *high confidence* that they will become more frequent in some  
2 regions at higher levels of global warming. The probability of compound flooding (storm surge, extreme  
3 rainfall and/or river flow) has increased in some locations, and will continue to increase due to both sea level  
4 rise and increases in heavy precipitation, including changes in precipitation intensity associated with TCs  
5 (*high confidence*). The land area affected by concurrent extremes has increased (*high confidence*).  
6 Concurrent extreme events at different locations, but possibly affecting similar sectors (e.g., critical crop-  
7 producing areas for global food supply) in different regions, will become more frequent with increasing  
8 global warming, in particular above 2°C of global warming (*high confidence*). {11.8, Box 11.3, Box 11.4}.

#### 9 **Low-Likelihood High-Impact (LLHI) Events Associated With Climate Extremes**

10 **The future occurrence of LLHI events linked to climate extremes is generally associated with *low***  
11 ***confidence*, but cannot be excluded, especially at global warming levels above 4°C.** Compound events,  
12 including concurrent extremes, are a factor increasing the probability of LLHI events (*high confidence*).  
13 With increasing global warming some compound events with low likelihood in past and current climate will  
14 become more frequent, and there is a higher chance of occurrence of historically unprecedented events and  
15 surprises (*high confidence*). However, even extreme events that do not have a particularly low probability in  
16 the present climate (at more than 1°C of global warming) can be perceived as surprises because of the pace  
17 of global warming (*high confidence*). {Box 11.2}

18  
19  
20

## 11.1 Framing

### 11.1.1 Introduction to the chapter

This chapter provides assessments of changes in weather and climate extremes (collectively referred to as extremes) framed in terms of the relevance to the Working Group II assessment. It assesses observed changes in extremes, their attribution to causes, and future projections, at three global warming levels: 1.5°C, 2°C, 4°C. This chapter is also one of the four “regional chapters” of the WGI report (along with Chapters 10 and 12 and the Atlas). Consequently, while it encompasses assessments of changes in extremes at global and continental scales to provide a large-scale context, it also addresses changes in extremes at regional scales.

Extremes are climatic impact-drivers (Annex VII: Glossary, see Chapter 12 for a comprehensive assessment). The IPCC risk framework (Chapter 1) articulates clearly that the exposure and vulnerability to climatic impact-drivers, such as extremes, modulate the risk of adverse impacts of these drivers, and that adaptation that reduces exposure and vulnerability will increase resilience resulting in a reduction in impacts. Nonetheless, changes in extremes lead to changes in impacts not only as a direct consequence of changes in their magnitude and frequency, but also through their influence on exposure and resilience.

The Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation (referred as the SREX report, IPCC, 2012) provided a comprehensive assessment on changes in extremes and how exposure and vulnerability to extremes determine the impacts and likelihood of disasters. Chapter 3 of that report (Seneviratne et al., 2012, hereafter also referred to as SREX Ch3) assessed physical aspects of extremes, and laid a foundation for the follow-up IPCC assessments. Several chapters of the WGI AR5 (IPCC AR5; IPCC, 2013) addressed climate extremes with respect to observed changes (Hartmann et al., 2013), model evaluation (Flato et al., 2013), attribution (Bindoff et al., 2013), and projected long-term changes (Collins et al., 2013). Assessments were also provided in the recent IPCC Special Reports on 1.5°C global warming (SR15, IPCC, 2018; Hoegh-Guldberg et al., 2018), on climate change and land (IPCC, 2019), and on oceans and the cryosphere (IPCC, 2019). These assessments are the starting point of the present assessment.

This chapter is structured as follows (Figure 11.1). This Section (11.1) provides the general framing and introduction to the chapter, highlighting key aspects that underlie the confidence and uncertainty in the assessment of changes in extremes, and introducing some main elements of the chapter. To provide readers a quick overview of past and future changes in extremes, a synthesis of global scale assessment for different types of extremes is included at the end of this Section (Tables 11.1 and 11.2). Section 11.2 introduces methodological aspects of research on climate extremes. Sections 11.3 to 11.7 assess past changes and their attribution to causes, and projected future changes in extremes, for different types of extremes, including temperature extremes, heavy precipitation and pluvial floods, river floods, droughts, and storms, in separate sections. Section 11.8 addresses compound events. Section 11.9 summarizes regional assessments of changes in temperature extremes, in precipitation extremes and in droughts by continents in tables. The chapter also includes several boxes and FAQs on more specific topics.

[START FIGURE 11.1 HERE]

**Figure 11.1:** Chapter 11 visual abstract of contents.

[END FIGURE 11.1 HERE]

### 11.1.2 What are extreme events and how are their changes studied?

Building on the SREX report and AR5, this Report defines an extreme weather event as “an event that is rare at a particular place and time of year” and an extreme climate event as “a pattern of extreme weather that



1 persists for some time, such as a season” (Annex VII: Glossary). The definitions of rare are wide ranging,  
2 depending on applications. Some studies consider an event as an extreme if it is unprecedented; on the other  
3 hand, other studies consider events that occur several times a year as moderate extreme events. Rarity of an  
4 event with a fixed magnitude also changes under human-induced climate change, making events that are  
5 unprecedented so far rather probable under present conditions, but unique in the observational record – and  
6 thus often considered as “surprises” (see Box 11.2).

7  
8 Various approaches are used to define extremes. These are generally based on the determination of relative  
9 (e.g. 90<sup>th</sup> percentile) or absolute (e.g. 35°C for a hot day) thresholds above which conditions are considered  
10 extremes. Changes in extremes can be examined from two perspectives, either focusing on changes in  
11 frequency of given extremes, or on changes in their intensity. These considerations in the definition of  
12 extremes are further addressed in Section 11.2.1.

### 13 14 15 *11.1.3 Types of extremes assessed in this chapter*

16  
17 The types of extremes assessed in this chapter include temperature extremes, heavy precipitation and pluvial  
18 floods, river floods, droughts, and storms. The drought assessment addresses meteorological droughts,  
19 agricultural and ecological droughts, and hydrological droughts (see Annex VII: Glossary). The storms  
20 assessment addresses tropical cyclones, extratropical cyclones, and severe convective storms. In addition,  
21 this chapter also assesses changes in compound events, that is, multivariate or concurrent extreme events,  
22 because of their relevance to impacts as well as the emergence of new literature on the subject. Most of the  
23 considered extremes were also assessed in the SREX and AR5. Compound events were not assessed in depth  
24 in past IPCC reports (SREX Ch3; Section 11.8). Marine-related extremes such as marine heat waves and  
25 extreme sea level, are assessed in Chapter 9 (Section 9.6.4 and Box 9.2) of this report.

26  
27 Extremes and related phenomena are of various spatial and temporal scales. Tornadoes have a spatial scale  
28 as small as less than 100 meters and a temporal scale as short as a few minutes. In contrast, a drought can last  
29 for multiple years, affecting vast regions. The level of complexity of the involved processes differs from one  
30 type of extreme to another, affecting our capability to detect, attribute and project changes in weather and  
31 climate extremes. Temperature and precipitation extremes studied in the literature are often based on  
32 extremes derived from daily values. Studies of events on longer time scales for both temperature or  
33 precipitation, or on sub-daily extremes, are scarcer, which generally limits the assessment for such events.  
34 Nevertheless, extremes on time scales different from daily are assessed for temperature extremes and heavy  
35 precipitation, when possible (Sections 11.3, 11.4). Droughts, as well as tropical and extratropical cyclones,  
36 are assessed as phenomena in general, not limited by their extreme forms, because these phenomena are  
37 relevant to impacts (Sections 11.6, 11.7). Both precipitation and wind extremes associated with storms are  
38 considered.

39  
40 Multiple concomitant extremes can lead to stronger impacts than those resulting from the same extremes had  
41 they happened in isolation. For this reason, the occurrence of multiple extremes that are multivariate and/or  
42 concurrent and/or happening in succession, also called “compound events” (SREX Ch3), are assessed in this  
43 chapter based on emerging literature on this topic (Section 11.8). Box 11.2 also provides an assessment on  
44 low-likelihood high-impact scenarios associated with extremes.

45  
46 The assessment of projected future changes in extremes is presented as function of different global warming  
47 levels (Section 11.2.4 and CC-Box 11.1). On the one hand, this provides traceability and comparison to the  
48 SR15 assessment (Hoegh-Guldberg et al., 2018, hereafter referred to as SR15 Ch3). On the other hand, this  
49 is useful for decision makers as actionable information, as much of the mitigation policy discussion and  
50 adaptation planning can be tied to the level of global warming. For example, regional changes in extremes,  
51 and thus their impacts, can be linked to global mitigation efforts. Additionally, there is also an advantage of  
52 separating uncertainty in future projections due to regional responses as function of global warming levels  
53 from other factors such as differences in global climate sensitivity and emission scenarios (CC-Box 11.1).  
54 However, information is also provided on the translation between information provided at global warming  
55 levels and for single emissions scenarios (CC-Box 11.1) to facilitate easier comparison with the AR5

1 assessment and with some analyses provided in other chapters as function of emissions scenarios.

2  
3 A global-scale synthesis of this chapter's assessments is provided in Section 11.1.7. In particular, Tables  
4 11.1 and 11.2 provide a synthesis for observed and attributed changes, and projected changes in extremes,  
5 respectively, at different global warming levels (1.5°C, 2°C, 4°C). Tables on regional-scale assessments for  
6 changes in temperature extremes, heavy precipitation and droughts, are provided in Section 11.9.

#### 7 8 9 **11.1.4 Effects of greenhouse gas and other external forcings on extremes**

10 SREX, AR5, and SR15 assessed that there is evidence from observations that some extremes have changed  
11 since the mid 20<sup>th</sup> century, that some of the changes are a result of anthropogenic influences, and that some  
12 observed changes are projected to continue into the future, while other changes are projected to emerge from  
13 natural climate variability under enhanced global warming (SREX Chapter 3, AR5 Chapter 10).

14  
15  
16 At the global scale but also at the regional scale to some extent, many of the changes in extremes are a direct  
17 consequence of enhanced radiative forcing, and the associated global warming and/or resultant increase in  
18 the water-holding capacity of the atmosphere, as well as changes in vertical stability and meridional  
19 temperature gradients that affect climate dynamics (see Box 11.1). Widespread observed and projected  
20 increases in the intensity and frequency of hot extremes, together with decreases in the intensity and  
21 frequency of cold extremes, are consistent with global and regional warming (Figure 11.2, Section 11.3).  
22 Extreme temperatures on land tend to increase more than the global mean temperature (Figure 11.2), due in  
23 large part to the land-sea contrast, and additionally to regional feedbacks in some regions (Section 11.1.6).  
24 Increases in the intensity of temperature extremes scale robustly and in general linearly with global warming  
25 across different geographical regions in projections up to 2100, with minimal dependence on emissions  
26 scenarios (Figures 11.3 and 11.A.1; Seneviratne et al., 2016; Wartenburger et al., 2017; Kharin et al., 2018;  
27 Section 11.2.4 and CC-Box 11.1). The frequency of hot temperature extremes (see Figure 11.6), the number  
28 of heat wave days and the length of heat wave seasons in various regions also scale well, but non-linearly  
29 (because of the threshold effect), with global mean temperatures (Wartenburger et al., 2017; Sun et al.,  
30 2018a).

31  
32 Changes in annual maximum one-day precipitation (Rx1day) are proportional to mean global surface  
33 temperature changes, at about 7% increase per 1°C temperature increase, that is, following the Clausius-  
34 Clapeyron relationship (Box 11.1), both in observations (Westra et al., 2013) and in future projections  
35 (Kharin et al., 2013) at the global scale. Extreme short-duration precipitation in North America also scales  
36 with global surface temperature (Li et al., 2018a; Prein et al., 2016b). At the local and regional scales,  
37 changes in extremes are also strongly modulated and controlled by regional forcings and feedback  
38 mechanisms (Section 11.1.6), whereby some regional forcings, for example, associated with changes in land  
39 cover and land or aerosol emissions, can have non-local or some (non-homogeneous) global-scale effects. In  
40 general, there is *high confidence* in changes in extremes due to global-scale thermodynamic processes (i.e.,  
41 global warming, mean moistening of the air) as the processes are well understood, while the confidence in  
42 those related to dynamic processes or regional and local forcing, including regional and local thermodynamic  
43 processes, is much lower due to multiple factors (see following sub-section and Box 11.1).

44  
45  
46 **[START FIGURE 11.2 HERE]**

47  
48 **Figure 11.2:** Time series of observed temperature anomalies for global average annual mean temperature (black), land  
49 average annual mean temperature (green), land average annual hottest daily maximum temperature (TXx,  
50 purple), and land average annual coldest daily minimum temperature (TNn, blue). Global and land mean  
51 temperature anomalies are relative to their 1850-1900 means based on the multi-product mean annual  
52 time series assessed in Section 2.3.1.1.3 (see text for references). TXx and TNn anomalies are relative to  
53 their respective 1961-1990 means and are based on the HadEX3 dataset (Dunn et al., 2020) using values  
54 for grid boxes with at least 90% temporal completeness over 1961-2018. Further details on data sources  
55 and processing are available in the chapter data table (Table 11.SM.9).

1  
2 **[END FIGURE 11.2 HERE]**  
3  
4

5 **[START FIGURE 11.3 HERE]**  
6  
7

8 **Figure 11.3:** Regional mean changes in annual hottest daily maximum temperature (TXx) for AR6 land regions and  
9 the global land, against changes in global mean surface air temperature (GSAT) as simulated by CMIP6  
10 models under different forcing scenarios SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. (a)  
11 shows individual models from the CMIP6 ensemble (grey), the multi-model median under three selected  
12 SSPs (colours), and the multi-model median (black). (b) to (l) show the multi-model-median for the  
13 pooled data for individual AR6 regions. Numbers in parantheses indicate the linear scaling between  
14 regional TXx and GSAT. The black line indicates the 1:1 reference scaling between TXx and GSAT. See  
15 Atlas.1.3.2 for the definition of regions. For details on the methods see Supplementary Material 11.SM.2.

16  
17  
18 **[END FIGURE 11.3 HERE]**  
19  
20

21 Since AR5, the attribution of extreme weather events, or the investigation of changes in the frequency and/or  
22 magnitude of individual and local- and regional-scale extreme weather events due to various drivers (see  
23 Cross-Working Group Box 1.1 (in Chapter 1) and Section 11.2.3) has provided evidence that greenhouse  
24 gases and other external forcings have affected individual extreme weather events. The events that have been  
25 studied are geographically uneven. A few events, for example, extreme rainfall events in the UK (Schaller et  
26 al., 2016; Vautard et al., 2016; Otto et al., 2018b) or heat waves in Australia (King et al., 2014; Perkins-  
27 Kirkpatrick et al., 2016; Lewis et al., 2017b), have spurred more studies than other events. Many highly  
28 impactful extreme weather events have not been studied in the event attribution framework. Studies in the  
29 developing world are also generally lacking. This is due to various reasons (Section 11.2) including lack of  
30 observational data, lack of reliable climate models, and lack of scientific capacity (Otto et al., 2020). While  
31 the events that have been studied are not representative of all extreme events that occurred and results from  
32 these studies may also be subject to selection bias, the large number of event attribution studies provide  
33 evidence that changes in the properties of these local and individual events are in line with expected  
34 consequences of human influence on the climate and can be attributed to external drivers (Section 11.9).  
35 Figure 11.4 summarizes assessments of observed changes in temperature extremes, in heavy precipitation  
36 and in droughts, and their attribution in a map form.

37  
38  
39 **[START FIGURE 11.4 HERE]**  
40

41 **Figure 11.4:** Overview of observed changes for cold, hot, and wet extremes and their potential human  
42 contribution. Shown are the direction of change and the confidence in 1) the observed changes in how  
43 cold and hot as well as wet extremes have already changed across the world and 2) in the contribution of  
44 whether human-induced climate change contributed in causing to these changes (attribution). In each  
45 region changes in extremes are indicated by colour (orange – increase in the type of extreme, blue –  
46 decrease, both colours – there are changes of opposing direction within the region the signal depends on  
47 the exact event definition, grey – there are no changes observed, and no fill – the data/evidence is too  
48 sparse to make an assessment). The squares and dots next to the symbol indicate the level of confidence  
49 for observing the trend and the human contribution, respectively. The more black dots/squares the higher  
50 the level of confidence. The information on this figure is based on regional assessment of the literature on  
51 observed trends, detection and attribution and event attribution in section 11.9.

52  
53 **[END FIGURE 11.4 HERE]**  
54  
55  
56  
57  
58  
59

1 [START BOX 11.1 HERE]

2  
3  
4 **BOX 11.1: Thermodynamic and dynamic changes in extremes across scales**

5  
6 Changes in weather and climate extremes are determined by local exchanges in heat, moisture, and other  
7 related quantities (thermodynamic changes) and those associated with atmospheric and oceanic motions  
8 (dynamic changes). While thermodynamic and dynamic processes are interconnected, considering them  
9 separately helps to disentangle the roles of different processes contributing to changes in climate extremes  
10 (e.g. Shepherd, 2014).

11  
12 **Temperature extremes**

13 An increase in the concentration of greenhouse gases in the atmosphere leads to the warming of tropospheric  
14 air and the Earth's surface. This direct thermodynamic effect leads to warmer temperatures everywhere with  
15 an increase in the frequency and intensity of warm extremes and a decrease in the frequency and intensity of  
16 cold extremes. The initial increase in temperature in turn leads to other thermodynamic responses and  
17 feedbacks affecting both the atmosphere and the surface. These include an increase in the water vapour  
18 content of the atmosphere (water vapour feedback, see Section 7.4.2.2) and a change in the vertical profile of  
19 temperature (e.g., lapse rate feedback, see Section 7.4.2.2). While the water vapour feedback always  
20 amplifies the initial temperature increases (positive feedback), the lapse rate feedback amplifies near-surface  
21 temperature increases (positive feedback) in mid- and high latitudes but reduces temperature increases  
22 (negative feedback) in tropical regions (Pithan and Mauritsen, 2014).

23  
24 Thermodynamic responses and feedbacks also occur through surface processes. For instance, observations  
25 and model simulations show that temperature increases, including extreme temperatures, are amplified in  
26 areas where seasonal snow cover is reduced due to decreases in surface albedo (see Section 11.3.1). In some  
27 mid-latitude areas, temperature increases are amplified by the higher atmospheric evaporative demand (Fu  
28 and Feng, 2014; Vicente-Serrano et al., 2020b) that results in a drying of soils in some regions (Section  
29 11.6), leading to increased sensible heat fluxes (soil-moisture temperature feedback, see Sections 11.1.6 and  
30 11.3.1). Other thermodynamic feedback processes include changes in the water-use efficiency of plants  
31 under enhanced atmospheric CO<sub>2</sub> concentrations that can reduce the overall transpiration, and thus also  
32 enhance temperature in projections (Sections 8.2.3.3, 11.1.6, 11.3, and 11.6).

33  
34 Changes in the spatial distribution of temperatures can also affect temperature extremes by modifying the  
35 characteristics of weather patterns (e.g., Suarez-Gutierrez et al., 2020). For example, a robust thermodynamic  
36 effect of polar amplification is a weakened north-south temperature gradient, which amplifies the warming  
37 of cold extremes in the Northern Hemisphere mid- and high latitudes because of the reduction of cold air  
38 advection (Holmes et al., 2015; Schneider et al., 2015; Gross et al., 2020). Much less robust is the dynamic  
39 effect of polar amplification (Section 7.4.4.1) and the reduced low-altitude meridional temperature gradient  
40 that has been linked to an increase in the persistence of weather patterns (e.g., heatwaves) and subsequent  
41 increases in temperature extremes (Francis and Vavrus, 2012; Coumou et al., 2015, 2018; Mann et al., 2017)  
42 (CC-Box 10.1).

43  
44 **Precipitation extremes**

45 Changes in temperature also control changes in water vapour through increases in evaporation and in the  
46 water-holding capacity of the atmosphere (Section 8.2.1). At the global scale, column-integrated water  
47 vapour content increases roughly following the Clausius-Clapeyron (C-C) relation, with an increase of  
48 approximately 7% for every degree celsius of global-mean surface warming (Section 8.2.1). Nonetheless, at  
49 regional scales, water vapour increases differ from this C-C rate due to several reasons (Section 8.2.2),  
50 including a change in weather regimes and limitations in moisture transport from the ocean, which warms  
51 more slowly than land (Byrne and O’Gorman, 2018). Observational studies (Fischer and Knutti, 2016; Sun et  
52 al., 2020) have shown the observed rate of increase of precipitation extremes is similar to the C-C scaling at  
53 the global scale. Climate model projections show that the increase in water vapour leads to robust increases  
54 in precipitation extremes everywhere, with a magnitude that varies between 4% and 8% per degree celsius of  
55 surface warming (thermodynamic contribution, Box 11.1, Figure 1b). At regional scales, climate models  
56 show that the dynamic contribution (Box 11.1, Figure 1c) can be substantial and strongly modify the

**Do Not Cite, Quote or Distribute**

11-15

Total pages: 345

1 projected rate of change of extreme precipitation (Box 11.1, Figure 1a) with large regions in the subtropics  
2 showing robust reductions and other areas (e.g., equatorial Pacific) showing robust amplifications (Box 11.1,  
3 Figure 1c). However, the dynamic contributions show large differences across models and are more  
4 uncertain than thermodynamic contributions (Shepherd, 2014; Trenberth et al., 2015; Pfahl et al., 2017; Box  
5 11.1, Figure 1c).

6  
7 Dynamic contributions can occur in response to changes in the vertical and horizontal distribution of  
8 temperature (thermodynamics) and can affect the frequency and intensity of synoptic and subsynoptic  
9 phenomena including tropical cyclones, extratropical cyclones, fronts, mesoscale-convective systems and  
10 thunderstorms. For example, the poleward shift and strengthening of the Southern Hemisphere mid-latitude  
11 storm tracks (Section 4.5.1) can modify the frequency/intensity of extreme precipitation. However, the  
12 precise way in which dynamic changes will affect precipitation extremes is unclear due to several competing  
13 effects (Shaw et al., 2016; Allan et al., 2020).

14  
15 Extreme precipitation can also be enhanced by dynamic responses and feedbacks occurring within storms  
16 that result from the extra latent heat released from the thermodynamic increases in moisture (Lackmann,  
17 2013; Willison et al., 2013; Marciano et al., 2015; Nie et al., 2018; Mizuta and Endo, 2020). The extra latent  
18 heat released within storms has been shown to increase precipitation extremes by strengthening convective  
19 updrafts and the intensity of the cyclonic circulation (e.g., Molnar et al., 2015; Nie et al., 2018), although  
20 weakening effects have also been found in mid-latitude cyclones (e.g., Kirshbaum et al., 2017). Additionally,  
21 the increase in latent heat can also suppress convection at larger scales due to atmospheric stabilization (Nie  
22 et al., 2018; Tandon et al., 2018; Kendon et al., 2019). As these dynamic effects result from feedback  
23 processes within storms where convective processes are crucial, their proper representation might require  
24 improving the horizontal/vertical resolution, the formulation of parameterizations, or both, in current climate  
25 models (i.e., Ban et al., 2015; Kendon et al., 2014; Meredith et al., 2015; Nie et al., 2018; Prein et al., 2015;  
26 Westra et al., 2014).

#### 27 28 29 **[START BOX 11.1, FIGURE 1 HERE]**

30  
31 **Box 11.1, Figure 1:** Multi-model (CMIP5) mean fractional changes (in % per degree of warming) for (a) annual  
32 maximum precipitation (Rx1day), (b) changes in Rx1day due to the thermodynamic contribution  
33 and (c) changes in Rx1day due to the dynamic contribution estimated as the difference between  
34 the total changes and the thermodynamic contribution. Changes were derived from a linear  
35 regression for the period 1950–2100. Uncertainty is represented using the simple approach: no  
36 overlay indicates regions with high model agreement, where  $\geq 80\%$  of models ( $n=22$ ) agree on sign  
37 of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models  
38 agree on sign of change. For more information on the simple approach, please refer to the Cross-  
39 Chapter Box Atlas 1. A detailed description of the estimation of dynamic and thermodynamic  
40 contributions is given in Pfahl et al. (2017). Adapted from (Pfahl et al., 2017), originally published  
41 in *Nature Climate Change*/ Springer Nature. Further details on data sources and processing are  
42 available in the chapter data table (Table 11.SM.9).

#### 43 44 **[END BOX 11.1, FIGURE 1 HERE]**

### 45 46 **Droughts**

47 Droughts are also affected by both thermodynamic and dynamic processes (Sections 8.2.3.3 and 11.6).  
48 Thermodynamic processes affect droughts by increasing atmospheric evaporative demand (Martin, 2018;  
49 Gebremeskel Haile et al., 2020; Vicente-Serrano et al., 2020b) through changes in air temperature, radiation,  
50 wind speed, and relative humidity. Dynamic processes affect droughts through changes in the occurrence,  
51 duration and intensity of weather anomalies, which are related to precipitation and the amount of sunlight  
52 (Section 11.6). While atmospheric evaporative demand increases with warming, regional changes in aridity  
53 are affected by increasing land-ocean warming contrast, vegetation feedbacks and responses to rising CO<sub>2</sub>  
54 concentrations and dynamic shifts in the location of the wet and dry parts of the atmospheric circulation in  
55 response to climate change as well as internal variability (Byrne and O’Gorman, 2015; Kumar et al., 2015;

Allan et al., 2020).

In summary, both thermodynamic and dynamic processes are involved in the changes of extremes in response to warming. Anthropogenic forcing (e.g., increases in greenhouse gas concentrations) directly affects thermodynamic variables, including overall increases in high temperatures and atmospheric evaporative demand, and regional changes in atmospheric moisture, which intensify heatwaves, droughts and heavy precipitation events when they occur (*high confidence*). Dynamic processes are often indirect responses to thermodynamic changes, are strongly affected by internal climate variability and are also less well understood. As such, there is *low confidence* in how dynamic changes affect the location and magnitude of extreme events in a warming climate.

[END BOX 11.1 HERE]

### 11.1.5 Effects of large-scale circulation on changes in extremes

Atmospheric large-scale circulation patterns and associated atmospheric dynamics are important determinants of the regional climate (Chapter 10). As a result, they are also important to the magnitude, frequency, and duration of extremes (Box 11.4). Aspects of changes in large-scale circulation patterns are assessed in Chapters 2, 3, 4, and 8 and representative atmospheric and oceanic modes are described in Annex IV. This subsection provides some general concepts, through a couple of examples, on why the uncertainty in the response of large-scale circulation patterns to external forcing can cascade to uncertainty in the response of extremes to external forcings. Details for specific types of extremes are covered in the relevant subsections. For example, the occurrence of the El Niño-Southern Oscillation (ENSO) influences precipitation regimes in many areas, favoring droughts in some regions and heavy rains in others (Box 11.4). The extent and strength of the Hadley circulation influences regions where tropical and extra-tropical cyclones occur, with important consequences for the characteristics of extreme precipitation, drought, and winds (Section 11.7). Changes in circulation patterns associated with land-ocean heat contrast, which affect the monsoon circulations (Section 8.4.2.4), lead to heavy precipitation along the coastal regions in East Asia (Freychet et al., 2015). As a result, changes in the spatial and/or temporal variability of the atmospheric circulation in response to warming affect characteristics of weather systems such as tropical cyclones (Sharmila and Walsh, 2018), storm tracks (Shaw et al., 2016), and atmospheric rivers (Waliser and Guan, 2017) (e.g. Section 11.7). Changes in weather systems come with changes in the frequency and intensity of extreme winds, extreme temperatures, and extreme precipitation, on the backdrop of thermodynamic responses of extremes to warming (Box 11.1). Floods are also affected by large-scale circulation modes, including ENSO, the North Atlantic Oscillation (NAO), the Atlantic Multi-decadal Variability (AMV), and the Pacific Decadal Variability (PDV) (Kundzewicz et al., 2018; Annex IV). Aerosol forcing, through changes in patterns of sea surface temperatures (SSTs), also affects circulation patterns and tropical cyclone activities (Takahashi et al., 2017).

Changes in atmospheric large-scale circulation due to external forcing are uncertain in general, but there are clear signals in some aspects (Chapter 2, 3, 4, and 8; Sections 2.3.1.4, 8.2.2.2). Among them, there has been a *very likely* widening of the Hadley circulation since the 1980s and the extratropical jets and cyclone tracks have *likely* been shifting poleward since the 1980s (Section 2.3.1.4). The poleward expansion affects drought occurrence in some regions (Section 11.6), and results in poleward shifts of tropical cyclones and storm tracks (Sections 11.7.1, 11.7.2). Although it is *very likely* that the amplitude of ENSO variability will not robustly change over the 21st century (Section 4.3.3.2), the frequency of extreme El Niños (Box 11.4), defined by precipitation threshold, is projected to increase with global warming (Section 6.5 of SROCC). This would have implications for projected changes in extreme events affected by ENSO, including droughts over wide areas (Section 11.6; Box 11.4) and tropical cyclones (Section 11.7.1). A case study is provided for extreme ENSOs in 2015/2016 in Box 11.4 to highlight the influence of ENSO on extremes.

In summary, large-scale atmospheric circulation patterns are important drivers for local and regional extremes. There is overall *low confidence* about future changes in the magnitude, frequency, and spatial distribution of these patterns, which contributes to uncertainty in projected responses of extremes, especially

1 in the near term.

### 2 3 4 **11.1.6 Effects of regional-scale processes and forcings and feedbacks on changes in extremes**

5  
6 At the local and regional scales, changes in extremes are strongly modulated by regional and local feedbacks  
7 (SRCCL, Jia et al., 2019; Seneviratne et al., 2013; Miralles et al., 2014; Lorenz et al., 2016; Vogel et al.,  
8 2017), changes in large-scale circulation patterns (11.1.5), and regional forcings such as changes in land use  
9 or aerosol concentrations (Chapters 3 and 7; Hirsch et al., 2017, 2018; Thiery et al., 2017; Wang et al.,  
10 2017f; Findell et al., 2017). In some cases, such responses may also include non-local effects (e.g., Persad  
11 and Caldeira, 2018; Miralles et al., 2019; de Vrese et al., 2016; Schumacher et al., 2019). Regional-scale  
12 forcing and feedbacks often affect temperature distributions asymmetrically, with generally higher effects for  
13 the hottest percentiles (Section 11.3).

14  
15 Land use can affect regional extremes, in particular hot extremes, in several ways (*high confidence*). This  
16 includes effects of land management (e.g. cropland intensification, irrigation, double cropping) and well as  
17 of land cover changes (deforestation) (Section 11.3.2; see also 11.6). Some of these processes are not well  
18 represented (e.g. effects of forest cover on diurnal temperature cycle) or not integrated (e.g. irrigation) in  
19 climate models (Sections 11.3.2, 11.3.3). Overall, the effects of land use forcing may be particularly relevant  
20 in the context of low-emissions scenarios, which include large land use modifications, for instance associated  
21 with the expansion of biofuels, biofuels with carbon capture and storage (BECCS), or re-afforestation to  
22 ensure negative emissions, as well as with the expansion of food production (e.g. SR15, Chapter 3; CC-Box  
23 5.1; van Vuuren et al., 2011, Hirsch et al., 2018). There are also effects on the water cycle through  
24 freshwater use (CC-Box 5.1; Section 11.6).

25  
26 Aerosol forcing also has a strong regional footprint associated with regional emissions, which affects  
27 temperature and precipitation extremes (*high confidence*; Sections 11.3, 11.4). From ca. the 1950s to 1980s,  
28 enhanced aerosol loadings led to regional cooling due to decreased global solar radiation (“global dimming”)  
29 which was followed by a phase of “global brightening” due to a reduction in aerosol loadings (Chapters 3  
30 and 7; Wild et al., 2005). King et al. (2016a) show that aerosol-induced cooling delayed the timing of a  
31 significant human contribution to record-breaking heat extremes in some regions. On the other hand, the  
32 decreased aerosol loading since the 1990s has led to an accelerated warming of hot extremes in some  
33 regions. Based on Earth System Model (ESM) simulations, Dong et al. (2017b) suggest that a substantial  
34 fraction of the warming of the annual hottest days in Western Europe since the mid-1990s has been due to  
35 decreases in aerosol concentrations in the region. Dong et al. (2016) also identify non-local effects of  
36 decreases in aerosol concentrations in Western Europe, which they estimate played a dominant role in the  
37 warming of the hottest daytime temperatures in Northeast Asia since the mid-1990s, via induced coupled  
38 atmosphere-land surface and cloud feedbacks, rather than a direct impact of anthropogenic aerosol changes  
39 on cloud condensation nuclei.

40  
41 In addition to regional forcings, regional feedback mechanisms can also substantially affect extremes (*high*  
42 *confidence*; Sections 11.3, 11.4, 11.6). In particular, soil moisture feedbacks play an important role for  
43 extremes in several mid-latitude regions, leading in particular to a marked additional warming of hot  
44 extremes compared to mean global warming (Seneviratne et al., 2016; Bathiany et al., 2018; Miralles et al.,  
45 2019), which is superimposed on the known land-sea contrast in mean warming (Vogel et al., 2017). Soil  
46 moisture-atmosphere feedbacks also affect drought development (Section 11.6). Additionally, effects of land  
47 surface conditions on circulation patterns have also been reported (Koster et al., 2016; Sato and Nakamura,  
48 2019). These regional feedbacks are also associated with substantial spread in models (Section 11.3), and  
49 contribute to the identified higher spread of regional projections of temperature extremes as function of  
50 global warming, compared with the spread resulting from the differences in projected global warming  
51 (global transient climate responses) in climate models (Seneviratne and Hauser, 2020). In addition, there are  
52 also feedbacks between soil moisture content and precipitation occurrence, generally characterized by  
53 negative spatial feedbacks and positive local feedbacks (Taylor et al., 2012; Guillod et al., 2015). Climate  
54 model projections suggest that these feedbacks are relevant for projected changes in heavy precipitation  
55 (Seneviratne et al., 2013), however, there is evidence that climate models do not capture the correct sign of

1 the soil moisture-precipitation feedbacks in several regions, in particular spatially and/or in some cases also  
2 temporally (Taylor et al., 2012; Moon et al., 2019). In the Northern Hemisphere high latitudes, the snow- and  
3 ice-albedo feedback, along with other factors, is projected to largely amplify temperature increases (e.g.,  
4 Pithan and Mauritsen, 2014), although the effect on temperature extremes is still unclear. It is also still  
5 unclear whether snow-albedo feedbacks in mountainous regions might have an effect on temperature and  
6 precipitation extremes (e.g. Gobiet et al., 2014), however these feedbacks play an important role in projected  
7 changes in high-latitude warming (Hall and Qu, 2006), and, in particular, in changes in cold extremes in  
8 these regions (Section 11.3).

9  
10 Finally, extreme events may also regionally amplify one another. This is, e.g., the case for heat waves and  
11 droughts, with high temperatures and stronger radiative forcing leading to drying tendencies on land due to  
12 increased evapotranspiration (Section 11.6), and drier soils then inducing decreased evapotranspiration and  
13 higher sensible heat flux and hot temperatures (Seneviratne et al., 2013; Miralles et al., 2014; Vogel et al.,  
14 2017; Zscheischler and Seneviratne, 2017; Zhou et al., 2019b; Kong et al., 2020; see Box 11.1, Section  
15 11.8).

16  
17 In summary, regional forcings and feedbacks, in particular associated with land use and aerosol forcings, and  
18 soil moisture-temperature, soil moisture-precipitation, and snow/ice-albedo-temperature feedbacks, play an  
19 important role in modulating regional changes in extremes. These can also lead to a higher warming of  
20 extreme temperatures compared to mean temperature (*high confidence*), and possibly cooling in some  
21 regions (*medium confidence*). However, there is only *medium confidence* in the representation of the  
22 associated processes in state-of-the-art Earth System Models.

### 23 24 25 **11.1.7 Global-scale synthesis**

26  
27 Tables 11.1 and 11.2 provide a synthesis for observed and attributed changes in extremes, and projected  
28 changes in extremes, respectively, at different levels of global warming. This synthesis assessment focuses  
29 on the more likely range of observed and projected changes. However, some low-likelihood high-impact  
30 scenarios can also be of high relevance as addressed in Box 11.2.

31  
32 Figure 11.5 provides a synthesis on the level of confidence in the attribution and projection of changes in  
33 extremes, building on the assessments from Tables 11.1 and 11.2. In the case where the physical processes  
34 underlying the changes in extremes in response to human forcing are well understood and the signal in the  
35 observations is still relatively weak, confidence in the projections would be higher than in the attribution  
36 because of an increase in the signal to noise ratio with higher global warming. On the other hand, when the  
37 observed signal is already strong and when observational evidence is consistent with model simulated  
38 responses, confidence in attribution may be higher than that in projections if certain physical processes could  
39 be expected to behave differently in a much warmer world and under much higher greenhouse gas forcing,  
40 and if such a behavior is poorly understood.

41  
42 Further synthesis figures for regional assessments are provided in Figure 11.4 (event attribution), Figure 11.6  
43 (projected change in hot temperature extremes) and Figure 11.7 (projected changes in precipitation  
44 extremes), and a synthesis on regional assessments for observed, attributed and projected changes in  
45 extremes is provided in Section 11.9 for all AR6 reference regions (See Chapter 1, section 1.4.5 and Figure  
46 1.18 for definition of AR6 regions).

47  
48 Confidence and likelihood of past changes and projected future changes at 2°C of global warming on the  
49 global scale. The information in this figure is based on Tables 11.1 and 11.2.

50 **[START FIGURE 11.5 HERE]**

51  
52  
53 **Figure 11.5:** Confidence and likelihood of past changes and projected future changes at 2°C of global warming on the  
54 global scale. The information in this figure is based on Tables 11.1 and 11.2.



1 [END FIGURE 11.5 HERE]

2  
3  
4 [START FIGURE 11.6 HERE]

5  
6  
7 **Figure 11.6:** Projected changes in the frequency of extreme temperature events under 1°C, 1.5°C, 2°C, 3°C, and 4°C  
8 global warming levels relative to the 1851-1900 baseline. Extreme temperatures are defined as the  
9 maximum daily temperatures that were exceeded on average once during a 10-year period (10-year event,  
10 blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results  
11 are shown for the global land and the AR6 regions. For each box plot, the horizontal line and the box  
12 represent the median and central 66% uncertainty range, respectively, of the frequency changes across the  
13 multi model ensemble, and the whiskers extend to the 90% uncertainty range. The dotted line indicates no  
14 change in frequency. The results are based on the multi-model ensemble from simulations of global  
15 climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6)  
16 under different SSP forcing scenarios. Adapted from (Li et al., 2020a). Further details on data sources and  
17 processing are available in the chapter data table (Table 11.SM.9).

18  
19  
20 [END FIGURE 11.6 HERE]

21  
22  
23 [START FIGURE 11.7 HERE]

24  
25  
26 **Figure 11.7:** Projected changes in the frequency of extreme precipitation events under 1°C, 1.5°C, 2°C, 3°C, and 4°C  
27 global warming levels relative to the 1951-1990 baseline. Extreme precipitation is defined as the  
28 maximum daily precipitation (Rx1day) that was exceeded on average once during a 10-year period (10-  
29 year event, blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base  
30 period. Results are shown for the global land and the AR6 regions. For each box plot, the horizontal line  
31 and the box represent the median and central 66% uncertainty range, respectively, of the frequency  
32 changes across the multi model ensemble, and the whiskers extend to the 90% uncertainty range. The  
33 dotted line indicates no change in frequency. The results are based on the multi-model ensemble from  
34 simulations of global climate models contributing to the sixth phase of the Coupled Model  
35 Intercomparison Project (CMIP6) under different SSP forcing scenarios. Adapted from (Li et al., 2020a).  
36 Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

37  
38  
39 [END FIGURE 11.7 HERE]

40  
41  
42 [START TABLE 11.1 HERE]

43  
44  
45 **Table 11.1:** Synthesis table on observed changes in extremes and contribution by human influences. Note that  
46 observed changes in marine extremes are assessed in the Cross-Chapter Box 9.1 in Chapter 9.

Phenomenon and direction of trend	Observed/detected trends since 1950 (for +0.5°C global warming or higher)	Human contribution to the observed trends since 1950 (for +0.5°C global warming or higher)
Warmer and/or more frequent hot days and nights over most land areas	<i>Virtually certain</i> on global scale {11.3}	<i>Extremely likely</i> main contributor on global scale {11.3}
Warmer and/or fewer cold days and nights over most land areas	<i>Continental-scale evidence:</i> Asia, Australasia, Europe, North America: <i>Very likely</i> Central and South America: <i>High confidence</i>	<i>Continental-scale evidence:</i> North America, Europe, Australasia, Asia: <i>Very likely</i> Central and South America: <i>High confidence</i>
Warm spells/heat waves; Increases in frequency or intensity over most land areas	Africa: <i>Medium confidence</i> {11.3, 11.9}	Africa: <i>Medium confidence</i> {11.3, 11.9}
Cold spells/cold waves; Decreases in frequency or		

intensity over most land areas		
Heavy precipitation events: increase in the frequency, intensity, and/or amount of heavy precipitation	<p><i>Likely</i> on global scale, over majority of land regions with good observational coverage {11.3}</p> <p><i>Continental-scale evidence:</i> Asia, Europe, North America: <i>Likely</i> Africa, Australasia, Central and South America: <i>Low confidence</i> {11.3, 11.9}</p>	<p><i>Likely</i> main contributor to the observed intensification of heavy precipitation in land regions on global scale. {11.3}</p> <p><i>Continental-scale evidence:</i> Asia, Europe, North America: <i>Likely</i> Africa, Australasia, Central and South America: <i>Low confidence</i> {11.3, 11.9}</p>
Agricultural and ecological drought events: Enhanced drying in dry season	<p><i>Medium confidence</i>, in predominant fraction of land area</p> <p>Observed decrease in water availability in the dry season due to increased evapotranspiration (driven by increased atmospheric evaporative demand) in a predominant fraction of the land area (<i>medium confidence</i>) {11.6}</p> <p>Increasing trends in agricultural and ecological droughts have been observed in AR6 regions on all continents (<i>medium confidence</i>) {11.6, 11.9}</p>	<p><i>Medium confidence</i>, in predominant fraction of land area</p> <p>Human contribution to decrease in water availability in the dry season in a predominant fraction of the land area (<i>medium confidence</i>) {11.6}</p>
Increase in precipitation associated with tropical cyclones	<i>Medium confidence</i> {11.7}	<i>High confidence</i> {11.7}
Increase in likelihood that a TC will be at major TC intensity (Cat. 3-5)	<i>Likely</i> {11.7}	<i>Medium confidence</i> {11.7}
Changes in frequency of rapidly intensifying tropical cyclones	<i>Likely</i> {11.7}	<i>Medium confidence</i> {11.7}
Poleward migration of tropical cyclones in the western Pacific	<i>Medium confidence</i> {11.7}	<i>Medium confidence</i> {11.7}
Decrease in TC forward motion over the USA	It is <i>likely</i> that TC translation speed has slowed over the USA since 1900. {11.7}	It is <i>more likely than not</i> that the slowdown of TC translation speed over the USA has contributions from anthropogenic forcing. {11.7}
Severe convective storms (tornadoes, hail, rainfall, wind, lightning)	<i>Low confidence</i> in past trends in hail and winds and tornado activity due to short length of high-quality data records. {11.7}	<i>Low confidence.</i> {11.7}
Increase in compound events	<p><i>Likely</i> increase in the probability of compound events.</p> <p><i>High confidence</i> that co-occurrent heat waves and droughts are becoming more frequent under enhanced greenhouse gas forcing at global scale.</p> <p><i>Medium confidence</i> that fire weather, i.e. compound hot, dry and windy events, have become more frequent in some regions.</p> <p><i>Medium confidence</i> that compound flooding risk has increased along the USA coastline. {11.8}</p>	<p><i>Likely</i> that human-induced climate change has increased the probability of compound events.</p> <p><i>High confidence</i> that human influence has increased the frequency of co-occurrent heat waves and droughts.</p> <p><i>Medium confidence</i> that human influence has increased fire weather occurrence in some regions.</p> <p><i>Low confidence</i> that human influences has contributed to changes in compound events leading to flooding. {11.8}</p>

--	--	--

[END TABLE 11.1 HERE]

[START TABLE 11.2 HERE]

**Table 11.2:** Synthesis table on projected changes in extremes. Note that projected changes in marine extremes are assessed in Chapter 9 and the Cross-chapter box 9.1 (marine heat waves). Assessments are provided compared to pre-industrial conditions.

Phenomenon and direction of trend	Projected changes at +1.5°C global warming	Projected changes at +2°C global warming	Projected changes at +4°C global warming
Warmer and/or more frequent hot days and nights over most land areas  Warmer and/or fewer cold days and nights over most land areas  Warm spells/heat waves; Increases in frequency or intensity over most land areas  Cold spells/cold waves: Decreases in frequency or intensity over most land areas	<i>Virtually certain</i> compared to pre-industrial on global scale.  <i>Extremely likely</i> on all continents  Highest increase of temperature of hottest days is projected in some mid-latitude and semi-arid regions, at about 1.5 times to twice the rate of global warming ( <i>high confidence</i> )  Highest increase of temperature of coldest days is projected in Arctic regions, at about three times the rate of global warming ( <i>high confidence</i> ) {11.3}  <i>Continental-scale projections:</i> <i>Extremely likely:</i> Africa, Asia, Australasia, Central and South America, Europe, North America {11.3, 11.9}}	<i>Virtually certain</i> compared to pre-industrial on global scale.  <i>Virtually certain</i> on all continents  Highest increase of temperature of hottest days is projected in some mid-latitude and semi-arid regions, at about 1.5 times to twice the rate of global warming ( <i>high confidence</i> )  Highest increase of temperature of coldest days is projected in Arctic regions, at about three times the rate of global warming ( <i>high confidence</i> ) {11.3}  <i>Continental-scale projections:</i> <i>Virtually certain:</i> Africa, Asia, Australasia, Central and South America, Europe, North America {11.3, 11.9}	<i>Virtually certain</i> compared to pre-industrial on global scale.  <i>Virtually certain</i> on all continents  Highest increase of temperature of hottest days is projected in some mid-latitude and semi-arid regions, at about 1.5 times to twice the rate of global warming ( <i>high confidence</i> )  Highest increase of temperature of coldest days is projected in Arctic regions, at about three times the rate of global warming ( <i>high confidence</i> ) {11.3}  <i>Continental-scale projections:</i> <i>Virtually certain:</i> Africa, Asia, Australasia, Central and South America, Europe, North America {11.3, 11.9}
Heavy precipitation events: increase in the frequency, intensity, and/or amount of heavy precipitation	<i>High confidence</i> that increases take place in most land regions {11.4}  <i>Very likely:</i> Asia, N. America <i>Likely:</i> Africa, Europe <i>High confidence:</i> Central and South America <i>Medium confidence:</i> Australasia {11.4, 11.9}	<i>Likely</i> that increases take place in most land regions {11.4}  <i>Extremely likely:</i> Asia, N. America <i>Very likely:</i> Africa, Europe <i>Likely:</i> Australasia, Central and South America {11.4, 11.9}	<i>Very likely</i> that increases take place in most land regions {11.4}  <i>Virtually certain:</i> Africa, Asia, N. America <i>Extremely likely:</i> Central and South America, Europe <i>Very likely:</i> Australasia {11.4, 11.9}

1  
2  
3  
4  
5  
6  
7  
8  
9  
10

<p>Agricultural and ecological droughts: Increases in intensity and/or duration of drought events</p>	<p><i>High confidence</i> over predominant fraction of land area</p> <p>Land area affected by increasing drought frequency and severity expands with increasing global warming (<i>high confidence</i>). {11.6, 11.9}</p> <p>Precipitation decreases is going to increase the severity of drought in some regions; atmospheric evaporative demand will continue to increase compared to pre-industrial conditions and lead to further increases in agricultural and ecological droughts due to increased evapotranspiration in some regions. (<i>high confidence</i>) {11.6, 11.9}</p>	<p><i>Likely</i> over predominant fraction of land area</p> <p>Land area affected by increasing drought frequency and severity expands with increasing global warming (<i>likely</i>). {11.6, 11.9}</p> <p>Precipitation decreases is going to increase the severity of drought in some regions; atmospheric evaporative demand will continue to increase compared to pre-industrial conditions and lead to further increases in agricultural and ecological droughts due to increased evapotranspiration in some regions. (<i>high confidence</i>) {11.6, 11.9}</p>	<p><i>Very likely</i> over predominant fraction of land area</p> <p>Land area affected by increasing drought frequency and severity expands with increasing global warming (<i>very likely</i>). {11.6, 11.9}</p> <p>Precipitation decreases is going to increase the severity of drought in several regions; atmospheric evaporative demand will continue to increase compared to pre-industrial conditions and lead to further increases in agricultural and ecological droughts due to increased evapotranspiration in several regions. (<i>high confidence</i>) {11.6, 11.9}</p>
<p>Increase in precipitation associated with tropical cyclones (TC)</p>	<p><i>High confidence</i> in a projected increase of TC rain rates at the global scale; the median projected rate of increase due to human emissions is about 11%. {11.7}</p> <p><i>Medium confidence</i> that rain rates will increase in every basin. {11.7}</p>	<p><i>High confidence</i> in a projected increase of TC rain rates at the global scale; the median projected rate of increase due to human emissions is about 14%. {11.7}</p> <p><i>Medium confidence</i> that rain rates will increase in every basin. {11.7}</p>	<p><i>High confidence</i> in a projected increase of TC rain rates at the global scale; the median projected rate of increase due to human emissions is about 28%. {11.7}</p> <p><i>Medium confidence</i> that rain rates will increase in every basin. {11.7}</p>
<p>Increase in mean tropical cyclone lifetime-maximum wind speed (intensity)</p>	<p><i>Medium confidence</i> {11.7}</p>	<p><i>High confidence</i> {11.7}</p>	<p><i>High confidence</i> {11.7}</p>
<p>Increase in likelihood that a TC will be at major TC intensity (Cat. 4-5)</p>	<p><i>High confidence</i> for an increase in the proportion of TCs that reach the strongest (Category 4-5) levels. The median projected increase in this proportion is about 10%. {11.7}</p>	<p><i>High confidence</i> for an increase in the proportion of TCs that reach the strongest (Category 4-5) levels. The median projected increase in this proportion is about 13%. {11.7}</p>	<p><i>High confidence</i> for an increase in the proportion of TCs that reach the strongest (Category 4-5) levels. The median projected increase in this proportion is about 20%. {11.7}</p>
<p>Severe convective storms</p>	<p>There is <i>medium confidence</i> that the frequency of severe convective storms increases in the spring with enhancement of convective available potential energy (CAPE), leading to extension of seasons of occurrence of severe convective storms. There is <i>high confidence</i> of future intensification of precipitation associated with severe convective storms. {11.7}</p>		
<p>Increase in compound events (frequency, intensity)</p>	<p><i>Likely</i> that probability of compound events will continue to increase with global warming.</p> <p><i>High confidence</i> that co-occurrent heat waves and droughts will continue to increase under higher levels of global warming, with higher frequency/intensity with every additional 0.5°C of global warming.</p> <p><i>High confidence</i> that fire weather, i.e. compound hot, dry and windy events, will become more frequent in some regions at higher levels of global warming.</p> <p><i>Medium confidence</i> that compound flooding at the coastal zone will increase under higher levels of global warming, with higher frequency/intensity with every additional 0.5°C of global warming. {11.8}</p>		

1 [END TABLE 11.2 HERE]

2  
3  
4 [START BOX 11.2 HERE]

5  
6  
7 **BOX 11.2: Low-likelihood high-impact changes in extremes**

8  
9 SREX (Chapter 3) assigned *low confidence* to low-probability high-impact (LLHI) events. Such events are  
10 often not anticipated and thus sometimes referred to as surprises. There are several types of LLHI events.  
11 Abrupt changes in mean climate are addressed in Chapter 4. Unanticipated LLHI events can either result  
12 from tipping points in the climate system (Section 1.4.4.3), such as the shutdown of the Atlantic  
13 thermohaline circulation (SROCC Ch6; Collins et al., 2019) or the drydown of the Amazonian rainforest  
14 (SR15 Ch3; Hoegh-Guldberg et al., 2018; Drijfhout et al. 2015), or from uncertainties in climate processes  
15 including climate feedbacks that may enhance or damp extremes either related to global or regional climate  
16 responses (Seneviratne et al., 2018b; Sutton, 2018). The *low confidence* does not by itself exclude the  
17 possibility of such events to occur, it is instead an indication of a poor state of knowledge. Such outcomes,  
18 while *unlikely*, could be associated with very high impacts, and are thus highly relevant from a risk  
19 perspective (see Chapter 1, Section 1.4.3, Box 11.4; Sutton, 2018, 2019). Alternatively, high impacts can  
20 occur when different extremes occur at the same time or in short succession at the same location or in several  
21 regions with shared vulnerability (e.g. food-basket regions Gaupp et al., 2019). These “compound events”  
22 are assessed in Section 11.8 and Box 11.4 provides a case-study example.

23  
24 The difficulties in determining the likelihood of occurrence and time frame of potential tipping points and  
25 LLHI events persist. However, new literature has emerged on unanticipated and low-probability high-impact  
26 events more generally. There are events that are sufficiently rare that they have not been observed in  
27 meteorological records, but whose occurrence is nonetheless plausible within the current state of the climate  
28 system, see examples below and McCollum et al. (2020). The rare nature of such events and the limited  
29 availability of relevant data makes it difficult to estimate their occurrence probability and thus gives little  
30 evidence on whether to include such hypothetical events in planning decisions and risk assessments. The  
31 estimation of such potential surprises is often limited to events that have historical analogues (including  
32 before the instrumental records began, Wetter et al., 2014), albeit the magnitude of the event may differ.  
33 Additionally, there is also a limitation of available resources to exhaust all plausible trajectories of the  
34 climate system. As a result, there will still be events that cannot be anticipated. These events can be surprises  
35 to many in that the events have not been experienced, although their occurrence could be inferred by  
36 statistical means or physical modelling approaches (Chen et al., 2017; van Oldenborgh et al., 2017;  
37 Harrington and Otto, 2018a). Another approach focusing on the estimation of low-probability events and of  
38 events whose likelihood of occurrence is unknown consists in using physical climate models to create a  
39 physically self-consistent storyline of plausible extreme events and assessing their impacts and driving  
40 factors in past (Section 11.2.3) or future conditions (11.2.4) (Cheng et al., 2018; Schaller et al., 2020;  
41 Shepherd, 2016; Shepherd et al., 2018; Sutton, 2018; Zappa and Shepherd, 2017; Wehrli et al., 2020;  
42 Hazeleger et al., 2015).

43  
44 In many parts of the world, observational data are limited to 50-60 years. This means that the chance to  
45 observe an extreme event that occurs once in several hundred or more years is small. Thus, when a very  
46 extreme event occurs, it becomes a surprise to many (Bao et al., 2017; McCollum et al., 2020), and very rare  
47 events are often associated with high impacts (van Oldenborgh et al., 2017; Philip et al., 2018b; Tozer et al.,  
48 2020). Attributing and projecting very rare events in a particular location by assessing their likelihood of  
49 occurrence within the same larger region and climate thus provides another way to make quantitative  
50 assessments regarding events that are extremely rare locally. Some examples of such events include for  
51 instance:

- 52 • Hurricane Harvey, that made landfall in Houston, TX in August 2017 (Section 11.7.1.4.)
- 53 • The 2010-2011 extreme floods in Queensland, Australia (Christidis et al., 2013a)
- 54 • The 2018 concurrent heat waves across the northern Hemisphere (Box 11.4)
- 55 • Tropical cyclone Idai in Mozambique (Cross-Chapter Box Disaster in WGII AR6 Chapter 4)

- The California fires in 2018 and 2019
- The 2019-2020 Australia fires (Cross-Chapter Box Disaster in WGII AR6 Chapter 4)

One factor making such events hard to anticipate is the fact that we now live in a non-stationary climate, and that the framework of reference for adaptation is continuously moving. As an example, the concurrent heat waves that occurred across the Northern Hemisphere in the summer of 2018 were considered very unusual and were indeed unprecedented given the total area that was concurrently affected (Toreti et al., 2019; Vogel et al., 2019; Drouard et al., 2019; Kornhuber et al., 2019); however, the probability of this event under 1°C global warming was found to be about 16% (Vogel et al., 2019), which is not particularly low. Similarly, the 2013 summer temperature over eastern China was the hottest on record at the time, but it had an estimated recurrence interval of about 4 years in the climate of 2013 (Sun et al., 2014). Furthermore, when other aspects of the risk, vulnerability, and exposure are historically high or have recently increased (see WGII, Chapter 16, Section 16.4), relatively moderate extremes can have very high impacts (Otto et al., 2015b; Philip et al., 2018b). As warming continues, the climate moves further away from its historical state with which we are familiar, resulting in an increased likelihood of unprecedented events and surprises. This is particularly the case under high global warming levels e.g. such as the climate of the late 21st century under high-emissions scenarios (above 4°C of global warming, CC-Box 11.1).

Another factor highlighted in Section 11.8 and Box 11.4 making events high-impact and difficult to anticipate is that several locations under moderate warming levels could be affected simultaneously, or very repeatedly by different types of extremes (Mora et al., 2018, Gaupp et al., 2019; Vogel et al., 2019). Box 11.4 shows that concurrent events at different locations, which can lead to major impacts across the world, can also result from the combination of anomalous circulation or natural variability (ENSO) patterns with amplification of resulting responses to human-induced global warming. Also multivariate extremes at single locations pose specific challenges to anticipation (Section 11.8), with low-likelihoods in the current climate but the probability of occurrence of such compound events strongly increasing with increasing global warming levels (Vogel et al., 2020a). Therefore, in order to estimate whether and at what level of global warming very high impacts arising from extremes would occur, the spatial extent of extremes and the potential of compounding extremes need to be assessed. Sections 11.3, 11.4, 11.7 and 11.8 highlight increasing evidence that temperature extremes, higher intensity precipitation accompanying tropical cyclones, and compound events such as dry/hot conditions conducive to wildfire or storm surges resulting from sea level rise and heavy precipitation events, pose widespread threats to societies already at relatively low warming levels. Studies have already shown that the probability for some recent extreme events is so small in the undisturbed world such that these events may not have been possible without human influence (Section 11.2.4). Box 11.2, Table 1, provides examples of projected changes in LLHI extremes (single extremes, compound events) of potential relevance for impact and adaptation assessments showing that today's very rare events can become commonplace in a warmer future.

In summary, the future occurrence of LLHI events linked to climate extremes is generally associated with *low confidence*, but cannot be excluded, especially at global warming levels above 4°C. Compound events, including concurrent extremes, are a factor increasing the probability of LLHI events (*high confidence*). With increasing global warming some compound events with low likelihood in past and current climate will become more frequent, and there is a higher chance of historically unprecedented events and surprises (*high confidence*). However, even extreme events that do not have a particularly low probability in the present climate (at more than 1°C of global warming) can be perceived as surprises because of the pace of global warming (*high confidence*).

**Box 11.2, Table 1:** Examples of changes in LLHI extreme conditions (single extremes, compound events) at different global warming levels

	+1°C (present-day)	+1.5°C	+2°C	+3°C and higher
Risk ratio for annual hottest daytime temperature (TXx) with 1% of probability under present-day warming (+1°C) (Kharin et	1	3.3 (i.e. 230% higher probability)	8.2 (i.e. 720% higher probability)	Not assessed

al., 2018): Global land				
Risk ratio for heavy precipitation events (Rx1day) with 1% of probability under present-day warming (+1°C) (Kharin et al., 2018): Global land	1	1.2 (i.e. 20% higher probability)	1.5 (i.e. 50% higher probability)	Not assessed
Risk ratio for 1- 5 day duration extreme floods with 1% of probability under present-day warming (+1°C) (Ali et al., 2019a) Indian subcontinent	Up to 3 in individual locations	Up to 5 in individual locations	2-6 in most locations	Up to 12 in individual locations (4°C)
Probability of “extremes extremes” hot days with 1/1000 probability at the end of 20 <sup>th</sup> century (Vogel et al., 2020a): Global land	~20 days over 20 years in most locations	about ~50 days in 20 years in most locations	about ~150 days in 20 years in most locations	about ~500 days in 20 years in most locations (3°C)
Probability of co-occurrence in the same week of hot days with 1/1000 probability and dry days with 1/1000 probability at the end of 20 <sup>th</sup> century (Vogel et al., 2020a): Amazon	0% probability	~1 week within 20 years	~4-5 weeks within 20 years	>9 weeks within 20 years (3°C)
Projected soil moisture drought duration per year (Samaniego et al., 2018): Mediterranean region	41 days (+46% compared to late 20 <sup>th</sup> century)	58 days (+107% compared to late 20 <sup>th</sup> century)	71 days (+154% compared to late 20 <sup>th</sup> century)	125 days (+346% compared to late 20 <sup>th</sup> century) (3°C)
Increase in days exposed to dangerous extreme heat (measured in Health Heat Index (HHI) (Sun et al., 2019c) global land	Not assessed, baseline is 1981-2000	1.6 times higher risk of experiencing heat > 40.6	2.3 times higher risk of experiencing heat > 40.6	~ 80% of land area exposed to dangerous heat, tropical regions 1/3 of the year (4°C)
Increase in regional mean fire season length (Sun et al., 2019c; Xu et al., 2020) global land	Not assessed, baseline is 1981-2000	6.2 days	9.5 days	~ 50 days (4°C)

1  
2  
3 **[END BOX 11.2 HERE]**  
4  
5

## 6 **11.2 Data and Methods**

7  
8 This section provides an assessment of observational data and methods used in the analysis and attribution of  
9 climate change specific to weather and climate extremes, and also introduces some concepts used in  
10 presenting future projections of extremes in the chapter. The main focus is on extreme events over land, as  
11 extremes in the ocean are assessed in Chapter 9 of this Report. Later sections (11.3-11.8) also provide  
12 additional assessments on relevant observational datasets and model validation specific for the type of  
13 extremes to be assessed. General background on climate modelling is provided in Chapters 4 and 10.  
14

### 15 **11.2.1 Definition of extremes**

16  
17 In the literature, an event is generally considered extreme if the value of a variable exceeds (or lies below) a  
18

1 threshold. The thresholds have been defined in different ways, leading to differences in the meaning of  
2 extremes that may share the same name. For example, two sets of frequency of hot/warm days have been  
3 used in the literature. One set counts the number of days when maximum daily temperature is above a  
4 relative threshold defined as the 90<sup>th</sup> or higher percentile of maximum daily temperature for the calendar day  
5 over a base period. An event based on such a definition can occur during any time of the year and the impact  
6 of such an event would differ depending on the season. The other set counts the number of days in which  
7 maximum daily temperature is above an absolute threshold such as 35°C, because exceedance of this  
8 temperature can sometimes cause health impacts (however, these impacts may depend on location and  
9 whether ecosystems and the population are adapted to such temperatures). While both types of hot extreme  
10 indices have been used to analyze changes in the frequency of hot/warm events, they represent different  
11 events that occur at different times of the year, possibly affected by different types of processes and  
12 mechanisms, and possibly also associated with different impacts.

13  
14 Changes in extremes have also been examined from two perspectives: changes in the frequency for a given  
15 magnitude of extremes or changes in the magnitude for a particular return period (frequency). Changes in the  
16 probability of extremes (e.g., temperature extremes) depend on the rarity of the extreme event that is  
17 assessed, with a larger change in probability associated with a rarer event (e.g., Kharin et al., 2018). On the  
18 other hand, changes in the magnitude represented by the return levels of the extreme events may not be as  
19 sensitive to the rarity of the event. While the answers to the two different questions are related, their  
20 relevance to different audiences may differ. Conclusions regarding the respective contribution of greenhouse  
21 gas forcing to changes in magnitude versus frequency of extremes may also differ (Otto et al., 2012).  
22 Correspondingly, the sensitivity of changes in extremes to increasing global warming is also dependent on  
23 the definition of the considered extremes. In the case of temperature extremes, changes in magnitude have  
24 been shown to often depend linearly on global surface temperature (Seneviratne et al., 2016; Wartenburger et  
25 al., 2017), while changes in frequency tend to be non-linear and can, for example, be exponential for  
26 increasing global warming levels (Fischer and Knutti, 2015; Kharin et al., 2018). When similar damage  
27 occurs once a fixed threshold is exceeded, it is more important to ask a question regarding changes in the  
28 frequency. But when the exceedance of this fixed threshold becomes a normal occurrence in the future, this  
29 can lead to a saturation in the change of probability (Harrington and Otto, 2018a). On the other hand, if the  
30 impact of an event increases with the intensity of the event, it would be more relevant to examine changes in  
31 the magnitude. Finally, adaptation to climate change might change the relevant thresholds over time,  
32 although such aspects are still rarely integrated in the assessment of projected changes in extremes. Framing,  
33 including how extremes are defined and how the questions are asked in the literature, is considered when  
34 forming the assessments of this chapter.

### 35 36 37 **11.2.2 Data**

38  
39 Studies of past and future changes in weather and climate extremes and in the mean state of the climate use  
40 the same original sources of weather and climate observations, including in-situ observations, remotely  
41 sensed data, and derived data products such as reanalyses. Chapter 2 (Section 2.3) and Chapter 10 (Section  
42 10.2) assess various aspects of these data sources and data products from the perspective of their general use  
43 and in the analysis of changes in the mean state of the climate in particular. Building on these previous  
44 chapters, this subsection highlights particular aspects that are related to extremes and that are most relevant  
45 to the assessment of this chapter. The SREX (Chapter 3, Seneviratne et al., 2012) and AR5 (Chapter 2,  
46 Hartmann et al., 2013) addressed critical issues regarding the quality and availability of observed data and  
47 their relevance for the assessment of changes in extremes.

48  
49 Extreme weather and climate events occur on time scales of hours (e.g., convective storms that produce  
50 heavy precipitation) to days (e.g., tropical cyclones, heat waves), to seasons and years (e.g., droughts). A  
51 robust determination of long-term changes in these events can have different requirements for the spatial and  
52 temporal scales and sample size of the data. In general, it is more difficult to determine long-term changes  
53 for events of fairly large temporal duration, such as “mega-droughts” that last several years or longer (e.g.,  
54 Ault et al. 2014), because of the limitations of the observational sample size. Literature that study changes in  
55 extreme precipitation and temperature often use indices representing specifics of extremes that are derived



1 from daily precipitation and temperature values. Station-based indices would have the same issues as those  
2 for the mean climate regarding the quality, availability, and homogeneity of the data. For the purpose of  
3 constructing regional information and/or for comparison with model outputs, such as model evaluation, and  
4 detection and attribution, these station-based indices are often interpolated onto regular grids. Two different  
5 approaches, involving two different orders of operation, have been used in producing such gridded datasets.  
6

7 In some cases, such as for the HadEX3 dataset (Dunn et al., 2020), indices of extremes are computed using  
8 time series directly derived from stations first and are then gridded over the space. As the indices are  
9 computed at the station level, the gridded data products represent point estimates of the indices averaged  
10 over the spatial scale of the grid box. In other instances, daily values of station observations are first gridded  
11 (e.g., Contractor et al., 2020), and the interpolated values can then be used to compute various indices by the  
12 users. Depending on the station density, values for extremes computed from data gridded this way represent  
13 extremes of spatial scales anywhere from the size of the grid box to a point. In regions with high station  
14 density (e.g., North America, Europe), the gridded values are closer to extremes of area means and are thus  
15 more appropriate for comparisons with extremes estimated from climate model output, which is often  
16 considered to represent areal means (Chen and Knutson, 2008; Gervais et al., 2014; Avila et al., 2015; Di  
17 Luca et al., 2020a). In regions with very limited station density (e.g., Africa), the gridded values are closer to  
18 point estimates of extremes. The difference in spatial scales among observational data products and model  
19 simulations needs to be carefully accounted for when interpreting the comparison among different data  
20 products. For example, the average annual maximum daily maximum temperature (TXx) over land  
21 computed from the original ERA-interim reanalysis (at 0.75° resolution) is about 0.4°C warmer than that  
22 computed when the ERA-interim dataset is upscaled to the resolution of 2.5° x 3.75° (Di Luca et al., 2020).  
23

24 Extreme indices computed from various reanalysis data products have been used in some studies, but  
25 reanalysis extreme statistics have not been rigorously compared to observations (Donat et al., 2016a).  
26 In general, changes in temperature extremes from various reanalyses were most consistent with gridded  
27 observations after about 1980, but larger differences were found during the pre-satellite era (Donat et al.,  
28 2014b). Overall, lower agreement across reanalysis datasets was found for extreme precipitation changes,  
29 although temporal and spatial correlations against observations were found to be still significant. In regions  
30 with sparse observations (e.g., Africa and parts of South America), there is generally less agreement for  
31 extreme precipitation between different reanalysis products, indicating a consequence of the lack of an  
32 observational constraint in these regions (Donat et al., 2014b, 2016a). More recent reanalyses, such as ERA5  
33 (Hersbach et al., 2020), seem to have improved over previous products, at least over some regions (e.g.,  
34 Mahto and Mishra, 2019; Gleixner et al., 2020; Sheridan et al., 2020). Caution is needed when reanalysis  
35 data products are used to provide additional information about past changes in these extremes in regions  
36 where observations are generally lacking.  
37

38 Satellite remote sensing data have been used to provide information about precipitation extremes because  
39 several products provide data at sub-daily resolution for precipitation (e.g., TRMM; Maggioni et al. 2016)  
40 and clouds (e.g., HIMAWARI; Bessho et al., 2016; Chen et al. 2019). However, satellites do not observe the  
41 primary atmospheric state variables directly and polar orbiting satellites do not observe any given place at all  
42 times. Hence, their utility as a substitute for high-frequency (i.e., daily) ground-based observations is limited.  
43 For instance, Timmermans et al. (2019) found little relationship between the timing of extreme daily and  
44 five-day precipitation in satellite and gridded station data products over the United States.  
45  
46

47 **[START BOX 11.3 HERE]**

### 48 **BOX 11.3: Extremes in paleoclimate archives compared to instrumental records**

49 Examining extremes in pre-instrumental information can help to put events occurring in the instrumental  
50 record (referred to as ‘observed’) in a longer-term context. This box focuses on extremes in the Common Era  
51 (CE, the last 2000 years), because there is generally higher confidence in pre-instrumental information  
52 gathered from the more recent archives from the Common Era than from earlier evidence. It addresses  
53 evidence of extreme events in paleo reconstructions, documentary evidence (such as grape harvest data,  
54 religious documents, newspapers, and logbooks) and model-based analyses, and whether observed extremes  
55

1 have or have not been exceeded in the Common Era. This box provides overviews of i) AR5 assessments  
2 and ii) types of evidence assessed here, evidence of iii) droughts, iv) temperature extremes, v) paleofloods,  
3 and vi) paleotempests, and vii) a summary.

4  
5 AR5 (Chapter 5, Masson-Delmotte et al., 2013) concluded with *high confidence* that droughts of greater  
6 magnitude and of longer duration than those observed in the instrumental period occurred in many regions  
7 during the preceding millennium. There was *high confidence* in evidence that floods during the past five  
8 centuries in northern and Central Europe, the western Mediterranean region, and eastern Asia were of a  
9 greater magnitude than those observed instrumentally, and *medium confidence* in evidence that floods in the  
10 near East, India and central North America were comparable to modern observed floods. While AR5  
11 assessed 20<sup>th</sup> century summer temperatures compared to those reconstructed in the Common Era, it did not  
12 assess shorter duration temperature extremes.

13  
14 Many factors affect confidence in information on pre-instrumental extremes. First, the geographical coverage  
15 of paleoclimate reconstructions of extremes is not spatially uniform (Smerdon and Pollack, 2016) and  
16 depends on both the availability of archives and records, which are environmentally dependent, and also the  
17 differing attention and focus from the scientific community. In Australia, for example, the paleoclimate  
18 network is sparser than for other regions, such as Asia, Europe and North America, and synthesised products  
19 rely on remote proxies and assumptions about the spatial coherence of precipitation between remote climates  
20 (Cook et al., 2016c; Freund et al., 2017). Second, pre-instrumental evidence of extremes may be focused on  
21 understanding archetypal extreme events, such as the climatic consequences of the 1815 eruption of Mount  
22 Tambora, Indonesia (Brohan et al., 2016; Veale and Endfield, 2016). These studies provide narrow evidence  
23 of extremes in response to specific forcings (Li, 2017) for specific epochs. Third, natural archives may  
24 provide information about extremes in one season only and may not represent all extremes of the same types.

25  
26 Evidence of shorter duration extreme event types, such as floods and tropical storms, is further restricted by  
27 the comparatively low chronological controls and temporal resolution (e.g., monthly, seasonal, yearly,  
28 multiple years) of most archives compared to the events (e.g., minutes to days). Natural archives may be  
29 sensitive only to intense environmental disturbances, and so only sporadically record short-duration or small  
30 spatial scale extremes. Interpreting sedimentary records as evidence of past short-duration extremes is also  
31 complex and requires a clear understanding of natural processes. For example, paleoflood reconstructions of  
32 flood recurrence and intensity produced from geological evidence (e.g., river and lake sediments),  
33 speleothems (Denniston and Luetscher, 2017), botanical evidence (e.g., flood damage to trees, or tree ring  
34 reconstructions), and floral and faunal evidence (e.g., diatom fossil assemblages) require understanding of  
35 sediment sources and flood mechanisms. Pre-instrumental records of tropical storm intensity and frequency  
36 (also called paleotempest records) derived from overwash deposits of coastal lake and marsh sediments are  
37 difficult to interpret. Many factors impact whether disturbances are deposited in archives (Muller et al.,  
38 2017) and deposits may provide sporadic and incomplete preservation histories (e.g., Tamura et al., 2018).

39  
40 Overall, the most complete pre-instrumental evidence of extremes occurs for long-duration, large-spatial-  
41 scale extremes, such as for multi-year meteorological droughts or seasonal- and regional-scale temperature  
42 extremes. Additionally, more precise insights into recent extremes emerge where multiple studies have been  
43 undertaken, compared to the confidence in extremes reported at single sites or in single studies, which may  
44 not necessarily be representative of large-scale changes, or for reconstructions that synthesise multiple  
45 proxies over large areas (e.g., drought atlases). Multiproxy synthesis products combine paleoclimate  
46 temperature reconstructions and cover sub-continental- to hemispheric-scale regions to provide continuous  
47 records of the Common Era (e.g. Ahmed et al., 2013; Neukom et al., 2014 for temperature).

48  
49 There is *high confidence* in the occurrence of long-duration and severe drought events during the Common  
50 Era for many locations, although their severity compared to recent drought events differs between locations  
51 and the lengths of reconstruction provided. Recent observed drought extremes in some regions (such as the  
52 Levant (Cook et al., 2016a), California in the United States (Cook et al., 2014; Griffin and Anchukaitis,  
53 2014), and the Andes (Domínguez-Castro et al., 2018)) do not have precedents within the multi-century  
54 periods reconstructed in these studies, in terms of duration and/or severity. In some regions (in Southwest  
55 North America (Asmerom et al., 2013; Cook et al., 2015), the Great Plains region (Cook et al., 2004), the

1 Middle East (Kaniewski et al., 2012), and China (Gou et al., 2015), recent drought extremes may have been  
2 exceeded in the Common Era. In further locations, there is conflicting evidence for the severity of pre-  
3 instrumental droughts compared to observed extremes, depending on the length of the reconstruction and the  
4 seasonal perspective provided (see Cook et al., 2016b; Freund et al., 2017 for Australia). There can also be  
5 differing conclusions for the severity, or even the occurrence, of specific individual pre-instrumental  
6 droughts when different evidence is compared (e.g., Büntgen et al., 2015; Wetter et al., 2014).

7  
8 There is *medium confidence* that the magnitude of large-scale, seasonal-scale extreme high temperatures in  
9 observed records exceed those reconstructed over the Common Era in some locations, such as Central  
10 Europe. In one example, multiple studies have examined the unusualness of present-day European summer  
11 temperature records in a long-term context, particularly in comparison to the exceptionally warm year of  
12 1540 CE in Central Europe. Several studies indicate recent extreme summers (2003 and 2010) in Europe  
13 have been unusually warm in the context of the last 500 years (Barriopedro et al., 2011; Wetter and Pfister,  
14 2013; Wetter et al., 2014; Orth et al., 2016a), or longer (Luterbacher et al., 2016). Others studies show  
15 summer temperatures in Central Europe in 1540 were warmer than the present-day (1966–2015) mean, but  
16 note that it is difficult to assess whether or not the 1540 summer was for its part warmer than observed  
17 record extreme temperatures (Orth et al., 2016a).

18  
19 There is *high confidence* that the magnitude of floods over the Common Era has exceeded observed records  
20 in some locations, including Central Europe and eastern Asia. Recent literature supports the AR5  
21 assessments (Masson-Delmotte et al., 2013) of floods. High temporally resolved records provide evidence,  
22 for example, of Common Era floods exceeding the probable maximum flood levels in the Upper Colorado  
23 River, USA (Greenbaum et al., 2014) and peak discharges that are double gauge levels along the middle  
24 Yellow River, China (Liu et al., 2014). Further studies demonstrate pre-instrumental or early instrumental  
25 differences in flood frequency compared to the instrumental period, including reconstructions of high and  
26 low flood frequency in the European Alps (e.g., Swierczynski et al., 2013; Amann et al., 2015) and  
27 Himalayas (Ballesteros Cánovas et al., 2017). The combination of extreme historical flood episodes  
28 determined from documentary evidence also increases confidence in the determination of flood frequency  
29 and magnitude, compared to using geomorphological archives alone (Kjeldsen et al., 2014). In regions, such  
30 as Europe and China, that have rich historical flood documents, there is strong evidence of high magnitude  
31 flood events over pre-instrumental periods (Benito et al., 2015; Kjeldsen et al., 2014; Macdonald and  
32 Sangster, 2017). A key feature of paleoflood records is variability in flood recurrence at centennial  
33 timescales (Wilhelm et al., 2019), although constraining climate-flood relationships remains challenging.  
34 Pre-instrumental floods often occurred in considerably different contexts in terms of land use, irrigation, and  
35 infrastructure, and may not provide direct insight into modern river systems, which further prevents long-  
36 term assessments of flood changes being made based on these sources.

37  
38 There is *medium confidence* that periods of both more and less tropical cyclone activity (frequency or  
39 intensity) than observed occurred over the Common Era in many regions. Paleotempest studies cover a  
40 limited number of locations that are predominantly coastal, and hence provide information on specific  
41 locations that cannot be extrapolated basin-wide (see Muller et al., 2017). In some locations, such as the Gulf  
42 of Mexico and the New England coast, similarly intense storms to those observed recently have occurred  
43 multiple times over centennial timescales (Donnelly et al., 2001; Bregy et al., 2018). Further research  
44 focused on the frequency of tropical storm activity. Extreme storms occurred considerably more frequently  
45 in particular periods of the Common Era, compared to the instrumental period in northeast Queensland,  
46 Australia (Nott et al., 2009; Haig et al., 2014), and the Gulf Coast (e.g., Brandon et al., 2013; Lin et al.,  
47 2014).

48  
49 The probability of finding an unprecedented extreme event increases with an increased length of past record-  
50 keeping, in the absence of longer-term trends. Thus, as a record is extended to the past based on paleo-  
51 reconstruction, there is a higher chance of very rare extreme events having occurred at some time prior to  
52 instrumental records. Such an occurrence is not, in itself, evidence of a change, or lack of a change, in the  
53 magnitude or the likelihood of extremes in the past or in the instrumental period at regional and local scales.  
54 Yet, the systematic collection of paleoclimate records over wide areas may provide evidence of changes in  
55 extremes. In one study, extended evidence of the last millennium from observational data and paleoclimate

1 reconstructions using tree rings indicates human activities affected the worldwide occurrence of droughts as  
2 early as the beginning of the 20<sup>th</sup> century (Marvel et al., 2019).

3  
4 In summary, there is *low confidence* in overall changes in extremes derived from paleo-archives. The most  
5 robust evidence is *high confidence* that high-duration and severe drought events occurred at many locations  
6 during the last 2000 years. There is also *high confidence* that high-magnitude flood events occurred at some  
7 locations during the last 2000 years, but overall changes in infrastructure and human water management  
8 make the comparison with present-day records difficult. But these isolated paleo-drought and paleo-flood  
9 events are not evidence of a change, or lack of a change, in the magnitude or the likelihood of relevant  
10 extremes.

11  
12 **[END BOX 11.3 HERE]**

### 13 14 15 **11.2.3 Attribution of extremes**

16  
17 Attribution science concerns the identification of causes for changes in characteristics of the climate system  
18 (e.g., trends, single extreme events). A general overview and summary of methods of attribution science is  
19 provided in the Cross-Working Group Box 1.1 (in Chapter 1). Trend detection using optimal fingerprinting  
20 methods is a well-established field, and has been assessed in the AR5 (Chapter 10, Bindoff et al., 2013), and  
21 Chapter 3 in this Report (Section 3.2.1). There are specific challenges when applying optimal fingerprinting  
22 to the detection and attribution of trends in extremes and on regional scales where the lower signal-to-noise  
23 ratio is a challenge. In particular, the method generally requires the data to follow a Normal (Gaussian)  
24 distribution, which is often not the case for extremes. Recent studies showed that extremes can, however, be  
25 transformed to a Gaussian distribution, for example by averaging over space, so that optimal fingerprinting  
26 techniques can still be used (Zhang et al., 2013; Wen et al., 2013; and Wan et al., 2019). Non-stationary  
27 extreme value distributions, which allow for the detailed detection and attribution of regional trends in  
28 temperature extremes, have also been used (Wang et al., 2017c).

29  
30 Apart from the detection and attribution of trends in extremes, new approaches have been developed to  
31 answer the question of whether and to what extent external drivers have altered the probability and intensity  
32 of an individual extreme event (NASEM, 2016). In AR5, there was an emerging consensus that the role of  
33 external drivers of climate change in specific extreme weather events could be estimated and quantified in  
34 principle, but related assessments were still confined to particular case studies, often using a single model,  
35 and typically focusing on high-impact events with a clear attributable signal.

36  
37 However, since AR5, the attribution of extreme weather events has emerged as a growing field of climate  
38 research with an increasing body of literature (see series of supplements to the annual State of the Climate  
39 report (Peterson et al., 2012, 2013b, Herring et al., 2014, 2015, 2016, 2018), including the number of  
40 approaches to examining extreme events (described in Easterling et al., 2016; Otto, 2017; Stott et al., 2016)).  
41 A commonly-used approach, often called the risk-based approach in the literature and referred to here as the  
42 “probability-based approach”, produces statements such as ‘anthropogenic climate change made this event  
43 type twice as likely’ or ‘anthropogenic climate change made this event 15% more intense’. This is done by  
44 estimating probability distributions of the index characterizing the event in today’s climate, as well as in a  
45 counterfactual climate, and either comparing intensities for a given occurrence probability (e.g., 1-in-100  
46 year event) or probabilities for a given magnitude (see FAQ 11.3). There are a number of different analytical  
47 methods encompassed in the probability-based approach building on observations and statistical analyses  
48 (e.g., van Oldenborgh et al., 2012), optimal fingerprint methods (Sun et al., 2014), regional climate and  
49 weather forecast models (e.g., Schaller et al., 2016), global climate models (GCMs) (e.g., Lewis and Karoly,  
50 2013), and large ensembles of atmosphere-only GCMs (e.g., Lott et al., 2013). A key component in any  
51 event attribution analysis is the level of conditioning on the state of the climate system. In the least  
52 conditional approach, the combined effect of the overall warming and changes in the large-scale atmospheric  
53 circulation are considered and often utilize fully coupled climate models (Sun et al., 2014). Other more  
54 conditional approaches involve prescribing certain aspects of the climate system. These range from  
55 prescribing the pattern of the surface ocean change at the time of the event (e.g. Hoerling et al., 2013, 2014),

1 often using AMIP-style global models, where the choice of sea surface temperature and ice patterns  
2 influences the attribution results (Sparrow et al., 2018), to prescribing the large-scale circulation of the  
3 atmosphere and using weather forecasting models or methods (e.g., Pall et al., 2017; Patricola and Wehner,  
4 2018; Wehner et al., 2018a). These highly conditional approaches have also been called “storylines”  
5 (Shepherd, 2016; Cross-Working Group Box 1.1 in Chapter 1) and can be useful when applied to extreme  
6 events that are too rare to otherwise analyse or where the specific atmospheric conditions were central to the  
7 impact. These methods are also used to enable the use of very-high-resolution simulations in cases where  
8 lower-resolution models do not simulate the regional atmospheric dynamics well (Shepherd, 2016; Shepherd  
9 et al., 2018). However, the imposed conditions limit an overall assessment of the anthropogenic influence on  
10 an event, as the fixed aspects of the analysis may also have been affected by climate change. For instance,  
11 the specified initial conditions in the highly conditional hindcast attribution approach often applied to  
12 tropical cyclones (e.g., Patricola and Wehner, 2018; Takayabu et al., 2015) permit only a conditional  
13 statement about the magnitude of the storm if similar large-scale meteorological patterns could have  
14 occurred in a world without climate change, thus precluding any attribution statement about the change in  
15 frequency if used in isolation. Combining conditional assessments of changes in the intensity with a multi-  
16 model approach does allow for the latter as well (Shepherd, 2016).

17  
18 The outcome of event attribution is dependent on the definition of the event (Leach et al., 2020), as well as  
19 the framing (Christidis et al., 2018; Jézéquel et al., 2018; Otto et al., 2016) and uncertainties in observations  
20 and modelling. Observational uncertainties arise both in estimating the magnitude of an event as well as its  
21 rarity (Angélil et al., 2017). Results of attribution studies can also be very sensitive to the choice of climate  
22 variables (Sippel and Otto, 2014; Wehner et al., 2016). Attribution statements are also dependent on the  
23 spatial (Uhe et al., 2016; Cattiaux and Ribes, 2018; Kirchmeier-Young et al., 2019) and temporal  
24 (Harrington, 2017; Leach et al., 2020) extent of event definitions, as events of different scales involve  
25 different processes (Zhang et al., 2020d) and large-scale averages generally yield higher attributable changes  
26 in magnitude or probability due to the smoothing out of the noise. In general, confidence in attribution  
27 statements for large-scale heat and lengthy extreme precipitation events have higher confidence than shorter  
28 and more localized events, such as extreme storms, an aspect also relevant for determining the emergence of  
29 signals in extremes or the confidence in projections (see also Cross-Chapter Box Atlas.1)

30  
31 The reliability of the representation of the event in question in the climate models used in a study is essential  
32 (Angélil et al., 2016; Herger et al., 2018). Extreme events characterized by atmospheric dynamics that stretch  
33 the capabilities of current-generation models (see Section 10.3.3.4, Shepherd, 2014; Woollings et al., 2018)  
34 limit the applicability of the probability-based approach of event attribution. The lack of model evaluation, in  
35 particular in early event attribution studies, has led to criticism of the emerging field of attribution science as  
36 a whole (Trenberth et al., 2015) and of individual studies (Angélil et al., 2017). In this regard, the storyline  
37 approach (Shepherd, 2016) provides an alternative option that does not depend on the model’s ability to  
38 represent the circulation reliably. In addition, several ways of quantifying statistical uncertainty (Paciorek et  
39 al., 2018) and model evaluation (Lott and Stott, 2016; Philip et al., 2018b, 2020) have been employed to  
40 evaluate the robustness of event attribution results. For the unconditional probability-based approach, multi-  
41 model and multi-approach (e.g., combining observational analyses and model experiments) methods have  
42 been used to improve the robustness of event attribution (Hauser et al., 2017; Otto et al., 2018a; Philip et al.,  
43 2018b, 2019, 2020; van Oldenborgh et al., 2018; Kew et al., 2019).

44  
45 In the regional tables provided in Section 11.9, the different lines of evidence from event attribution studies  
46 and trend attributions are assessed alongside one another to provide an assessment of the human contribution  
47 to observed changes in extremes in all AR6 regions .

#### 48 49 50 **11.2.4 Projecting changes in extremes as a function of global warming levels**

51  
52 The most important quantity used to characterize past and future climate change is global warming relative  
53 to its pre-industrial level. On the one hand, changes in global warming are linked quasi-linearly to global  
54 cumulative CO<sub>2</sub> emissions (IPCC, 2013). On the other hand, changes in regional climate, including many  
55 types of extremes, scale quasi-linearly with changes in global warming, often independently of the

1 underlying emissions scenarios (SR15 Ch3; Seneviratne et al., 2016; Wartenburger et al., 2017; Matthews et  
2 al., 2017; Tebaldi and Knutti 2018, Sun et al., 2018a, Kharin et al., 2018, Beusch et al., 2020b; Li et al.,  
3 2020). Finally, the use of global warming levels in the context of global policy documents (in particular the  
4 2015 Paris Agreement, UNFCCC 2015), implies that information on changes in the climate system, and in  
5 particular extremes, as a function of global warming are of particular policy relevance. Cross-Chapter Box  
6 11.1 provides an overview on the translation between information at global warming levels (GWLs) and  
7 scenarios.

8  
9 The assessment of projections of future changes in extremes as function of GWL has an advantage in  
10 separating uncertainty associated with the global warming response (see Chapter 4) from the uncertainty  
11 resulting from the regional climate response as a function of GWLs (Seneviratne and Hauser, 2020). If the  
12 interest is in the projection of regional changes at certain GWLs, such as those defined by the Paris  
13 Agreement, projections based on time periods and emission scenarios have unnecessarily larger uncertainty  
14 due to differences in model global transient climate responses. To take advantage of this feature and to  
15 provide easy comparison with SR15, assessments of projected changes in this chapter are largely provided in  
16 relation to future GWLs, with a focus on changes at +1.5°C, +2°C, and +4°C of global warming above pre-  
17 industrial levels (e.g. Tables 11.1, 11.2 and regional tables in Section 11.9). These encompass a scenario  
18 compatible with the aim of the Paris Agreement (+1.5°C), a scenario slightly overshooting the aims of the  
19 Paris Agreement (+2°C), and a “worst-case” scenario with no mitigation (+4°C). The CC-Box 11.1 provides  
20 a background on the GWL sampling approach used in the AR6, both for the computation of GWL  
21 projections from ESMs contributing to the 6<sup>th</sup> Phase of the Coupled Model Intercomparison Project (CMIP6)  
22 as well as for the mapping of existing scenario-based literature for CMIP6 and the 5<sup>th</sup> Phase of CMIP  
23 (CMIP5) to assessments as function of GWLs (see also Section 11.9. and Table 11.3 for an example).

24  
25 While regional changes in many types of extremes do scale robustly with global surface temperature,  
26 generally irrespective of emission scenarios (Section 11.1.4; Figures 11.3, 11.6, 11.7; CC-Box 11.1), effects  
27 of local forcing can distort this relation. In particular, emission scenarios with the same radiative forcing can  
28 have different regional extreme precipitation responses resulting from different aerosol forcing (Wang et al.,  
29 2017d). Another example is related to forcing from land use and land cover changes (Section 11.1.6).  
30 Climate models often either overestimate or underestimate observed changes in annual maximum daily  
31 maximum temperature depending on the region and considered models (Donat et al., 2017; Vautard et al.,  
32 1999). Part of the discrepancies may be due to the lack of representation of some land forcings, in particular  
33 crop intensification and irrigation (Mueller et al., 2016b; Thiery et al., 2017; Findell et al., 2017; Thiery et  
34 al., 2020). Since these local forcings are not represented and their future changes are difficult to project,  
35 these can be important caveats when using GWL scaling to project future changes for these regions.  
36 However, these caveats also apply to the use of scenario-based projections.

37  
38 SR15 (Chapter 3) assessed different climate responses at +1.5°C of global warming, including transient  
39 climate responses, short-term stabilization responses, and long-term equilibrium stabilization responses, and  
40 their implications for future projections of different extremes. Indeed, the temporal dimension, that is, when  
41 the given GWL occurs, also matters for projections, in particular beyond the 21st century and for some  
42 climate variables with large inertia (e.g., sea level rise and associated extremes). Nonetheless, for  
43 assessments focused on conditions within the next decades and for the main extremes considered in this  
44 chapter, derived projections are relatively insensitive to details of climate scenarios and can be well  
45 estimated based on transient simulations (CC-Box 11.1; see also SR15).

46  
47 An important question is the identification of the GWL at which a given change in a climate extreme can  
48 begin to emerge from climate noise. Figure 11.8 displays analyses of the GWLs at which emergence in hot  
49 extremes (20-year return values of TXx, TXx\_20yr) and heavy precipitation (20-year return values of  
50 Rx1day, Rx1day\_20yr) is identified in AR6 regions for the whole CMIP5 and CMIP6 ensembles). Overall,  
51 signals for extremes emerge very early for TXx\_20yr, already below 0.2°C in many regions (Fig. 11.8a,b),  
52 and at around 0.5°C in most regions. This is consistent with conclusions from the SR15 Ch3 for less-rare  
53 temperature extremes (TXx on the yearly time scale), which shows that a difference as small as 0.5°C of  
54 global warming, e.g. between +1.5°C and +2°C of global warming, leads to detectable differences in  
55 temperature extremes in TXx in most WGI AR6 regions in CMIP5 projections (e.g., Wartenburger et al.,

2017; Seneviratne et al., 2018b). The GWL emergence for Rx1day\_20yr is also largely consistent with analyses for less-extreme heavy precipitation events (Rx5day on the yearly time scale) in the SR15 (see Chapter 3).

To some extent, analyses as functions of GWLs replace the time axis with a global surface temperature axis. Nonetheless, information on the timing of given changes in extremes is obviously also relevant. Regarding this information, that is, the time frame at which given global warming levels are reached, the readers are referred to Chapter 4 (Section 4.6; see also CC-Box 11.1). Figure 11.5 provides a synthesis of attributed and projected changes in extremes as function of GWLs (see also Figs. 11.3, 11.6, and 11.7 for regional analyses).

[START FIGURE 11.8 HERE]

**Figure 11.8:** Global and regional-scale emergence of changes in temperature (a) and precipitation (b) extremes for the globe (glob.), global oceans (oc.), global lands (land), and the AR6 regions. Colours indicate the multi-model mean global warming level at which the difference in 20-year means of the annual maximum daily maximum temperature (TXx) and the annual maximum daily precipitation (Rx1day) become significantly different from their respective mean values during the 1851–1900 base period. Results are based on simulations from the CMIP5 and CMIP6 multi-model ensembles. See Atlas.1.3.2 for the definition of regions. Adapted from Seneviratne and Hauser, 2020) under the terms of the Creative Commons Attribution license.

[END FIGURE 11.8 HERE]

[START CROSS-CHAPTER BOX 11.1 HERE]

#### **Cross-Chapter Box 11.1: Translating between regional information at global warming levels vs scenarios for end users**

**Contributors:** Erich Fischer (Switzerland), Mathias Hauser (Switzerland), Sonia I. Seneviratne (Switzerland), Richard Betts (UK), José M. Gutiérrez (Spain), Richard G. Jones (UK), June-Yi Lee (Republic of Korea), Malte Meinshausen (Australia/Germany), Friederike Otto (UK/Germany), Izidine Pinto (Mozambique), Roshanka Ranasinghe (The Netherlands/Sri Lanka/Australia), Joeri Rogelj (Germany/Belgium), Bjørn Samset (Norway), Claudia Tebaldi (USA), Laurent Terray (France)

#### **Background**

Traditionally, projections of climate variables are summarized and communicated as function of time and scenario. Recently, quantifying global and regional climate at specific global warming levels (GWLs) has become widespread, motivated by the inclusion of explicit GWLs in the long-term temperature goal of the Paris Agreement (Section 1.6.2). GWLs, expressed as changes in global surface temperature relative to the 1850-1900 period (see CCBox 2.3), are used in the SR15 and in the assessment of Reasons for Concerns in the WGII reports (see also CCBox 12.1). CCB 11.1, Figure 1 illustrates how the assessment of the climate response at GWLs relates to the uncertainty in scenarios regarding the timing of the respective GWLs, as well as to the uncertainty in the associated regional climate responses, including extremes and other climatic impact-drivers (CIDs). For many (but not all) climate variables and CIDs the response pattern for a given GWL is consistent across different scenarios (Chapters 1, 4, 9, 11 and Atlas). GWLs are defined as long-term means (e.g. 20-year averages) compared to the pre-industrial period, are commonly used in the literature and were also underlying main assessments of SR15 (Chapter 3).

[START CROSS-CHAPTER BOX 11.1, FIGURE 1 HERE]

**Cross-Chapter Box 11.1, Figure 1:** Schematic representation of relationship between emission scenarios, global

warming levels (GWLs), regional climate responses, and impacts. The illustration shows the implied uncertainty problem associated with differentiating between 1.5, 2°C, and other GWLs. Focusing on GWL raises questions associated with emissions pathways to get to these temperatures (scenarios), as well as questions associated with regional climate responses and the associated impacts at the corresponding GWL (the impacts question). Adapted from (James et al., 2017) and (Rogelj, 2013) under the terms of the Creative Commons Attribution license.

**[END CROSS-CHAPTER BOX 11.1, FIGURE 1 HERE]**

Numerous studies have compared the regional response to anthropogenic forcing at GWLs in annual and seasonal mean values and extremes of different climate and impact variables across different multi-model ensembles and/or different scenarios (e.g. Frieler et al., 2012; Schewe et al., 2014; Schleussner et al., 2016; Seneviratne et al., 2016; Wartenburger et al., 2017; Dosio and Fischer, 2018; Tebaldi et al., 2020; (Herger et al., 2015; Betts et al., 2018; Samset et al., 2019), see Sections 4.6.1, 8.5.3, 9.3.1, 9.5, 9.6.3, 10.4.3 and 11.2.4 for further details). The regional response patterns at given GWLs have been found to be consistent across different scenarios for many climate variables (CC-Box 11.1 Fig.2) (Pendergrass et al., 2015; Seneviratne et al., 2016; Wartenburger et al., 2017; Seneviratne and Hauser, 2020). The consistency tends to be higher for temperature-related variables than for variables in the hydrological cycle or variables characterizing atmospheric dynamics, and for intermediate to high emission scenarios than for low-emission scenarios (e.g. for mean precipitation in the RCP2.6 scenario: Pendergrass et al., 2015; Wartenburger et al., 2017). Nonetheless, CCB 11.1 Figure 2 illustrates that even for mean precipitation, which is known to be forcing-dependent (Section 4.6.1 and Section 8.5.3), scenario differences in the response pattern at a given GWL are smaller than model uncertainty and internal variability in many regions (Herger et al., 2015). The response pattern is further found to be broadly consistent between models that reach a GWL relatively early and those that reach it later under a given SSP (see CC Box 11.1 Fig.2 g, h)

**[START CROSS-CHAPTER 11.1, FIGURE 2 HERE]**

**Cross-Chapter Box 11.1, Figure 2:** (a-c) CMIP6 multi-model mean precipitation change at 2°C GWL (20-yr mean) in three different SSP scenarios relative to 1850-1900. All models reaching the corresponding GWL in the corresponding scenario are averaged. The number of models averaged across is shown at the top right of the panel. The maps for the other two SSP scenarios SSP1-1.9 (five models only) and SSP3-7.0 (not shown) are consistent. (d-f) Same as (a-c) but for annual mean temperature. (g) Annual mean temperature change at 2°C in CMIP6 models with high warming rate reaching the GWL in the corresponding scenario before the earliest year of the assessed very likely range (section 4.3.4) (h) Climate response at 2°C GWL across all SSP1-1.9, SSP2-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5 in all other models not shown in (g). The good agreement of (g) and (h) demonstrate that the mean temperature response at 2°C is not sensitive to the rate of warming and thereby the GSAT warming of the respective models in 2081-2100. Uncertainty is represented using the advanced approach: No overlay indicates regions with robust signal, where  $\geq 66\%$  of models show change greater than variability threshold and  $\geq 80\%$  of all models agree on sign of change; diagonal lines indicate regions with no change or no robust signal, where  $< 66\%$  of models show a change greater than the variability threshold; crossed lines indicate regions with conflicting signal, where  $\geq 66\%$  of models show change greater than variability threshold and  $< 80\%$  of all models agree on sign of change. For more information on the advanced approach, please refer to the Cross-Chapter Box Atlas.1.



**[END CROSS-CHAPTER BOX 11.1, FIGURE 2 HERE]**

In contrast to linear pattern scaling (Mitchell, 2003; Collins et al., 2013a), the use of GWLs as a dimension of integration does not require linearity in the response of a climate variable. It is thus even useful for metrics which do not show a linear response, such as the frequency of heat extremes over land and oceans (Fischer and Knutti, 2015; Perkins-Kirkpatrick and Gibson, 2017; Frölicher et al., 2018; Kharin et al., 2018) if the relationship of the variable of interest to the GWL is scenario independent. The latter means that the response is independent of the pathway and relative contribution of various radiative forcings. For some more complex indices like warm-spell duration or for regions with strong aerosol changes, discrepancies can be larger (Wang et al., 2017d; King et al., 2018; Tebaldi et al., 2020) (see also subsection below on GWLs vs scenarios for further caveats).

The limited scenario dependence of the GWL-based response for many variables implies that the regional response to emissions scenarios can be split in almost independent contributions of 1) the transient global warming response to scenarios (see Chapter 4), and 2) the regional response as function of a given GWL, which has also been referred to as “regional climate sensitivity” (Seneviratne and Hauser, 2020). This property has also been used to develop regionally-resolved emulators for global climate models, using global surface temperature as input (Beusch et al., 2020; Tebaldi et al., 2020). Analyses of the CMIP6 and CMIP5 multi-model ensembles shows that the GWL-based responses are very similar for temperature and precipitation extremes across the ensembles (Li et al., 2020a; Seneviratne and Hauser, 2020; Wehner, 2020). This is despite their difference in global warming response (Chapter 4), confirming a substantial decoupling between the two responses (global warming vs GWL-based regional response) for these variables. Thus, the GWL approach isolates the uncertainty in the regional climate response from the global warming uncertainty induced by scenario, global mean model response and internal variability (CCB Figure 1).

**Mapping between GWL- and scenario-based responses in model analyses**

To map scenario-based climate projections into changes at specific GWLs, first, all individual ESM simulations that reach a certain GWL are identified. Second, the climate response patterns at the respective GWL are calculated using an approach termed here “GWL-sampling approach” – sometimes also referred to as epoch analysis, time shift, or time sampling approach –, taking into account all models and scenarios (CCB Figure 3). Note that the range of years when a given GWL is reached in the CMIP6 ensemble is different from the AR6 assessed range of projected global surface temperature (Table 4.5; Section 4.3.4). The latter further takes into account different lines of evidence, including the assessed observed warming between pre-industrial and present day, information from observational constraints on CMIP6, and emulators using the assessed transient climate response (TCR) and equilibrium climate sensitivity (ECS) ranges (Section 4.3.4). Hence the Chapter 4 assessed range (Table 4.5) is the reference to determine when a given GWL is *likely* reached under given scenarios, while the mapping between scenarios/time frames and GWLs is used to assess the respective regional responses happening at these time frames (which also allows to account for the global surface temperature assessment rather than using scenarios analyses directly from CMIP6 output).

In the model-based assessment of Chapters 4, 8, 10, 11, 12 and the Atlas, the estimation of changes at GWLs are generally defined as the 20-year time period in which the mean global surface air temperature (GSAT; CCBox 2.3) first exceeds a certain anomaly relative to 1850-1900 (for simulations that start after 1850, relative to all years up to 1900 CCB Figure 3). The years when each individual model reaches a given GWL for CMIP6 and CMIP5 can be found in Hauser et al. (2021). The changes at given GWLs are identified for each ensemble member (for all scenarios) individually. Thereby, a given GWL is potentially reached a few years earlier or later in different realizations of the same model due to internal variability, but the temperature averaged across the 20-year period analysed in any simulation is consistent with the GWL. Instead of blending the information from the different scenarios, the Interactive Atlas can be used to compare the GWL spatial patterns and timings across the different scenarios (see Section Atlas 1.3.1).

**[START CROSS-CHAPTER BOX 11.1, FIGURE 3 HERE]**

**Cross-Chapter Box 11.1, Figure 3:** Illustration of the AR6 GWL sampling approach to derive the timing and the response at a given GWL for the case of CMIP6 data. For the mapping of scenarios/time slices into GWLs for CMIP6, please refer to Table 4.2. Respective numbers for the CMIP6 multi-model experiment are provided in the Chapter 11 Supplementary Material (11.SM.1). Note that the time frames used to derive the GWL time slices can also include different number of years (e.g. 30 years for some analyses).

**[END CROSS-CHAPTER BOX 11.1, FIGURE 3 HERE]****Mapping between GWL- and scenario-based responses for literature**

A large fraction of the literature considers scenario-based analyses for given time slices. When GWL-based information is required instead, an approximated mapping of the multi-model mean can be derived based on the known GWL in the given experiments for a particular time period. As a rough approximation, CMIP6 multi-model mean projections for the near-term (2021-2040) correspond to changes at about 1.5°C, and projections for the high-end scenario (SSP5-8.5) for the long-term (2081-2100) correspond to about 4-5°C of global warming (see Table 4.2 for changes in the CMIP6 ensemble and the Chapter 11 Supplementary Material (11.SM.1) and Hauser (2021) for details on other time periods and CMIP5). These approximated changes are for instance used for some of the GWL-based assessments provided in the Chapter 11 regional tables (Section 11.9; Table 11.3) when literature based on scenario projections is used to assess estimated changes at given GWLs.

**GWLs vs scenarios**

The use of scenarios remains a key element to inform mitigation decisions (Chapter 1, CCB1.4), to assess which emission pathways are consistent with a certain GWL (CCB1.4 Figure 1), to estimate when certain GWLs are reached (Section 4.3.4), and to assess for which variables it is meaningful to use GWLs as a dimension of integration. The use of scenarios is also essential for variables whose climate response strongly depends on the contribution of radiative forcing (e.g. aerosols) and land use and land management changes, and are time and warming rate dependent (e.g. sea level rise), or differ between transient and quasi-equilibrium states. Furthermore, the use of concentration or emission-driven scenario simulations is required if regional climate assessments need to account for the uncertainty in GSAT changes or climate-carbon feedbacks.

Forcing dependence of the GWL response is found for global mean precipitation (Section 8.4.3), but less for regional patterns of mean precipitation changes (CC-Box 11.1, Fig. 2). Limited dependence is found for extremes, as highlighted above. In the cryosphere, elements that are quick to respond to warming like sea ice area, permafrost, and snow show little scenario dependence (Chapter 9.3.1.1, 9.5.2.3, 9.5.3.3), whereas slow-responding variables such as ice volumes of glaciers and ice sheets respond with a substantial delay and due to their inertia, the response depends on when a certain GWL is reached. This also applies to some extent for sea level rise where, for example, the contributions of melting glaciers and ice sheets depend on the pathway followed to reach a given GWL (Chapter 9.6.3.4).

In addition to the lagged effect, the climate response at a given GWL may differ before and after a period of overshoot, for example in the Atlantic Meridional Overturning Circulation (e.g. Palter et al. 2018). Finally, as assessed in IPCC SR15, there is a difference in the response even for temperature-related variables if a GWL is reached in a rapidly warming transient state or in an equilibrium state when the land-sea warming contrast is less pronounced (e.g. King et al. 2020). However, in this report GWLs are used in the context of projections for the 21<sup>st</sup> century when the climate response is mostly not in equilibrium and where projections for many variables are less dependent on the pathway than for projections beyond 2100 (Section 9.6.3.4).

## Key conclusions on assessments based on GWLs

GWL-based projections can inform society and policymakers on how climate would change under GWLs consistent with the aims of the Paris Agreement (stabilization at 1.5°C/well below 2°C), as well as on the consequences of missing these aims and reaching GWLs of 3°C or 4°C by the end of the century. The AR6 assessment shows that every bit of global warming matters and that changes in global warming of 0.5°C lead to statistically significant changes in mean climate and climate extremes on global scale and for large regions (Sections 4.6.2, 11.2.4, 11.3, 11.4, 11.6, 11.9; Figs 11.8, 11.9, Atlas, Interactive Atlas), as also assessed in the IPCC SR15.

[END CROSS-CHAPTER BOX 11.1 HERE]

### 11.3 Temperature extremes

This section assesses changes in temperature extremes at global, continental and regional scales. The main focus is on the changes in the magnitude and frequency of moderate extreme temperatures (those that occur several times a year) to very extreme temperatures (those that occur once-in-10-years or longer) of time scales from a day to a season, though there is a strong emphasis on the daily scale where literature is most concentrated.

#### 11.3.1 Mechanisms and drivers

The SREX (IPCC, 2012) and AR5 (IPCC, 2014) concluded that greenhouse gas forcing is the dominant factor for the increases in the intensity, frequency, and duration of warm extremes and the decrease in those of cold extremes. This general global-scale warming is modulated by large-scale atmospheric circulation patterns, as well as by feedbacks such as soil moisture-evapotranspiration-temperature and snow/ice-albedo-temperature feedbacks, and local forcings such as land use change or changes in aerosol concentrations at the regional and local scales (Box 11.1, Sections 11.1.5, 11.1.6). Therefore, changes in temperature extremes at regional and local scales can have heterogeneous spatial distributions. Changes in the magnitudes (or intensities) of extreme temperatures are often larger than changes in global surface temperature, because of larger warming on land than on the ocean surface (2.3.1.1) and feedbacks, though they are of similar magnitude to changes in the local mean temperature (Fig 11.2).

Extreme temperature events are associated with large-scale meteorological patterns (Grotjahn et al., 2016). Quasi-stationary anticyclonic circulation anomalies or atmospheric blocking events are linked to temperature extremes in many regions, such as in Australia (Parker et al., 2014; Perkins-Kirkpatrick et al., 2016), Europe (Brunner et al., 2017, 2018; Schaller et al., 2018), Eurasia (Yao et al., 2017), Asia (Chen et al., 2016; Ratnam et al., 2016; Rohini et al., 2016), and North America (Yu et al., 2018, 2019b; Zhang and Luo, 2019). Mid-latitude planetary wave modulations affect short-duration temperature extremes such as heat waves (Perkins, 2015; Kornhuber et al., 2020). The large-scale modes of variability (Annex VI) affect the strength, frequency, and persistence of these meteorological patterns and, hence, temperature extremes. For example, cold and warm extremes in the mid-latitudes are associated with atmospheric circulation patterns such as the Pacific-North American (PNA) pattern, as well as atmosphere-ocean coupled modes such as Pacific Decadal Variability (PDV), the North Atlantic Oscillation (NAO), and Atlantic Multidecadal Variability (AMV) (Kamae et al., 2014; Johnson et al., 2018; Ruprich-Robert et al., 2018; Yu et al., 2018, 2019a; Müller et al., 2020; Section 11.1.5). Changes in the modes of variability in response to warming would therefore affect temperature extremes (Clark and Brown, 2013; Horton et al., 2015). The level of confidence in those changes, both in the observations and in future projections, varies, affecting the level of confidence in changes in temperature extremes in different regions. As highlighted in Chapters 2-4 of this Report, it is *likely* that there have been observational changes in the extratropical jets and mid-latitude jet meandering (Section 2.3.1.4.3; Cross-Chapter Box 10.1). There is *low confidence* in possible effects of Arctic warming

1 on mid-latitude temperature extremes (Cross-Chapter Box 10.1). A large portion of the multi-decadal  
2 changes in extreme temperature remains after the removal of the effect of these modes of variability and can  
3 be attributed to human influence (Kamae et al., 2017b; Wan et al., 2019). Thus, global warming dominates  
4 changes in temperature extremes at the regional scale and it is *very unlikely* that dynamic responses to  
5 greenhouse-gas induced warming would alter the direction of these changes.  
6

7 Land-atmosphere feedbacks strongly modulate regional- and local-scale changes in temperature extremes  
8 (*high confidence*; Section 11.1.6; Seneviratne et al., 2013; Lemordant et al., 2016; Donat et al., 2017;  
9 Sillmann et al., 2017b; Hirsch et al., 2019). This effect is particularly notable in mid-latitude regions where  
10 the drying of soil moisture amplifies high temperatures, in particular through increases in sensible heat flux  
11 (Whan et al., 2015; Douville et al., 2016; Vogel et al., 2017). Land-atmosphere feedbacks amplifying  
12 temperature extremes also include boundary-layer feedbacks and effects on atmospheric circulation  
13 (Miralles et al., 2014a; Schumacher et al., 2019). Soil moisture-temperature feedbacks affect past and  
14 present-day heat waves in observations and model simulations, both locally (Miralles et al. 2014; Hauser et  
15 al. 2016; Meehl et al. 2016; Wehrli et al., 2019; Cowan et al., 2016) and beyond the regions of feedback  
16 occurrence through changes in regional circulation patterns (Koster et al., 2016; Sato and Nakamura, 2019;  
17 Stéfanon et al., 2014). The uncertainty due to the representation of land-atmosphere feedbacks in ESMs is a  
18 cause of discrepancy between observations and simulations (Clark et al., 2006; Mueller and Seneviratne,  
19 2014; Meehl et al., 2016). The decrease of plant transpiration or the increase of stomata resistance under  
20 enhanced CO<sub>2</sub> concentrations is a direct CO<sub>2</sub> forcing of land temperatures (warming due to reduced  
21 evaporative cooling), which contributes to higher warming on land (Lemordant et al., 2016; Vicente-Serrano  
22 et al., 2020c). The snow/ice-albedo feedback plays an important role in amplifying temperature variability in  
23 the high latitudes (Diro et al. 2018) and can be the largest contributor to the rapid warming of cold extremes  
24 in the mid- and high latitudes of the Northern Hemisphere (Gross et al., 2020).  
25

26 Regional external forcings, including land-use changes and emissions of anthropogenic aerosols, play an  
27 important role in the changes of temperature extremes in some regions (*high confidence*, Section 11.1.6).  
28 Deforestation may have contributed to about one third of the warming of hot extremes in some mid-latitude  
29 regions since the pre-industrial time (Lejeune et al., 2018). Aspects of agricultural practice, including no-till  
30 farming, irrigation, and overall cropland intensification, may cool hot temperature extremes (Davin et al.,  
31 2014; Mueller et al., 2016b). For instance, cropland intensification has been suggested to be responsible for a  
32 cooling of the highest temperature percentiles in the US Midwest (Mueller et al., 2016b). Irrigation has been  
33 shown to be responsible for a cooling of hot temperature extremes of up to 1-2°C in many mid-latitude  
34 regions in the present climate (Thiery et al., 2017; Thiery et al., 2020), a process not represented in most of  
35 state-of-the-art ESMs (CMIP5, CMIP6). Double cropping may have led to increased hot extremes in the  
36 inter-cropping season in part of China (Jeong et al., 2014). Rapid increases in summertime warming in  
37 western Europe and northeast Asia since the 1980s are linked to a reduction in anthropogenic aerosol  
38 precursor emissions over Europe (Dong et al., 2016, 2017; Nabat et al., 2014), in addition to the effect of  
39 increased greenhouse gas forcing (see also Chapter 10, Section 10.1.3.1). This effect of aerosols on  
40 temperature-related extremes is also noted for declines in short-lived anthropogenic aerosol emissions over  
41 North America (Mascioli et al., 2016). On the local scale, the urban heat island (UHI) effect results in higher  
42 temperatures in urban areas than in their surrounding regions and contributes to warming in regions of rapid  
43 urbanization, in particular for night-time temperature extremes (Box 10.3; Phelan et al., 2015; Chapman et  
44 al., 2017; Sun et al., 2019). But these local and regional forcings are generally not (well-) represented in the  
45 CMIP5 and CMIP6 simulations (see also Section 11.3.3), contributing to uncertainty in model simulated  
46 changes.  
47

48 In summary, greenhouse gas forcing is the dominant driver leading to the warming of temperature extremes.  
49 At regional scales, changes in temperature extremes are modulated by changes in large-scale patterns and  
50 modes of variability, feedbacks including soil moisture-evapotranspiration-temperature or snow/ice-albedo-  
51 temperature feedbacks, and local and regional forcings such as land use and land cover changes, or aerosol  
52 concentrations, and decadal and multidecadal natural variability. This leads to heterogeneity in regional  
53 changes and their associated uncertainties (*high confidence*). Urbanization has exacerbated the effects of  
54 global warming in cities, in particular for night-time temperature extremes (*high confidence*).  
55

### 11.3.2 Observed trends

The SREX (IPCC, 2012) reported a *very likely* decrease in the number of cold days and nights and increase in the number of warm days and nights at the global scale. Confidence in trends was assessed as regionally variable (*low to medium confidence*) due to either a lack of observations or varying signals in sub-regions.

Since SREX (IPCC, 2012) and AR5 (IPCC, 2014), many regional-scale studies have examined trends in temperature extremes using different metrics that are based on daily temperatures, such as the CCI/WCRP/JCOMM Expert Team on Climate Change Detection and Indices (ETCCDI) indices (Dunn et al., 2020). The additional observational records, along with a stronger warming signal, show very clearly that changes observed at the time of AR5 (IPCC, 2014) continued, providing strengthened evidence of an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes. While the magnitude of the observed trends in temperature-related extremes varies depending on the region, spatial and temporal scales, and metric assessed, evidence of a warming effect is overwhelming, robust, and consistent. In particular, an increase in the intensity and frequency of hot extremes is almost always associated with an increase in the hottest temperatures and in the number of heatwave days. It is also the case for changes in cold extremes. For this reason, and to simplify the presentation, the phrase “increase in the intensity and frequency of hot extremes” is used to represent, collectively, an increase in the magnitude of extreme day and/or night temperatures, in the number of warm days and/or nights, and in the number of heat wave days. Changes in cold extremes are assessed similarly.

On the global scale, evidence of an increase in the number of warm days and nights and a decrease in the number of cold days and nights, and an increase in the coldest and hottest extreme temperatures is very robust and consistent among all variables. Figure 11.2 displays timeseries of globally-averaged annual maximum daily maximum (TXx) and annual minimum daily minimum temperature (TNn) on land. Warming of land mean TXx is similar to the mean land warming, which is about 45% higher than global warming (Section 2.3.1). Warming of land mean TNn is even higher, with about 3°C of warming since 1960 (Figure 11.2). Figure 11.9 shows maps of linear trends over 1960-2018 in the annual maximum daily maximum (TXx), the annual minimum daily minimum temperature (TNn), and frequency of warm days (TX90p). The maps for TXx and TNn show trends consistent with overall warming in most regions, with a particularly high warming of TXx in Europe and north-western South America, and a particularly high warming of TNn in the Arctic. Consistent with the observed warming in global surface temperature (2.3.1.2) and the observed trends in TXx and TNn, the frequency of TX90p has increased while that of cold nights (TN10p) has decreased since the 1950s: Nearly all land regions showed statistically significant decreases in TN10p (Alexander, 2016; Dunn et al., 2020), though trends in TX90p are variable with some decreases in southern South America, mainly during austral summer (Rusticucci et al., 2017). A decrease in the number of cold spell days is also observed over nearly all land surface areas (Easterling et al., 2016) and in the northern mid-latitudes in particular (van Oldenborgh et al., 2019). These observed changes are also consistent when a new global land surface daily air temperature dataset is analyzed (Zhang et al., 2019c). Consistent warming trends in temperature extremes globally, and in most land areas, over the past century are also found in a range of observation-based data sets (Fischer and Knutti, 2014; Donat et al., 2016a; Dunn et al., 2020), with the extremes related to daily minimum temperatures changing faster than those related to daily maximum temperatures (Dunn et al., 2020) (Fig. 11.2). Seasonal variations in trends in temperature-related extremes have been demonstrated. A warming in warm-season temperature extremes is detected, even during the “slower surface global warming” period from the late 1990s to early 2010s (Cross-Chapter Box 3.1) (Kamae et al., 2014; Seneviratne et al., 2014; Imada et al., 2017). Many studies of past changes in temperature extremes for particular regions or countries show trends consistent with this global picture, as summarized below and in Tables 11.4, 11.7, 11.10, 11.13, 11.16 and 11.19 in Section 11.9.

[START FIGURE 11.9 HERE]

**Figure 11.9:** Linear trends over 1960-2018 in the annual maximum daily maximum temperature (TXx, a), the annual minimum daily minimum temperature (TNn, b), and the annual number of days when daily maximum

1 temperature exceeds its 90<sup>th</sup> percentile from a base period of 1961-1990 (TX90p, c), based on the  
2 HadEX3 data set (Dunn et al., 2020). Linear trends are calculated only for grid points with at least 66% of  
3 the annual values over the period and which extend to at least 2009. Areas without sufficient data are  
4 shown in grey. No overlay indicates regions where the trends are significant at  $p = 0.1$  level. Crosses  
5 indicate regions where trends are not significant. For details on the methods see Supplementary Material  
6 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table  
7 11.SM.9).

8  
9 **[END FIGURE 11.9 HERE]**

10  
11  
12 In Africa (Table 11.4), while it is difficult to assess changes in temperature extremes in parts of the continent  
13 because of a lack of data, evidence of an increase in the intensity and frequency of hot extremes and decrease  
14 in the intensity and frequency of cold extremes is clear and robust in regions where data are available. These  
15 include an increase in the frequency of warm days and nights and a decrease in the frequency of cold days  
16 and nights with *high confidence* (Donat et al., 2013b, 2014b; Kruger and Sekele, 2013; Chaney et al., 2014;  
17 Filahi et al., 2016; Moron et al., 2016; Ringard et al., 2016; Barry et al., 2018; Gebrechorkos et al., 2018) and  
18 an increase in heat waves (Russo et al., 2016; Ceccherini et al., 2017). The increase in TNn is more notable  
19 than in TXx (Figure 11.9). Cold spells occasionally strike subtropical areas, but are *likely* to have decreased  
20 in frequency (Barry et al., 2018). The frequency of cold events has *likely* decreased in South Africa (Song et  
21 al., 2014; Kruger and Nxumalo, 2017), North Africa (Driouech et al., 2021; Filahi et al., 2016), and the  
22 Sahara (Donat et al., 2016a). Over the whole continent, there is *medium confidence* in an increase in the  
23 intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes; it is  
24 *likely* that similar changes have also occurred in areas with poor data coverage, as warming is widespread  
25 and as projected future changes are similar over all regions (11.3.5).

26  
27 In Asia (Table 11.7), there is very *robust evidence* for a *very likely* increase in the intensity and frequency of  
28 hot extremes and decrease in the intensity and frequency of cold extremes in recent decades. This is clear in  
29 global studies (e.g. Alexander, 2016; Dunn et al., 2020), as well as in numerous regional studies (Table  
30 11.7). The area fraction with extreme warmth in Asia increased during 1951–2016 (Imada et al., 2018). The  
31 frequency of warm extremes increased and the frequency of cold extremes decreased in East Asia (Zhou et  
32 al., 2016a; Chen and Zhai, 2017; Yin et al., 2017; Lee et al., 2018c; Qian et al., 2019) and west Asia (Acar  
33 Deniz and Gönençgil, 2015; Erlat and Türkeş, 2016; Imada et al., 2017; Rahimi et al., 2018; Rahimi and  
34 Hejabi, 2018) with *high confidence*. The duration of heat extremes has also lengthened in some regions, for  
35 example, in southern China (Luo and Lau, 2016), but there is *medium confidence* of heat extremes increasing  
36 in frequency in South Asia (AlSarmi and Washington, 2014; Sheikh et al., 2015; Mazdiyasi et al., 2017;  
37 Zahid et al., 2017; Nasim et al., 2018; Khan et al., 2019; Roy, 2019). Warming trends in daily temperature  
38 extremes indices have also been observed in central Asia (Hu et al., 2016; Feng et al., 2018), the Hindu Kush  
39 Himalaya (Sun et al., 2017), and Southeast Asia (Supari et al., 2017; Cheong et al., 2018). The intensity and  
40 frequency of cold spells in all Asian regions have been decreasing since the beginning of the 20th century  
41 (*high confidence*) (Sheikh et al., 2015; Donat et al., 2016a; Dong et al., 2018; van Oldenborgh et al., 2019).

42  
43 In Australasia (Table 11.10), there is very *robust evidence* for *very likely* increases in the number of warm  
44 days and warm nights and decrease in the number of cold days and cold nights since 1950 (Lewis and King,  
45 2015; Jakob and Walland, 2016; Alexander and Arblaster, 2017). The increase in extreme minimum  
46 temperatures occurs in all seasons over most of Australia and typically exceeds the increase in extreme  
47 maximum temperatures (Wang et al., 2013b; Jakob and Walland, 2016). However, some parts of southern  
48 Australia have shown stable or increased numbers of frost days since the 1980s (Dittus et al., 2014) (see also  
49 Section 11.3.4). Similar positive trends in extreme minimum and maximum temperatures have been  
50 observed in New Zealand, in particular in the autumn-winter seasons, although they generally show higher  
51 spatial variability (Caloiero, 2017). In the tropical Western Pacific region, spatially coherent warming trends  
52 in maximum and minimum temperature extremes have been reported for the period of 1951–2011 (Whan et  
53 al., 2014; McGree et al., 2019).

54  
55 In Central and South America (Table 11.13), there is *high confidence* that observed hot extremes (TN90p,  
56 TX90p) have increased and cold extremes (TN10p, TX10p) have decreased over recent decades, though

1 trends vary among different extremes types, datasets, and regions (Dereczynski et al., 2020; Dittus et al.,  
2 2016; Dunn et al., 2020; Meseguer-Ruiz et al., 2018; Olmo et al., 2020; Rusticucci et al., 2017; Salvador and  
3 de Brito, 2018; Skansi et al., 2013). An increase in the intensity and frequency of heatwave events was also  
4 observed between 1961 and 2014, in an area covering most of South America (Ceccherini et al., 2016;  
5 Geirinhas et al., 2018). However, there is *medium confidence* that warm extremes (TXx and TX90p) have  
6 decreased in the last decades over the central region of SES during austral summer (Tencer, B.; Rusticucci,  
7 2012; Skansi et al., 2013; Rusticucci et al., 2017; Wu and Polvani, 2017). There is *medium confidence* that  
8 TNn extremes are increasing faster than TXx extremes, with the largest warming rates observed over  
9 Northeast Brazil (NEB) and North South America (NSA) for cold nights (Skansi et al., 2013).

10  
11 In Europe (Table 11.16), there is very robust evidence for a *very likely* increase in maximum temperatures  
12 and the frequency of heat waves. The increase in the magnitude and frequency of high maximum  
13 temperatures has been observed consistently across regions including in central (Twardosz and Kossowska-  
14 Cezak, 2013; Christidis et al., 2015; Lorenz et al., 2019) and southern Europe (Croitoru and Piticar, 2013; El  
15 Kenawy et al., 2013; Christidis et al., 2015; Nastos and Kapsomenakis, 2015; Fioravanti et al., 2016; Ruml et  
16 al., 2017). In northern Europe, a strong increase in extreme winter warming events has been observed  
17 (Matthes et al., 2015; Vikhamar-Schuler et al., 2016). Temperature observations for wintertime cold spells  
18 show a long-term decreasing frequency in Europe (Brunner et al., 2018; van Oldenborgh et al., 2019), and  
19 typical cold spells such as that observed during the 2009/2010 winter had an occurrence probability that is  
20 twice smaller currently than if climate change had not occurred (Christiansen et al., 2018).

21  
22 In North America (Table 11.19), there is very robust evidence for a *very likely* increase in the intensity and  
23 frequency of hot extremes and decrease in the intensity and frequency of cold extremes for the whole  
24 continent, though there are substantial spatial and seasonal variations in the trends. Minimum temperatures  
25 display warming consistently across the continent, while there are more contrasting trends in the annual  
26 maximum daily temperatures in parts of the USA (Figure 11.9) (Lee et al., 2014; van Oldenborgh et al.,  
27 2019; Dunn et al., 2020). In Canada, there is a clear increase in the intensity and frequency of hot extremes  
28 and decrease in the intensity and frequency of cold extremes (Vincent et al., 2018). In Mexico, a clear  
29 warming trend in TNn was found, particularly in the northern arid region (Montero-Martínez et al., 2018).  
30 The number of warm days has increased and the number of cold days has decreased (García-Cueto et al.,  
31 2019). Cold spells have undergone a reduction in magnitude and intensity in all regions of North America  
32 (Bennett and Walsh, 2015; Donat et al., 2016a; Grotjahn et al., 2016; Vose et al., 2017; García-Cueto et al.,  
33 2019; van Oldenborgh et al., 2019).

34  
35 Extreme heat events have increased around the Arctic since 1979, particularly over Arctic North America  
36 and Greenland (Matthes et al., 2015; Dobricic et al., 2020), which is consistent with summer melt (9.4.1).  
37 Observations north of 60°N show increases in wintertime warm days and nights over 1979-2015, while cold  
38 days and nights declined (Sui et al., 2017). Extreme heat days are particularly strong in winter, with  
39 observations showing the warmest mid-winter temperatures at the North Pole rising at twice the rate of mean  
40 temperature (Moore, 2016), as well as increases in Arctic winter warm days ( $T > -10^{\circ}\text{C}$ ) (Vikhamar-Schuler et  
41 al., 2016; Graham et al., 2017). Arctic annual minimum temperatures have increased at about three times the  
42 rate of global surface temperature since the 1960s (Figs. 11.2, 11.9), consistent with the observed mean cold  
43 season (October-May) warming of  $3.1^{\circ}\text{C}$  in the region (Atlas 11.2).

44  
45 Trends in some measures of heat waves are also observed at the global scale. Globally-averaged heat wave  
46 intensity, heat wave duration, and the number of heat wave days have significantly increased from 1950-  
47 2011 (Perkins, 2015). There are some regional differences in trends in characteristics of heat waves with  
48 significant increases reported in Europe (Russo et al., 2015; Forzieri et al., 2016; Sánchez-Benítez et al.,  
49 2020) and Australia (CSIRO and BOM, 2016; Alexander and Arblaster, 2017). In Africa, there is *medium*  
50 *confidence* that heat waves, regardless of the definition, have been becoming more frequent, longer-lasting,  
51 and hotter over more than three decades (Fontaine et al., 2013; Mouhamed et al., 2013; Ceccherini et al.,  
52 2016, 2017; Forzieri et al., 2016; Moron et al., 2016; Russo et al., 2016). The majority of heat wave  
53 characteristics examined in China between 1961-2014 show increases in heat wave days, consistent with  
54 warming (You et al., 2017; Xie et al., 2020). Increases in the frequency and duration of heat waves are also  
55 observed in Mongolia (Erdenebat and Sato, 2016) and India (Ratnam et al., 2016; Rohini et al., 2016). In the

1 UK, the lengths of short heat waves have increased since the 1970s, while the lengths of long heat waves  
2 (over 10 days) have decreased over some stations in the southeast of England (Sanderson et al., 2017b). In  
3 Central and South America, there are increases in the frequency of heat waves (Barros et al., 2015;  
4 Bitencourt et al., 2016; Ceccherini et al., 2016; Piticar, 2018), although decreases in Excess Heat Factor  
5 (EHF), which is a metric for heat wave intensity, are observed in South America in data derived from  
6 HadGHCND (Cavanaugh and Shen, 2015).

7  
8 In summary, it is *virtually certain* that there has been an increase in the number of warm days and nights and  
9 a decrease in the number of cold days and nights on the global scale since 1950. Both the coldest extremes  
10 and hottest extremes display increasing temperatures. It is *very likely* that these changes have also occurred  
11 at the regional scale in Europe, Australasia, Asia, and North America. It is *virtually certain* that there has  
12 been increases in the intensity and duration of heat waves and in the number of heat wave days at the global  
13 scale. These trends *likely* occur in Europe, Asia, and Australia. There is *medium confidence* in similar  
14 changes in temperature extremes in Africa and *high confidence* in South America; the lower confidence is  
15 due to reduced data availability and fewer studies. Annual minimum temperatures on land have increased  
16 about three times more than global surface temperature since the 1960s, with particularly strong warming in  
17 the Arctic (*high confidence*).

### 18 19 20 **11.3.3 Model evaluation**

21  
22 AR5 assessed that CMIP3 and CMIP5 models generally captured the observed spatial distributions of the  
23 mean state and that the inter-model range of simulated temperature extremes was similar to the spread  
24 estimated from different observational datasets; the models generally captured trends in the second half of  
25 the 20th century for indices of extreme temperature, although they tended to overestimate trends in hot  
26 extremes and underestimate trends in cold extremes (Flato et al., 2013). Post-AR5 studies on the CMIP5  
27 models' performance in simulating mean and changes in temperature extremes continue to support the AR5  
28 assessment (Fischer and Knutti, 2014; Sillmann et al., 2014; Ringard et al., 2016; Borodina et al., 2017b;  
29 Donat et al., 2017; Di Luca et al., 2020a). Over Africa, the observed warming in temperature extremes is  
30 captured by CMIP5 models, although it is underestimated in west and central Africa (Sherwood et al., 2014;  
31 Diedhiou et al., 2018). Over East Asia, the CMIP5 ensemble performs well in reproducing the observed  
32 trend in temperature extremes averaged over China (Dong et al., 2015). Over Australia, the multi-model  
33 mean performs better than individual models in capturing observed trends in gridded station based ETCCDI  
34 temperature indices (Alexander and Arblaster, 2017).

35  
36 Initial analyses of CMIP6 simulations (Chen et al., 2020a; Di Luca et al., 2020b; Kim et al., 2020; Li et al.,  
37 2020a; Thorarinsdottir et al., 2020; Wehner et al., 2020) indicate the CMIP6 models perform similarly to the  
38 CMIP5 models regarding biases in hot and cold extremes. In general, CMIP5 and CMIP6 historical  
39 simulations are similar in their performance in simulating the observed climatology of extreme temperatures  
40 (*high confidence*). The general warm bias in hot extremes and cold bias in cold extremes reported for CMIP5  
41 models (Kharin et al., 2013; Sillmann et al., 2013a) remain in CMIP6 models (Di Luca et al., 2020b).  
42 However, there is some evidence that CMIP6 models better represent some of the underlying processes  
43 leading to extreme temperatures, such as seasonal and diurnal variability and synoptic-scale variability (Di  
44 Luca et al., 2020b). Whether these improvements are sufficient to enhance our understanding of past changes  
45 or to reduce uncertainties in future projections remains unclear. The relative error estimates in the simulation  
46 of various indices of temperature extremes in the available CMIP6 models show that no single model  
47 performs the best on all indices and the multi-model ensemble seems to out-perform any individual model  
48 due to its reduction in systematic bias (Kim et al., 2020). Figure 11.10 show errors in the 1979-2014 average  
49 annual TXx and annual TNn simulated by available CMIP6 models in comparison with HadEX3 and ERA5  
50 (Li et al., 2020; Kim et al., 2020, Wehner et al., 2020). While the magnitude of the model error depends on  
51 the reference data set, the model evaluations drawn from different reference data sets are quite similar. In  
52 general, models reproduce the spatial patterns and magnitudes of both cold and hot temperature extremes  
53 quite well. There are also systematic biases. Hot extremes tend to be too cool in mountainous and high-  
54 latitude regions, but too warm in the eastern United States and South America. For cold extremes, CMIP6  
55 models are too cool, except in northeastern Eurasia and the southern mid-latitudes. Errors in seasonal mean



1 temperatures are uncorrelated with errors in extreme temperatures and are often of opposite sign (Wehner et  
2 al., 2020).

3  
4  
5 **[START FIGURE 11.10 HERE]**

6  
7 **Figure 11.10:** Multi-model mean bias in temperature extremes (°C) for the period 1979-2014, calculated as the  
8 difference between the CMIP6 multi-model mean and the average of observations from the values  
9 available in HadEX3 for (a) the annual hottest temperature (TXx) and (b) the annual coldest temperature  
10 (TNn). Areas without sufficient data are shown in grey. Adapted from Wehner et al. (2020) under the  
11 terms of the Creative Commons Attribution license. Further details on data sources and processing are  
12 available in the chapter data table (Table 11.SM.9).

13  
14 **[END FIGURE 11.10 HERE]**

15  
16  
17 Atmospheric model (AMIP) simulations are often used in event attribution studies to assess the influence of  
18 global warming on observed temperature-related extremes. These simulations typically capture the observed  
19 trends in temperature extremes, though some regional features, such as the lack of warming in daytime warm  
20 temperature extremes over South America and parts of North America, are not reproduced in the model  
21 simulations (Dittus et al., 2018), possibly due to internal variability, deficiencies in local surface processes,  
22 or forcings that are not represented in the SSTs. Additionally, the AMIP models assessed tend to produce  
23 overly persistent heat wave events. This bias in the duration of the events does not impact the reliability of  
24 the models' positive trends (Freychet et al., 2018).

25  
26 Several regional climate models (RCMs) have also been evaluated in terms of their performance in  
27 simulating the climatology of extremes in various regions of the Coordinated Regional Downscaling  
28 Experiment (CORDEX) (Giorgi et al., 2009), especially in East Asia (Ji and Kang, 2015; Yu et al., 2015;  
29 Park et al., 2016; Bucchignani et al., 2017; Gao et al., 2017a; Niu et al., 2018; Sun et al., 2018b; Wang et al.,  
30 2018a), Europe (Cardoso et al., 2019; Gaertner et al., 2018; Jacob et al., 2020; Kim et al., 2020; Lorenz et  
31 al., 2019; Smiatek et al., 2016; Vautard et al., 2013; Vautard et al., 2020b), and Africa (Kim et al., 2014b;  
32 Diallo et al., 2015; Dosio, 2017; Samouly et al., 2018; Mostafa et al., 2019). Compared to GCMs, RCM  
33 simulations show an added value in simulating temperature-related extremes, though this depends on  
34 topographical complexity and the parameters employed (see Section 10.3.3). The improvement with  
35 resolution is noted in East Asia (Park et al., 2016; Zhou et al., 2016b; Shi et al., 2017; Hui et al., 2018).  
36 However, in the European CORDEX ensemble, different aerosol climatologies with various degrees of  
37 complexity were used in projections (Bartók et al., 2017; Lorenz et al., 2019) and the land surface models  
38 used in the RCMs do not account for physiological CO<sub>2</sub> effects on photosynthesis leading to enhanced water-  
39 use efficiency and decreased evapotranspiration (Schwingshackl et al., 2019), which could lead to biases in  
40 the representation of temperature extremes in these projections (Boé et al., 2020). In addition, there are key  
41 cold biases in temperature extremes over areas with complex topography (Niu et al., 2018). Over North  
42 America, 12 RCMs were evaluated over the ARCTIC-CORDEX region (Diaconescu et al., 2018). Models  
43 were able to simulate well climate indices related to mean air temperature and hot extremes over most of the  
44 Canadian Arctic, with the exception of the Yukon region where models displayed the largest biases related to  
45 topographic effects. Two RCMs were evaluated against observed extremes indices over North America over  
46 the period 1989–2009, with a cool bias in minimum temperature extremes shown in both RCMs (Whan and  
47 Zwiers, 2016). The most significant biases are found in TXx and TNn, with fewer differences in the  
48 simulation of annual minimum daily maximum temperature (TXn) and annual maximum daily minimum  
49 temperature (TNx) in central and western North America. Over Central and South America, maximum  
50 temperatures from the Eta RCM are generally underestimated, although hot days, warm nights, and heat  
51 waves are increasing in the period 1961-1990, in agreement with observations (Chou et al., 2014b; Tencer et  
52 al., 2016; Bozkurt et al., 2019).

53  
54 Some land forcings are not well represented in climate models. As highlighted in the IPCC SRCCL Ch2,  
55 there is *high agreement* that temperate deforestation leads to summer warming and winter cooling (Bright et  
56 al., 2017; Zhao and Jackson, 2014; Gálós et al., 2011, 2013; Wickham et al., 2013; Ahlswede and Thomas,

2017; Anderson-Teixeira et al., 2012; Anderson et al., 2011; Chen et al., 2012; Strandberg and Kjellström, 2019), which has substantially contributed to the warming of hot extremes in the northern mid-latitudes over the course of the 20<sup>th</sup> century (Lejeune et al., 2018) and in recent years (Strandberg and Kjellström, 2019). However, observed forest effects on the seasonal and diurnal cycle of temperature are not well captured in several ESMs: while observations show a cooling effect of forest cover compared to non-forest vegetation during daytime (Li et al., 2015), in particular in arid, temperate, and tropical regions (Alkama and Cescatti, 2016), several ESMs simulate a warming of daytime temperatures for regions with forest vs non-forest cover (Lejeune et al., 2017). Also irrigation effects, which can lead to regional cooling of temperature extremes, are generally not integrated in current-generations of ESMs (Section 11.3.1).

In summary, there is *high confidence* that climate models can reproduce the mean state and overall warming of temperature extremes observed globally and in most regions, although the magnitude of the trends may differ. The ability of models to capture observed trends in temperature-related extremes depends on the metric evaluated, the way indices are calculated, and the time periods and spatial scales considered. Regional climate models add value in simulating temperature-related extremes over GCMs in some regions. Some land forcings on temperature extremes are not well captured (effects of deforestation) or generally not represented (irrigation) in ESMs.

#### 11.3.4 Detection and attribution, event attribution

SREX (IPCC, 2012) assessed that it is *likely* anthropogenic influences have led to the warming of extreme daily minimum and maximum temperatures at the global scale. AR5 concluded that human influence has *very likely* contributed to the observed changes in the intensity and frequency of daily temperature extremes on the global scale in the second half of the 20th century (IPCC, 2014). With regard to individual, or regionally- or locally-specific events, AR5 concluded that it is *likely* human influence has substantially increased the probability of occurrence of heat waves in some locations.

Studies since AR5 continue to attribute the observed increase in the frequency or intensity of hot extremes and the observed decrease in the frequency or intensity of cold extremes to human influence, dominated by anthropogenic greenhouse gas emissions, on global and continental scales, and for many AR6 regions. These include attribution of changes in the magnitude of annual TXx, TNx, TXn, and TNn, based on different observational data sets including, HadEX2 and HadEX3, CMIP5 and CMIP6 simulations, and different statistical methods (Kim et al., 2016; Wang et al., 2017c; Seong et al., 2020). As is the case for an increase in mean temperature (3.3.1), an increase in extreme temperature is mostly due to greenhouse gas forcing, offset by aerosol forcing. The aerosols' cooling effect is clearly detectable over Europe and Asia (Seong et al., 2020). As much as 75% of the moderate daily hot extremes (above 99.9<sup>th</sup> percentile) over land are due to anthropogenic warming (Fischer and Knutti, 2015). New results are found to be more robust due to the extended period that improves the signal-to-noise ratio. The effect of anthropogenic forcing is clearly detectable and attributable in the observed changes in these indicators of temperature extremes, even at country and sub-country scales, such as in Canada (Wan et al., 2019). Changes in the number of warm nights, warm days, cold nights, and cold days, and other indicators such as the Warm Spell Duration Index (WSDI), are also attributed to anthropogenic influence (Hu et al., 2020; Christidis and Stott, 2016).

Regional studies, including for Asia (Dong et al., 2018; Lu et al., 2018), Australia (Alexander and Arblaster, 2017), and Europe (Christidis and Stott, 2016), found similar results. A clear anthropogenic signal is also found in the trends in the Combined Extreme Index (CEI) for North America, Asia, Australia, and Europe (Dittus et al., 2016). While various studies have described increasing trends in several heat wave metrics (HWD, HWA, EHF, etc.) in different regions (e.g., Bandyopadhyay et al., 2016; Cowan et al., 2014; Sanderson et al., 2017), few recent studies have explicitly attributed these changes to causes; most of them stated that observed trends are consistent with anthropogenic warming. The detected anthropogenic signals are clearly separable from the response to natural forcing, and the results are generally insensitive to the use of different model samples, as well as different data availability, indicating robust attribution. Studies of monthly, seasonal, and annual records in various regions (Kendon, 2014; Lewis and King, 2015; Bador et al., 2016; Meehl et al., 2016; Zhou et al., 2019a) and globally (King, 2017) show an increase in the breaking of

1 hot records and a decrease in the breaking of cold records (King, 2017). Changes in anthropogenically-  
2 attributable record-breaking rates are noted to be largest over the Northern Hemisphere land areas (Shiogama  
3 et al., 2016). Yin and Sun (2018) found clear evidence of an anthropogenic signal in the changes in the  
4 number of frost and icing days, when multiple model simulations were used. In some key wheat-producing  
5 regions of southern Australia, increases in frost days or frost season length have been reported (Dittus et al.,  
6 2014; Crimp et al., 2016); these changes are linked to decreases in rainfall, cloud-cover, and subtropical  
7 ridge strength, despite an overall increase in regional mean temperatures (Dittus et al., 2014; Pepler et al.,  
8 2018).

9  
10 A significant advance since AR5 has been a large number of studies focusing on extreme temperature events  
11 at monthly and seasonal scales, using various extreme event attribution methods. Diffenbaugh et al. (2017)  
12 found anthropogenic warming has increased the severity and probability of the hottest month over >80% of  
13 the available observational area on the global scale. Christidis and Stott (2014) provide clear evidence that  
14 warm events have become more probable because of anthropogenic forcings. Sun et al. (2014) found human  
15 influence has caused a more than 60-fold increase in the probability of the extreme warm 2013 summer in  
16 eastern China since the 1950s. Human influence is found to have increased the probability of the historically  
17 hottest summers in many regions of the world, both in terms of mean temperature (Mueller et al., 2016a) and  
18 wet-bulb globe temperature (WBGT) (Li et al., 2017a). In most regions of the Northern Hemisphere,  
19 changes in the probability of extreme summer average WBGT were found to be about an order of magnitude  
20 larger than changes in the probability of extreme hot summers estimated by surface air temperature (Li et al.,  
21 2017a). In addition to these generalised, global-scale approaches, extreme event studies have found an  
22 attributable increase in the probability of hot annual and seasonal temperatures in many locations, including  
23 Australia (Knutson et al., 2014a; Lewis and Karoly, 2014), China (Sun et al., 2014; Sparrow et al., 2018;  
24 Zhou et al., 2020), Korea (Kim et al., 2018c) and Europe (King et al., 2015b).

25  
26 There have also been many extreme event attribution studies that examined short duration temperature  
27 extremes, including daily temperatures, temperature indices, and heat wave metrics. Examples of these  
28 events from different regions are summarised in various annual Explaining Extreme Events supplements of  
29 the Bulletin of the American Meteorological Society (Peterson et al., 2012, 2013b, Herring et al., 2014,  
30 2015, 2016, 2018, 2019, 2020), including a number of approaches to examine extreme events (described in  
31 Easterling et al., 2016; Otto, 2017; Stott et al., 2016). Several studies of recent events from 2016 onwards  
32 have determined an infinite risk ratio (fraction of attributable risk (FAR) of 1), indicating the occurrence  
33 probability for such events is close to zero in model simulations without anthropogenic influences (see  
34 Herring et al., 2018, 2019, 2020; Imada et al., 2019; Vogel et al., 2019). Though it is difficult to accurately  
35 estimate the lower bound of the uncertainty range of the FAR in these cases (Paciorek et al., 2018), the fact  
36 that those events are so far outside the envelop of the models with only natural forcing indicates that it is  
37 *extremely unlikely* for those events to occur without human influence.

38  
39 Studies that focused on the attributable signal in observed cold extreme events show human influence  
40 reducing the probability of those events. Individual attribution studies on the extremely cold winter of 2011  
41 in Europe (Peterson et al., 2012), in the eastern US during 2014 and 2015 (Trenary et al., 2015, 2016; Wolter  
42 et al., 2015; Bellprat et al., 2016), in the cold spring of 2013 in the United Kingdom (Christidis et al., 2014),  
43 and of 2016 in eastern China (Qian et al., 2018; Sun et al., 2018b) all showed a reduced probability due to  
44 human influence on the climate. An exception is the study of Grose et al. (2018), who found an increase in  
45 the probability of the severe western Australian frost of 2016 due to anthropogenically-driven changes in  
46 circulation patterns that drive cold outbreaks and frost probability.

47  
48 Different event attribution studies can produce a wide range of changes in the probability of event  
49 occurrence because of different framing. The temperature event definition itself plays a crucial role in the  
50 attributable signal (Fischer and Knutti, 2015; Kirchmeier-Young et al., 2019). Large-scale, longer-duration  
51 events tend to have notably larger attributable risk ratios (Angélil et al., 2014, 2018; Uhe et al., 2016;  
52 Harrington, 2017; Kirchmeier-Young et al., 2019), as natural variability is smaller. While uncertainty in the  
53 best estimates of the risk ratios may be large, their lower bounds can be quite insensitive to uncertainties in  
54 observations or model descriptions, thus increasing confidence in conservative attribution statements (Jeon et  
55 al., 2016).

1  
2 The relative strength of anthropogenic influences on temperature extremes is regionally variable, in part due  
3 to differences in changes in atmospheric circulation, land surface feedbacks, and other external drivers like  
4 aerosols. For example, in the Mediterranean and over western Europe, risk ratios on the order of 100 have  
5 been found (Kew et al., 2019; Vautard et al., 2020a), whereas in the US, changes are much less pronounced.  
6 This is probably a reflection of the land-surface feedback enhanced extreme 1930s temperatures that reduce  
7 the rarity of recent extremes, in addition to the definition of the events and framing of attribution analyses  
8 (e.g., spatial and temporal scales considered). Local forcing may mask or enhance the warming effect of  
9 greenhouse gases. In India, short-lived aerosols or an increase in irrigation may be masking the warming  
10 effect of greenhouse gases (Wehner et al., 2018c). Irrigation and crop intensification have been shown to  
11 lead to a cooling in some regions, in particular in North America, Europe, and India (Mueller et al., 2016b;  
12 Thiery et al., 2017, 2020; Chen and Dirmeyer, 2019), (*high confidence*). Deforestation has contributed about  
13 one third of the total warming of hot extremes in some mid-latitude regions since pre-industrial times  
14 (Lejeune et al., 2018). Despite all of these differences, and larger uncertainties at the regional scale, nearly  
15 all studies demonstrated that human influence has contributed to an increase in the frequency or intensity of  
16 hot extremes and to a decrease in the frequency or intensity of cold extremes.

17  
18 In summary, long-term changes in various aspects of long- and short-duration extreme temperatures,  
19 including intensity, frequency, and duration have been detected in observations and attributed to human  
20 influence at global and continental scales. It is *extremely likely* that human influence is the main contributor  
21 to the observed increase in the intensity and frequency of hot extremes and the observed decrease in the  
22 intensity and frequency of cold extremes on the global scale. It is *very likely* that this applies on continental  
23 scales as well. Some specific recent hot extreme events would have been *extremely unlikely* to occur without  
24 human influence on the climate system. Changes in aerosol concentrations have affected trends in hot  
25 extremes in some regions, with the presence of aerosols leading to attenuated warming, in particular from  
26 1950-1980. Crop intensification, irrigation and no-till farming have attenuated increases in summer hot  
27 extremes in some regions, such as central North America (*medium confidence*).

### 30 11.3.5 Projections

31  
32 AR5 (Chapter 12, Collins et al., 2013a) concluded it is *virtually certain* there will be more frequent hot  
33 extremes and fewer cold extremes at the global scale and over most land areas in a future warmer climate  
34 and it is *very likely* heat waves will occur with a higher frequency and longer duration. SR15 (Chapter 3,  
35 Hoegh-Guldberg et al., 2018) assessment on projected changes in hot extremes at 1.5°C and 2°C global  
36 warming is consistent with the AR5 assessment, concluding it is *very likely* a global warming of 2°C, when  
37 compared with a 1.5°C warming, would lead to more frequent and more intense hot extremes on land, as  
38 well as to longer warm spells, affecting many densely-inhabited regions. SR15 also assessed it is *very likely*  
39 the strongest increases in the frequency of hot extremes are projected for the rarest events, while cold  
40 extremes will become less intense and less frequent and cold spells will be shorter.

41  
42 New studies since AR5 and SR15 confirm these assessments. New literature since AR5 includes projections  
43 of temperature-related extremes in relation to changes in mean temperatures, projections based on CMIP6  
44 simulations, projections based on stabilized global warming levels, and the use of new metrics. Constraints  
45 for the projected changes in hot extremes were also provided (Borodina et al., 2017b; Sippel et al., 2017b;  
46 Vogel et al., 2017). Overall, projected changes in the magnitude of extreme temperatures over land are larger  
47 than changes in global mean temperature, over mid-latitude land regions in particular (Figures 11.3, 11.11),  
48 (Fischer et al., 2014; Seneviratne et al., 2016; Sanderson et al., 2017a; Wehner et al., 2018b; Di Luca et al.,  
49 2020a). Large warming in hot and cold extremes will occur even at the 1.5°C global warming level (Figure  
50 11.11). At this level, widespread significant changes at the grid-box level occur for different temperature  
51 indices (Aerenson et al., 2018). In agreement with CMIP5 projections, CMIP6 simulations show that a 0.5°C  
52 increment in global warming will significantly increase the intensity and frequency of hot extremes and  
53 decrease the intensity and frequency of cold extremes on the global scale (Figures 11.6, 11.8, 11.12). It takes  
54 less than half of a degree for the changes in TXx to emerge above the level of natural variability (Figure  
55 11.8) and the 66% ranges of the land medians of the 10-year or 50-year TXx events do not overlap between

1 1.0°C and 1.5°C in the CMIP6 multi-model ensemble simulations (Figure 11.6, Li et al., 2020).

2  
3  
4 **[START FIGURE 11.11 HERE]**

5  
6  
7 **Figure 11.11:** Projected changes in (a-c) annual maximum temperature (TXx) and (d-f) annual minimum temperature (TNn) at 1.5°C, 2°C, and 4°C of global warming compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-model ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The numbers in the top right indicate the number of simulations included. Uncertainty is represented using the simple approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models agree on sign of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

17  
18 **[END FIGURE 11.11 HERE]**

19  
20  
21 Projected warming is larger for TNn and exhibits strong equator-to-pole amplification similar to the warming of boreal winter mean temperatures. The warming of TXx is more uniform over land and does not exhibit this behaviour (Figure 11.11). The warming of temperature extremes on global and regional scales tends to scale linearly with global warming (Section 11.1.4) (Fischer et al., 2014; Seneviratne et al., 2016, Wartenburger et al., 2017; Li et al., 2020; see also SR15, Chapter 3). In the mid-latitudes, the rate of warming of hot extremes can be as large as twice the rate of global warming (Figure 11.11). In the Arctic winter, the rate of warming of the temperature of the coldest nights is about three times the rate of global warming (Appendix Figure 11.A.1). Projected changes in temperature extremes can deviate from projected changes in annual mean warming in the same regions (Figure 11.3, Figs. 11.A.1 and 11.A.2, Di Luca et al., 2020a; Wehner, 2020) due to the additional processes that control the response of regional extremes, including, in particular, soil moisture-evapotranspiration-temperature feedbacks for hot extremes in the mid-latitudes and subtropical regions, and snow/ice-albedo-temperature feedbacks in high-latitude regions.

33  
34  
35 **[START FIGURE 11.12 HERE]**

36  
37 **Figure 11.12:** Projected changes in the intensity of extreme temperature events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1851-1900 baseline. Extreme temperature events are defined as the daily maximum temperatures (TXx) that were exceeded on average once during a 10-year period (10-year event, blue) and that once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the intensity changes across the space, and the whiskers extend to the 90% uncertainty range. The results are based on the multi-model ensemble median estimated from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Adapted from (Li et al., 2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

48  
49 **[END FIGURE 11.12 HERE]**

50  
51  
52 The probability of exceeding a certain hot extreme threshold will increase, while those for cold extreme will decrease with global warming (Mueller et al., 2016a; Lewis et al., 2017b; Suarez-Gutierrez et al., 2020b). The changes tend to scale nonlinearly with the level of global warming, with larger changes for more rare events (Section 11.2.4 and CCB 11.11; Figure. 11.6 and 11.12; e.g. Fischer and Knutti, 2015, Kharin et al., 2018; Li et al., 2020). For example, the CMIP5 ensemble projects the frequency of the present-day climate 20-year hottest daily temperature to increase by 80% at the 1.5°C global warming level and by 180% at the 2.0°C global warming level, and the frequency of the present-day climate 100-year hottest daily temperature

1 to increase by 200% and more than 700% at the 1.5°C and 2.0°C warming levels, respectively (Kharin et al.,  
2 2018). CMIP6 simulations project similar changes (Li et al., 2020a).

3  
4 Tebaldi and Wehner (2018) showed that at the middle of the 21<sup>st</sup> century, 66% of the land surface area would  
5 experience the present-day 20-year return values of TX<sub>x</sub> and the running 3-day average of the daily  
6 maximum temperature every other year on average under the RCP8.5 scenario, as opposed to only 34%  
7 under RCP4.5. By the end of the century, these area fractions increase to 92% and 62%, respectively. Such  
8 nonlinearities in the characteristics of future regional extremes are shown, for instance, for Europe (Lionello  
9 and Scarascia, 2020; Spinoni et al., 2018a; Dosio and Fischer, 2018), Asia (Guo et al., 2017; Harrington and  
10 Otto, 2018b; King et al., 2018), and Australia (Lewis et al., 2017a) under various global warming thresholds.  
11 The non-linear increase in fixed-threshold indices (e.g., percentile-based for a given reference period or  
12 based on an absolute threshold) as a function of global warming is consistent with a linear warming of the  
13 absolute temperature of the temperature extremes (e.g., Whan et al., 2015). Compared to the historical  
14 climate, warming will result in strong increases in heat wave area, duration, and magnitude (Vogel et al.,  
15 2020b). These changes are mostly due to the increase in mean seasonal temperature, rather than changes in  
16 temperature variability, though the latter can have an effect in some regions (Di Luca et al., 2020a; Suarez-  
17 Gutierrez et al., 2020a; Brown, 2020).

18  
19 Projections of temperature-related extremes in RCMs in the CORDEX regions demonstrate robust increases  
20 under future scenarios and can provide information on finer spatial scales than GCMs (e.g. Coppola et al.,  
21 2021). Five RCMs in the CORDEX-East Asia region project decreases in the 20-year return values of  
22 temperature extremes (summer maxima), with models that exhibit warm biases projecting stronger warming  
23 (Park and Min, 2018). Similarly, in the African domain, future increases in TX<sub>90p</sub> and TN<sub>90p</sub> are projected  
24 (Dosio, 2017; Mostafa et al., 2019). This regional-scale analysis provides fine scale information, such as  
25 distinguishing the increase in TX<sub>90p</sub> over sub-equatorial Africa (Democratic Republic of Congo, Angola  
26 and Zambia) with values over the Gulf of Guinea, Central African Republic, South Sudan, and Ethiopia.  
27 Empirical-statistical downscaling has also been used to produce more robust estimates for future heat waves  
28 compared to RCMs based on large multi-model ensembles (Furrer et al., 2010; Keellings and Waylen, 2014;  
29 Wang et al., 2015; Benestad et al., 2018).

30  
31 In all continental regions, including Africa (Table 11.4), Asia (Table 11.7), Australasia (Table 11.10),  
32 Central and South America (Table 11.13), Europe (Table 11.16), North America (Table 11.19) and at the  
33 continental scale, it is *very likely* the intensity and frequency of hot extremes will increase and the intensity  
34 and frequency of cold extremes will decrease compared with the 1995-2014 baseline, even under 1.5°C  
35 global warming, and those changes are *virtually certain* to occur under 4°C global warming. At the regional  
36 scale and for almost all AR6 regions, it is *likely* the intensity and frequency of hot extremes will increase and  
37 the intensity and frequency of cold extremes will decrease compared with the 1995-2014 baseline, even  
38 under 1.5°C global warming and those changes will *virtually certain* to occur under 4°C global warming.  
39 Exceptions include lower confidence in the projected decrease in the intensity and frequency of cold  
40 extremes compared with the 1995-2014 baseline under 1.5°C of global warming (*medium confidence*) and  
41 4°C of global warming (*very likely*) in North Central America, Central North America, and Western North  
42 America.

43  
44 In Africa (Table 11.4), evidence includes increases in the intensity and frequency of hot extremes, such as  
45 warm days, warm nights, and heat waves, and decreases in the intensity and frequency of cold extremes,  
46 such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and CORDEX  
47 simulations (Giorgi et al., 2014; Engelbrecht et al., 2015; Lelieveld et al., 2016; Russo et al., 2016; Dosio,  
48 2017; Bathiany et al., 2018; Mba et al., 2018; Nangombe et al., 2018; Weber et al., 2018; Kruger et al., 2019;  
49 Coppola et al., 2021; Li et al., 2020). Cold spells are projected to decrease under all RCPs and even at low  
50 warming levels in West and Central Africa (Diedhiou et al., 2018) and the number of cold days is projected  
51 to decrease in East Africa (Ongoma et al., 2018b).

52  
53 In Asia (Table 11.7), evidence includes increases in the intensity and frequency of hot extremes, such as  
54 warm days, warm nights, and heat waves, and decreases in the intensity and frequency of cold extremes,  
55 such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and CORDEX

1 simulations (Gao et al., 2018; Han et al., 2018; Li et al., 2019b; Pal and Eltahir, 2016; Shin et al., 2018;  
2 Sillmann et al., 2013b; Singh and Goyal, 2016; Sui et al., 2018; Xu et al., 2017; Zhang et al., 2015c; Zhao et  
3 al., 2015; Zhou et al., 2014; Zhu et al., 2020). More intense heat waves of longer durations and occurring at a  
4 higher frequency are projected over India (Murari et al., 2015; Mishra et al., 2017) and Pakistan (Nasim et  
5 al., 2018). Future mid-latitude warm extremes, similar to those experienced during the 2010 event, are  
6 projected to become more extreme, with temperature extremes increasing potentially by 8.4°C (RCP8.5)  
7 over northwest Asia (van der Schrier et al., 2018). Over WSB, ESB and RFE, an increase in extreme heat  
8 durations is expected in all scenarios (Sillmann et al., 2013b; Kattsov et al., 2017; Reyer et al., 2017). In the  
9 MENA regions (ARP, WCA), extreme temperatures could increase by almost 7°C by 2100 under RCP8.5  
10 (Lelieveld et al., 2016).

11  
12 In Australasia (Table 11.10), evidence includes increases in the intensity and frequency of hot extremes, such  
13 as warm days, warm nights, and heat waves, and decreases in the intensity and frequency of cold extremes,  
14 such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and CORDEX  
15 simulations (Coppola et al., 2021; Alexander and Arblaster, 2017; CSIRO and BOM, 2015; Herold et al.,  
16 2018; Lewis et al., 2017a; Evans et al., 2020). Over most of Australia, increases in the intensity and  
17 frequency of hot extremes are projected to be predominantly driven by the long-term increase in mean  
18 temperatures (Di Luca et al., 2020a). Future projections indicate a decrease in the number of frost days  
19 regardless of the region and season considered (Alexander and Arblaster, 2017; Herold et al., 2018).

20  
21 In Central and South America (Table 11.13), evidence includes increases in the intensity and frequency of  
22 hot extremes, such as warm days, warm nights, and heat waves, and decreases in the intensity and frequency  
23 of cold extremes, such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and  
24 CORDEX simulations (Chou et al., 2014a; Cabré et al., 2016; López-Franca et al., 2016; Stennett-Brown et  
25 al., 2017; Li et al., 2020a; Coppola et al., 2021b; Vichot-Llano et al., 2021). Over SES during the austral  
26 summer, the increase in the frequency of TN90p is larger than that projected for TX90p, consistent with  
27 observed past changes (López-Franca et al., 2016). Under RCP8.5, the number of heat wave days are  
28 projected to increase for the intra-Americas region for the end of the 21st century (Angeles-Malaspina et al.,  
29 2018). A general decrease in the frequency of cold spells and frost days is projected as indicated by several  
30 indices based on minimum temperature (López-Franca et al., 2016).

31  
32 In Europe (Table 11.16), evidence includes increases in the intensity and frequency of hot extremes, such as  
33 warm days, warm nights, and heat waves, and decreases in the intensity and frequency of cold extremes,  
34 such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and CORDEX  
35 simulations (Coppola et al., 2021; Cardoso et al., 2019; Jacob et al., 2018; Lau and Nath, 2014; Lhotka et al.,  
36 2018; Lionello and Scarascia, 2020; Molina et al., 2020; Ozturk et al., 2015; Rasmijn et al., 2018; Russo et  
37 al., 2015; Schoetter et al., 2015; Suarez-Gutierrez et al., 2018; Vogel et al., 2017; Winter et al., 2017; Li et  
38 al., 2020). Increases in heat waves are greater over the southern Mediterranean and Scandinavia (Forzieri et  
39 al., 2016; Abaurrea et al., 2018; Dosio and Fischer, 2018; Rohat et al., 2019). The biggest increases in the  
40 number of heat wave days are expected for southern European cities (Guerreiro et al., 2018a; Junk et al.,  
41 2019), and Central European cities will see the biggest increases in maximum heat wave temperatures  
42 (Guerreiro et al., 2018a).

43  
44 In North America (Table 11.19), evidence includes increases in the intensity and frequency of hot extremes,  
45 such as warm days, warm nights, and heat waves, and decreases in the intensity and frequency of cold  
46 extremes, such as cold days and cold nights, over the continent as projected by CMIP5, CMIP6, and  
47 CORDEX simulations (Li et al., 2020; Coppola et al., 2021; Alexandru, 2018; Grotjahn et al., 2016; Li et al.,  
48 2018a; Vose et al., 2017a; Yang et al., 2018a; Zhang et al., 2019d). Projections of temperature extremes for  
49 the end of the 21st century show that warm days and nights are *very likely* to increase and cold days and  
50 nights are *very likely* to decrease in all regions. There is *medium confidence* in large increases in warm days  
51 and warm nights in summer, particularly over the United States, and in large decreases in cold days in  
52 Canada in fall and winter (Li et al., 2020; Coppola et al., 2021; Alexandru, 2018; Grotjahn et al., 2016; Li et  
53 al., 2018a; Vose et al., 2017a; Yang et al., 2018a; Zhang et al., 2019d). Minimum winter temperatures are  
54 projected to rise faster than mean winter temperatures (Underwood et al., 2017). Projections for the end of  
55 the century under RCP8.5 showed the 4-day cold spell that happens on average once every 5 years is

1 projected to warm by more than 10 °C and CMIP5 models do not project current 1-in-20 year annual  
2 minimum temperature extremes to recur over much of the continent (Wuebbles et al., 2014).  
3

4 In summary, it is *virtually certain* that further increases in the intensity and frequency of hot extremes and  
5 decreases in the intensity and frequency of cold extremes will occur throughout the 21<sup>st</sup> century and around  
6 the world. It is *virtually certain* the number of hot days and hot nights and the length, frequency, and/or  
7 intensity of warm spells or heat waves compared to 1995-2014 will increase over most land areas. In most  
8 regions, changes in the magnitude of temperature extremes are proportional to global warming levels (*high*  
9 *confidence*). The highest increase of temperature of hottest days is projected in some mid-latitude and semi-  
10 arid regions, at about 1.5 time to twice the rate of global warming (high confidence). The highest increase of  
11 temperature of coldest days is projected in Arctic regions, at about three times the rate of global warming  
12 (high confidence). The probability of temperature extremes generally increases non-linearly with increasing  
13 global warming levels (*high confidence*). Confidence in assessments depends on the spatial and temporal  
14 scales of the extreme in question, with *high confidence* in projections of temperature-related extremes at  
15 global and continental scales for daily to seasonal scales. There is *high confidence* that, on land, the  
16 magnitude of temperature extremes increases more strongly than global mean temperature.  
17

## 18 19 **11.4 Heavy precipitation**

20  
21 This section assesses changes in heavy precipitation at global and regional scales. The main focus is on  
22 extreme precipitation at a daily scale where literature is most concentrated, though extremes of shorter (sub-  
23 daily) and longer (five-day or more) durations are also assessed to the extent the literature allows.  
24

### 25 26 **11.4.1 Mechanisms and drivers**

27  
28 SREX (Chapter 3, Seneviratne et al., 2012) assessed changes in heavy precipitation in the context of the  
29 effects of thermodynamic and dynamic changes. Box 11.1 assesses thermodynamic and dynamic changes in  
30 a warming world to aid the understanding of changes in observations and projections in some extremes and  
31 the sources of uncertainties (See also Chapter 8, Section 8.2.3.2). In general, warming increases the  
32 atmospheric water-holding capacity following the Clausius-Clapeyron (C-C) relation. This thermodynamic  
33 effect results in an increase in extreme precipitation at a similar rate at the global scale. On a regional scale,  
34 changes in extreme precipitation are further modulated by dynamic changes (Box 11.1).  
35

36 Large-scale modes of variability, such as the North Atlantic Oscillation (NAO), El Niño-Southern  
37 Oscillation (ENSO), Atlantic Multidecadal Variability (AMV), and Pacific Decadal Variability (PDV)  
38 (Annex VI), modulate precipitation extremes through changes in environmental conditions or embedded  
39 storms (Section 8.3.2). Latent heating can invigorate these storms (Nie et al., 2018; Zhang et al., 2019g);  
40 changes in dynamics can increase precipitation intensity above that expected from the C-C scaling rate  
41 (8.2.3.2, Box 11.1, and Section 11.7). Additionally, the efficiency of converting atmospheric moisture into  
42 precipitation can change as a result of cloud microphysical adjustment to warming, resulting in changes in  
43 the characteristics of extreme precipitation; but changes in precipitation efficiency in a warming world are  
44 highly uncertain (Sui et al., 2020).  
45

46 It is difficult to separate the effect of global warming from internal variability in the observed changes in the  
47 modes of variability (Section 2.4). Future projections of modes of variability are highly uncertain (Section  
48 4.3.3), resulting in uncertainty in regional projections of extreme precipitation. Future warming may amplify  
49 monsoonal extreme precipitation. Changes in extreme storms, including tropical/extratropical cyclones and  
50 severe convective storms, result in changes in extreme precipitation (Section 11.7). Also, changes in sea  
51 surface temperatures (SSTs) alter land-sea contrast, leading to changes in precipitation extremes near coastal  
52 regions. For example, the projected larger SST increase near the coasts of East Asia and India can result in  
53 heavier rainfall near these coastal areas from tropical cyclones (Mei and Xie, 2016) or torrential rains  
54 (Manda et al., 2014). The warming in the western Indian Ocean is associated with increases in moisture  
55 surges on the low-level monsoon westerlies towards the Indian subcontinent, which may lead to an increase



1 in the occurrence of precipitation extremes over central India (Krishnan et al., 2016; Roxy et al., 2017).

2  
3 Decreases in atmospheric aerosols results in warming and thus an increase in extreme precipitation (Samset  
4 et al., 2018; Sillmann et al., 2019). Changes in atmospheric aerosols also result in dynamic changes such as  
5 changes in tropical cyclones (Takahashi et al., 2017; Strong et al., 2018). Uncertainty in the projections of  
6 future aerosol emissions results in additional uncertainty in the heavy precipitation projections of the 21<sup>st</sup>  
7 century (Lin et al., 2016).

8  
9 There has been new evidence of the effect of local land use and land cover change on heavy precipitation.  
10 There is a growing set of literature linking increases in heavy precipitation in urban centres to urbanization  
11 (Argüeso et al., 2016; Zhang et al., 2019f). Urbanization intensifies extreme precipitation, especially in the  
12 afternoon and early evening, over the urban area and its downwind region (*medium confidence*) (Box 10.3).  
13 There are four possible mechanisms: a) increases in atmospheric moisture due to horizontal convergence of  
14 air associated with the urban heat island effect (Shastri et al., 2015; Argüeso et al., 2016); b) increases in  
15 condensation due to urban aerosol emissions (Han et al., 2011; Sarangi et al., 2017); c) aerosol pollution that  
16 impacts cloud microphysics (Schmid and Niyogi, 2017) (Box 8.1); and d) urban structures that impede  
17 atmospheric motion (Ganeshan and Murtugudde, 2015; Paul et al., 2018; Shepherd, 2013). Other local  
18 forcing, including reservoirs (Woldemichael et al., 2012), irrigation (Devanand et al., 2019), or large-scale  
19 land use and land cover change (Odoulami et al., 2019), can also affect local extreme precipitation.

20  
21 In summary, precipitation extremes are controlled by both thermodynamic and dynamic processes.  
22 Warming-induced thermodynamic change results in an increase in extreme precipitation, at a rate that  
23 closely follows the Clausius-Clapeyron relationship at the global scale (*high confidence*). The effects of  
24 warming-induced changes in dynamic drivers on extreme precipitation are more complicated, difficult to  
25 quantify and are an uncertain aspect of projections. Precipitation extremes are also affected by forcings other  
26 than changes in greenhouse gases, including changes in aerosols, land use and land cover change, and  
27 urbanization (*medium confidence*).

#### 28 29 30 **11.4.2 Observed Trends**

31  
32 Both SREX (Chapter 3, Seneviratne et al., 2012) and AR5 (IPCC, 2014 Chapter 2) concluded it was *likely*  
33 the number of heavy precipitation events over land had increased in more regions than it had decreased,  
34 though there were wide regional and seasonal variations, and trends in many locations were not statistically  
35 significant. This assessment has been strengthened with multiple studies finding robust evidence of the  
36 intensification of extreme precipitation at global and continental scales, regardless of spatial and temporal  
37 coverage of observations and the methods of data processing and analysis.

38  
39 The average annual maximum precipitation amount in a day (Rx1day) has significantly increased since the  
40 mid-20<sup>th</sup> century over land (Du et al., 2019; Dunn et al., 2020) and in the humid and dry regions of the globe  
41 (Dunn et al., 2020). The percentage of observing stations with statistically significant increases in Rx1day is  
42 larger than expected by chance, while the percentage of stations with statistically significant decreases is  
43 smaller than expected by chance, over the global land as a whole and over North America, Europe, and Asia  
44 (Figure 11.13, Sun et al., 2020) and over global monsoon regions (Zhang and Zhou, 2019) where data  
45 coverage is relatively good. The addition of the past decade of observational data shows a more robust  
46 increase in Rx1day over the global land region (Sun et al., 2020). Light, moderate, and heavy daily  
47 precipitation has all intensified in a gridded daily precipitation data set (Contractor et al., 2020). Daily mean  
48 precipitation intensities have increased since the mid-20th century in a majority of land regions (*high*  
49 *confidence*, Section 8.3.1.3). The probability of precipitation exceeding 50 mm/day increased during 1961-  
50 2018 (Benestad et al., 2019). The globally averaged annual fraction of precipitation from days in the top 5%  
51 (R95pTOT) has also significantly increased (Dunn et al., 2020). The increase in the magnitude of Rx1day in  
52 the 20<sup>th</sup> century is estimated to be at a rate consistent with C-C scaling with respect to global mean  
53 temperature (Fischer and Knutti, 2016; Sun et al., 2020). Studies on past changes in extreme precipitation of  
54 durations longer than a day are more limited, though there are some studies examining long-term trends in  
55 annual maximum five-day precipitation (Rx5day). On global and continental scales, long-term changes in

1 Rx5day are similar to those of Rx1day in many aspects (Zhang and Zhou 2019; Sun et al., 2020). As  
2 discussed below, at the regional scale, changes in Rx5day are also similar to those of Rx1day where there are  
3 analyses of changes in both Rx1day and Rx5day.

4  
5 Overall, there is a lack of systematic analysis of long-term trends in sub-daily extreme precipitation at the  
6 global scale. Often, sub-daily precipitation data have only sporadic spatial coverage and are of limited  
7 length. Additionally, the available data records are far shorter than needed for a robust quantification of past  
8 changes in sub-daily extreme precipitation (Li et al., 2018b). Despite these limitations, there are studies in  
9 regions of almost all continents that generally indicate intensification of sub-daily extreme precipitation,  
10 although *confidence* in an overall increase at the global scale remains *very low*. Studies include an increase in  
11 extreme sub-daily rainfall in summer over South Africa (Sen Roy and Rouault, 2013), annually in Australia  
12 (Guerreiro et al., 2018b), over 23 urban locations in India (Ali and Mishra, 2018), in Peninsular Malaysia  
13 (Syafrina et al., 2015), and in eastern China in the summer season during 1971-2013 (Xiao et al., 2016). In  
14 some regions in Italy (Arnone et al., 2013; Libertino et al., 2019) and in the US during 1950-2011 (Barbero  
15 et al., 2017), there is also an increase. In general, an increase in sub-daily heavy precipitation results in an  
16 increase in pluvial floods over smaller watersheds (Ghausi and Ghosh, 2020).

17  
18 There is a considerable body of literature examining scaling of sub-daily precipitation extremes, conditional  
19 on day-to-day air or dew-point temperatures (Westra et al., 2014; Fowler et al., 2021). This scaling, termed  
20 apparent scaling (Fowler et al., 2020) is robust when different methodologies are used in different regions,  
21 ranging between the C-C and two-times the C-C rate (e.g. Burdanowitz et al., 2019; Formayer and Fritz,  
22 2017; Lenderink et al., 2017). This is confirmed when sub-daily precipitation data collected from multiple  
23 continents (Lewis et al., 2019a) are analysed in a consistent manner using different methods (Ali et al.,  
24 2021). It has been hoped that apparent scaling might be used to help understand past and future changes in  
25 extreme sub-daily precipitation. However, apparent scaling samples multiple synoptic weather states, mixing  
26 thermodynamic and dynamic factors that are not directly relevant for climate change responses (8.2.3.2)  
27 (Prein et al., 2016b; Bao et al., 2017; Zhang et al., 2017c; Drobinski et al., 2018; Sun et al., 2019d). The  
28 spatial pattern of apparent scaling is different from those of projected changes over Australia (Bao et al.,  
29 2017) and North America (Sun et al., 2019) in regional climate model simulations. It thus remains difficult to  
30 use the knowledge about apparent scaling to infer past and future changes in extreme sub-daily precipitation  
31 according to observed and projected changes in local temperature.

32  
33 In Africa (Table 11.5), evidence shows an increase in extreme daily precipitation for the late half of the 20<sup>th</sup>  
34 century over the continent where data are available; there is a larger percentage of stations showing  
35 significant increases in extreme daily precipitation than decreases (Sun et al., 2020). There are increases in  
36 different metrics relevant to extreme precipitation in various regions of the continent (Chaney et al., 2014;  
37 Harrison et al., 2019; Dunn et al., 2020; Sun et al., 2020). There is an increase in extreme precipitation  
38 events in southern Africa (Weldon and Reason, 2014; Kruger et al., 2019) and a general increase in heavy  
39 precipitation over East Africa, the Greater Horn of Africa (Omondi et al., 2014). Over sub-Saharan Africa,  
40 increases in the frequency and intensity of extreme precipitation have been observed over the well-gauged  
41 areas during 1950-2013; however, this covers only 15% of the total area of sub-Saharan Africa (Harrison et  
42 al., 2019). *Confidence* about the increase in extreme precipitation for some regions where observations are  
43 more abundant is *medium*, but for Africa as whole, it is *low* because of a general lack of continent-wide  
44 systematic analysis, the sporadic nature of available precipitation data over the continent, and spatially non-  
45 homogenous trends in places where data are available (Donat et al., 2014a; Mathbout et al., 2018; Alexander  
46 et al., 2019; Funk et al., 2020)

47  
48 In Asia (Table 11.8), there is *robust evidence* that extreme precipitation has increased since the 1950s (*high*  
49 *confidence*), however this is dominated by high spatial variability. Increases in Rx1day and Rx5day during  
50 1950-2018 are found over two thirds of stations and the percentage of stations with statistically significant  
51 trends is significantly larger than can be expected by chance (Sun et al., 2020, also Fig 11.13). An increase  
52 in extreme precipitation has also been observed in various regional studies based on different metrics of  
53 extreme precipitation and different spatial and temporal coverage of the data. These include an increase in  
54 daily precipitation extremes over central Asia (Hu et al., 2016), most of South Asia (Zahid and Rasul, 2012;  
55 Pai et al., 2015; Sheikh et al., 2015; Adnan et al., 2016; Malik et al., 2016; Dimri et al., 2017; Priya et al.,

1 2017; Roxy et al., 2017; Hunt et al., 2018; Kim et al., 2019; Wester et al., 2019), the Arabian Peninsula  
2 (Rahimi and Fatemi, 2019; Almazroui and Saeed, 2020; Atif et al., 2020), Southeast Asia (Siswanto et al.,  
3 2015; Supari et al., 2017; Cheong et al., 2018); the northwest Himalaya (Malik et al., 2016), parts of east  
4 Asia (Nayak et al., 2017; Baek et al., 2017; Ye and Li, 2017), the western Himalayas since the 1950s (Ridley  
5 et al., 2013; Dimri et al., 2015; Madhura et al., 2015), WSB, ESB and RFE (Donat et al., 2016a) and a  
6 decrease was found over the eastern Himalayas (Sheikh et al., 2015; Talchabhadel et al., 2018). Increases  
7 have been observed over Jakarta (Siswanto et al., 2015), but Rx1day over most parts of the Maritime  
8 Continent has decreased (Villafuerte and Matsumoto, 2015). Trends in extreme precipitation over China are  
9 mixed with increases and decreases (Fu et al., 2013a; Jiang et al., 2013; Ma et al., 2015; Yin et al., 2015;  
10 Xiao et al., 2016) and are not significant over China as whole (Li et al., 2018c; Ge et al., 2017; Hu et al.,  
11 2016; Jiang et al., 2013; Liu et al., 2019b; Chen et al., 2021; Deng et al., 2018; He and Zhai, 2018; Tao et al.,  
12 2018). With few exceptions, most Southeast Asian countries have experienced an increase in rainfall  
13 intensity, but with a reduced number of wet days (Donat et al., 2016a; Cheong et al., 2018; Naveendrakumar  
14 et al., 2019), though large differences in trends exists if the trends are estimated from different datasets  
15 including gauge-based, remotely-sensed, and reanalysis over a relatively short period (Kim et al. 2019).  
16 There is a significant increase in heavy rainfall ( $>100 \text{ mm day}^{-1}$ ) and a significant decrease in moderate  
17 rainfall ( $5\text{--}100 \text{ mm day}^{-1}$ ) in central India during the South Asian monsoon season (Deshpande et al., 2016;  
18 Roxy et al., 2017).

19  
20 In Australasia (Table 11.11), available evidence has not shown an increase or a decrease in heavy  
21 precipitation over Australasia as a whole (*medium confidence*), but heavy precipitation tends to increase over  
22 northern Australia (particularly the northwest) and decrease over the eastern and southern regions (e.g.,  
23 Jakob and Walland, 2016; Dey et al., 2018; Guerreiro et al., 2018; Dunn et al., 2020; Sun et al., 2020).  
24 Available studies that used long-term observations since the mid-20th century showed nearly as many  
25 stations with an increase as those with a decrease in heavy precipitation (Jakob and Walland, 2016) or  
26 slightly more stations with a decrease than with an increase in Rx1day and Rx5day (Sun et al., 2020), or  
27 strong differences in Rx1day trends with increases over northern Australia and central Australia in general  
28 but mostly decreases over southern Australia and eastern Australia (Dunn et al., 2020). Over New Zealand,  
29 decreases are observed for moderate-heavy precipitation events, but there are no significant trends for very  
30 heavy events (more than 64 mm in a day) for the period 1951-2012. The number of stations with an increase  
31 in very wet days is similar to that with a decrease during 1960-2019 (MfE and Stats NZ, 2020). Overall,  
32 there is *low confidence* in trends in the frequency of heavy rain days with mostly decreases over New  
33 Zealand (Caloiero, 2015; Harrington and Renwick, 2014).

34  
35 In Central and South America (Table 11.14), evidence shows an increase in extreme precipitation, but in  
36 general there is *low confidence*; while continent-wide analyses produced wetting trends, trends are not  
37 robust. Rx1day increased at more stations than it decreased in South America between 1950-2018 (Sun et al.,  
38 2020). Over 1950-2010, both Rx5day and R99p increased over large regions of South America, including  
39 NWS, NSA, and SES (Skansi et al., 2013). There are large regional differences. A decrease in daily extreme  
40 precipitation is observed in northeastern Brazil (Bezerra et al., 2018; Dereczynski et al., 2020; Skansi et al.,  
41 2013). Trends in extreme precipitation indices were not statistically significant over the period 1947-2012  
42 within the São Francisco River basin in the Brazilian semi-arid region (Bezerra et al., 2018). An increase in  
43 extreme rainfall is observed in AMZ with *medium confidence* (Skansi et al., 2013) and in SES with *high*  
44 *confidence* (Ávila et al., 2016; Barros et al., 2015; Lovino et al., 2018; Skansi et al., 2013; Wu and Polvani,  
45 2017; Dereczynski et al., 2020; Valverde and Marengo, 2014). Among all sub-regions, SES shows the  
46 highest rate of increase for rainfall extremes, followed by AMZ (Skansi et al., 2013). Increases in the  
47 intensity of heavy daily rainfall events have been observed in the southern Pacific and in the Titicaca basin  
48 (Huerta and Lavado-Casimiro, 2020; Skansi et al., 2013). In SCA trends in annual precipitation are generally  
49 not significant, although small (but significant) increases are found in Guatemala, El Salvador, and Panama  
50 (Hidalgo et al., 2017). Small positive trends were found in multiple extreme precipitation indices over the  
51 Caribbean region over a short time period (1986-2010) (Stephenson et al., 2014; McLean et al., 2015)

52  
53 In Europe (Table 11.17), there is robust evidence that the magnitude and intensity of extreme precipitation  
54 has *very likely* increased since the 1950s. There is a significant increase in Rx1day and Rx5day during 1950-  
55 2018 in Europe as whole (Sun et al., 2020, also Figure 11.13). The number of stations with increases far

1 exceeds those with decreases in the frequency of daily rainfall exceeding its 90<sup>th</sup> or 95<sup>th</sup> percentile in century-  
2 long series (Cioffi et al., 2015). The 5-, 10-, and 20-year events of one-day and five-day precipitation during  
3 1951-1960 became more common since the 1950s (van den Besselaar et al., 2013). There can be large  
4 discrepancies among studies and regions and seasons (Croitoru et al., 2013; Willems, 2013; Casanueva et al.,  
5 2014; Roth et al., 2014; Fischer et al., 2015); evidence for increasing extreme precipitation is more  
6 frequently observed for summer and winter, but not in other seasons (Madsen et al., 2014; Helama et al.,  
7 2018). An increase is observed in central Europe (Volosciuk et al., 2016; Zeder and Fischer, 2020), and in  
8 Romania (Croitoru et al., 2016). Trends in the Mediterranean region are in general not spatially (Reale and  
9 Lionello, 2013), with decreases in the western Mediterranean and some increases in the eastern  
10 Mediterranean (Rajczak et al., 2013; Casanueva et al., 2014; de Lima et al., 2015; Gajić-Čapka et al., 2015;  
11 Sunyer et al., 2015; Pedron et al., 2017; Serrano-Notivoli et al., 2018; Ribes et al., 2019). In the Netherlands,  
12 the total precipitation contributed from extremes higher than the 99<sup>th</sup> percentile doubles per degree C  
13 increase in warming (Myhre et al., 2019), though extreme rainfall trends in northern Europe may differ in  
14 different seasons (Irannezhad et al., 2017).

15  
16 In North America (Table 11.20), there is robust evidence that the magnitude and intensity of extreme  
17 precipitation has *very likely* increased since the 1950s. Both Rx1day and Rx5day have significantly increased  
18 in North America during 1950-2018 (Sun et al., 2020, also Figure 11.13). There is, however, regional  
19 diversity. In Canada, there is a lack of detectable trends in observed annual maximum daily (or shorter  
20 duration) precipitation (Shephard et al., 2014; Mekis et al., 2015; Vincent et al., 2018). In the United States,  
21 there is an overall increase in one-day heavy precipitation, both in terms of intensity and frequency (Sun et  
22 al., 2020; Donat et al., 2013; Huang et al., 2017; Villarini et al., 2012; Easterling et al., 2017; Wu, 2015;  
23 Howarth et al., 2019), except for the southern part of the US (Hoerling et al., 2016) where internal variability  
24 may have played a substantial role in the lack of observed increases. In Mexico, increases are observed in  
25 R10mm and R95p (Donat et al., 2016a), very wet days over the cities (García-Cueto et al., 2019) and in  
26 PRCPTOT and Rx1day (Donat et al., 2016b).

27  
28 In Small Islands, there is a lack of evidence showing changes in heavy precipitation overall. There were  
29 increases in extreme precipitation in Tobago from 1985–2015 (Stephenson et al., 2014; Dookie et al., 2019)  
30 and decreases in southwestern French Polynesia and the southern subtropics (*low confidence*; Atlas.10; Table  
31 11.5). Extreme precipitation leading to flooding in the small islands has been attributed in part to TCs, as  
32 well as being influenced by ENSO (Khouakhi et al., 2016; Hoegh-Guldberg et al., 2018) (Box 11.5).

### 33 34 35 [START FIGURE 11.13 HERE]

36  
37 **Figure 11.13:** Signs and significance of the observed trends in annual maximum daily precipitation (Rx1day) during  
38 1950–2018 at 8345 stations with sufficient data. (a) Percentage of stations with statistically significant  
39 trends in Rx1day; green dots show positive trends and brown dots negative trends. Box-and-whisker plots  
40 indicate the expected percentage of stations with significant trends due to chance estimated from 1000  
41 bootstrap realizations under a no-trend null hypothesis. The boxes mark the median, 25<sup>th</sup> percentile, and  
42 75<sup>th</sup> percentile. The upper and lower whiskers show the 97.5<sup>th</sup> and the 2.5<sup>th</sup> percentiles, respectively. Maps  
43 of stations with positive (b) and negative (c) trends. The light color indicates stations with non-significant  
44 trends and the dark color stations with significant trends. Significance is determined by a two-tailed test  
45 conducted at the 5% level. Adapted from Sun et al., (2020). © American Meteorological Society. Used  
46 with permission. Further details on data sources and processing are available in the chapter data table  
47 (Table 11.SM.9).

### 48 49 [END FIGURE 11.13 HERE]

50  
51  
52 In summary, the frequency and intensity of heavy precipitation have *likely* increased at the global scale over  
53 a majority of land regions with good observational coverage. Since 1950, the annual maximum amount of  
54 precipitation falling in a day or over five consecutive days has *likely* increased over land regions with  
55 sufficient observational coverage for assessment, with increases in more regions than there are decreases.  
56 Heavy precipitation has *likely* increased on the continental scale over three continents, including North  
57 America, Europe, and Asia where observational data are more abundant. There is *very low confidence* about

1 changes in sub-daily extreme precipitation due to a limited number of studies and the data used in these  
2 studies are often limited.

### 5 **11.4.3 Model evaluation**

6  
7 The evaluation of the skill of climate models to simulate heavy precipitation extremes is challenging due to a  
8 number of factors, including the lack of reliable observations and the spatial scale mismatch between  
9 simulated and observed data (Avila et al., 2015, Alexander et al., 2019). Simulated precipitation represents  
10 areal means, but station-based observations are conducted at point locations and are often sparse. The areal-  
11 reduction factor, the ratio between pointwise station estimates of extreme precipitation and extremes of the  
12 areal mean, can be as large as 130% at CMIP6 resolutions (~100km) (Gervais et al., 2014). Hence, the order  
13 in which gridded station based extreme values are constructed (i.e., if the extreme values are extracted at the  
14 station first and then gridded or if the daily station values are gridded and then the extreme values are  
15 extracted) represents different spatial scales of extreme precipitation and needs to be taken into account in  
16 model evaluation (Wehner et al. 2020). This aspect has been considered in some studies. Reanalysis products  
17 are used in place of station observations for their spatial completeness as well as spatial-scale comparability  
18 (Sillmann et al., 2013a; Kim et al., 2020; Li et al., 2020). However, reanalyses share similar  
19 parameterizations to the models themselves, reducing the objectivity of the comparison.

20  
21 Different generations of the Coupled Model Intercomparison Project (CMIP) models have improved over  
22 time, though quite modestly (Flato et al., 2013; Watterson et al., 2014). Improvements in the representation  
23 of the magnitude of the ETCCDI indices in CMIP5 over CMIP3 (Sillmann et al., 2013a; Chen and Sun,  
24 2015a) have been attributed to higher resolution as higher-resolution models represent smaller areas at  
25 individual grid boxes. Additionally, the spatial distribution of extreme rainfall simulated by high-resolution  
26 models (CMIP5 median resolution ~ 180 × 96) is generally more comparable to observations (Sillmann et al.,  
27 2013b; Kusunoki, 2017, 2018b; Scher et al., 2017) as these models tend to produce more realistic storms  
28 compared to coarser models (11.7.2). Higher horizontal resolution alone improves simulation of extreme  
29 precipitation in some models (Wehner et al., 2014; Kusunoki, 2017, 2018), but this is insufficient in other  
30 models (Bador et al., 2020) as model parameterization also plays a significant role (Wu et al., 2020a). A  
31 simple comparison of climatology may not fully reflect the improvements of the new models that have more  
32 comprehensive formulations of processes (Di Luca et al., 2015). Dittus et al. (2016) found that many of the  
33 eight CMIP5 models they evaluated reproduced the observed increase in the difference between areas  
34 experiencing an extreme high (90%) and an extreme low (10%) proportion of the annual total precipitation  
35 from heavy precipitation (R95p/PRCPTOT) for Northern Hemisphere regions. Additionally, CMIP5 models  
36 reproduced the relation between changes in extreme and non-extreme precipitation: an increase in extreme  
37 precipitation is at the cost of a decrease in non-extreme precipitation (Thackeray et al., 2018), a characteristic  
38 found in the observational record (Gu and Adler, 2018).

39  
40 CMIP6 models perform reasonably well in capturing large-scale features of precipitation extremes, including  
41 intense precipitation extremes in the intertropical convergence zone (ITCZ), and weak precipitation extremes  
42 in dry areas in the tropical regions (Li et al., 2020) but a double-ITCZ bias over the equatorial central and  
43 eastern Pacific that appeared in CMIP5 models remains (3.3.2.1). There are also regional biases in the  
44 magnitude of precipitation extremes (Kim et al., 2020). The models also have difficulties in reproducing  
45 detailed regional patterns of extreme precipitation such as over the northeast US (Agel and Barlow, 2020),  
46 though they performed better for summer extremes over the US (Akinsanola et al., 2020). The comparison  
47 between climatologies in the observations and in model simulations shows that the CMIP6 and CMIP5  
48 models that have similar horizontal resolutions also have similar model evaluation scores and their error  
49 patterns are highly correlated (Wehner et al., 2020). In general, extreme precipitation in CMIP6 models tends  
50 to be somewhat larger than in CMIP5 models (Li et al., 2020a), reflecting smaller spatial scales of extreme  
51 precipitation represented by slightly higher resolution models (Gervais et al., 2014). This is confirmed by  
52 Kim et al. (2020), who showed that Rx1day and Rx5day simulated by CMIP6 models tend to be closer to  
53 point estimates of HadEX3 data (Dunn et al., 2020) than those simulated by CMIP5. Figure 11.14 shows the  
54 multi-model ensemble bias in mean Rx1day over the period 1979-2014 from 21 available CMIP6 models  
55 when compared with observations and reanalyses. Measured by global land root mean square error, the

1 model performance is generally consistent across different observed/reanalysis data products for the extreme  
2 precipitation metric (Figure 11.14). The magnitude of extreme area-mean precipitation simulated by the  
3 CMIP6 models is consistently smaller than the point estimates of HadEX3, but the model values are more  
4 comparable to those of areal-mean values (Figure 11.14) of the ERA5 reanalysis or REGEN (Contractor et  
5 al., 2020b). Taylor-plot-based performance metrics reveal strong similarities in the patterns of extreme  
6 precipitation errors over land regions between CMIP5 and CMIP6 (Srivastava et al., 2020; Wehner et al.,  
7 2020) and between annual mean precipitation errors and Rx1day errors for both generations of models  
8 (Wehner et al., 2020).

9  
10 In general, there is *high confidence* that historical simulations by CMIP5 and CMIP6 models of similar  
11 horizontal resolutions are interchangeable in their performance in simulating the observed climatology of  
12 extreme precipitation.

13  
14  
15 **[START FIGURE 11.14 HERE]**

16  
17 **Figure 11.14:** Multi-model mean bias in annual maximum daily precipitation (Rx1day, %) for the period 1979-2014,  
18 calculated as the difference between the CMIP6 multi-model mean and the average of available  
19 observational or reanalysis products including (a) ERA5, (b) HadEX3, and (c) and REGEN. Bias is  
20 expressed as the percent error relative to the long-term mean of the respective observational data  
21 products. Brown indicates that models are too dry, while green indicates that they are too wet. Areas  
22 without sufficient observational data are shown in grey. Adapted from Wehner et al. (2020) under the  
23 terms of the Creative Commons Attribution license. Further details on data sources and processing are  
24 available in the chapter data table (Table 11.SM.9).

25  
26 **[END FIGURE 11.14 HERE]**

27  
28  
29 Studies using regional climate models (RCMs), for example, CORDEX (Giorgi et al., 2009) over Africa  
30 (Dosio et al., 2015; Klutse et al., 2016; Pinto et al., 2016; Gibba et al., 2019), Australia, East Asia (Park et  
31 al., 2016), Europe (Prein et al., 2016a; Fantini et al., 2018), and parts of North America (Diaconescu et al.,  
32 2018) suggest that extreme rainfall events are better captured in RCMs compared to their host GCMs due to  
33 their ability to address regional characteristics, for example, topography and coastlines. However, CORDEX  
34 simulations do not show good skill over south Asia for heavy precipitation and do not add value with respect  
35 to their GCM source of boundary conditions (Mishra et al., 2014a; Singh et al., 2017b). The evaluation of  
36 models in simulating regional processes is discussed in detail in Chapter 10 (Section 10.3.3.4). The high-  
37 resolution simulation of mid-latitude winter extreme precipitation over land is of similar magnitude to point  
38 observations. Simulation of summer extreme precipitation has a high bias when compared with observations  
39 at the same spatial scale. Simulated extreme precipitation in the tropics also appears to be too large,  
40 indicating possible deficiencies in the parameterization of cumulus convection at this resolution. Indeed,  
41 precipitation distributions at both daily and sub-daily time scales are much improved with a convection-  
42 permitting model (Belušić et al., 2020) over west Africa (Berthou et al., 2019b), East Africa (Finney et al.,  
43 2019), North America and Canada (Cannon and Innocenti, 2019; Innocenti et al., 2019) and over Belgium in  
44 Europe (Vanden Broucke et al., 2019).

45  
46 In summary, there is *high confidence* in the ability of models to capture the large-scale spatial distribution of  
47 precipitation extremes over land. The magnitude and frequency of extreme precipitation simulated by  
48 CMIP6 models are similar to those simulated by CMIP5 models (*high confidence*).

#### 51 **11.4.4 Detection and attribution, event attribution**

52  
53 Both SREX (Chapter 3, Seneviratne et al., 2012) and AR5 (Chapter 10, IPCC, 2014) concluded with *medium*  
54 *confidence* that anthropogenic forcing has contributed to a global-scale intensification of heavy precipitation  
55 over the second half of the 20th century. These assessments were based on the evidence of anthropogenic  
56 influence on aspects of the global hydrological cycle, in particular, the human contribution to the warming-

1 induced observed increase in atmospheric moisture that leads to an increase in heavy precipitation, and  
2 limited evidence of anthropogenic influence on extreme precipitation of durations of one and five days.  
3

4 Since AR5 there has been new and robust evidence and improved understanding of human influence on  
5 extreme precipitation. In particular, detection and attribution analyses have provided consistent and robust  
6 evidence of human influence on extreme precipitation of one- and five-day durations at global to continental  
7 scales. The observed increases in Rx1day and Rx5day over the Northern Hemisphere land area during 1951-  
8 2005 can be attributed to the effect of combined anthropogenic forcing, including greenhouse gases and  
9 anthropogenic aerosols, as simulated by CMIP5 models and the rate of intensification with regard to  
10 warming is consistent with C-C scaling (Zhang et al., 2013). This is confirmed to be robust when an  
11 additional nine years of observational data and the CMIP6 model simulations were used (Paik et al., 2020;  
12 CCB3.2, Figure 1). Additionally, the influence of greenhouse gases is attributed as the dominant contributor  
13 to the observed intensification. The global average of Rx1day in the observations is consistent with  
14 simulations by both CMIP5 and CMIP6 models under anthropogenic forcing, but not under natural forcing  
15 (CCB3.2, Figure 1). The observed increase in the fraction of annual total precipitation falling into the top 5<sup>th</sup>  
16 or top 1<sup>st</sup> percentiles of daily precipitation can also be attributed to human influence at the global scale (Dong  
17 et al., 2020). CMIP5 models were able to capture the fraction of land experiencing a strong intensification of  
18 heavy precipitation during 1960-2010 under anthropogenic forcing, but not in unforced simulations (Fischer  
19 et al., 2014)). But the models underestimated the observed trends (Borodina et al., 2017a). Human influence  
20 also significantly contributed to the historical changes in record-breaking one-day precipitation (Shiogama et  
21 al., 2016). There is also limited evidence of the influences of natural forcing. Substantial reductions in  
22 Rx5day and SDII (daily precipitation intensity) over the global summer monsoon regions occurred during  
23 1957-2000 after explosive volcanic eruptions (Paik and Min, 2018). The reduction in post-volcanic eruption  
24 extreme precipitation in the simulations is closely linked to the decrease in mean precipitation, for which  
25 both thermodynamic effects (moisture reduction due to surface cooling) and dynamic effects (monsoon  
26 circulation weakening) play important roles.  
27

28 There has been new evidence of human influence on extreme precipitation at continental scales, including  
29 the detection of the combined effect of greenhouse gases and aerosol forcing on Rx1day and Rx5day over  
30 North America, Eurasia, and mid-latitude land regions (Zhang et al., 2013) and of greenhouse gas forcing in  
31 Rx1day and Rx5day in the mid-to-high latitudes, western and eastern Eurasia, and the global dry regions  
32 (Paik et al., 2020). These findings are corroborated by the detection of human influence in the fraction of  
33 extreme precipitation in the total precipitation over Asia, Europe, and North America (Dong et al., 2020).  
34 Human influence was found to have contributed to the increase in frequency and intensity of regional  
35 precipitation extremes in North America during 1961-2010, based on both optimal fingerprinting and event  
36 attribution approaches (Kirchmeier-Young and Zhang, 2020). Tabari et al. (2020) found the observed  
37 latitudinal increase in extreme precipitation over Europe to be consistent with model-simulated responses to  
38 anthropogenic forcing.  
39

40 Evidence of human influence on extreme precipitation at regional scales is more limited and less robust. In  
41 northwest Australia, the increase in extreme rainfall since 1950 can be related to increased monsoonal flow  
42 due to increased aerosol emissions, but cannot be attributed to an increase in greenhouse gases (Dey et al.,  
43 2019a). Anthropogenic influence on extreme precipitation in China was detected in one study (Li et al.,  
44 2017), but it was not detected in another study (Li et al., 2018e) using different detection and data-processing  
45 procedures, indicating the lack of robustness in the detection results. A still weak signal-to-noise ratio seems  
46 to be the main cause for the lack of robustness, as detection would become robust 20 years in the future (Li  
47 et al., 2018e). Krishnan et al. (2016) attributed the observed increase in heavy rain events (intensity > 100  
48 mm/day) in the post-1950s over central India to the combined effects of greenhouse gases, aerosols, land use  
49 and land cover changes, and rapid warming of the equatorial Indian Ocean SSTs. Roxy et al. (2017) and  
50 Devanand et al. (2019) showed the increase in widespread extremes over the South Asian Monsoon during  
51 1950-2015 is due to the combined impacts of the warming of the Western Indian Ocean (Arabian Sea) and  
52 the intensification of irrigation water management over India  
53

54 Anthropogenic influence may have affected the large-scale meteorological processes necessary for extreme  
55 precipitation and the localized thermodynamic and dynamic processes, both contributing to changes in

1 extreme precipitation events. Several new methods have been proposed to disentangle these effects by either  
2 conditioning on the circulation state or attributing analogues. In particular, the extremely wet winter of  
3 2013/2014 in the UK can be attributed, approximately to the same degree, to both temperature-induced  
4 increases in saturation vapour pressure and changes in the large-scale circulation (Vautard et al., 2016; Yiou  
5 et al., 2017). There are multiple cases indicating that very extreme precipitation may increase at a rate more  
6 than the C-C rate (6-7%/ °C) (Pall et al., 2017; Risser and Wehner, 2017; van der Wiel et al., 2017; van  
7 Oldenborgh et al., 2017; Wang et al., 2018).

8  
9 Event attribution studies found an influence of anthropogenic activities on the probability or magnitude of  
10 observed extreme precipitation events, including European winters (Schaller et al., 2016; Otto et al., 2018b),  
11 extreme 2014 precipitation over the northern Mediterranean (Vautard et al., 2015), parts of the US for  
12 individual events (Knutson et al., 2014b; Szeto et al., 2015; Eden et al., 2016; van Oldenborgh et al., 2017),  
13 extreme rainfall in 2014 over Northland, New Zealand (Rosier et al., 2016) or China (Burke et al., 2016; Sun  
14 and Miao, 2018; Yuan et al., 2018b; Zhou et al., 2018). For other heavy rainfall events, however, studies  
15 identified a lack of evidence about anthropogenic influences (Imada et al., 2013; Schaller et al., 2014; Otto et  
16 al., 2015c; Siswanto et al., 2015). There are also studies whose results are inconclusive because of limited  
17 reliable simulations (Christidis et al., 2013b; Angélil et al., 2016). Overall, both the spatial and temporal  
18 scales on which extreme precipitation events are defined are important for attribution; events defined on  
19 larger scales have larger signal-to-noise ratios and thus the signal is more readily detectable. At the current  
20 level of global warming, there is a strong enough signal to be detectable for large-scale extreme precipitation  
21 events, but the chance to detect such signals for smaller-scale events becomes smaller (Kirchmeier-Young et  
22 al., 2019).

23  
24 In summary, most of the observed intensification of heavy precipitation over land regions is *likely* due to  
25 anthropogenic influence, for which greenhouse gases emissions are the main contributor. New and robust  
26 evidence since AR5 includes attribution of the observed increase in annual maximum one-day and five-day  
27 precipitation and in the fraction of annual precipitation due to heavy events to human influence. It also  
28 includes a larger fraction of land showing enhanced extreme precipitation and a larger probability of record-  
29 breaking one-day precipitation than expected by chance, both of which can only be explained when  
30 anthropogenic greenhouse gas forcing is considered. Human influence has contributed to the intensification  
31 of heavy precipitation in three continents where observational data are more abundant, including North  
32 America, Europe and Asia (*high confidence*). On the spatial scale of AR6 regions, *evidence* of human  
33 influence on extreme precipitation is *limited*, but new evidence is emerging; in particular, studies attributing  
34 individual heavy precipitation events found that human influence was a significant driver of the events,  
35 particularly in the winter season.

#### 36 37 38 **11.4.5 Projections**

39  
40 AR5 concluded it is *very likely* that extreme precipitation events will be more frequent and more intense over  
41 most of the mid-latitude land masses and wet tropics in a warmer world (Collins et al., 2013). Post-AR5  
42 studies provide more and *robust evidence* to support the previous assessments. These include an observed  
43 increase in extreme precipitation (11.4.3) and human causes of past changes (11.4.4), as well as projections  
44 based on either GCM and/or RCM simulations. CMIP5 models project the rate of increase in Rx1day with  
45 warming is independent of the forcing scenario (Pendergrass et al., 2015, Chapter 8, Section 8.5.3.1) or  
46 forcing mechanism (Sillmann et al., 2017). This is confirmed in CMIP6 simulations (Li et al., 2020, and  
47 Sillmann et al., 2019). In particular, for extreme precipitation that occurs once a year or less frequently, the  
48 magnitudes of the rates of change per 1°C change in global mean temperature are similar regardless of  
49 whether the temperature change is caused by increases in CO<sub>2</sub>, CH<sub>4</sub>, solar forcing, or SO<sub>4</sub> (Sillmann et al.,  
50 2019). In some models, CESM1 in particular, the extreme precipitation response to warming may follow a  
51 quadratic relation (Pendergrass et al., 2019). Figure 11.15 shows changes in the 10-year and 50-year return  
52 values of Rx1day at different warming levels as simulated by the CMIP6 models. The median value of the  
53 scaling over land, across all SSP scenarios and all models, is close to 7%/°C for the 50-year return value of  
54 Rx1day. It is just slightly smaller for the 10-year and 50-year return values of Rx5day (Li et al., 2020a). The  
55 90% ranges of the multi-model ensemble changes across all land grid boxes in the 50-yr return values for



Rx1day and Rx5day do not overlap between 1.5°C and 2°C warming levels (Li et al., 2020), indicating that a small increment such as 0.5°C in global warming can result in a significant increase in extreme precipitation. Projected long-period Rx1day return value changes are larger than changes in mean Rx1day and increase with increasing rarity (Pendergrass, 2018; Mizuta and Endo, 2020; Wehner, 2020). The rate of change of moderate extreme precipitation may depend more on the forcing agent, similar to the mean precipitation response to warming (Lin et al., 2016, 2018a). Thus, there is *high confidence* that extreme precipitation that occurs once a year or less frequently increases proportionally to the amount of surface warming and the rate of change in precipitation is not dependent on the underlying forcing agents of warming.

[START FIGURE 11.15 HERE]

**Figure 11.15:** Projected changes in the intensity of extreme precipitation events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1851-1900 baseline. Extreme precipitation events are defined as the daily precipitation (Rx1day) that was exceeded on average once during a 10-year period (10-year event, blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the intensity changes across the space, and the whiskers extend to the 90% uncertainty range. The results are based on the multi-model ensemble median estimated from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Based on (Li et al., 2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

[END FIGURE 11.15 HERE]

The spatial patterns of the projected changes across different warming levels are quite similar, as shown in Figure 11.16 and confirmed by near-linear scaling between extreme precipitation and global warming levels at regional scales (Seneviratne and Hauser, 2020). Internal variability modulates changes in heavy rainfall (Wood and Ludwig, 2020), resulting in different changes in different regions (Seneviratne and Hauser, 2020). Extreme precipitation nearly always increases across land areas with larger increases at higher global warming levels, except in very few regions, such as southern Europe around the Mediterranean Basin in some seasons. The *very likely* ranges of the multi-model ensemble changes across all land grid boxes in the 50-yr return values for Rx1day and Rx5day between 1.5°C and 1°C warming levels are above zero for all continents except Europe, with *likely* range above zero over Europe (Li et al., 2020). Decreases in extreme precipitation are confined mostly to subtropical ocean areas and are highly correlated to decreases in mean precipitation due to storm track shifts. These subtropical decreases can extend to nearby land areas in individual realizations.

Projected increases in the probability of extreme precipitation of fixed magnitudes are non-linear and show larger increases for more rare events (Figures 11.7 and 11.15, Fischer and Knutti, 2015, Li et al., 2020, Kharin et al., 2018). CMIP5-model-projected increases in the probability of high (99<sup>th</sup> and 99.9<sup>th</sup>) percentile precipitation between 1.5°C and 2°C warming scenarios are consistent with what can be expected based on observed changes (Fischer and Knutti, 2015), providing confidence in the projections. CMIP5 model simulations show that the frequency for present-day climate 20-year extreme precipitation is projected to increase by 10% at the 1.5°C global warming level and by 22% at the 2.0°C global warming level, while the increase in the frequency for present-day climate 100-year extreme precipitation is projected to increase by 20% and more than 45% at the 1.5°C and 2.0°C warming levels, respectively (Kharin et al., 2018). CMIP6 simulations with SSP scenarios show the frequency of 10-year and 50-year events will be approximately doubled and tripled, respectively, at a very high warming level of 4°C (Figure 11.7, Li et al., 2020).

The number of studies on the projections of extreme hourly precipitation are limited. The ability of GCMs to simulate hourly precipitation extremes is limited (Morrison et al., 2019) and very few modelling centres archive sub-daily and hourly precipitation prior to CMIP6 experiments. RCM simulations project an increase in extreme sub-daily precipitation in North America (Li et al., 2019a) and over Sweden (Olsson and Foster,

1 2013), but these models still do not explicitly resolve convective processes that are important for properly  
2 simulating extreme sub-daily precipitation. Simulations by RCMs that explicitly resolve convective  
3 processes (convection-permitting models) are limited in length and only available in a few regions because  
4 of high computing costs. Yet, a majority of the available convection-permitting simulations project increases  
5 in the intensities of extreme sub-daily precipitation events with the amount similar to or higher than the C-C  
6 scaling rate (Ban et al., 2015; Helsen et al., 2020; Kendon et al., 2014, 2019; Prein et al., 2016b; Fowler et  
7 al., 2020). An increase is projected in extreme sub-daily precipitation over Africa (Kendon et al., 2019); over  
8 East Africa (Finney et al., 2020) and West Africa (Berthou et al., 2019a; Fitzpatrick et al., 2020), even for  
9 areas where parameterized RCMs project a decrease; in Europe (Chan et al., 2020 and Hodnebrog et al.,  
10 2019); as well as in the continental US (Prein et al., 2016). Overall, available evidence, while limited, points  
11 to an increase in extreme sub-daily precipitation in the future. Studies on future changes in extreme  
12 precipitation for a month or longer are limited. One study projects an increase in extreme monthly  
13 precipitation in Japan under 4°C global warming for around 80% of stations in the summer (Hatsuzuka and  
14 Sato, 2019).

15  
16 In Africa (Table 11.5), extreme precipitation will *likely* increase under warming levels of 2°C or below  
17 (compared to pre-industrial values) and *very likely* increase at higher warming levels. Simulations by  
18 CMIP5, CMIP6 and CORDEX regional models project an increase in daily extreme precipitation between  
19 1.5°C and 2.0°C warming levels. The pattern of change in heavy precipitation under different scenarios or  
20 warming levels is similar with larger increases for higher warming levels (e.g., Nikulin et al., 2018; Li et al.,  
21 2020). With increases in warming, extreme precipitation is projected to increase in the majority of land  
22 regions in Africa (Mtongori et al., 2016; Pfahl et al., 2017; Diedhiou et al., 2018; Dunning et al., 2018;  
23 Akinyemi and Abiodun, 2019; Giorgi et al., 2019). Over southern Africa, heavy precipitation will *likely*  
24 increase by the end of the 21<sup>st</sup> century under RCP 8.5 (Dosio, 2016; Pinto et al., 2016; Abiodun et al., 2017;  
25 Dosio et al., 2019). However, heavy rainfall amounts are projected to decrease over western South Africa  
26 (Pinto et al., 2018) as a result of a projected decrease in the frequency of the prevailing westerly winds south  
27 of the continent that translates into fewer cold fronts and closed mid-latitudes cyclones (Engelbrecht et al.,  
28 2009; Pinto et al., 2018). Heavy precipitation will *likely* increase by the end of the century under RCP8.5 in  
29 West Africa (Diallo et al., 2016; Dosio, 2016; Sylla et al., 2016; Abiodun et al., 2017; Akinsanola and Zhou,  
30 2018; Dosio et al., 2019) and is projected to increase (*medium confidence*) in central Africa (Fotso-Nguemo  
31 et al., 2018, 2019; Sonkoué et al., 2019) and eastern Africa (Thierry et al., 2016; Ongoma et al., 2018a). In  
32 northeast and central east Africa, extreme precipitation intensity is projected to increase across CMIP5,  
33 CMIP6 and CORDEX-CORE (*high confidence*) in most areas annually (Coppola et al., 2021a), but the  
34 trends differ from season to season in all future scenarios (Dosio et al., 2019). In northern Africa, there is *low*  
35 *confidence* in the projected changes in heavy precipitation, either due to a lack of agreement among studies  
36 on the sign of changes (Sillmann et al., 2013a; Giorgi et al., 2014) or due to insufficient evidence.

37  
38 In Asia (Table 11.8), extreme precipitation will *likely* increase at global warming levels of 2°C and below,  
39 but *very likely* increase at higher warming levels for the region as whole. The CMIP6 multi-model median  
40 projects an increase in the 10- and 50-yr return values of Rx1day and Rx5day over more than 95% of  
41 regions, even at the 2°C warming level, with larger increases at higher warming levels, independent of  
42 emission scenarios (Li et al., 2020, also Figure 11.7). CMIP5 models produced similar projections. Both  
43 heavy rainfall and rainfall intensity are projected to increase (Endo et al., 2017; Guo et al., 2016, 2018; Han  
44 et al., 2018; Kim et al., 2018; Xu et al., 2016; Zhou et al., 2014). A half-degree difference in warming  
45 between the 1.5°C and 2.0°C warming levels can result in a detectable increase in extreme precipitation over  
46 the region (Li et al., 2020), in the Asian-Australian monsoon region (Chevuturi et al., 2018), and over South  
47 Asia and China (Lee et al., 2018b; Li et al., 2018f). While there are regional differences, extreme  
48 precipitation is projected to increase in almost all sub-regions, though there can be spatial heterogeneity  
49 within sub-regions, such as in India (Shashikanth et al., 2018) and Southeast Asia (Ohba and Sugimoto,  
50 2019). In East and Southeast Asia, there is *high confidence* that extreme precipitation is projected to intensify  
51 (Guo et al., 2018; Li et al., 2018a; Seo et al., 2014; Sui et al., 2018; Wang et al., 2017b, 2017c; Xu et al.,  
52 2016; Zhou et al., 2014, Nayak et al., 2017; Mandapaka and Lo, 2018; Raghavan et al., 2018; Tangang et al.,  
53 2018; Supari et al., 2020). Extreme daily precipitation is also projected to increase in South Asia  
54 (Shashikanth et al., 2018; Han et al., 2018; Xu et al., 2017). The extreme precipitation indices, including  
55 Rx5day, R95p, and days of heavy precipitation (i.e., R10mm), are all projected to increase under the RCP4.5

1 and RCP8.5 scenarios in central and northern Asia (Xu et al., 2017; Han et al., 2018). A general wetting  
2 across the whole Tibetan Plateau and the Himalaya is projected, with increases in heavy precipitation in the  
3 21st century (Zhou et al., 2014; Zhang et al., 2015c; Gao et al., 2018; Palazzi et al., 2013; Rajbhandari et al.,  
4 2015; Wu et al., 2017; Paltan et al., 2018). Agreement in projected changes by different models is low in  
5 regions of complex topography such as Hindu-Kush-Himalaya (Wester et al., 2019), but CMIP5, CMIP6 and  
6 CORDEX-CORE simulations consistently project an increase in heavy precipitation in higher latitude areas  
7 (WSB, ESB, RFE) (Coppola et al., 2021a) (*high confidence*).

8  
9 In Australasia (Table 11.11), most CMIP5 models project an increase in Rx1day under RCP4.5 and RCP8.5  
10 scenarios for the late 21<sup>st</sup> century (CSIRO and BOM, 2015; Alexander and Arblaster, 2017; Grose et al.,  
11 2020) and the CMIP6 multi-model median projects an increase in the 10- and 50-yr return values of Rx1day  
12 and Rx5day at a rate between 5-6% per degree celsius of near-surface global mean warming (Li et al., 2020,  
13 also Figure 11.7). Yet, there is large uncertainty in the increase because projected changes in dynamic  
14 processes lead to a decrease in Rx1day that can offset the thermodynamic increase over a large portion of  
15 the region (Pfahl et al., 2017, see also Box 11.1 Figure 1). Projected changes in moderate extreme  
16 precipitation (the 99<sup>th</sup> percentile of daily precipitation) by RCMs under RCP8.5 for 2070-2099 are mixed,  
17 with more regions showing decreases than increases (Evans et al., 2020). It is *likely* that daily rainfall  
18 extremes such as Rx1day will increase at the continental scale for global warming levels at or above 3°C,  
19 daily rainfall extremes are projected to increase at the 2.0°C global warming level (*medium confidence*), and  
20 there is *low confidence* in changes at the 1.5°C. Projected changes show important regional differences with  
21 *very likely* increases over NAU (Alexander and Arblaster, 2017; Herold et al., 2018; Grose et al., 2020) and  
22 NZ (MfE, 2018) where projected dynamic contributions are small (Pfahl et al., 2017), see also Box 11.1  
23 Figure 1) and *medium confidence* on increases over central, eastern, and southern Australia where dynamic  
24 contributions are substantial and can affect local phenomena (CSIRO and BOM, 2015; Pepler et al., 2016;  
25 Bell et al., 2019; Dowdy et al., 2019).

26  
27 In Central and South America (Table 11.14), extreme precipitation will *likely* increase at global warming  
28 levels of 2°C and below, but *very likely* increase at higher warming levels for the region as whole. A larger  
29 increase in global surface temperature leads to a larger increase in extreme precipitation, independent of  
30 emission scenarios (Li et al., 2020a). But there are regional differences in the projection and projected  
31 changes for more moderate extreme precipitation are also more uncertain. Extreme precipitation, represented  
32 by the R<sub>50mm</sub> and R<sub>90p</sub> extreme indices, is projected to increase on the eastern coast of SCA, but to decrease  
33 along the Pacific coasts of El Salvador and Guatemala (Imbach et al., 2018). Chou et al. (2014) and Giorgi et  
34 al. (2014) projected an increase in extreme precipitation over southeastern South America and the Amazon.  
35 Projected changes in moderate extreme precipitation represented by the 99<sup>th</sup> percentile of daily precipitation  
36 by different models under different emission scenarios, even at high warming levels, are mixed, with  
37 increases projected for all regions by the CORDEX-CORE and CMIP5 simulations, but increases for some  
38 regions and decreases for other regions by CMIP6 simulations (Coppola et al., 2021a). Extreme precipitation  
39 is projected to increase in the La Plata basin (Cavalcanti et al., 2015; Carril et al., 2016). Taylor et al. (2018)  
40 projected a decrease in days with intense rainfall in the Caribbean under 2°C global warming by the 2050s  
41 under RCP4.5 relative to 1971-2000.

42  
43 In Europe (Table 11.17), extreme precipitation will *likely* increase at global warming levels of 2°C and  
44 below, but *very likely* increase for higher warming levels for the region as whole. The CMIP6 multi-model  
45 median projects an increase in the 10- and 50-yr return values of Rx1day and Rx5day over a majority of the  
46 region at the 2°C global warming level, with more than 95% of the region showing an increase at higher  
47 warming levels (Li et al., 2020, also Figure 11.7). The most intense precipitation events observed today in  
48 Europe are projected to almost double in occurrence for each degree celsius of further global warming  
49 (Myhre et al., 2019). Extreme precipitation is projected to increase in both boreal winter and summer over  
50 Europe (Madsen et al., 2014; OB et al., 2015; Nissen and Ulbrich, 2017). There are regional differences,  
51 with decreases or no change for the southern part of Europe, such as the southern Mediterranean (Lionello  
52 and Scarascia, 2020; Trambly and Somot, 2018; Coppola et al., 2020), uncertain changes over central  
53 Europe (Argüeso et al., 2012; Croitoru et al., 2013; Rajczak et al., 2013; Casanueva et al., 2014; Patarčić et  
54 al., 2014; Paxian et al., 2014; Roth et al., 2014; Fischer and Knutti, 2015; Monjo et al., 2016) and a strong  
55 increase in the remaining parts, including the Alps region (Gobiet et al., 2014; Donnelly et al., 2017),

1 particularly in winter (Fischer et al., 2015), and northern Europe. In a 3°C warmer world, there will be a  
2 robust increase in extreme rainfall over 80% of land areas in northern Europe (Madsen et al., 2014; Donnelly  
3 et al., 2017; Cardell et al., 2020).

4  
5 In North America (Table 11.20), the intensity and frequency of extreme precipitation will *likely* increase at  
6 the global warming levels of 2°C and below and *very likely* increase at higher warming levels. An increase is  
7 projected by CMIP6 model simulations (Li et al., 2020) and by previous model generations (Easterling et al.,  
8 2017; Wu, 2015; Zhang et al. 2018f; Innocenti et al., 2019b), as well as by RCMs (Coppola et al., 2020).  
9 Projections of extreme precipitation over the southern portion of the continent and over Mexico in particular  
10 are more uncertain, with decreases possible (Alexandru, 2018; Sillmann et al., 2013b; Coppola et al., 2020).

### 11 12 13 [START FIGURE 11.16 HERE]

14  
15 **Figure 11.16:** Projected changes in annual maximum daily precipitation at (a) 1.5°C, (b) 2°C, and (c) 4°C of global  
16 warming compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-  
17 model ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The  
18 numbers on the top right indicate the number of simulations included. Uncertainty is represented using  
19 the simple approach: no overlay indicates regions with high model agreement, where ≥80% of models  
20 agree on sign of change; diagonal lines indicate regions with low model agreement, where <80% of  
21 models agree on sign of change. For more information on the simple approach, please refer to the Cross-  
22 Chapter Box Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on  
23 data sources and processing are available in the chapter data table (Table 11.SM.9).

### 24 25 26 [END FIGURE 11.16 HERE]

27  
28  
29 In summary, heavy precipitation will generally become more frequent and more intense with additional  
30 global warming. At global warming levels of 4°C relative to the pre-industrial, very rare (e.g., 1 in 10 or  
31 more years) heavy precipitation events would become more frequent and more intense than in the recent  
32 past, on the global scale (*virtually certain*), and in all continents and AR6 regions: The increase in frequency  
33 and intensity is *extremely likely* for most continents and *very likely* for most AR6 regions. The likelihood is  
34 lower at lower global warming levels and for less-rare heavy precipitation events. At the global scale, the  
35 intensification of heavy precipitation will follow the rate of increase in the maximum amount of moisture  
36 that the atmosphere can hold as it warms (*high confidence*), of about 7% per °C of global warming. The  
37 increase in the frequency of heavy precipitation events will accelerate with more warming and will be higher  
38 for rarer events (*high confidence*), with 10-year and 50-year events to be approximately double and triple,  
39 respectively, at the 4°C warming level. Increases in the intensity of extreme precipitation events at regional  
40 scales will depend on the amount of regional warming as well as changes in atmospheric circulation and  
41 storm dynamics leading to regional differences in the rate of heavy precipitation changes (*high confidence*).

## 42 43 44 11.5 Floods

45  
46 Floods are the inundation of normally dry land and are classified into types (e.g., pluvial floods, flash floods,  
47 river floods, groundwater floods, surge floods, coastal floods) depending on the space and time scales and  
48 the major factors and processes involved (Chapter 8, Section 8.2.3.2, Nied et al., 2014; Aerts et al., 2018).  
49 Flooded area is difficult to measure or quantify and, for this reason, many of the existing studies on changes  
50 in floods focus on streamflow. Thus, this section assesses changes in flow as a proxy for river floods, in  
51 addition to some types of flash floods. Pluvial and urban floods, types of flash floods resulting from the  
52 precipitation intensity exceeding the capacity of natural and artificial drainage systems, are directly linked to  
53 extreme precipitation. Because of this link, changes in extreme precipitation are the main proxy for inferring  
54 changes in pluvial and urban floods (see also Section 12.4, REF Chapter 12), assuming there is no additional  
55 change in the surface condition. Changes in these types of floods are not assessed in this section, but can be  
56 inferred from the assessment of changes in heavy precipitation in Section 11.4. Coastal floods due to extreme  
57 sea levels and flood changes at regional scales are assessed in Chapter 12 (12.4).

### 11.5.1 Mechanisms and drivers

Since AR5, the number of studies on understanding how floods may have changed and will change in the future has substantially increased. Floods are a complex interplay of hydrology, climate, and human management, and the relative importance of these factors is different for different flood types and regions.

In addition to the amount and intensity of precipitation, the main factors for river floods include antecedent soil moisture (Paschalis et al., 2014; Berghuijs et al., 2016; Grillakis et al., 2016; Woldemeskel and Sharma, 2016) and snow water-equivalent in cold regions (Sikorska et al., 2015; Berghuijs et al., 2016). Other factors are also important, including stream morphology (Borga et al., 2014; Slater et al., 2015), river and catchment engineering (Pisaniello et al., 2012; Nakayama and Shankman, 2013; Kim and Sanders, 2016), land-use and land-cover characteristics (Aich et al., 2016; Rogger et al., 2017) and changes (Knighton et al., 2019), and feedbacks between climate, soil, snow, vegetation, etc. (Hall et al., 2014; Ortega et al., 2014; Berghuijs et al., 2016; Buttle et al., 2016; Teufel et al., 2019). Water regulation and management have, in general, increased resilience to flooding (Formetta and Feyen, 2019), masking effects of an increase in extreme precipitation on flood probability in some regions, even though they do not eliminate very extreme floods (Vicente-Serrano et al., 2017). This means that an increase in precipitation extremes may not always result in an increase in river floods (Sharma et al., 2018; Do et al., 2020). Yet, as very extreme precipitation can become a dominant factor for river floods, there can then be some correspondence in the changes in very extreme precipitation and river floods (Ivancic and Shaw, 2015; Wasko and Sharma, 2017; Wasko and Nathan, 2019). This has been observed in the western Mediterranean (Llasat et al., 2016), in China (Zhang et al., 2015a) and in the US (Peterson et al., 2013a; Berghuijs et al., 2016; Slater and Villarini, 2016).

In regions with a seasonal snow cover, snowmelt is the main cause of extreme river flooding over large areas (Pall et al., 2019). Extensive snowmelt combined with heavy and/or long-duration precipitation can cause significant floods (Li et al., 2019b; Krug et al., 2020). Changes in floods in these regions can be uncertain because of the compounding and competing effects of the responses of snow and rain to warming that affect snowpack size: warming results in an increase in precipitation, but also a reduction in the time period of snowfall accumulation (Teufel et al., 2019). An increase in atmospheric CO<sub>2</sub> enhances water-use efficiency by plants (Roderick et al., 2015; Milly and Dunne, 2016; Swann et al., 2016; Swann, 2018); this could reduce evapotranspiration and contribute to the maintenance of soil moisture and streamflow levels under enhanced atmospheric CO<sub>2</sub> concentrations (Yang et al., 2019). This mechanism would suggest an increase in the magnitude of some floods in the future (Kooperman et al., 2018). But this effect is uncertain as an increase in leaf area index and vegetation coverage could also result in overall larger water consumption (Mátyás and Sun, 2014; Mankin et al., 2019; Teuling et al., 2019), and there are also other CO<sub>2</sub>-related mechanisms that come into play (Chapter 5, CC Box 5.1).

Various factors, such as extreme precipitation (Cho et al., 2016; Archer and Fowler, 2018), glacier lake outbursts (Schneider et al., 2014; Schwanghart et al., 2016), or dam breaks (Biscarini et al., 2016) can cause flash floods. Very intense rainfall, along with a high fraction of impervious surfaces can result in flash floods in urban areas (Hettiarachchi et al., 2018). Because of this direct connection, changes in very intense precipitation can translate to changes in urban flood potential (Rosenzweig et al., 2018), though there can be a spectrum of urban flood responses to this flood potential (Smith et al., 2013), as many factors such as the overland flow rate and the design of urban (Falconer et al., 2009) and storm water drainage systems (Maksimović et al., 2009) can play an important role. Nevertheless, changes in extreme precipitation are the main proxy for inferring changes in some types of flash floods, which are addressed in Chapter 12 (Section 12.4)), given the relation between extreme precipitation and pluvial floods, the very limited literature on urban and pluvial floods (e.g., Skougaard Kaspersen et al., 2017), and limitations of existing methodologies for assessing changes in floods (Archer et al., 2016).

In summary, there is not always a one-to-one correspondence between an extreme precipitation event and a flood event, or between changes in extreme precipitation and changes in floods, because floods are affected by many factors in addition to heavy precipitation (*high confidence*). Changes in extreme precipitation may

1 be used as a proxy to infer changes in some types of flash floods that are more directly related to extreme  
2 precipitation (*high confidence*).  
3  
4

### 5 **11.5.2 Observed trends**

6

7 The SREX (Seneviratne et al., 2012) assessed *low confidence* for observed changes in the magnitude or  
8 frequency of floods at the global scale. This assessment was confirmed by the AR5 report (Hartmann et al.,  
9 2013). The SR15 (Hoegh-Guldberg et al., 2018) found increases in flood frequency and extreme streamflow  
10 in some regions, but decreases in other regions. While the number of studies on flood trends has increased  
11 since the AR5 report, and there were also new analyses after the release of SR15 (Berghuijs et al., 2017;  
12 Blöschl et al., 2019; Gudmundsson et al., 2019), hydrological literature on observed flood changes is  
13 heterogeneous, focusing at regional and sub-regional basin scales, making it difficult to synthesise at the  
14 global and sometimes regional scales. The vast majority of studies focus on river floods using streamflow as  
15 a proxy, with limited attention to urban floods. Streamflow measurements are not evenly distributed over  
16 space, with gaps in spatial coverage, and their coverage in many regions of Africa, South America, and parts  
17 of Asia is poor (e.g. Do et al., 2017), leading to difficulties in detecting long-term changes in floods (Slater  
18 and Villarini, 2017). See also Chapter 8, Section 8.3.1.5.  
19

20 Peak flow trends are characterized by high regional variability and lack overall statistical significance of a  
21 decrease or an increase over the globe as a whole. Of more than 3500 streamflow stations in the US, central  
22 and northern Europe, Africa, Brazil, and Australia, 7.1% stations showed a significant increase and 11.9%  
23 stations showed a significant decrease in annual maximum peak flow during 1961-2005 (Do et al., 2017).  
24 This is in direct contrast to the global and continental scale intensification of short-duration extreme  
25 precipitation (11.4.2). There may be some consistency over large regions (see Gudmundsson et al., 2019), in  
26 high streamflows (> 90<sup>th</sup> percentile), including a decrease in some regions (e.g., in the Mediterranean) and an  
27 increase in others (e.g., northern Asia), but gauge coverage is often limited. On a continental scale, a  
28 decrease seems to dominate in Africa (Tramblay et al., 2020) and Australia (Ishak et al., 2013; Wasko and  
29 Nathan, 2019), an increase in the Amazon (Barichivich et al., 2018), and trends are spatially variable in other  
30 continents (Do et al., 2017; Hodgkins et al., 2017; Bai et al., 2016; Zhang et al., 2015b). In Europe, flow  
31 trends have large spatial differences (Hall et al., 2014; Mediero et al., 2015; Kundzewicz et al., 2018;  
32 Mangini et al., 2018), but there appears to be a pattern of increase in northwestern Europe and a decrease in  
33 southern and eastern Europe in annual peak flow during 1960-2000 (Blöschl et al., 2019). In North America,  
34 peak flow has increased in the northeast US and decreased in the southwest US (Peterson et al., 2013a;  
35 Armstrong et al., 2014; Mallakpour and Villarini, 2015; Archfield et al., 2016; Burn and Whitfield, 2016;  
36 Wehner et al., 2017; Neri et al., 2019). There are important changes in the seasonality of peak flows in  
37 regions where snowmelt dominates, such as northern North America (Burn and Whitfield, 2016; Dudley et  
38 al., 2017) and northern Europe (Blöschl et al., 2017), corresponding to strong winter and spring warming.  
39

40 In summary, the seasonality of floods has changed in cold regions where snowmelt dominates the flow  
41 regime in response to warming (*high confidence*). *Confidence* about peak flow trends over past decades on  
42 the global scale is *low*, but there are regions experiencing increases, including parts of Asia, southern South  
43 America, the northeast USA, northwestern Europe, and the Amazon, and regions experiencing decreases,  
44 including parts of the Mediterranean, Australia, Africa, and the southwestern USA.  
45  
46

### 47 **11.5.3 Model evaluation**

48

49 Hydrological models used to simulate floods are structurally diverse (Dankers et al., 2014; Mateo et al.,  
50 2017; Şen, 2018), often requiring extensive calibration since sub-grid processes and land-surface properties  
51 need to be parameterized, irrespective of the spatial resolutions (Döll et al., 2016; Krysanova et al., 2017).  
52 The data that are used to drive and calibrate the models are usually of coarse resolution, necessitating the use  
53 of a wide variety of downscaling techniques (Muerth et al., 2013). This adds uncertainty not only to the  
54 models, but also to the reliability of the calibrations. The quality of the flood simulations also depends on the  
55 spatial scale, as flood processes are different for catchments of different sizes. It is more difficult to replicate

1 flood processes for large basins, as water management and water use are often more complex for these  
2 basins.

3  
4 Studies that use different regional hydrological models show large spread in flood simulations (Dankers et  
5 al., 2014; Roudier et al., 2016; Trigg et al., 2016; Krysanova et al., 2017). Regional models reproduce  
6 moderate and high flows (0.02 – 0.1 flow annual exceedance probabilities) reasonably well, but there are  
7 large biases for the most extreme flows (0 - 0.02 annual flow exceedance probability), independent of the  
8 climatic and physiographic characteristics of the basins (Huang et al., 2017)). Global-scale hydrological  
9 models have even more challenges, as they struggle to reproduce the magnitude of the flood hazard (Trigg et  
10 al., 2016). Additionally, the ensemble mean of multiple models does not perform better than individual  
11 models (Zaherpour et al., 2018).

12  
13 The use of hydrological models for assessing changes in floods, especially for future projections, adds  
14 another dimension of uncertainty on top of uncertainty in the driving climate projections, including emission  
15 scenarios, and uncertainty in the driving climate models (both RCMs and GCMs) (Arnell and Gosling, 2016;  
16 Hundercha et al., 2016; Krysanova et al., 2017). The differences in hydrological models (Roudier et al., 2016;  
17 Thober et al., 2018), as well as post-processing of climate model output for the hydrological models (Muerth  
18 et al., 2013; Maier et al., 2018), both add to uncertainty for flood projections.

19  
20 In summary, there is *medium confidence* that simulations for the most extreme flows by regional  
21 hydrological models can have large biases. Global-scale hydrological models still struggle with reproducing  
22 the magnitude of floods. Projections of future floods are hampered by these difficulties and cascading  
23 uncertainties, including uncertainties in emission scenarios and the climate models that generate inputs.

#### 24 25 26 **11.5.4 Attribution**

27  
28 There are very few studies focused on the attribution of long-term changes in floods, but there are studies on  
29 changes in flood events. Most of the studies focus on flash floods and urban floods, which are closely related  
30 to intense precipitation events (Hannaford, 2015). In other cases, event attribution focused on runoff using  
31 hydrological models, and examples include river basins in the UK (Schaller et al., 2016; Kay et al., 2018)  
32 (See Section 11.4.4), the Okavango river in Africa (Wolski et al., 2014), and the Brahmaputra in Bangladesh  
33 (Philip et al., 2019). Findings about anthropogenic influences vary between different regions and basins. For  
34 some flood events, the probability of high floods in the current climate is lower than in a climate without an  
35 anthropogenic influence (Wolski et al., 2014), while in other cases anthropogenic influence leads to more  
36 intense floods (Cho et al., 2016; Pall et al., 2017; van der Wiel et al., 2017; Philip et al., 2018a; Teufel et al.,  
37 2019). Factors such as land cover change and river management can also increase the probability of high  
38 floods (Ji et al., 2020). These, along with model uncertainties and the lack of studies overall, suggest a *low*  
39 *confidence* in general statements to attribute changes in flood events to anthropogenic climate change. Some  
40 individual regions have been well studied, which allows for *high confidence* in the attribution of increased  
41 flooding in these cases (Section 11.9 table). For example, flooding in the UK following increased winter  
42 precipitation (Schaller et al., 2016; Kay et al., 2018) can be attributed to anthropogenic climate change  
43 (Schaller et al., 2016; Vautard et al., 2016; Yiou et al., 2017; Otto et al., 2018b).

44  
45 Attributing changes in heavy precipitation to anthropogenic activities (Section 11.4.4) cannot be readily  
46 translated to attributing changes in floods to human activities, because precipitation is only one of the  
47 multiple factors, albeit an important one, that affect floods. For example, Teufel et al. (2017) showed that  
48 while human influence increased the odds of the flood-producing rainfall for the 2013 Alberta flood in  
49 Canada, it was not detected to have influenced the probability of the flood itself. Schaller et al. (2016)  
50 showed human influence on the increase in the probability of heavy precipitation translated linearly into an  
51 increase in the resulting river flow of the Thames in winter 2014, but its contribution to the inundation was  
52 inconclusive.

53  
54 Gudmundsson et al. (2021) compared the spatial pattern of the observed regional trends in high river flows  
55 (> 90<sup>th</sup> percentile) over 1971-2010 with those simulated by global hydrological models driven by outputs of

1 climate models under all historical forcing and with pre-industrial climate model simulations. They found  
2 complex spatial patterns of extreme river flow trends. They also found the observed spatial patterns of trends  
3 can be reproduced only if anthropogenic climate change is considered and that simulated effects of water and  
4 land management cannot reproduce the observed spatial pattern of trends. As there is only one study and  
5 multiple caveats, including relatively poor observational data coverage, there is *low confidence* about human  
6 influence on the changes in high river flows on the global scale.  
7

8 In summary there is *low confidence* in the human influence on the changes in high river flows on the global  
9 scale. *Confidence* is in general *low* in attributing changes in the probability or magnitude of flood events to  
10 human influence because of a limited number of studies and differences in the results of these studies, and  
11 large modelling uncertainties.  
12

### 13 **11.5.5 Future projections**

14 The SREX report (Chapter 3, Seneviratne et al., 2012) stressed the low availability of studies on flood  
15 projections under different emission scenarios and concluded there was *low confidence* in projections of  
16 flood events given the complexity of the mechanisms driving floods at the regional scale. The AR5 WGII  
17 report (Chapter 3, Jimenez Cisneros et al., 2014) assessed with *medium confidence* the pattern of future flood  
18 changes, including flood hazards increasing over about half of the globe (parts of southern and Southeast  
19 Asia, tropical Africa, northeast Eurasia, and South America) and flood hazards decreasing in other parts of  
20 the world, despite uncertainties in GCMs and their coupling to hydrological models. SR15 (Chapter 3,  
21 Hoegh-Guldberg et al., 2018) assessed with *medium confidence* that global warming of 2°C would lead to an  
22 expansion of the fraction of global area affected by flood hazards, compared to conditions at 1.5°C of global  
23 warming, as a consequence of changes in heavy precipitation.  
24

25 The majority of new studies that produce future flood projections based on hydrological models do not  
26 typically consider aspects that are also important to actual flood severity or damages, such as flood  
27 prevention measures (Neumann et al., 2015; Şen, 2018), flood control policies (Barraqué, 2017), and future  
28 changes in land cover (see also Chapter 8, Section 8.4.1.5). At the global scale, Alfieri et al. (2017a) used  
29 downscaled projections from seven GCMs as input to drive a hydrodynamic model. They found successive  
30 increases in the frequency of high floods in all continents except Europe, associated with increasing levels of  
31 global warming (1.5°C, 2°C, 4°C). These results are supported by Paltan et al. (2018), who applied a  
32 simplified runoff aggregation model forced by outputs from four GCMs. Huang et al. (2018b) used three  
33 hydrological models forced with bias-adjusted outputs from four GCMs to produce projections for four river  
34 basins including the Rhine, Upper Mississippi, Upper Yellow, and Upper Niger under 1.5°C, 2°C, and 3°C  
35 global warming. This study found diverse projections for different basins, including a shift towards earlier  
36 flooding for the Rhine and the Upper Mississippi, a substantial increase in flood frequency in the Rhine only  
37 under the 1.5°C and 2°C scenarios, and a decrease in flood frequency in the Upper Mississippi under all  
38 scenarios.  
39

40 At the continental and regional scales, the projected changes in floods are uneven in different parts of the  
41 world, but there is a larger fraction of regions with an increase than with a decrease over the 21st century  
42 (Hirabayashi et al., 2013; Dankers et al., 2014; Arnell and Gosling, 2016; Döll et al., 2018). These results  
43 suggest *medium confidence* in flood trends at the global scale, but *low confidence* in projected regional  
44 changes. Increases in flood frequency or magnitude are identified for southeastern and northern Asia and  
45 India (high agreement across studies), eastern and tropical Africa, and the high latitudes of North America  
46 (*medium agreement*), while decreasing frequency or magnitude is found for central and eastern Europe and  
47 the Mediterranean (*high confidence*), and parts of South America, southern and central North America, and  
48 southwest Africa (Hirabayashi et al., 2013; Dankers et al., 2014; Arnell and Gosling, 2016; Döll et al., 2018).  
49 Over South America, most studies based on global and regional hydrological models show an increase in the  
50 magnitude and frequency of high flows in the western Amazon (Sorribas et al., 2016; Langerwisch et al.,  
51 2013; Guimberteau et al., 2013; Zulkafli et al., 2016) and the Andes (Hirabayashi et al., 2013; Bozkurt et al.,  
52 2018). Chapter 12, Section 12.4, provides a detailed assessment of regional flood projections.  
53  
54  
55



1 In summary, global hydrological models project a larger fraction of land areas to be affected by an increase  
2 in river floods than by a decrease in river floods (*medium confidence*). There is *medium confidence* that river  
3 floods will increase in the western Amazon, the Andes, and southeastern and northern Asia. Regional  
4 changes in river floods are more uncertain than changes in pluvial floods because complex hydrological  
5 processes and forcings, including land cover change and human water management, are involved.  
6  
7

## 8 **11.6 Droughts**

9

10 Droughts refer to periods of time with substantially below-average moisture conditions, usually covering  
11 large areas, during which limitations in water availability result in negative impacts for various components  
12 of natural systems and economic sectors (Wilhite and Pulwarty, 2017; Ault, 2020). Depending on the  
13 variables used to characterize it and the systems or sectors being impacted, drought may be classified in  
14 different types (Figure 8.6; Table 11.A.1) such as **meteorological** (precipitation deficits), **agricultural** (e.g.,  
15 crop yield reductions or failure, often related to soil moisture deficits), **ecological** (related to plant water  
16 stress that causes e.g., tree mortality), or **hydrological** droughts (e.g., water shortage in streams or storages  
17 such as reservoirs, lakes, lagoons, and groundwater) (See Annex VII: Glossary). The distinction of drought  
18 types is not absolute as drought can affect different sub-domains of the Earth system concomitantly, but  
19 sometimes also asynchronously, including propagation from one drought type to another (Brunner and  
20 Tallaksen, 2019). Because of this, drought cannot be characterized using a single universal definition (Lloyd-  
21 Hughes, 2014) or directly measured based on a single variable (SREX Chapter 3; Wilhite and Pulwarty,  
22 2017). Drought can happen on a wide range of timescales - from "flash droughts" on a scale of weeks, and  
23 characterized by a sudden onset and rapid intensification of drought conditions (Hunt et al., 2014; Otkin et  
24 al., 2018; Pendergrass et al., 2020) to multi-year or decadal rainfall deficits (sometimes termed  
25 "megadroughts"; Annex VII: Glossary) (Ault et al., 2014; Cook et al., 2016b; Garreaud et al., 2017).  
26 Droughts are often analysed using indices that are measures of drought severity, duration and frequency  
27 (Table 11.A.1; Chapter 8, Sections 8.3.1.6, 8.4.1.6, Chapter 12, Sections 12.3.2.6 and 12.3.2.7). There are  
28 many drought indices published in the scientific literature, as also highlighted in the IPCC SREX report  
29 (SREX Chapter 3). These can range from anomalies in single variables (e.g., precipitation, soil moisture,  
30 runoff, evapotranspiration) to indices combining different atmospheric variables.  
31

32 This assessment is focused on changes in physical conditions and metrics of direct relevance to droughts  
33 (Table 11.A.1): a) precipitation deficits, b) excess of atmospheric evaporative demand (AED), c) soil  
34 moisture deficits, d) hydrological deficits, and e) atmospheric-based indices combining precipitation and  
35 AED. In the regional tables (Section 11.9), the assessment is structured by drought types, addressing i)  
36 meteorological, ii) agricultural and ecological, and iii) hydrological droughts. Note that the latter two  
37 assessments are directly informing the Chapter 12 assessment on projected regional changes in these climatic  
38 impact-drivers (Chapter 12, Section 12.4). The text refers to AR6 regions acronyms (Section 11.9, see  
39 Chapter 1, Section 1.4.5) when referring to changes in AR6 regions.  
40  
41

### 42 **11.6.1 Mechanisms and drivers**

43

44 Similar to many other extreme events, droughts occur as a combination of thermodynamic and dynamic  
45 processes (Box 11.1). Thermodynamic processes contributing to drought, which are modified by greenhouse  
46 gas forcing both at global and regional scales, are mostly related to heat and moisture exchanges and also  
47 partly modulated by plant coverage and physiology. They affect, for instance, atmospheric humidity,  
48 temperature, and radiation, which in turn affect precipitation and/or evapotranspiration in some regions and  
49 time frames. On the other hand, dynamic processes are particularly important to explain drought variability  
50 on different time scales, from a few weeks (flash droughts) to multiannual (megadroughts). There is *low*  
51 *confidence* in the effects of greenhouse gas forcing on changes in atmospheric dynamic (Chapter 2, Section  
52 2.4; Chapter 4, Section 4.3.3), and, hence, on associated changes in drought occurrence. Thermodynamic  
53 processes are thus the main driver of drought changes in a warming climate (*high confidence*).  
54  
55

### 11.6.1.1 Precipitation deficits

Lack of precipitation is generally the main factor controlling drought onset. There is *high confidence* that atmospheric dynamics, which varies on interannual, decadal and longer time scales, is the dominant contributor to variations in precipitation deficits in the majority of the world regions (Dai, 2013; Seager and Hoerling, 2014; Miralles et al., 2014b; Burgman and Jang, 2015; Dong and Dai, 2015; Schubert et al., 2016; Raymond et al., 2018; Baek et al., 2019; Drumond et al., 2019; Herrera-Estrada et al., 2019; Gimeno et al., 2020; Mishra, 2020). Precipitation deficits are driven by dynamic mechanisms taking place on different spatial scales, including synoptic processes –atmospheric rivers and extratropical cyclones, blocking and ridges (Section 11.7; Sousa et al., 2017), dominant large-scale circulation patterns (Kingston et al., 2015), and global ocean-atmosphere coupled patterns such as IPO, AMO and ENSO (Dai and Zhao, 2017). These various mechanisms occur on different scales, are not independent, and substantially interact with one another. Also regional moisture recycling and land-atmosphere feedbacks play an important role for some precipitation anomalies (see below).

There is *high confidence* that land-atmosphere feedbacks play a substantial or dominant role in affecting precipitation deficits in some regions (SREX, Chapter 3; Gimeno et al., 2012; Guillod et al., 2015; Haslinger et al., 2019; Herrera-Estrada et al., 2019; Koster et al., 2011; Santanello Jr. et al., 2018; Taylor et al., 2012; Tuttle and Salvucci, 2016). The sign of the feedbacks can be either positive or negative, as well as local or non-local (Taylor et al., 2012; Guillod et al., 2015; Tuttle and Salvucci, 2016). ESMs tend to underestimate non-local negative soil moisture-precipitation feedbacks (Taylor et al., 2012) and also show high variations in their representation in some regions (Berg et al., 2017a). Soil moisture-precipitation feedbacks contribute to changes in precipitation in climate model projections in some regions, but ESMs display substantial uncertainties in their representation, and there is thus only *low confidence* in these contributions (Berg et al., 2017a; Vogel et al., 2017, 2018).

### 11.6.1.2 Atmospheric evaporative demand

Atmospheric evaporative demand (AED) quantifies the maximum amount of actual evapotranspiration (ET) that can happen from land surfaces if they are not limited by water availability (Table 11.A.1). AED is affected by both radiative and aerodynamic components. For this reason, the atmospheric dryness, often quantified with the relative humidity or the vapor pressure deficit (VPD), is not equivalent to the AED, as other variables are also highly relevant, including solar radiation and wind speed (Hobbins et al., 2012; McVicar et al., 2012b; Sheffield et al., 2012). AED can be estimated using different methods (McMahon et al., 2013). Methods solely based on air temperature (e.g. Hargreaves, Thornthwaite) usually overestimate it in terms of magnitude and temporal trends (Sheffield et al., 2012), in particular in the context of substantial background warming. Physically-based combination methods such as the Penman-Monteith equation are more adequate and recommended since 1998 by the Food and Agriculture Organization (Pereira et al., 2015). For this reason, the assessment of this chapter, when considering atmospheric-based drought indices, only includes AED estimates using the latter (see also Section 11.9). AED is generally higher than ET, since it represents an upper bound for it. Hence, an AED increase does not necessarily lead to increased ET (Milly and Dunne, 2016), in particular under drought conditions given soil moisture limitation (Bonan et al., 2014; Berg et al., 2016; Konings et al., 2017; Stocker et al., 2018). In general, AED is highest in regions where ET is lowest (e.g., desert areas), further illustrating the decoupling between the two variables under limited soil moisture.

The influence of AED on drought depends on the drought type, background climate, the environmental conditions and the moisture availability (Hobbins et al., 2016, 2017; Vicente-Serrano et al., 2020b). This influence also includes effects not related to increased ET. Under low soil moisture conditions, increased AED increases plant stress, enhancing the severity of agricultural and ecological droughts (Williams et al., 2013; Allen et al., 2015; McDowell et al., 2016; Grossiord et al., 2020). Moreover, high VPD impacts overall plant physiology; it affects the leaf and xylem safety margins, and decreases the sap velocity and plant hydraulic conductance (Fontes et al., 2018). VPD also affects the plant metabolism of carbon and if prolonged, it may cause plant mortality via carbon starvation (Breshears et al., 2013; Hartmann, 2015).

1 Drought projections based exclusively on AED metrics overestimate changes in soil moisture and runoff  
2 deficits. Nevertheless, AED also directly impacts hydrological drought, as ET from surface waters is not  
3 limited (Wurbs and Ayala, 2014; Friedrich et al., 2018; Hogeboom et al., 2018; Xiao et al., 2018a), and this  
4 effect increases under climate change projections (Wang et al., 2018c; Althoff et al., 2020). In addition, high  
5 AED increases crop water consumptions in irrigated lands (García-Garizábal et al., 2014), contributing to  
6 intensifying hydrological droughts downstream (Fazel et al., 2017; Vicente-Serrano et al., 2017).

7  
8 On subseasonal to decadal scales, temporal variations in AED are strongly controlled by circulation  
9 variability (Williams et al., 2014; Chai et al., 2018; Martens et al., 2018), but thermodynamic processes also  
10 play a fundamental role and under human-induced climate change dominate the changes in AED.

11 Atmospheric warming due to increased atmospheric CO<sub>2</sub> concentrations increases AED by means of  
12 enhanced VPD in the absence of other influences (Scheff and Frierson, 2015). Indeed, because of the greater  
13 warming over land than over oceans (Chapter 2, Section 2.3.1.1; Section 11.3), the saturation pressure of  
14 water vapor increases more over land than over oceans; oceanic air masses advected over land thus contain  
15 insufficient water vapour to keep pace with the greater increase in saturation vapour pressure over land  
16 (Sherwood and Fu, 2014; Byrne and O’Gorman, 2018; Findell et al., 2019). Land-atmosphere feedbacks are  
17 also important in affecting atmospheric moisture content and temperature, with resulting effects on relative  
18 humidity and VPD (Berg et al., 2016; Haslinger et al., 2019; Zhou et al., 2019; Box 11.1).

### 19 20 21 *11.6.1.3 Soil moisture deficits*

22  
23 Soil moisture shows an important correlation with precipitation variability (Khong et al., 2015; Seager et al.,  
24 2019), but ET also plays a substantial role in further depleting moisture from soils, in particular in humid  
25 regions during periods of precipitation deficits (Padrón et al., 2020; Teuling et al., 2013). In addition, soil  
26 moisture plays a role in drought self-intensification under dry conditions in which ET is decreased and leads  
27 to higher AED (Miralles et al., 2019), an effect that can also contribute to trigger “flash droughts” (Otkin  
28 et al., 2016, 2018; DeAngelis et al., 2020; Pendergrass et al., 2020). If soil moisture becomes limited, ET is  
29 reduced, which on one hand may decrease the rate of soil drying, but on the other hand can lead to further  
30 atmospheric dryness through various feedback loops (Seneviratne et al., 2010; Miralles et al., 2014a, 2019;  
31 Teuling, 2018; Vogel et al., 2018; Zhou et al., 2019b; Liu et al., 2020). The process is complex since  
32 vegetation cover plays a role in modulating albedo and in providing access to deeper stores of water (both in  
33 the soil and groundwater), and changes in land cover and in plant phenology may alter ET (Sterling et al.,  
34 2013; Woodward et al., 2014; Frank et al., 2015; Döll et al., 2016; Ukkola et al., 2016; Trancoso et al., 2017;  
35 Hao et al., 2019; Lian et al., 2020). Snow depth has strong and direct impacts on soil moisture in many  
36 systems (Gergel et al., 2017; Williams et al., 2020).

37  
38 Soil moisture directly affects plant water stress and ET. Soil moisture is the primary factor that controls  
39 xylem hydraulic conductance, i.e. plant water uptake in plants (Sperry et al., 2016; Hayat et al., 2019; Chen  
40 et al., 2020d). For this reason, soil moisture deficits are the main driver of xylem embolism, the primary  
41 mechanism of plant mortality (Anderegg et al., 2012, 2016; Rowland et al., 2015). Also carbon assimilation  
42 by plants strongly depend on soil moisture (Hartzell et al., 2017), with implications for carbon starvation and  
43 plant dying if soil moisture deficits are prolonged (Sevanto et al., 2014). These mechanisms explain that  
44 soil moisture deficits are usually more relevant than AED excess to explain gross primary production  
45 anomalies and vegetation stress, mostly in sub-humid and semi-arid regions (Stocker et al. 2018; Liu et al.,  
46 2020b). CO<sub>2</sub> concentrations are shown to potentially decrease plant ET and increase plant water-use  
47 efficiency, affecting soil moisture levels, although this effect interacts with other CO<sub>2</sub> physiological and  
48 radiative effects (Section 11.6.5.2; Chapter 5, CC Box 5.1), and has less relevance under low soil moisture  
49 (Morgan et al., 2011; Xu et al., 2016b; Nackley et al., 2018; Dikšaitytė et al., 2019). ESMs represent both  
50 surface (ca. 10cm) and total column soil moisture, whereby total soil moisture is of more direct relevance for  
51 root water uptake, in particular by trees. There is evidence that surface soil moisture projections are  
52 substantially drier than total soil moisture projections, and may thus overestimate drying of relevance for  
53 most vegetation (Berg et al., 2017b).

#### 11.6.1.4 Hydrological deficits

Drivers of streamflow and surface water deficits are complex and strongly depend on the hydrological system analysed (e.g., streamflows in the headwaters, medium course of the rivers, groundwater, highly regulated hydrological basins). Soil hydrological processes, which control the propagation of meteorological droughts throughout different parts of the hydrological cycle (Van Loon and Van Lanen, 2012), are spatially and temporally complex (Herrera-Estrada et al., 2017; Huang et al., 2017c) and difficult to quantify (Van Lanen et al., 2016; Apurv et al., 2017; Caillouet et al., 2017; Konapala and Mishra, 2017; Hasan et al., 2019). The physiographic characteristics of the basins also affect how droughts propagate throughout the hydrological cycle (Van Loon and Van Lanen, 2012; Van Lanen et al., 2013; Van Loon, 2015; Konapala and Mishra, 2020; Valiya Veetil and Mishra, 2020). In addition, the assessment of groundwater deficits is very difficult given the complexity of processes that involve natural and human-driven feedbacks and interactions with the climate system (Taylor et al., 2013). Streamflow and surface water deficits are affected by land cover, groundwater and soil characteristics (Van Lanen et al., 2013; Van Loon and Laaha, 2015; Barker et al., 2016; Tjiedeman et al., 2018), as well as human activities (water management and demand, damming) and land use changes (He et al., 2017; Jehanzaib et al., 2020; Van Loon et al., 2016; Veldkamp et al., 2017; Wu et al., 2018; Xu et al., 2019b; Section 11.6.4.3). Finally, snow and glaciers are relevant for water resources in some regions. For instance, warming affects snowpack levels (Dierauer et al., 2019; Huning and AghaKouchak, 2020), as well as the timing of snow melt, thus potentially affecting the seasonality and magnitude of low flows (Barnhart et al., 2016).

#### 11.6.1.5 Atmospheric-based drought indices

Given difficulties of drought quantification and data constraints, atmospheric-based drought indices combining both precipitation and AED have been developed, as they can be derived from meteorological data that is available in most regions with few exceptions. These demand/supply indices are not intended to be metrics of soil moisture, streamflow or vegetation water stress. Because of their reliance on precipitation and AED, they are mostly related to the actual water balance in humid regions, in which ET is not limited by soil moisture and tends towards AED. In water-limited regions and in dry periods everywhere, they constitute an upper bound for overall water-balance deficits (e.g. of surface waters) but are also related to conditions conducive to vegetation stress, particularly under soil moisture limitation (Section 11.6.1.2).

Although there are many atmospheric-based drought indices, two are assessed in this chapter: the Palmer Drought Severity Index (PDSI) and the Standardized Precipitation Evapotranspiration Index (SPEI). The PDSI has been widely used to monitor and quantify drought severity (Dai et al., 2018), but is affected by some constraints (SREX Chapter 3; Mukherjee et al., 2018). Although the calculation of the PDSI is based on a soil water budget, the PDSI is essentially a climate drought index that mostly responds to the precipitation and the AED (van der Schrier et al., 2013; Vicente-Serrano et al., 2015; Dai et al., 2018). The SPEI also combines precipitation and AED, being equally sensitive to these two variables (Vicente-Serrano et al., 2015). The SPEI is more sensitive to AED than the PDSI (Cook et al., 2014a; Vicente-Serrano et al., 2015), although under humid and normal precipitation conditions, the effects of AED on the SPEI are small (Tomas-Burguera et al., 2020). Given the limitations associated with temperature-based AED estimates (Section 11.6.1.2), only studies using the Penman-Monteith-based SPEI and PDSI (hereafter SPEI-PM and PDSI-PM) are considered in this assessment and in the regional tables in Section 11.9.

#### 11.6.1.6 Relation of assessed variables and metrics for changes in different drought types

This chapter assesses changes in meteorological drought, agricultural and ecological droughts, and hydrological droughts. Precipitation-based indices are used for the estimation of changes in meteorological droughts, such as the Standardized Precipitation Index (SPI) and the Consecutive Dry Days (CDD). Changes in total soil moisture and soil moisture-based drought events are used for the estimation of changes in agricultural and ecological droughts, complemented by changes in surface soil moisture, water-balance estimates (precipitation minus ET), and SPEI-PM and PDSI-PM. For hydrological droughts, changes in low

1 flows are assessed, sometimes complemented by changes in mean streamflow.

2  
3 In summary, different drought types exist and they are associated with different impacts and respond  
4 differently to increasing greenhouse gas concentrations. Precipitation deficits and changes in  
5 evapotranspiration govern net water availability. A lack of sufficient soil moisture, sometimes amplified by  
6 increased atmospheric evaporative demand, result in agricultural and ecological drought. Lack of runoff and  
7 surface water result in hydrological drought. Drought events are both the result of dynamic and/or  
8 thermodynamic processes, with thermodynamic processes being the main driver of drought changes under  
9 human-induced climate change (*high confidence*).

## 10 11 12 **11.6.2 Observed trends**

13  
14 Evidence on observed drought trends at the time of the SREX (Chapter 3) and AR5 (Chapter 2) was limited.  
15 SREX concluded that “There is *medium confidence* that since the 1950s some regions of the world have  
16 experienced a trend to more intense and longer droughts, in particular in southern Europe and West Africa,  
17 but in some regions droughts have become less frequent, less intense, or shorter, for example, in central  
18 North America and northwestern Australia”. The assessment at the time did not distinguish between different  
19 drought types. This chapter includes numerous updates on observed drought trends, associated with  
20 extensive new literature and longer datasets since the AR5.

### 21 22 23 **11.6.2.1 Precipitation deficits**

24  
25 Strong precipitation deficits have been recorded in recent decades in the Amazon (2005, 2010), southwestern  
26 China (2009-2010), southwestern North America (2011-2014), Australia (1997-2009), California (2014), the  
27 middle East (2012-2016), Chile (2010-2015), the Great Horn of Africa (2011), among others (van Dijk et al.,  
28 2013; Mann and Gleick, 2015; Rowell et al., 2015; Marengo and Espinoza, 2016; Dai and Zhao, 2017;  
29 Garreaud et al., 2017, 2020; Marengo et al., 2017; Brito et al., 2018; Cook et al., 2018). Global studies  
30 generally show no significant trends in SPI time series (Orlowsky and Seneviratne, 2013; Spinoni et al.,  
31 2014), and in derived drought frequency and severity data (Spinoni et al., 2019), with very few regional  
32 exceptions (Figure 11.17 and Section 11.9). Long-term decreases in precipitation are found in some AR6  
33 regions in Africa (CAF, ESAF), and several regions in South America (NES, SAM, SWS, SSA) (Section  
34 11.9). Evidence of precipitation-based drying trends is also found in Western Africa (WAF), consistent with  
35 studies based on CDD trends (Chaney et al., 2014; Donat et al., 2014b; Barry et al., 2018; Dunn et al.,  
36 2020)(Figure 11.17), however there is a partial recovery of the rainfall trends since the 1980s in this region  
37 (Chapter 10, 10.4.2.1). Some AR6 regions show a decrease in meteorological drought, including NAU,  
38 CAU, NEU and CNA (Section 11.9). Other regions do not show substantial trends in long-term  
39 meteorological drought, or display mixed signals depending on the considered time frame and subregions,  
40 such as in Southern Australia (SAU; Gallant et al., 2013; Delworth and Zeng, 2014; Alexander and  
41 Arblaster, 2017; Spinoni et al., 2019; Dunn et al., 2020; Rauniyar and Power, 2020) and the Mediterranean  
42 (MED; Camuffo et al., 2013; Gudmundsson and Seneviratne, 2016; Spinoni et al., 2017; Stagge et al., 2017;  
43 Caloiero et al., 2018; Peña-Angulo et al., 2020) (see also Section 11.9 and Atlas 8.2).

### 44 45 46 **11.6.2.2 Atmospheric evaporative demand**

47  
48 In several regions, AED increases have intensified recent drought events (Williams et al., 2014, 2020; Seager  
49 et al., 2015b; Basara et al., 2019; García-Herrera et al., 2019), enhanced vegetation stress (Allen et al., 2015;  
50 Sanginés de Cárcer et al., 2018; Yuan et al., 2019), or contributed to the depletion of soil moisture or runoff  
51 through enhanced ET (Teuling et al., 2013; Padrón et al., 2020) (*high confidence*). Trends in pan evaporation  
52 measurements and Penman-Monteith AED estimates provide an indication of possible trends in the influence  
53 of AED on drought. Given the observed global temperature increases (Chapter 2; Section 2.3.1.1; Section  
54 11.3) and dominant decrease in relative humidity over land areas (Simmons et al., 2010; Willett et al., 2014),  
55 VPD has increased globally (Barkhordarian et al., 2019; Yuan et al., 2019). Pan evaporation has increased as

1 a consequence of VPD changes in several AR6 regions such as East Asia (EAS; Li et al., 2013; Sun et al.,  
2 2018; Yang et al., 2018a), West Central Europe (WCE; Mozny et al., 2020), MED; Azorin-Molina et al.,  
3 2015) and Central and Southern Australia (CAU, SAU; Stephens et al., 2018). Nevertheless, there is an  
4 important regional variability in observed trends, and in other AR6 regions pan evaporation has decreased  
5 (e.g. in North Central America, NCA (Breña-Naranjo et al., 2016) and in the Tibetan Plateau, TIB (Zhang et  
6 al., 2018a)). Physical models also show an important regional diversity, with an increase in New Zealand  
7 (NZ; Salinger, 2013) and the Mediterranean (MED; Gocic and Trajkovic, 2014; Azorin-Molina et al., 2015;  
8 Piticar et al., 2016), a decrease in SAS (Jhajharia et al., 2015), and strong spatial variability in North  
9 America (Seager et al., 2015b). This variability is driven by the role of other meteorological variables  
10 affecting AED. Changes in solar radiation as a consequence of solar dimming and brightening may affect  
11 trends (Kambezidis et al., 2012; Sanchez-Lorenzo et al., 2015; Wang and Yang, 2014; Chapter 7, Section  
12 7.2.2.2). Wind speed is also relevant (McVicar et al., 2012a), and studies suggest a reduction of the wind  
13 speed in some regions (Zhang et al., 2019h) that could compensate the role of the VPD increase.  
14 Nevertheless, the VPD trend seems to dominate the overall AED trends, compared to the effects of trends in  
15 wind speed and solar radiation (Wang et al., 2012; Park Williams et al., 2017; Vicente-Serrano et al., 2020b).

### 16 17 18 11.6.2.3 Soil moisture deficits

19  
20 There are limited long-term measurements of soil moisture from ground observations (Dorigo et al., 2011;  
21 Qiu et al., 2016; Quiring et al., 2016), which impedes their use in the analysis of trends. Among the few  
22 existing observational studies covering at least two decades, several studies have investigated trends in  
23 ground soil moisture in East Asia (Section 11.9; (Chen and Sun, 2015b; Liu et al., 2015; Qiu et al., 2016)).  
24 Alternatively, microwave-based satellite measurements of surface soil moisture have also been used to  
25 analyse trends (Dorigo et al., 2012; Jia et al., 2018). Although there is regional evidence that microwave-  
26 based soil moisture estimates can capture well drying trends in comparison with ground soil moisture  
27 observations (Jia et al., 2018), there is only *medium confidence* in the derived trends, since satellite soil  
28 moisture data are affected by inhomogeneities (Dorigo et al., 2015; Rodell et al., 2018; Preimesberger et al.,  
29 2020). Furthermore, microwave-based satellites only sense surface soil moisture, which differs from root-  
30 zone soil moisture (Berg et al., 2017b), although relationships can be derived between the two (Brocca et al.,  
31 2011). Several studies have also analysed long-term soil moisture timeseries from observations-driven land-  
32 surface or hydrological models, including land-based reanalysis products (Albergel et al., 2013; Jia et al.,  
33 2018; Gu et al., 2019b; Markonis et al., 2021). Such models have also been used to assess changes in land  
34 water availability, estimated as precipitation minus ET, which is equal to the sum of soil moisture and runoff  
35 (Greve et al., 2014; Padrón et al., 2020).

36  
37 Overall, evidence from global studies suggests that several land regions have been affected by increased soil  
38 drying or water-balance in past decades, despite some spread among products (Albergel et al., 2013; Greve  
39 et al., 2014; Gu et al., 2019b; Padrón et al., 2020). Drying has not only occurred in dry regions, but also in  
40 humid regions (Greve et al., 2014). Some studies have specifically addressed changes in soil moisture at  
41 regional scale (Section 11.9). For AR6 regions, several studies suggest an increase in the frequency and areal  
42 extent of soil moisture deficits, with examples in East Asia (EAS; Cheng et al., 2015; Qin et al., 2015; Jia et  
43 al., 2018), Western and Central Europe (WCE; Trnka et al., 2015b), and the Mediterranean (MED; Hanel et  
44 al., 2018; Moravec et al., 2019; Markonis et al., 2021). Nonetheless, some analyses also show no long-term  
45 trends in soil drying in some AR6 regions, e.g. in Eastern (ENA; Park Williams et al., 2017) and Central  
46 North America (CNA; Seager et al., 2019), as well as in North-Eastern Africa (NEAF; Kew et al., 2021).  
47 The soil moisture drying trends identified in both global and regional studies are generally related to  
48 increases in ET (associated with higher AED) rather than decreases in precipitation, as identified on global  
49 land for trends in water-balance in the dry season (Padrón et al., 2020), as well as for some regions (Teuling  
50 et al., 2013; Cheng et al., 2015; Trnka et al., 2015a; Van Der Linden et al., 2019; Li et al., 2020c).

51  
52 Evidence from observed or observations-derived trends in soil moisture and precipitation minus ET, are  
53 combined with evidence from SPEI and PDSI-PM studies to derive regional assessments of changes in  
54 agricultural and ecological droughts (Section 11.9). This assessment is summarized in Section 11.6.2.6.

55

#### 11.6.2.4 Hydrological deficits

There is evidence based on streamflow records of increased hydrological droughts in East Asia (Zhang et al., 2018b) and southern Africa (Gudmundsson et al., 2019). In areas of Western and Central Europe and of Northern Europe, there is no evidence of changes in the severity of hydrological droughts since 1950 based on flow reconstructions (Caillouet et al., 2017; Barker et al., 2019) and observations (Vicente-Serrano et al., 2019). In the Mediterranean region, there is *high confidence* in hydrological drought intensification (Giuntoli et al., 2013; Gudmundsson et al., 2019; Lorenzo-Lacruz et al., 2013; Masseroni et al., 2020; Section 11.9). In Southeastern South America there is a decrease in the severity of hydrological droughts (Rivera and Penalba, 2018). In North America, depending on the methods, datasets and study periods, there are differences between studies that suggest an increase (Shukla et al., 2015; Udall and Overpeck, 2017) vs a decrease in hydrological drought frequency (Mo and Lettenmaier, 2018), but in general there is strong spatial variability (Poshtiri and Pal, 2016). Streamflow observation reference networks of near-natural catchments have also been used to isolate the effect of climate trends on hydrological drought trends in a few regions, but these show limited trends in Northern Europe and Western and Central Europe (Stahl et al., 2010; Bard et al., 2015; Harrigan et al., 2018), North America (Dudley et al., 2020) and most of Australia with the exception of Eastern and Southern Australia (Zhang et al., 2016c). Given the low availability of observations, there are few studies analysing trends of drought severity in the groundwater. Nevertheless, some studies suggest a noticeable response of groundwater droughts to climate variability (Lorenzo-Lacruz et al., 2017) and increased drought frequency and severity associated with warming, probably as a consequence of enhanced ET induced by higher AED (Maxwell and Condon, 2016). This is supported by studies in Northern Europe (Bloomfield et al., 2019) and North America (Condon et al., 2020).

#### 11.6.2.5 Atmospheric-based drought indices

Globally, trends in SPEI-PM and PDSI-PM suggest slightly higher increases of drought frequency and severity in regions affected by drying over the last decades in comparison to the SPI (Dai and Zhao, 2017; Spinoni et al., 2019; Song et al., 2020), mainly in regions of West and Southern Africa, the Mediterranean and East Asia (Figure 11.17), which is consistent with observed soil moisture trends (Section 11.6.2.3). These indices suggest that AED has contributed to increase the severity of agricultural and ecological droughts compared to meteorological droughts (García-Herrera et al., 2019; Williams et al., 2020), reduce soil moisture during the dry season (Padrón et al., 2020), increase plant water stress (Allen et al., 2015; Grossiord et al., 2020; Solander et al., 2020) and trigger more severe forest fires (Abatzoglou and Williams, 2016; Turco et al., 2019; Nolan et al., 2020). A number of regional studies based on these drought indices have also shown stronger drying trends in comparison to trends in precipitation-based indices in the following AR6 regions (see also 11.9): NSA (Fu et al., 2013b; Marengo and Espinoza, 2016), SCA (Hidalgo et al., 2017), WCA (Tabari and Aghajanjloo, 2013; Sharafati et al., 2020), SAS (Niranjan Kumar et al., 2013), NEAF (Zeke et al., 2017), WSAF (Edossa et al., 2016), NWN and NEN (Bonsal et al., 2013), EAS (Yu et al., 2014; Chen and Sun, 2015b; Li et al., 2020b; Liang et al., 2020; Wu et al., 2020b) and MED (Kelley et al., 2015; Stagge et al., 2017; González-Hidalgo et al., 2018; Mathbout et al., 2018a).

#### [START FIGURE 11.17 HERE]

**Figure 11.17:** Observed linear trend for (a) consecutive dry days (CDD) during 1960-2018, (b) standardized precipitation index (SPI) and (c) standardized precipitation-evapotranspiration index (SPEI) during 1951-2016. CDD data are from the HadEx3 dataset (Dunn et al., 2020), trend calculation of CDD as in Figure 11.9. Drought severity is estimated using 12-month SPI (SPI-12) and 12-month SPEI (SPEI-12). SPI and SPEI datasets are from Spinoni et al. (2019). The threshold to identify drought episodes was set at -1 SPI/SPEI units. Areas without sufficient data are shown in grey. No overlay indicates regions where the trends are significant at  $p = 0.1$  level. Crosses indicate regions where trends are not significant. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1 **[END FIGURE 11.17 HERE]**

2  
3  
4 *11.6.2.6 Synthesis for different drought types*

5  
6 Few AR6 regions show observed increases in meteorological drought (Section 11.9), mostly in Africa and  
7 South America (NES: *high confidence*; WAF, CAF, ESAF, SAM, SWS, SSA, SAS: *medium confidence*); a  
8 few others show a decrease (WSB, ESB, NAU, CAU, NEU, CNA: *medium confidence*). There are stronger  
9 signals indicating observed increases in agricultural and ecological drought (Section 11.9), which highlights  
10 the role of increased ET, driven by increased AED, for these trends (Sections 11.6.2.3, 11.6.2.5). Past  
11 increases in agricultural and ecological droughts are found on all continents and several regions (WAF, CAF,  
12 WSAF, ESAF, WCA, ECA, EAS, SAU, MED, WCE, NES: *medium confidence*), while decreases are found  
13 only in one AR6 region (NAU: *medium confidence*). The more limited availability of datasets makes it more  
14 difficult to assess historical trends in hydrological drought at regional scale (Section 11.9). Increasing (MED:  
15 *high confidence*; WAF, EAS, SAU: *medium confidence*) and decreasing (NEU, SES: *medium confidence*)  
16 trends in hydrological droughts have only been observed in a few regions.

17  
18 In summary, there is *high confidence* that AED has increased on average on continents, contributing to  
19 increased ET and resulting water stress during periods with precipitation deficits, in particular during dry  
20 seasons. There is *medium confidence* in increases in precipitation deficits in a few regions of Africa and  
21 South America. Based on multiple evidence, there is *medium confidence* that agricultural and ecological  
22 droughts have increased in several regions on all continents (WAF, CAF, WSAF, ESAF, WCA, ECA, EAS,  
23 SAU, MED, WCE, NES: *medium confidence*), while there is only *medium confidence* in decreases in one  
24 AR6 region (NAU). More frequent hydrological droughts are found in fewer regions (MED: *high*  
25 *confidence*; WAF, EAS, SAU: *medium confidence*).

26  
27  
28 **11.6.3 Model evaluation**

29  
30 *11.6.3.1 Precipitation deficits*

31  
32 ESMS generally show limited performance and large spread in identifying precipitation deficits and  
33 associated long-term trends in comparison with observations (Nasrollahi et al., 2015). Meteorological  
34 drought trends in the CMIP5 ensemble showed substantial disagreements compared with observations  
35 (Orlowsky and Seneviratne, 2013; Knutson and Zeng, 2018) including a tendency to overestimate drying, in  
36 particular in mid- to high latitudes (Knutson and Zeng, 2018). CMIP6 models display a better performance in  
37 reproducing long-term precipitation trends or seasonal dynamics in some studies in southern South America  
38 (Rivera and Arnould, 2020), East Asia (Xin et al., 2020), southern Asia (Gusain et al., 2020), and  
39 southwestern Europe (Peña-Angulo et al., 2020b), but there is still too *limited evidence* to allow for an  
40 assessment of possible differences in performance between CMIP5 and CMIP6. Furthermore, ESMS are  
41 generally found to underestimate the severity of precipitation deficits and the dry day frequencies in  
42 comparison to observations (Fantini et al., 2018; Ukkola et al., 2018). This is probably related to  
43 shortcomings in the simulation of persistent weather events in the mid-latitudes (Chapter 10, Section  
44 10.3.3.3). In addition, ESMS also show a tendency to underestimate precipitation-based drought persistence  
45 at monthly to decadal time scales (Ault et al., 2014; Moon et al., 2018). The overall inter-model spread in  
46 the projected frequency of precipitation deficits is also substantial (Touma et al., 2015; Zhao et al., 2016;  
47 Engström and Keellings, 2018). Moreover, there are spatial differences in the spread, which is higher in the  
48 regions where enhanced drought conditions are projected and under high-emission scenarios (Orlowsky and  
49 Seneviratne, 2013). Nonetheless, some event attribution studies have concluded that droughts at regional  
50 scales can be adequately simulated by some climate models (Schaller et al., 2016; Otto et al., 2018c).

51  
52  
53 *11.6.3.2 Atmospheric evaporative demand*

54  
55 There is only limited evidence on the evaluation of AED in state-of-the-art ESMS, which is performed on



externally computed AED based on model output (Scheff and Frierson, 2015; Liu and Sun, 2016, 2017). An evaluation of average AED in 17 CMIP5 ESMs for 1981-1999 based on potential evaporation show that the models' spatial patterns resemble the observations, but that the magnitude of potential evaporation displays strong divergence among models globally and regionally (Scheff and Frierson, 2015). The evaluation of AED in 12 CMIP5 ESMs with pan evaporation observations in East Asia for 1961-2000 (Liu and Sun, 2016, 2017) show that the ESMs capture seasonal cycles well, but that regional AED averages are underestimated due to biases in the meteorological variables controlling the aerodynamic and radiative components of AED. CMIP5 ESMs also show a strong underestimation of atmospheric drying trends compared to reanalysis data (Douville and Plazzotta, 2017).

### 11.6.3.3 Soil moisture deficits

The performance of climate models for representing soil moisture deficits shows more uncertainty than for precipitation deficits since in addition to the uncertainties related to cloud and precipitation processes, there is uncertainty related to the representation of complex soil hydrological and boundary-layer processes (Van Den Hurk et al., 2011; Lu et al., 2019; Quintana-Seguí et al., 2020). A limitation is also the lack of observations, and in particular soil moisture, in most regions (Section 11.6.2.3), and the paucity of land surface property data to parameterize land surface models, in particular soil types, soil properties and depth (Xia et al., 2015). The spatial resolution of models is an additional limitation since the representation of some land-atmosphere feedbacks and topographic effects requires detailed resolution (Nicolai-Shaw et al., 2015; Van Der Linden et al., 2019). Beside climate models, also land surface and hydrological models are used to derive historical and projected trends in soil moisture and related land water variables (Albergel et al., 2013; Cheng et al., 2015; Gu et al., 2019b; Padrón et al., 2020; Markonis et al., 2021; Pokhrel et al., 2021).

Overall, there are contrasting results on the performance of land surface models and climate models in representing soil moisture. Some studies suggest that soil moisture anomalies are well captured by land surface models driven with observation-based forcing (Dirmeyer et al., 2006; Albergel et al., 2013; Xia et al., 2014; Balsamo et al., 2015; Reichle et al., 2017; Spennemann et al., 2020), but other studies report limited agreement in the representation of interannual soil moisture variability (Stillman et al., 2016; Yuan and Quiring, 2017; Ford and Quiring, 2019) and noticeable seasonal differences in model skill (Xia et al., 2014, 2015) in some regions. Models with good skill can nonetheless display biases in absolute soil moisture (Xia et al., 2014; Gu et al., 2019a), but these are not necessarily of relevance for the simulation of surface water fluxes and drought anomalies (Koster et al., 2009). There is also substantial intermodel spread (Albergel et al. 2013), particularly for the root-zone soil moisture (Berg et al., 2017b).

Regarding the performance of regional and global climate models, an evaluation of an ensemble of RCM simulations for Europe (Stegehuis et al., 2013) shows that these models display too strong drying in early summer, resulting in an excessive decrease of latent heat fluxes, with potential implications for more severe droughts in dry environments (Teuling, 2018; Van Der Linden et al., 2019). Compared with a range of observational ET estimates, CMIP5 models show an overestimation of ET on annual scale, but an ET underestimation in boreal summer in many North-Hemisphere mid-latitude regions, also suggesting a tendency towards excessive soil drying (Mueller and Seneviratne, 2014), consistent with identified biases in soil moisture-temperature coupling (Donat et al., 2018; Vogel et al., 2018; Selten et al., 2020). Land surface models used in ESM display a bias in their representation of the sensitivity of interannual land carbon uptake to soil moisture conditions, which appears related to a limited range of soil moisture variations compared to observations (Humphrey et al., 2018).

For future projections, the spread of soil moisture outputs among different ESMs is more important than internal variability and scenario uncertainty, and the bias is strongly related to the sign of the projected change (Ukkola et al., 2018; Lu et al., 2019; Selten et al., 2020). CMIP5 ESMs that project more drying and warming in mid-latitude regions show a substantial bias in soil moisture-temperature coupling (Donat et al., 2018; Vogel et al., 2018). Although CMIP6 and CMIP5 simulations for soil moisture changes are overall similar, some differences are found in projections in a few regions (Cook et al., 2020)(see also Section 11.9).

1 There is still *limited evidence* to assess whether there are substantial differences in model performance in the  
2 two ensembles, but improvements in modeling aspects relevant for soil moisture have been reported for  
3 precipitation (11.6.3.2), and a better performance has been found in CMIP6 for the representation of long-  
4 term trends in soil moisture in the continental USA (Yuan et al., 2021). Despite the mentioned model  
5 limitations, the representation of soil moisture processes in ESMs uses physical and biological understanding  
6 of the underlying processes, which can represent well the temporal anomalies associated with temporal  
7 variability and trends in climate. In summary, there is *medium confidence* in the representation of soil  
8 moisture deficits in ESMs and related land surface and hydrological models.  
9

#### 10 11 *11.6.3.4 Hydrological deficits*

12  
13 Streamflow and groundwater are not directly simulated by ESMs, which only simulate runoff, but they are  
14 generally represented in hydrological models (Prudhomme et al., 2014; Giuntoli et al., 2015), which are  
15 typically driven in a stand-alone manner by observed or simulated climate forcing. The simulation of  
16 hydrological deficits is much more problematic than the simulation of mean streamflow or peak flows  
17 (Fundel et al., 2013; Stoelzle et al., 2013; Velázquez et al., 2013; Staudinger et al., 2015), since models tend  
18 to be too responsive to the climate forcing and do not satisfactorily capture low flows (Tallaksen and Stahl,  
19 2014). Simulations of hydrological drought metrics show uncertainties related to the contribution of both  
20 GCMs and hydrological models (Bosshard et al., 2013; Giuntoli et al., 2015; Samaniego et al., 2017; Vetter  
21 et al., 2017), but hydrological models forced by the same climate input data also show a large spread (Van  
22 Huijgevoort et al., 2013; Ukkola et al., 2018). At the catchment scale, the hydrological model uncertainty is  
23 higher than both GCM and downscaling uncertainty (Vidal et al., 2016), and the hydrological models show  
24 issues in representing drought propagation throughout the hydrological cycle (Barella-Ortiz and Quintana  
25 Seguí, 2019). A study on the evaluation of streamflow droughts in seven global (hydrological and land  
26 surface) models compared with observations in near-natural catchments of Europe showed a substantial  
27 spread among models, an overestimation of the number of drought events, and an underestimation of drought  
28 duration and drought-affected area (Tallaksen and Stahl, 2014).  
29

#### 30 31 *11.6.3.5 Atmospheric-based drought indices*

32  
33 A number of studies have analysed the ability of models to capture drought severity and trends based on  
34 climatic drought indices. Given the limitations of ESMs in reproducing the dynamic of precipitation deficits  
35 and AED (11.6.3.1, 11.6.3.2), atmospheric-based drought indices derived from ESM data for these two  
36 variables are also affected by uncertainties and biases. A comparison of historical trends in PDSI-PM for  
37 1950-2014 derived from CMIP3 and CMIP5 with respective estimates derived from observations (Dai and  
38 Zhao, 2017) show a similar behaviour at global scale (long-term decrease), but low spatial agreement in the  
39 trends except in a few regions (Mediterranean, South Asia, northwestern US). In future projections there is  
40 an important spread in PDSI-PM and SPEI-PM among different models (Cook et al., 2014a).  
41

#### 42 43 *11.6.3.6 Synthesis for different drought types*

44  
45 The performance of ESMs used to assess changes in variables related to meteorological droughts,  
46 agricultural and ecological droughts, and hydrological droughts, show the presence of biases and  
47 uncertainties compared to observations, but there is *medium confidence* in their overall performance for  
48 assessing drought projections given process understanding. Given the substantial inter-model spread  
49 documented for all related variables, the consideration of multi-model projections increases the confidence  
50 of model-based assessments, with only *low confidence* in assessments based on single models.  
51

52 In summary, the evaluation of ESMs, land surface and hydrological models for the simulation of droughts is  
53 complex, due to the regional scale of drought trends, their overall low signal-to-noise ratio, and the lack of  
54 observations in several regions, in particular for soil moisture and streamflow. There is *medium confidence*  
55 in the ability of ESMs to simulate trends and anomalies in precipitation deficits and AED, and also *medium*

1 *confidence* in the ability of ESMs and hydrological models to simulate trends and anomalies in soil moisture  
2 and streamflow deficits, on global and regional scales.  
3  
4

#### 5 **11.6.4 Detection and attribution, event attribution**

##### 6 **11.6.4.1 Precipitation deficits**

7  
8  
9 There are only two AR6 regions in which there is at least *medium confidence* that human-induced climate  
10 change has contributed to changes in meteorological droughts (Section 11.9). In South-western South  
11 America (SSW), there is *medium confidence* that human-induced climate change has contributed to an  
12 increase in meteorological droughts (Boisier et al., 2016; Garreaud et al., 2020), while in Northern Europe  
13 (NEU), there is *medium confidence* that it has contributed to a decrease in meteorological droughts  
14 (Gudmundsson and Seneviratne, 2016) (Section 11.9). In other AR6 regions, there is inconclusive evidence  
15 in the attribution of long-term trends, but a human contribution to single meteorological events or  
16 subregional trends has been identified in some instances (Section 11.9; see also below). In the Mediterranean  
17 (MED) region, some studies have identified a precipitation decline or increase in meteorological drought  
18 probability for time frames since the early or mid 20<sup>th</sup> century and a possible human contribution to these  
19 trends (Hoerling et al., 2012; Gudmundsson and Seneviratne, 2016; Knutson and Zeng, 2018), also on  
20 subregional scale in Syria from 1930 to 2010 (Kelley et al., 2015). On the contrary, other studies have not  
21 identified precipitation and meteorological drought trends in the region for the long-term (Camuffo et al.,  
22 2013; Paulo et al., 2016; Vicente-Serrano et al., 2021) and also from the mid 20<sup>th</sup> century (Narrant and  
23 Douguédroit, 2006; Stagge et al., 2017). There is evidence of substantial internal variability in long-term  
24 precipitation trends in the region (Section 11.6.2.1), which limits the attribution of human influence on  
25 variability and trends of meteorological droughts from observational records (Kelley et al., 2012; Peña-  
26 Angulo et al., 2020b). In addition, there are important subregional trends showing mixed signals (MedECC,  
27 2020)(Section 11.9). The evidence thus leads to an assessment of *low confidence* in the attribution of  
28 observed short-term changes in meteorological droughts in the region (Section 11.9). In North America, the  
29 human influence on precipitation deficits is complex (Wehner et al., 2017), with *low confidence* in the  
30 attribution of long-term changes in meteorological drought in AR6 regions (Lehner et al., 2018; Section  
31 11.9). In Africa there is *low confidence* that human influence has contributed to the observed long-term  
32 meteorological drought increase in Western Africa (Section 11.9; Chapter 10, Section 10.6.2). There is *low*  
33 *confidence* in the attribution of the observed increasing trends in meteorological drought in Eastern Southern  
34 Africa, but evidence that human-induced climate change has affected recent meteorological drought events  
35 in the region (11.9).  
36

37 Attribution studies for recent meteorological drought events are available for various regions. In Central and  
38 Western Europe, a multi-method and multi-model attribution study on the 2015 Central European drought  
39 did not find conclusive evidence for whether human-induced climate change was a driver of the rainfall  
40 deficit, as the results depended on model and method used (Hauser et al., 2017). In the Mediterranean region,  
41 a human contribution was found in the case of the 2014 meteorological drought in the southern Levant based  
42 on a single-model study (Bergaoui et al., 2015). In Africa, there is some evidence of a contribution of human  
43 emissions to single meteorological drought events, such as the 2015-2017 southern African drought (Funk et  
44 al., 2018a; Yuan et al., 2018a; Pascale et al., 2020), and the three-year 2015-2017 drought in the western  
45 Cape Town region of South Africa (Otto et al., 2018c). An attributable signal was not found in droughts that  
46 occurred in different years with different spatial extents in the last decade in Northern and Southern East  
47 Africa (Marthews et al., 2015; Uhe et al., 2017; Otto et al., 2018a; Philip et al., 2018b; Kew et al., 2021).  
48 However, an attributable increase in 2011 long rain failure was identified (Lott et al., 2013). Further studies  
49 have attributed some African meteorological drought events to large-scale modes of variability, such as the  
50 strong 2015 El Niño (Philip et al., 2018; Box 11.4) and increased SSTs overall (Funk et al., 2015b, 2018b).  
51 Natural variability was dominant in the California droughts of 2011/12-2013/14 (Seager et al., 2015a). In  
52 Asia, no climate change signal was found in the record dry spell over Singapore-Malaysia in 2014 (Mcbride  
53 et al., 2015) or the drought in central southwest Asia in 2013/2014 (Barlow and Hoell, 2015). Nevertheless,  
54 the South East Asia drought of 2015 has been attributed to anthropogenic warming effects (Shiogama et al.,  
55 2020). Recent droughts occurring in South America, specifically in the southern Amazon region in 2010

1 (Shiogama et al., 2013) and in Northeast South America in 2014 (Otto et al., 2015) and 2016 (Martins et al.,  
2 2018) were not attributed to anthropogenic climate change. Nevertheless, the central Chile drought between  
3 2010 and 2018 has been suggested to be partly associated to global warming (Boisier et al., 2016; Garreaud  
4 et al., 2020). The 2013 New Zealand meteorological drought was attributed to human influence by  
5 Harrington et al. (2014, 2016) based on fully coupled CMIP5 models, but, no corresponding change in the  
6 dry end of simulated precipitation from a stand-alone atmospheric model was found by Angélil et al. (2017).  
7

8 Event attribution studies also highlight a complex interplay of anthropogenic and non-anthropogenic  
9 climatological factors for some events. For example, anthropogenic warming contributed to the 2014 drought  
10 in North Eastern-Africa by increasing east African and west Pacific temperatures, and increasing the gradient  
11 between standardized western and central Pacific SSTs causing reduced rainfall (Funk et al., 2015b). As  
12 different methodologies, models and data sources have been used for the attribution of precipitation deficits,  
13 Angélil et al. (2017) reexamined several events using a single analytical approach and climate model and  
14 observational datasets. Their results showed a disagreement in the original anthropogenic attribution in a  
15 number of precipitation deficit events, which increased uncertainty in the attribution of meteorological  
16 droughts events.  
17

#### 19 11.6.4.2 Soil moisture deficits

20  
21 There is a growing number of studies on the detection and attribution of long-term changes in soil moisture  
22 deficits. Mueller and Zhang (2016) concluded that anthropogenic forcing contributed significantly to an  
23 increase in the land surface area affected by soil moisture deficits, which can be reproduced by CMIP5  
24 models only if anthropogenic forcings are involved. A similar assessment was provided globally by Gu et al.  
25 (2019b) also using CMIP5 models. Padrón et al. (2019) analyzed long-term reconstructed and CMIP5  
26 simulated dry season water availability, defined as precipitation minus ET (i.e., equivalent to soil moisture  
27 and runoff availability), and found that patterns of changes in dry-season deficits in the recent three last  
28 decades can only be explained by anthropogenic forcing and are mostly related to changes in ET. Similarly  
29 Williams et al. (2020) concluded human-induced climate change contributed to the strong soil moisture  
30 deficits recorded in the last two decades in western North America through VPD increases associated with  
31 higher air temperatures and lower air humidity. There are few studies analysing the attribution of particular  
32 episodes of soil moisture deficits to anthropogenic influence. Nevertheless, the available modeling studies  
33 coincide in supporting an anthropogenic attribution associated with more extreme temperatures, exacerbating  
34 AED and increasing ET, and thus depleting soil moisture, as observed in southern Europe in 2017 (García-  
35 Herrera et al., 2019) and in Australia in 2018 (Lewis et al., 2019b) and 2019 (van Oldenborgh et al., 2021),  
36 the latter event having strong implications in the propagation of widespread mega-fires (Nolan et al., 2020).  
37

#### 39 11.6.4.3 Hydrological deficits

40  
41 It is often difficult to separate the role of climate trends from changes in land use, water management and  
42 demand for changes in hydrological deficits, especially on regional scale. However, a global study based on  
43 a recent multi-model experiment with global hydrological models and covering several AR6 regions  
44 suggests a dominant role of anthropogenic radiative forcing for trends in low, mean and high flows, while  
45 simulated effects of water and land management do not suffice to reproduce the observed spatial pattern of  
46 trends (Gudmundsson et al., 2021). Regional studies also suggest that climate trends have been dominant  
47 compared to land use and human water management for explaining trends in hydrological droughts in some  
48 regions, for instance in Ethiopia (Fenta et al., 2017), in China (Xie et al., 2015), and in North America for the  
49 Missouri and Colorado basins, as well as in California (Shukla et al., 2015; Udall and Overpeck, 2017;  
50 Ficklin et al., 2018; Xiao et al., 2018a; Glas et al., 2019; Martin et al., 2020; Milly and Dunne, 2020).  
51

52 In other regions the influence of human water uses can be more important to explain hydrological drought  
53 trends (Liu et al., 2016b; Mohammed and Scholz, 2016). There is *medium confidence* that human-induced  
54 climate change has contributed to an increase of hydrological droughts in the Mediterranean (Giuntoli et al.,  
55 2013; Vicente-Serrano et al., 2014; Gudmundsson et al., 2017), but also *medium confidence* that changes in

land use and terrestrial water management contributed to these trends as well (Teuling et al., 2019; Vicente-Serrano et al., 2019; Section 11.9). A global study with a single hydrological model estimated that human water consumption has intensified the magnitude of hydrological droughts by 20%-40% over the last 50 years, and that the human water use contribution to hydrological droughts was more important than climatic factors in the Mediterranean, and the central US, as well as in parts of Brazil (Wada et al., 2013). However, Gudmundsson et al. (2021) concluded that the contribution of human water use is smaller than that of anthropogenic climate change to explain spatial differences in the trends of low flows based on a multi-model analysis. There is still *limited evidence* and thus *low confidence* in assessing these trends at the scale of single regions, with few exceptions (Section 11.9).

#### 11.6.4.4 Atmospheric-based drought indices

Different studies using atmospheric-based drought indices suggest an attributable anthropogenic signal, characterized by the increased frequency and severity of droughts (Cook et al., 2018), associated to increased AED (Section 11.6.4.2). The majority of studies are based on the PDSI-PM. Williams et al. (2015) and Griffin and Anchukaitis (2014) concluded that increased AED has had an increased contribution to drought severity over the last decades, and played a dominant role in the intensification of the 2012-2014 drought in California. The same temporal pattern and physical mechanism was stressed by Li et al. (2017) in Central Asia. Marvel et al. (2019) compared tree ring-based reconstructions of the PDSI-PM over the past millennium with PDSI-PM estimates based on output from CMIP5 models, suggesting a contribution of greenhouse gas forcing to the changes since the beginning of the 20<sup>th</sup> century, although characterized with temporal differences that could be driven by temporal variations in the aerosol forcing, in agreement with the dominant external forcings of aridification at global scale between 1950 and 2014 (Bonfils et al., 2020). In the Mediterranean region there is *medium confidence* of drying attributable to anthropogenic forcing as a consequence of the strong AED increase (Gocic and Trajkovic, 2014; Liuzzo et al., 2014; Azorin-Molina et al., 2015; Maček et al., 2018), which has enhanced the severity of drought events (Vicente-Serrano et al., 2014; Stagge et al., 2017; González-Hidalgo et al., 2018). In particular, this effect was identified to be the main driver of the intensification of the 2017 drought that affected southwestern Europe, and was attributed to the human forcing (García-Herrera et al., 2019). Nangombe et al. (2020) and Zhang et al. (2020) concluded from differences between precipitation and AED that anthropogenic forcing contributed to 2018 droughts that affected southern Africa and southeastern China, respectively, principally as consequence of the high AED that characterised these two events.

#### 11.6.4.5 Synthesis for different drought types

The regional evidence on attribution for single AR6 regions generally shows *low confidence* for a human contribution to observed trends in meteorological droughts at regional scale, with few exceptions (Section 11.9). There is *medium confidence* that human influence has contributed to changes in agricultural and ecological droughts and has led to an increase in the overall affected land area. At regional scales, there is *medium confidence* in a contribution of human-induced climate change to increases in agricultural and ecological droughts in the Mediterranean (MED) and Western North America (WNA) (Section 11.9). There is *medium confidence* that human-induced climate change has contributed to an increase in hydrological droughts in the Mediterranean region, but also *medium confidence* in contributions from other human influences, including water management and land use (Section 11.9). Several meteorological and agricultural and ecological drought events have been attributed to human-induced climate change, even in regions where no long-term changes are detected (*medium confidence*). However, a lack of attribution to human-induced climate change has also been shown for some events (*medium confidence*).

In summary, human influence has contributed to changes in water availability during the dry season over land areas, including decreases over several regions due to increases in evapotranspiration (*medium confidence*). The increases in evapotranspiration have been driven by increases in atmospheric evaporative demand induced by increased temperature, decreased relative humidity and increased net radiation over affected land areas (*high confidence*). There is *low confidence* that human influence has affected trends in

1 meteorological droughts in most regions, but *medium confidence* that they have contributed to the severity of  
2 some single events. There is *medium confidence* that human-induced climate change has contributed to  
3 increasing trends in the probability or intensity of recent agricultural and ecological droughts, leading to an  
4 increase of the affected land area. Human-induced climate change has contributed to global-scale change in  
5 low flow, but human water management and land use changes are also important drivers (*medium*  
6 *confidence*).

### 9 **11.6.5 Projections**

10 SREX (Chapter 3) assessed with *medium confidence* projections of increased drought severity in some  
11 regions, including southern Europe and the Mediterranean, central Europe, Central America and Mexico,  
12 northeast Brazil, and southern Africa, and *low confidence* elsewhere given large inter-model spread. AR5  
13 (Chapters 11 and 12) also assessed large uncertainties in drought projections at the regional and global  
14 scales. The assessment of drought mechanisms under future climate change scenarios depends on the model  
15 used (Section 11.6.3). Moreover, uncertainties in drought projections are affected by the consideration of  
16 plant physiological responses to increasing atmospheric CO<sub>2</sub> (Greve et al., 2019; Mankin et al., 2019; Milly  
17 and Dunne, 2016; Yang et al., 2020; Chapter 5, Cross-Chapter Box 5.1), the role of soil moisture-atmosphere  
18 feedbacks for changes in water-balance and aridity (Berg et al., 2016; Zhou et al., 2021), and statistical  
19 issues related to considered drought time scales (Vicente-Serrano et al., 2020a). Nonetheless, the extensive  
20 literature available since AR5 allows a substantially more robust assessment of projected changes in  
21 droughts, also subdivided in different drought types (meteorological drought, agricultural and ecological  
22 drought, and hydrological drought). This includes assessments of projected changes in droughts, including  
23 changes at 1.5°C, 2°C and 4°C of global warming, for all AR6 regions (Section 11.9). Projected changes  
24 show increases in drought frequency and intensity in several regions as function of global warming (*high*  
25 *confidence*). There are also substantial increases in drought hazard probability from 1.5°C to 2°C global  
26 warming as well as for further additional increments of global warming (Figs. 11.18 and 11.19) (*high*  
27 *confidence*). These findings are based both on CMIP5 and CMIP6 analyses (Section 11.9; Greve et al., 2018;  
28 Wartenburger et al., 2017; Xu et al., 2019a), and strengthen the conclusions of the SR15 Ch3.

#### 32 **11.6.5.1 Precipitation deficits**

33 Studies based on CMIP5, CMIP6 and CORDEX projections show a consistent signal in the sign and spatial  
34 pattern of projections of precipitation deficits. Global studies based on these multi-model ensemble  
35 projections (Orlowsky and Seneviratne, 2013; Martin, 2018; Spinoni et al., 2020; Ukkola et al., 2020;  
36 Coppola et al., 2021b) show particularly strong signal-to-noise ratios for increasing meteorological droughts  
37 in the following AR6 regions: MED, ESAF, WSAF, SAU, CAU, NCA, SCA, NSA and NES (Section 11.9).  
38 There is also substantial evidence of changes in meteorological droughts at 1.5°C vs 2°C of global warming  
39 from global studies (Wartenburger et al., 2017; Xu et al., 2019a). The patterns of projected changes in mean  
40 precipitation are consistent with the changes in the drought duration, but they are not consistent with the  
41 changes in drought intensity (Ukkola et al., 2020). In general, CMIP6 projections suggest a stronger increase  
42 of the probability of precipitation deficits than CMIP5 projections (Cook et al., 2020; Ukkola et al., 2020).  
43 Projections for the number of CDDs in CMIP6 (Figure 11.19) for different levels of global warming relative  
44 to 1850-1900 show similar spatial patterns as projected precipitation deficits. The robustness of the patterns  
45 in projected precipitation deficits identified in the global studies is also consistent with results from regional  
46 studies (Giorgi et al., 2014; Marengo and Espinoza, 2016; Pinto et al., 2016; Huang et al., 2018a; Maure et  
47 al., 2018; Nangombe et al., 2018; Tabari and Willems, 2018; Abiodun et al., 2019; Dosio et al., 2019).

48 In Africa, a strong increase in the length of dry spells (CDD) is projected for 4°C of global warming over  
49 most of the continent with the exception of central and eastern Africa (Giorgi et al., 2014; Han et al., 2019;  
50 Sillmann et al., 2013; Section 11.9). In West Africa, a strong reduction of precipitation is projected  
51 (Sillmann et al., 2013a; Diallo et al., 2016; Akinsanola and Zhou, 2018; Han et al., 2019; Todzo et al., 2020)  
52 at 4°C of global warming, and CDD would increase with stronger global warming levels (Klutse et al.,  
53 2018). The regions most strongly affected are Southern Africa (ESAF, WSAF; (Nangombe et al., 2018);  
54  
55

1 Abiodun et al., 2019) and Northern Africa (part of MED region), with increases in meteorological droughts  
2 already at 1.5°C of global warming, and further increases with increasing global warming (Section 11.9).  
3 CDD is projected to increase more in the southern Mediterranean (northern Africa) than in the northern part  
4 of the Mediterranean region (Lionello and Scarascia, 2020).

5  
6 In Asia, most AR6 regions show *low confidence* in projected changes in meteorological droughts at 1.5°C  
7 and 2°C of global warming, with a few regions displaying a decrease in meteorological droughts at 4°C of  
8 global warming (RAR, ESB, RFE, ECA; *medium confidence*), although there is a projected increase in  
9 meteorological droughts in Southeast Asia (SEA) at 4°C (*medium confidence*) (Section 11.9). In Southeast  
10 Asia, an increasing frequency of precipitation deficits is projected as a consequence of an increasing  
11 frequency of extreme El Niño (Cai et al., 2014a, 2015, 2018).

12  
13 In central America, projections suggest an increase in mid-summer meteorological drought (Imbach et al.,  
14 2018) and increased CDD (Nakaegawa et al., 2013; Chou et al., 2014a; Giorgi et al., 2014). In the Amazon,  
15 there is also a projected increase in dryness (Marengo and Espinoza, 2016), which is the combination of a  
16 projected increase in the frequency and geographic extent of meteorological drought in the eastern Amazon,  
17 and an opposite trend in the West (Duffy et al., 2015). In southwestern South America, there is a projected  
18 increase of the CDD (Chou et al., 2014a; Giorgi et al., 2014) and in Chile, drying is projected to prevail  
19 (Boisier et al., 2018). In the South America monsoon region, an increase in CDD is projected (Chou et al.,  
20 2014a; Giorgi et al., 2014), but a decrease is projected in southeastern and southern South America (Giorgi et  
21 al., 2014). In Central America, mid summer meteorological drought is projected to intensify during 2071-  
22 2095 for the RCP8.5 scenario (Corrales-Suastegui et al., 2019).

23  
24 An increase in the frequency, duration and intensity of meteorological droughts is projected in southwest,  
25 south and east Australia (Kirono et al., 2020; Shi et al., 2020). In Canada and most of the USA, and based on  
26 the SPI, Swain and Hayhoe (2015) identified drier summer conditions in projections over most of the region,  
27 and there is a consistent signal toward an increase in duration and intensity of droughts in southern North  
28 America (Pascale et al., 2016; Escalante-Sandoval and Nuñez-Garcia, 2017). In California, more  
29 precipitation variability is projected, characterised by increased frequency of consecutive drought and humid  
30 periods (Swain et al., 2018).

31  
32 Substantial increases in meteorological drought are projected in Europe, in particular in the Mediterranean  
33 region already at 1.5°C of global warming (Section 11.9). In southern Europe, model projections display a  
34 consistent drying among models (Russo et al., 2013; Hertig and Trambly, 2017; Guerreiro et al., 2018a;  
35 Raymond et al., 2019). In Western and Central Europe there is some spread in CMIP5 projections, with  
36 some models projecting very strong drying and others close to no trend (Vogel et al., 2018), although CDD  
37 is projected to increase in CMIP5 projections under the RCP 8.5 scenario (Hari et al., 2020). The overall  
38 evidence suggests an increase in meteorological drought at 4°C in the WCE region (*medium confidence*;  
39 Section 11.9).

40  
41 Overall, based on both global and regional studies, several hot spot regions are identified displaying more  
42 frequent and severe meteorological droughts with increasing with global warming, including several AR6  
43 regions at 1.5°C (WSAF, ESAF, SAU, MED, NES) and 2°C of global warming (WSAF, ESAF, EAU, SAU,  
44 MED, NCA, SCA, NSA, NES) (Section 11.9). At 4°C of global warming, there is also confidence in  
45 increases in meteorological droughts in further regions (WAF, WCE, ENA, CAR, NWS, SAM, SWS, SSA;  
46 Section 11.9), showing a geographical expansion of meteorological drought with increasing global warming.  
47 Only few regions are projected to have less intense or frequent meteorological droughts (Section 11.9).

#### 50 11.6.5.2 Atmospheric evaporative demand

51  
52 Effects of AED on droughts in future projections is under debate. CMIP5 models project an AED increase  
53 over the majority of the world with increasing global warming, mostly as a consequence of strong VPD  
54 increases (Scheff and Frierson, 2015; Vicente-Serrano et al., 2020b). However, ET is projected to increase  
55 less than AED in many regions, due to plant physiological responses related to i) CO<sub>2</sub> effects on plant

1 photosynthesis and ii) soil moisture control on ET.

2  
3 Several studies suggest that increasing atmospheric CO<sub>2</sub> could lead to reduced leaf stomatal conductance,  
4 which would increase water-use efficiency and reduce plant water needs, thus limiting ET (Chapter 5, Cross-  
5 Chapter Box 5.1; Greve et al., 2017; Lemordant et al., 2018; Milly and Dunne, 2016; Roderick et al., 2015;  
6 Scheff et al., 2017; Swann, 2018; Swann et al., 2016). The implementation of a CO<sub>2</sub>-dependent land resistance  
7 parameter has been suggested for the estimation of AED (Yang et al., 2019). Nevertheless, there are other  
8 relevant mechanisms, as soil moisture deficits and VPD also play an important role in the control of the leaf  
9 stomatal conductance (Xu et al., 2016b; Menezes-Silva et al., 2019; Grossiord et al., 2020) and a number of  
10 ecophysiological and anatomical processes affect the response of plant physiology under higher atmospheric  
11 CO<sub>2</sub> concentrations (Mankin et al., 2019; Menezes-Silva et al., 2019; Chapter 5, Cross-Chapter Box 5.1).  
12 The benefits of the atmospheric CO<sub>2</sub> for plant stress and agricultural and ecological droughts would be  
13 minimal precisely during dry periods given stomatal closure in response to limited soil moisture (Allen et al.,  
14 2015; Xu et al., 2016b). In addition, CO<sub>2</sub> effects on plant stomatal conductance could not entirely  
15 compensate the increased demand associated to warming (Liu and Sun, 2017); in large tropical and  
16 subtropical regions (e.g. southern Africa, the Amazon, the Mediterranean and southern North America),  
17 AED is projected to increase even considering the possible CO<sub>2</sub> effects on the land resistance (Vicente-  
18 Serrano et al., 2020b). Moreover, these CO<sub>2</sub> effects would not affect the direct evaporation from soils and  
19 water bodies, which is very relevant in the reservoirs of warm areas (Friedrich et al., 2018). Because of these  
20 uncertainties, there is *low confidence* whether increased CO<sub>2</sub>-induced water-use efficiency in vegetation will  
21 substantially reduce global plant transpiration and will diminish the frequency and severity of soil moisture  
22 and streamflow deficits associated with the radiative effect of higher CO<sub>2</sub> concentrations (Chapter 5, CC Box  
23 5.1).  
24

25 Another mechanism reducing the ET response to increased AED in projections is the control of soil moisture  
26 limitations on ET, which leads to reduced stomatal conductance under water stress (Berg and Sheffield,  
27 2018; Stocker et al., 2018; Zhou et al., 2021). This response may be further amplified through VPD-induced  
28 decreases in stomatal conductance (Anderegg et al., 2020). However, the decreased stomatal conductance in  
29 response to both soil moisture limitation and enhanced CO<sub>2</sub> would further enhance AED (Sherwood and Fu,  
30 2014; Berg et al., 2016; Teuling, 2018; Miralles et al., 2019), whereby the overall effects on AED in ESMs  
31 are found to be of similar magnitude for soil moisture limitation and CO<sub>2</sub> physiological effects on stomatal  
32 conductance (Berg et al., 2016). Increased AED is thus both a driver and a feedback with respect to changes  
33 in ET, complicating the interpretation of its role on drought changes with increasing CO<sub>2</sub> concentrations and  
34 global warming.  
35

### 36 37 11.6.5.3 Soil moisture deficits

38  
39 Areas with projected soil moisture decreases do not fully coincide with areas with projected precipitation  
40 decreases, although there is substantial consistency in the respective patterns (Dirmeyer et al., 2013; Berg  
41 and Sheffield, 2018). There are, however, more regions affected by increased soil moisture deficits (Figure  
42 11.19) than precipitation deficits (CC-Box 11.1, Figures 2a,b,c), as a consequence of enhanced AED and the  
43 associated increased ET, as highlighted by some studies (Dai et al., 2018; Orłowsky and Seneviratne, 2013;  
44 Chapter 8, Section 8.2.2.1). Moisture in the top soil layer is projected to decrease more than precipitation at  
45 all warming levels (Lu et al., 2019), extending the regions affected by severe soil moisture deficits over most  
46 of south and central Europe (Lehner et al., 2017; Ruosteenoja et al., 2018; Samaniego et al., 2018; Van Der  
47 Linden et al., 2019), southern North America (Cook et al., 2019), South America (Orłowsky and  
48 Seneviratne, 2013), southern Africa (Lu et al., 2019), East Africa (Rowell et al., 2015), southern Australia  
49 (Kirono et al., 2020), India (Mishra et al., 2014b) and East Asia (Cheng et al., 2015) (Figure 11.19).  
50 Projected changes in total soil moisture display less widespread drying than those for surface soil moisture  
51 (Berg et al., 2017b), but still more than for precipitation (CC-Box 11.1, Figures 2a,b,c). The severity of  
52 droughts based on surface soil moisture in future projections is stronger than projections based on  
53 precipitation and runoff (Dai et al., 2018; Vicente-Serrano et al., 2020a). Nevertheless, in many parts of the  
54 world in which soil moisture is projected to decrease, the signal to noise ratio among models is low and only  
55 in the Mediterranean, Europe, the southwestern United States, and southern Africa the projections show a



1 high signal to noise ratio in soil moisture projections (Lu et al., 2019; (Figure 11.19). Increases in soil  
 2 moisture deficits are found to be statistically significant at regional scale in the Mediterranean region,  
 3 Southern Africa and Western South America for changes as small as 0.5°C in global warming, based on  
 4 differences between +1.5°C and +2°C of global warming (Wartenburger et al., 2017). Several other regions  
 5 are affected when considering changes in droughts for higher changes in global warming (Figure 11.19;  
 6 Section 11.9). Seasonal projections of drought frequency for boreal winter (DJF) and summer (JJA), from  
 7 CMIP6 multimodel ensemble for 1.5°C, 2°C and 4°C global warming levels, show contrasting trends (Fig  
 8 11.19). In the boreal winter in the Northern Hemisphere, the areas affected by drying show high agreement  
 9 with those characterized by increase in meteorological drought projections (Chapter 8, Figure 8.14; Chapter  
 10 12, Figure 12.4). On the contrary, in the boreal summer the drought frequency increases worldwide in  
 11 comparison to meteorological drought projections, with large areas of the Northern Hemisphere displaying a  
 12 high signal to noise ratio (low spread between models). This stresses the dominant influence of ET (as a result  
 13 of increased AED) in intensifying agricultural and ecological droughts in the warm season in many locations,  
 14 including mid- to high latitudes.

15  
 16 Increased soil moisture limitation and associated changes in droughts are projected to lead to increased  
 17 vegetation stress affecting the global land carbon sink in ESM projections (Green et al., 2019), with  
 18 implications for projected global warming (Cross-Chapter Box 5). There is *high confidence* that the global  
 19 land sink will become less efficient due to soil moisture limitations and associated agricultural and  
 20 ecological drought conditions in some regions in higher emission scenarios specially under global warming  
 21 levels above 4°C ; however, there is *low confidence* on how these water cycle feedbacks will play out in  
 22 lower emission scenarios (at 2°C global warming or lower) (Cross-Chapter Box5.1).

23  
 24  
 25 **[START FIGURE 11.18 HERE]**

26  
 27 **Figure 11.18:** Projected changes in the intensity (a) and frequency (b) of drought under 1°C, 1.5°C, 2°C, 3°C, and 4°C  
 28 global warming levels relative to the 1850-1900 baseline. Summaries are computed for the AR6 regions  
 29 in which there is at least medium confidence in increase in agriculture/ ecological drought at the 2°C  
 30 warming level (“drying regions”), including W. North-America, C. North-America, N. Central-America,  
 31 S. Central-America, N. South-America, N. E. South-America, South-American-Monsoon, S.W.South-  
 32 America, S.South-America, West & Central-Europe, Mediterranean, W.Southern-Africa, E.Southern-  
 33 Africa, Madagascar, E.Australia, S.Australia (c). A drought event is defined as a 10-year drought event  
 34 whose annual mean soil moisture was below its 10th percentile from the 1850-1900 base period. For each  
 35 box plot, the horizontal line and the box represent the median and central 66% uncertainty range,  
 36 respectively, of the frequency or the intensity changes across the multi-model ensemble, and the whiskers  
 37 extend to the 90% uncertainty range. The line of zero in (a) indicates no change in intensity, while the  
 38 line of one in (b) indicates no change in frequency. The results are based on the multi-model ensemble  
 39 estimated from simulations of global climate models contributing to the sixth phase of the Coupled Model  
 40 Intercomparison Project (CMIP6) under different SSP forcing scenarios. Intensity changes in (a) are  
 41 expressed as standard deviations of the interannual variability in the period 1850-1900 of the  
 42 corresponding model. For details on the methods see Supplementary Material 11.SM.2. Further details on  
 43 data sources and processing are available in the chapter data table (Table 11.SM.9).

44  
 45 **[END FIGURE 11.18 HERE]**

#### 46 47 48 *11.6.5.4 Hydrological deficits*

49  
 50 Some studies support wetting tendencies as a response to a warmer climate when considering globally-  
 51 averaged changes in runoff over land (Roderick et al., 2015; Greve et al., 2017; Yang et al., 2018e), and  
 52 streamflow projections respond to enhanced CO<sub>2</sub> concentrations in CMIP5 models (Yang et al., 2019).  
 53 Nevertheless, when focusing regionally on low-runoff periods, model projections also show an increase of  
 54 hydrological droughts in large world regions (Wanders and Van Lanen, 2015; Dai et al., 2018; Vicente-  
 55 Serrano et al., 2020a). In general, the frequency of hydrological deficits is projected to increase over most of  
 56 the continents, although with regionally and seasonally differentiated effects (Section 11.9), with *medium*  
 57 *confidence* of increase in the following AR6 regions: WCE, MED, SAU, WCA, WNA, SCA, NSA, SAM,

**Do Not Cite, Quote or Distribute**

1 SWS, SSA, WSAF, ESAF and MDG (Section 11.9; Cook et al., 2019; Forzieri et al., 2014; Giuntoli et al.,  
2 2015; Marx et al., 2018; Prudhomme et al., 2014; Roudier et al., 2016; Wanders and Van Lanen, 2015; Zhao  
3 et al., 2020). There are, however, large uncertainties related to the hydrological/impact model used  
4 (Prudhomme et al., 2014; Schewe et al., 2014; Gosling et al., 2017), limited signal-to-noise ratio (due to  
5 model spread) in several regions (Giuntoli et al., 2015), and also uncertainties in the projection of future  
6 human activities including water demand and land cover changes, which may represent more than 50% of  
7 the projected changes in hydrological droughts in some regions (Wanders and Wada, 2015).

8  
9 Regions dependent on mountainous snowpack as a temporary reservoir may be affected by severe  
10 hydrological droughts in a warmer world. In the southern European Alps, both winter and summer low flows  
11 are projected to be more severe, with a 25% decrease in the 2050s (Vidal et al., 2016). In the western United  
12 States, a 22% reduction in winter snow water equivalent is projected at around 2°C of global warming with a  
13 further decrease of a 70% reduction at 4°C global warming (Rhoades et al., 2018). This decline would cause  
14 less predictable hydrological droughts in snowmelt-dominated areas of North America (Livneh and Badger,  
15 2020). The exact magnitude of the influence of higher temperatures on snow-related droughts is, however,  
16 difficult to estimate (Mote et al., 2016), since the streamflow changes could affect the timing of peak  
17 streamflows but not necessarily their magnitude. In addition, projected changes in hydrological droughts  
18 downstream of declining glaciers can be very complex to assess (Chapter 9, see also SROCC).

#### 19 20 21 *11.6.5.5 Atmospheric-based drought indices*

22  
23 Studies show a stronger drying in projections based on atmospheric-based drought indices compared to ESM  
24 projections of changes in soil moisture (Berg and Sheffield, 2018) and runoff (Yang et al., 2019). It has been  
25 suggested that this difference is due to physiological CO<sub>2</sub> effects (Greve et al., 2019; Lemordant et al., 2018;  
26 Milly and Dunne, 2016; Roderick et al., 2015; Scheff, 2018; Swann, 2018; Swann et al., 2016; Yang et al.,  
27 2020; Section 11.6.5.2). Nonetheless, there is evidence that differences in projections between atmospheric-  
28 based drought indices and water-balance metrics from ESMs are not alone due to CO<sub>2</sub>-plant effects (Berg et  
29 al., 2016; Scheff et al., 2021), and can be also related to the fact that AED is an upper bound for ET in dry  
30 regions and conditions (Section 11.6.1.2) and that soil moisture stress limits increases in ET in projections  
31 (Berg et al., 2016; Zhou et al., 2021; Section 11.6.5.2). Atmospheric-based indices show in general more  
32 drying than total column soil moisture (Berg and Sheffield, 2018; Cook et al., 2020; Scheff et al., 2021), but  
33 are more consistent with projected increases in surface soil moisture deficits (Dirmeyer et al., 2013; Dai et  
34 al., 2018; Lu et al., 2019; Cook et al., 2020; Vicente-Serrano et al., 2020a).

35  
36 Atmospheric-based drought indices are not metrics of soil moisture or runoff (11.6.1.5) so their projections  
37 may not necessarily reflect the same trend of online simulated soil moisture and runoff. Independently of  
38 effects on the land water balance, atmospheric-based drought indices will reflect the potential vegetation  
39 stress resulting from deficits between available water and enhanced AED, even in conditions with no or only  
40 low ET. Under dry conditions, the enhanced AED associated to the human forcing would increase plant  
41 water stress (Brodribb et al., 2020), with effects on widespread forest dieback and mortality (Anderegg et al.,  
42 2013; Williams et al., 2013; Allen et al., 2015; McDowell and Allen, 2015; McDowell et al., 2016, 2020),  
43 and stronger risk of megafires (Flannigan et al., 2016; Podschwit et al., 2018; Clarke and Evans, 2019;  
44 Varela et al., 2019). For these reasons, there is *high confidence* that the future projections of enhanced  
45 drought severity showed by the PDSI-PM and the SPEI-PM are representative of more frequent and severe  
46 plant stress episodes and more severe agricultural and ecological drought impacts in some regions.

47  
48 Global tendencies towards more severe and frequent agricultural and ecological drought conditions are  
49 identified in future projections when focusing on atmospheric-based drought indices such as the PDSI-PM or  
50 the SPEI-PM. They expand the spatial extent of drought conditions compared to meteorological drought to  
51 most of North America, Europe, Africa, Central and East Asia and southern Australia (Cook et al., 2014a;  
52 Chen and Sun, 2017b, 2017a; Zhao and Dai, 2017; Gao et al., 2017b; Lehner et al., 2017; Dai et al., 2018;  
53 Naumann et al., 2018; Potopová et al., 2018; Vicente-Serrano et al., 2020a; Gu et al., 2020; Dai, 2021).  
54 Projections in PDSI-PM and SPEI-PM are used in complement to changes in total soil moisture for the  
55 assessed projected changes in agricultural and ecological drought (Section 11.9).

### 11.6.5.6 Synthesis for different drought types

The tables in Section 11.9 provide assessed projected changes in meteorological drought, agricultural and ecological drought, and hydrological droughts. The assessment shows that several regions will be affected by more severe agricultural and ecological droughts even if global warming is stabilized at well below 2°C, and 1.5°C, within the bounds of the Paris Agreement (*high confidence*). The most affected regions include WCE, MED, EAU, SAU, SCA, NSA, SAM, SWS, SSA, NCA, CAN, WSAF, ESAF and MDG (*medium confidence*). At 4°C of global warming, even more regions would be affected by agricultural and ecological droughts (WCE, MED, CAU, EAU, SAU, WCA, EAS, SCA, CAR, NSA, NES, SAM, SWS, SSA, NCA, CAN, ENA, WNA, WSAF, ESAF and MDG). NEAF, SAS are also projected to experience less agricultural and ecological drought with global warming (*medium confidence*). Projected changes in meteorological droughts are overall less extended but also affect several AR6 regions, at 1.5°C and 2°C (MED, EAU, SAU, SCA, NSA, NCA, WSAF, ESAF, MDG) and 4°C of global warming (WCE, MED, EAU, SAU, SEA, SCA, CAR, NWS, NSA, NES, SAM, SWS, SSA, NCA, ENA, WAF, WSAF, ESAF, MDG). Several regions are also projected to be affected by more hydrological droughts at 1.5°C and 2°C (WCE, MED, WNA, WSAF, ESAF) and 4°C of global warming (NEU, WCE, EEU, MED, SAU, WCA, SCA, NSA, SAM, SWS, SSA, WNA, WSAF, ESAF, MDG). To illustrate the changes in both intensity and frequency of drought in the regions where strongest changes are projected, Figure 11.18 displays changes in the intensity and frequency of soil moisture drought under different global warming levels (1.5°C, 2°C, 4°C) relative to the 1851-1900 baseline based on CMIP6 simulations under different SSP forcing scenarios. The 90% uncertainty ranges for the projected changes in both intensity and frequency are above zero, indicating significant increase in both intensity and frequency of drought in these regions as whole.

In summary, the land area affected by increasing drought frequency and severity expands with increasing global warming (*high confidence*). New evidence strengthens the SR15 conclusion that even relatively small incremental increases in global warming (+0.5°C) cause a worsening of droughts in some regions (*high confidence*). Several regions will be affected by more frequent and severe agricultural and ecological droughts even if global warming is stabilized at 1.5-2°C (*high confidence*). The most affected regions include WCE, MED, EAU, SAU, SCA, NSA, SAM, SWS, SSA, NCA, CAN, WSAF, ESAF and MDG (*medium confidence*). At 4°C of global warming, even more regions would be affected by agricultural and ecological droughts (WCE, MED, CAU, EAU, SAU, WCA, EAS, SCA, CAR, NSA, NES, SAM, SWS, SSA, NCA, CAN, ENA, WNA, WSAF, ESAF and MDG). Some regions are also projected to experience less agricultural and ecological drought with global warming (*medium confidence*; (NEAF, SAS)). There is *high confidence* that the projected increases in agricultural and ecological droughts are strongly affected by AED increases in a warming climate, although ET increases are projected to be smaller than those in AED due to soil moisture limitations and CO<sub>2</sub> effects on leaf stomatal conductance. Enhanced atmospheric CO<sub>2</sub> concentrations lead to enhanced water-use efficiency in plants (*medium confidence*), but there is *low confidence* that it can ameliorate agricultural and ecological droughts, or hydrological droughts, at higher global warming levels characterized by limited soil moisture and enhanced AED.

Projected changes in meteorological droughts are overall less extended than for agricultural and ecological droughts, but also affect several AR6 regions, even at 1.5°C and 2°C of global warming. Several regions are also projected to be more strongly affected by hydrological droughts with increasing global warming (NEU, WCE, EEU, MED, SAU, WCA, SCA, NSA, SAM, SWS, SSA, WNA, WSAF, ESAF, MDG). Increased soil moisture limitation and associated changes in droughts are projected to lead to increased vegetation stress in many regions, with implications for the global land carbon sink (CC-Box 5). There is *high confidence* that the global land sink will become less efficient due to soil moisture limitations and associated drought conditions in some regions in higher emission scenarios specially under global warming levels above 4°C ; however, there is *low confidence* on how these water cycle feedbacks will play out in lower emission scenarios (at 2°C global warming or lower) (Cross-Chapter Box5.1).

[START FIGURE 11.19 HERE]

1  
2 **Figure 11.19:** Projected changes in (a-c) the number of consecutive dry days (CDD), (d-f) annual mean soil moisture  
3 over the total column, and (g-l) the frequency and intensity of one-in-ten year soil moisture drought for  
4 the June-to-August and December-to-February seasons at 1.5°C, 2°C, and 4°C of global warming  
5 compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-model  
6 ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The numbers in  
7 the top right indicate the number of simulations included. Uncertainty is represented using the simple  
8 approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models agree on sign  
9 of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on  
10 sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box  
11 Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources  
12 and processing are available in the chapter data table (Table 11.SM.9).

13  
14  
15 **[END FIGURE 11.19 HERE]**

## 16 17 18 **11.7 Extreme storms**

19  
20 Extreme storms, such as tropical cyclones (TCs), extratropical cyclones (ETCs), and severe convective  
21 storms often have substantial societal impacts. Quantifying the effect of climate change on extreme storms is  
22 challenging, partly because extreme storms are rare, short-lived, and local, and individual events are largely  
23 influenced by stochastic variability. The high degree of random variability makes detection and attribution of  
24 extreme storm trends more uncertain than detection and attribution of trends in other aspects of the  
25 environment in which the storms evolve (e.g., larger-scale temperature trends). Projecting changes in  
26 extreme storms is also challenging because of constraints in the models' ability to accurately represent the  
27 small-scale physical processes that can drive these changes. Despite the challenges, progress has been made  
28 since AR5.

29  
30 SREX (Chapter 3) concluded that there is *low confidence* in observed long-term (40 years or more) trends in  
31 TC intensity, frequency, and duration, and any observed trends in phenomena such as tornadoes and hail; it  
32 is *likely* that extratropical storm tracks have shifted poleward in both the Northern and Southern  
33 Hemispheres and that heavy rainfalls and mean maximum wind speeds associated with TCs will increase  
34 with continued greenhouse gas (GHG) warming; it is *likely* that the global frequency of TCs will either  
35 decrease or remain essentially unchanged, while it is *more likely than not* that the frequency of the most  
36 intense storms will increase substantially in some ocean basins; there is *low confidence* in projections of  
37 small-scale phenomena such as tornadoes and hail storms; and there is *medium confidence* that there will be  
38 a reduced frequency and a poleward shift of mid-latitude cyclones due to future anthropogenic climate  
39 change.

40  
41 Since SREX, several IPCC assessments also assessed storms. AR5 (Chapter 2, Hartmann et al., 2013)  
42 assessment with *low confidence* observed long-term trends in TC metrics, but revised the statement from  
43 SREX to state that it is *virtually certain* that there are increasing trends in North Atlantic TC activity since  
44 the 1970s, with *medium confidence* that anthropogenic aerosol forcing has contributed to these trends. AR5  
45 concluded that it is *likely* that TC precipitation and mean intensity will increase and *more likely than not* that  
46 the frequency of the strongest storms increases with continued GHG warming. *Confidence* in projected  
47 trends in overall TC frequency remained *low*. *Confidence* in observed and projected trends in hail storm and  
48 tornado events also remained *low*. SROCC (Chapter 6, Collins et al., 2019) assessed past and projected TCs  
49 and ETCs supporting the conclusions of AR5 with some additional detail. Literature subsequent to AR5 adds  
50 support to the likelihood of increasing trends in TC intensity, precipitation, and frequency of the most intense  
51 storms, while some newer studies have added uncertainty to projected trends in overall frequency. A  
52 growing body literature since AR5 on the poleward migration of TCs led to a new assessment in SROCC of  
53 *low confidence* that the migration in the western North Pacific represents a detectable climate change  
54 contribution from anthropogenic forcing. SR15 (Chapter 3, Hoegh-Guldberg et al., 2018) essentially  
55 confirmed the AR5 assessment of TCs and ETCs, adding that heavy precipitation associated with TCs is  
56 projected to be higher at 2°C compared to 1.5°C global warming (*medium confidence*).

1  
2 SREX, AR5, SROCC, and SR15, do not provide assessments of the atmospheric rivers and SROCC and  
3 SR15 do not assess severe convective storms and extreme winds. This section assesses the state of  
4 knowledge on the four phenomena of TCs, ETCs, severe convective storms, and extreme winds.  
5 Atmospheric rivers are addressed in Chapter 8. In this respect, this assessment closely mirrors the SROCC  
6 assessment of TCs and ETCs, while updating SREX and AR5 assessments of severe convective storms and  
7 extreme winds.  
8  
9

## 10 **11.7.1 Tropical cyclones**

### 11 *11.7.1.1 Mechanisms and drivers*

12  
13  
14 The genesis, development, and tracks of TCs depend on conditions of the larger-scale circulations of the  
15 atmosphere and ocean (Christensen et al., 2013). Large-scale atmospheric circulations (Annex VI), such as  
16 the Hadley and Walker circulations and the monsoon circulations, and internal variability acting on various  
17 time-scales, from intra-seasonal (e.g., the Madden-Julian and Boreal Summer Intraseasonal oscillations  
18 (MJO, BSISO), and equatorial waves) and inter-annual (e.g., the El Niño-Southern Oscillation (ENSO) and  
19 Pacific and Atlantic Meridional Modes (PMM, AMM)), to inter-decadal (e.g., Atlantic Multidecadal  
20 Variability and Pacific Decadal Variability (PDV)) can all significantly affect TCs. This broad range of  
21 natural variability makes detection of anthropogenic effects difficult, and uncertainties in the projected  
22 changes of these modes of variability increase uncertainty in the projected changes in TC activity. Aerosol  
23 forcing also affects SST patterns and cloud microphysics, and it is *likely* that observed changes in TC activity  
24 are partly caused by changes in aerosol forcing (Evan et al., 2011; Ting et al., 2015; Sobel et al., 2016, 2019;  
25 Takahashi et al., 2017; Zhao et al., 2018; Reed et al., 2019). Among possible changes from these drivers,  
26 there is *medium confidence* that the Hadley cell has widened and will continue to widen in the future  
27 (Chapter 2.3, 3.3, 4.5). This *likely* causes latitudinal shifts of TC tracks (Sharmila and Walsh, 2018).  
28 Regional TC activity changes are also strongly affected by projected changes in SST warming patterns  
29 (Yoshida et al., 2017), which are highly uncertain (Chapter 4, 9).  
30

### 31 *11.7.1.2 Observed trends*

32  
33  
34 Identifying past trends in TC metrics remains a challenge due to the heterogeneous character of the historical  
35 instrumental data, which are known as “best-track” data (Schreck et al., 2014). There is *low confidence* in  
36 most reported long-term (multidecadal to centennial) trends in TC frequency- or intensity-based metrics due  
37 to changes in the technology used to collect the best-track data. This should not be interpreted as implying  
38 that no physical (real) trends exist, but rather as indicating that either the quality or the temporal length of the  
39 data is not adequate to provide robust trend detection statements, particularly in the presence of multidecadal  
40 variability.  
41

42 There are previous and ongoing efforts to homogenize the best-track data (Elsner et al., 2008; Kossin et al.,  
43 2013, 2020; Choy et al., 2015; Landsea, 2015; Emanuel et al., 2018) and there is substantial literature that  
44 finds positive trends in intensity-related metrics in the best-track during the “satellite period”, which is  
45 generally limited to the past ~40 years (Kang and Elsner, 2012; Kishtawal et al., 2012; Kossin et al., 2013,  
46 2020; Mei and Xie, 2016; Zhao et al., 2018; Tauvale and Tsuboki, 2019). When best-track trends are tested  
47 using homogenized data, the intensity trends generally remain positive, but are smaller in amplitude (Kossin  
48 et al., 2013; Holland and Bruyère, 2014). Kossin et al. (2020) extended the homogenized TC intensity record  
49 to the period 1979–2017 and identified significant global increases in major TC exceedance probability of  
50 about 6% per decade. In addition to trends in TC intensity, there is evidence that TC intensification rates and  
51 the frequency of rapid intensification events have increased within the satellite era (Kishtawal et al. 2012;  
52 Balaguru et al., 2018; Bhatia et al., 2018). The increase in intensification rates is found in the best-track as  
53 well as the homogenized intensity data.  
54

55 A subset of the best-track data corresponding to hurricanes that have directly impacted the United States

1 since 1900 is considered to be reliable, and shows no trend in the frequency of U.S. landfall events (Knutson  
 2 et al., 2019). However, in this period since 1900, an increasing trend in normalized U.S. hurricane damage,  
 3 which accounts for temporal changes in exposed wealth (Grinsted et al., 2019), and a decreasing trend in TC  
 4 translation speed over the U.S. (Kossin, 2019) have been identified. A similarly reliable subset of the data  
 5 representing TC landfall frequency over Australia shows a decreasing trend in eastern Australia since the  
 6 1800s (Callaghan and Power, 2011), as well as in other parts of Australia since 1982 (Chand et al., 2019;  
 7 Knutson et al., 2019), and a paleoclimate proxy reconstruction shows that recent levels of TC interactions  
 8 along parts of the Australian coastline are the lowest in the past 550–1,500 years (Haig et al., 2014). Existing  
 9 TC datasets show substantial interdecadal variations in basin-wide TC frequency and intensity in the western  
 10 North Pacific, but a statistically significant northwestward shift in the western North Pacific TC tracks since  
 11 the 1980s (Lee et al., 2020b). In the case of the North Indian Ocean, analyses of trends are highly dependent  
 12 on the details of each analysis (e.g., pre- and/or post-monsoon season period, or Bay of Bengal and/or  
 13 Arabian Sea region). The most consistent trends are an increase in the occurrence of the most intense TCs  
 14 and a decrease in the overall TC frequency, in particular in the Bay of Bengal (Sahoo and Bhaskaran, 2016;  
 15 Balaji et al., 2018; Singh et al., 2019; Baburaj et al., 2020). In the South Indian Ocean (SIO), an increase in  
 16 the occurrence of the most intense TCs has been noted, however there are well-known data quality issues  
 17 there (Kuleshov et al., 2010; Fitchett, 2018). When the SIO data are homogenized, a significant increase is  
 18 found in the fractional proportion of global category 3-5 TC estimates to all category 1-5 estimates (Kossin  
 19 et al., 2020).

20  
 21 As with all confined regional analyses of TC frequency, it is generally unclear whether any identified  
 22 changes are due to a basin-wide change in TC frequency, or to systematic track shifts (or both). From an  
 23 impacts perspective, however, these changes over land are highly relevant and emphasize that large-scale  
 24 modifications in TC behaviour can have a broad spectrum of impacts on a regional scale.

25  
 26 Subsequent to AR5, two metrics that are argued to be comparatively less sensitive to data issues than  
 27 frequency- and intensity-based metrics have been analysed. Trends in these metrics have been identified over  
 28 the past ~70 years or more (Knutson et al., 2019). The first metric, the mean latitude where TCs reach their  
 29 peak intensity, exhibits a global and regional poleward migration during the satellite period (Kossin et al.,  
 30 2014). The poleward migration can influence TC hazard exposure and risk (Kossin et al., 2016a) and is  
 31 consistent with the independently-observed expansion of the tropics (Lucas et al., 2014). The migration has  
 32 been linked to changes in the Hadley circulation (Altman et al., 2018; Sharmila and Walsh, 2018; Studholme  
 33 and Gulev, 2018). The migration is also apparent in the mean locations where TCs exhibit eyes (Knapp et al.,  
 34 2018), which is when TCs are most intense. Part of the Northern Hemisphere poleward migration is due to  
 35 inter-basin changes in TC frequency (Kossin et al., 2014, 2016b, Moon et al., 2015, 2016) and the trends, as  
 36 expected, can be sensitive to the time period chosen (Tennille and Ellis, 2017; Kossin, 2018; Song and  
 37 Klotzbach, 2018) and to subsetting of the data by intensity (Zhan and Wang, 2017). The poleward migration  
 38 is particularly pronounced and well-documented in the western North Pacific basin (Kossin et al., 2016a;  
 39 Oey and Chou, 2016; Liang et al., 2017; Nakamura et al., 2017; Altman et al., 2018; Daloz and Camargo,  
 40 2018; Sun et al., 2019b; Lee et al., 2020b; Yamaguchi and Maeda, 2020a; Kubota et al., 2021).

41  
 42  
 43 **[START FIGURE 11.20 HERE]**

44  
 45 **Figure 11.20:** Summary schematic of past and projected changes in tropical cyclone (TC), extratropical cyclone (ETC),  
 46 atmospheric river (AR), and severe convective storm (SCS) behaviour. Global changes (blue shading)  
 47 from top to bottom: 1) Increased mean and maximum rain-rates in TCs, ETCs, and ARs [past (*low*  
 48 *confidence* due to lack of reliable data) & projected (*high confidence*)]. 2) Increased proportion of  
 49 stronger TCs [past (*medium confidence*) & projected (*high confidence*)]. 3) Decrease or no change in  
 50 global frequency of TC genesis [past (*low confidence* due to lack of reliable data) & projected (*medium*  
 51 *confidence*)]. 4) Increased and decreased ETC wind-speed, depending on the region, as storm-tracks  
 52 change [past (*low confidence* due to lack of reliable data) & projected (*medium confidence*)]. Regional  
 53 changes, from left to right: 1) Poleward TC migration in the western North Pacific and subsequent  
 54 changes in TC exposure [past (*medium confidence*) & projected (*medium confidence*)]. 2) Slowdown of  
 55 TC forward translation speed over the contiguous US and subsequent increase in TC rainfall [past  
 56 (*medium confidence*) & projected (*low confidence* due to lack of directed studies)]. 3) Increase in mean

1 and maximum SCS rain-rate and increase in springtime SCS frequency and season length over the  
2 contiguous US [past (*low confidence* due to lack of reliable data) & projected (*medium confidence*)].

3  
4 **[END FIGURE 11.20 HERE]**

5  
6  
7 A second metric that is argued to be comparatively less sensitive to data issues than frequency- and intensity-  
8 based metrics is TC translation speed (Kossin, 2018), which exhibits a global slowdown in the best-track  
9 data over the period 1949-2016. TC translation speed is a measure of the speed at which TCs move across  
10 the Earth's surface and is very closely related to local rainfall amounts (i.e., a slower translation speed causes  
11 greater local rainfall). TC translation speed also affects structural wind damage and coastal storm surge by  
12 changing the hazard event duration. The slowdown is observed in the best-track data from all basins except  
13 the Northern Indian Ocean and is also found in a number of regions where TCs interact directly with land.  
14 The slowing trends identified in the best-track data by Kossin (2018) have been argued to be largely due to  
15 data heterogeneity. Moon et al. (2019) and Lanzante (2019) provide evidence that meridional TC track shifts  
16 project onto the slowing trends and argue that these shifts are due to the introduction of satellite data. Kossin  
17 (2019) provides evidence that the slowing trend is real by focusing on Atlantic TC track data over the  
18 contiguous United States in the 118-year period 1900–2017, which are generally considered reliable. In this  
19 period, mean TC translation speed has decreased by 17%. The slowing TC translation speed is expected to  
20 increase local rainfall amounts, which would increase coastal and inland flooding. In combination with  
21 slowing translation speed, abrupt TC track direction changes – that can be associated with track “meanders”  
22 or “stalls” – have become increasingly common along the North American coast since the mid-20th century,  
23 leading to more rainfall in the region (Hall and Kossin, 2019).

24  
25 In summary, there is mounting evidence that a variety of TC characteristics have changed over various time  
26 periods. It is *likely* that the proportion of major TC intensities and the frequency of rapid intensification  
27 events have both increased globally over the past 40 years. It is *very likely* that the average location where  
28 TCs reach their peak wind-intensity has migrated poleward in the western North Pacific Ocean since the  
29 1940s. It is *likely* that TC translation speed has slowed over the U.S. since 1900.

30  
31  
32 *11.7.1.3 Model evaluation*

33  
34 Accurate projections of future TC activity have two principal requirements: accurate representation of  
35 changes in the relevant environmental factors (e.g., SSTs) that can affect TC activity, and accurate  
36 representation of actual TC activity in given environmental conditions. In particular, models' capacity to  
37 reproduce historical trends or interannual variabilities of TC activity is relevant to the confidence in future  
38 projections. One test of the models is to evaluate their ability to reproduce the dependency of the TC  
39 statistics in the different basins in the real world, in addition to their capability of reproducing atmospheric  
40 and ocean environmental conditions. For the evaluation of projections of TC-relevant environmental  
41 variables, AR5 confidence statements were based on global surface temperature and moisture, but not on the  
42 detailed regional structure of SST and atmospheric circulation changes such as steering flows and vertical  
43 shear, which affect characteristics of TCs (genesis, intensity, tracks, etc.). Various aspects of TC metrics are  
44 used to evaluate how capable models are of simulating present-day TC climatologies and variability (e.g. TC  
45 frequency, wind-intensity, precipitation, size, tracks, and their seasonal and interannual changes) (Camargo  
46 and Wing, 2016; Knutson et al., 2019, 2020; Walsh et al., 2015). Other examples of TC  
47 climatology/variability metrics are spatial distributions of TC occurrence and genesis (Walsh et al., 2015),  
48 seasonal cycles and interannual variability of basin-wide activity (Zhao et al., 2009; Shaevitz et al., 2014;  
49 Kodama et al., 2015; Murakami et al., 2015; Yamada et al., 2017) or landfalling activity (Lok and Chan,  
50 2018), as well as newly developed process-diagnostics designed specifically for TCs in climate models (Kim  
51 et al., 2018a; Wing et al., 2019; Moon et al., 2020).

52  
53 Confidence in the projection of intense TCs, such as those of Category 4-5, generally becomes higher as the  
54 resolution of the models becomes higher. CMIP5/6 class climate models (~100-200 km grid spacing) cannot  
55 simulate TCs of Category 4-5 intensity. They do simulate storms of relatively high vorticity that are at best

1 described as “TC-like”, but metrics like storm counts are highly dependent on tracking algorithms (Wehner  
2 et al., 2015; Zarzycki and Ullrich, 2017; Roberts et al., 2020a). High-resolution global climate models (~10-  
3 60 km grid spacing) as used in HighResMIP (Haarsma et al., 2016; Roberts et al., 2020a) begin to capture  
4 some structures of TCs more realistically, as well as produce intense TCs of Category 4-5 despite the effects  
5 of parameterized deep cumulus convection processes (Murakami et al., 2015; Wehner et al., 2015; Yamada  
6 et al., 2017; Roberts et al., 2018; Moon et al., 2020). Convection-permitting models (~1-10 km grid-  
7 spacing), such as used in some dynamical downscaling studies, provide further realism with capturing eye  
8 wall structures (Tsuboki et al., 2015). Model characteristics besides resolution, especially details of  
9 convective parameterization, can influence a model’s ability to simulate intense TCs (Reed and Jablonowski,  
10 2011; Zhao et al., 2012; He and Posselt, 2015; Kim et al., 2018a; Zhang and Wang, 2018; Camargo et al.,  
11 2020). However, models’ dynamical cores and other physics also affect simulated TC properties (Reed et al.,  
12 2015; Vidale et al., 2021). Both wide-area regional and global convection-permitting models without the  
13 need for parameterized convection are becoming more useful for TC regional model projection studies  
14 (Tsuboki et al., 2015; Kanada et al., 2017a; Gutmann et al., 2018) and global model projection studies (Satoh  
15 et al., 2015, 2017; Yamada et al., 2017), as they capture more realistic eye-wall structures of TCs (Kinter et  
16 al., 2013) and are becoming more useful for investigating changes in TC structures (Kanada et al., 2013;  
17 Yamada et al., 2017). Large ensemble simulations of global climate models with 60 km grid spacing provide  
18 TC statistics that allow more reliable detection of changes in the projections, which are not well captured in  
19 any single experiment (Yoshida et al., 2017; Yamaguchi et al., 2020). Variable resolution global models  
20 offer an alternative to regional models for individual TC or basin-wide simulations (Yanase et al., 2012;  
21 Zarzycki et al., 2014; Harris et al., 2016; Reed et al., 2020; Stansfield et al., 2020). Computationally less  
22 intense than equivalent uniform resolution global models, they also do not require lateral boundary  
23 conditions, thus reducing this source of error (Hashimoto et al., 2016). Confidence in the projection of TC  
24 statistics and properties is increased by the higher-resolution models with more realistic simulations.

25  
26 Operational forecasting models also reproduce TCs and their use for climate projection studies shows  
27 promise. However, there is limited application for future projections as they are specifically developed for  
28 operational purposes and TC climatology is not necessarily well evaluated. Intercomparison of operational  
29 models indicates that enhancement of horizontal resolution can provide more credible projections of TCs  
30 (Nakano et al., 2017). Likewise, high-resolution climate models show promise as TC forecast tools (Zarzycki  
31 and Jablonowski, 2015; Reed et al., 2020), further narrowing the continuum of weather and climate models  
32 and increasing confidence in projections of future TC behaviour. However, higher horizontal resolution does  
33 not necessarily lead to an improved TC climatology (Camargo et al., 2020).

34  
35 Atmosphere-ocean interaction is an important process in TC evolution. Atmosphere-ocean coupled models  
36 are generally better than atmosphere-only models at capturing realistic processes related to TCs (Murakami  
37 et al., 2015; Ogata et al., 2015, 2016; Zarzycki, 2016; Kanada et al., 2017b; Scoccimarro et al., 2017),  
38 although the basin-scale SST biases commonly found in atmosphere-ocean models can introduce substantial  
39 errors in the simulated TC number (Hsu et al., 2019). Higher-resolution ocean models improve the  
40 simulation of TCs by reducing the SST climatology bias (Li and Sriviver, 2018; Roberts et al., 2020a). Coarse  
41 resolution atmospheric models may degrade coupled model performance as well. For example, in a case  
42 study of Hurricane Harvey, Trenberth et al. (2018) suggested that the lack of realistic hurricane activity  
43 within coupled climate models hampers the models’ ability to simulate SST and ocean heat content and their  
44 changes.

45  
46 Even with higher-resolution atmosphere-ocean coupled models, TC projection studies still rely on  
47 assumptions in experimental design that introduce uncertainties. Computational constraints often limit the  
48 number of simulations, resulting in relatively small ensemble sizes and incomplete analyses of possible  
49 future SST magnitude and pattern changes (Zhao and Held, 2011; Knutson et al., 2013a). Uncertainties in  
50 aerosol forcing also are reflected in TC projection uncertainty (Wang et al., 2014).

51  
52 Regional climate models (RCM) with grid spacing around 15-50 km can be used to study the projection of  
53 TCs. RCMs are run with lateral and surface boundary conditions, which are specified by the atmospheric  
54 state and SSTs simulated by GCMs. Various combinations of the lateral and surface boundary conditions can  
55 be chosen for RCM studies, and uncertainties in the projection can be further examined in general. They are



1 used for studying changes in TC characteristics in a specific area, such as Vietnam (Redmond et al., 2015)  
2 and the Philippines (Gallo et al., 2019).

3  
4 Less computationally-expensive downscaling approaches that allow larger ensembles and long-term studies  
5 are also used in the projection of TCs (Emanuel et al., 2006; Lee et al., 2018a). A statistical–dynamical TC  
6 downscaling method requires assumptions of the rate of seeding of random initial disturbances, which are  
7 generally assumed to not change with climate change (Emanuel et al., 2008; Emanuel, 2013). The results  
8 with the downscaling approach might depend on the assumptions which are required for the simplification of  
9 the methods.

10  
11 In summary, various types of models are useful to study climate changes of TCs, and there is no unique  
12 solution for choosing a model type. However, higher-resolution models generally capture TC properties  
13 more realistically (*high confidence*). In particular, models with horizontal resolutions of 10–60 km are  
14 capable of reproducing strong TCs with Category 4–5 and those of 1–10 km are capable of the eyewall  
15 structure of TCs. Uncertainties in TC simulations come from details of the model configuration of both  
16 dynamical and physical processes. Models with realistic atmosphere–ocean interactions are generally better  
17 than atmosphere-only models at reproducing realistic TC evolutions (*high confidence*).

#### 18 19 20 11.7.1.4 Detection and attribution, event attribution

21  
22 There is general agreement in the literature that anthropogenic greenhouse gases and aerosols have  
23 measurably affected observed oceanic and atmospheric variability in TC-prone regions (see Chapter 3). This  
24 underpinned the SROCC assessment of *medium confidence* that humans have contributed to the observed  
25 increase in Atlantic hurricane activity since the 1970s (Chapter 5, Bindoff et al., 2013). Literature subsequent  
26 to the AR5 lends further support to this statement (Knutson et al., 2019). However, there is still no consensus  
27 on the relative magnitude of human and natural influences on past changes in Atlantic hurricane activity, and  
28 particularly on which factor has dominated the observed increase (Ting et al., 2015) and it remains uncertain  
29 whether past changes in Atlantic TC activity are outside the range of natural variability. A recent result using  
30 high-resolution dynamical model experiments suggested that the observed spatial contrast in TC trends  
31 cannot be explained only by multi-decadal natural variability, and that external forcing plays an important  
32 role (Murakami et al., 2020). Observational evidence for significant global increases in the proportion of  
33 major TC intensities (Kossin et al., 2020) is consistent with both theory and numerical modeling simulations,  
34 which generally indicate an increase in mean TC peak intensity and the proportion of very intense TCs in a  
35 warming world (Knutson et al., 2015, 2020, Walsh et al., 2015, 2016). In addition, high-resolution coupled  
36 model simulations provide support that natural variability alone is *unlikely* to explain the magnitude of the  
37 observed increase in TC intensification rates and upward TC intensity trend in the Atlantic basin since the  
38 early 1980s (Bhatia et al., 2019; Murakami et al. 2020).

39  
40 The cause of the observed slowdown in TC translation speed is not yet clear. Yamaguchi et al. (2020) used  
41 large ensemble simulations to argue that part of the slowdown is due to actual latitudinal shifts of TC tracks,  
42 rather than data artefacts, in addition to atmospheric circulation changes, while Zhang et al. (2020a) used  
43 large ensemble simulations to show that anthropogenic forcing can lead to a robust slowdown, particularly  
44 outside of the tropics at higher latitudes. Yamaguchi and Maeda (2020b) found a significant slowdown in the  
45 western North Pacific over the past 40 years and attributed the slowdown to a combination of natural  
46 variability and global warming. The slowing trend since 1900 over the U.S. is robust and significant after  
47 removing multidecadal variability from the time series (Kossin, 2019). Among the hypotheses discussed is  
48 the physical linkage between warming and slowing circulation (Held and Soden 2006, see also Section  
49 8.2.2.2), with expectations of Arctic amplification and weakening circulation patterns through weakening  
50 meridional temperature gradients (Coumou et al., 2018; see also Cross-Chapter Box 10.1), or through  
51 changes in planetary wave dynamics (Mann et al., 2017). The tropics expansion and the poleward shift of the  
52 mid-latitude westerlies associated with warming is also suggested for the reason of the slowdown (Zhang et  
53 al., 2020a). However, the connection of these mechanisms to the slowdown has not been robustly shown yet.  
54 Furthermore, slowing trends have not been unambiguously observed in circulation patterns that steer TCs  
55 such as the Walker and Hadley circulations (Section 2.3.1.4), although these circulations generally slow

1 down in numerical simulations under global warming (Sections 4.5.1.6 and 8.4.2.2).

2  
3 The observed poleward trend in western North Pacific TCs remains significant after accounting for the  
4 known modes of dominant interannual to decadal variability in the region (Kossin et al., 2016a), and is also  
5 found in CMIP5 model-simulated TCs (in the recent historical period 1980–2005), although it is weaker than  
6 observed and is not statistically significant (Kossin et al., 2016a). However, the trend is significant in 21st  
7 century CMIP5 projections under the RCP8.5 scenario, with a similar spatial pattern and magnitude to the  
8 past observed changes in that basin over the period 1945–2016, supporting a possible anthropogenic GHG  
9 contribution to the observed trends (Knutson et al., 2019; Kossin et al., 2016a).

10  
11 The recent active TC seasons in some basins have been studied to determine whether there is anthropogenic  
12 influence. For 2015, Murakami et al. (2017) explored the unusually high TC frequency near Hawaii and in  
13 the eastern Pacific basin. Zhang et al. (2016) considered unusually high Accumulated Cyclone Energy  
14 (ACE) in the western North Pacific. Yang et al. (2018) and Yamada et al. (2019) looked at TC intensification  
15 in the western North Pacific. These studies suggest that the anomalous TC activity in 2015 was not solely  
16 explained by the effect of an extreme El Niño (see BOX 11.3), that there was also an anthropogenic  
17 contribution, mainly through the effects of SSTs in subtropical regions. In the post-monsoon seasons of 2014  
18 and 2015, tropical storms with lifetime maximum winds greater than  $46 \text{ m s}^{-1}$  were first observed over the  
19 Arabian Sea, and Murakami et al. (2017b) showed that the probability of late-season severe tropical storms is  
20 increased by anthropogenic forcing compared to the preindustrial era. Murakami et al. (2018) concluded that  
21 the active 2017 Atlantic hurricane season was mainly caused by pronounced SSTs in the tropical North  
22 Atlantic and that these types of seasonal events will intensify with projected anthropogenic forcing. The  
23 trans-basin SST change, which might be driven by anthropogenic aerosol forcing, also affects TC activity.  
24 Takahashi et al. (2017) suggested that a decrease in sulfate aerosol emissions caused about half of the  
25 observed decreasing trends in TC genesis frequency in the south-eastern region of the western North Pacific  
26 during 1992–2011.

27  
28 Event attribution is used in case studies of TCs to test whether the severities of recent intense TCs are  
29 explained without anthropogenic effects. In a case study of Hurricane Sandy (2012), Lackmann (2015) found  
30 no statistically significant impact of anthropogenic climate change on storm intensity, while projections in a  
31 warmer world showed significant strengthening. On the other hand, Magnusson et al. (2014) found that in  
32 ECMWF simulations, the simulated cyclone depth and intensity, as well as precipitation, were larger when  
33 the model was driven by the warmer actual SSTs than the climatological average SSTs. In super typhoon  
34 Haiyan, which struck the Philippines on 8 November 2013, Takayabu et al. (2015) took an event attribution  
35 approach with cloud system-resolving ( $\sim 1\text{km}$ ) downscaling ensemble experiments to evaluate the  
36 anthropogenic effect on typhoons, and showed that the intensity of the simulated worst case storm in the  
37 actual conditions was stronger than that in a hypothetical condition without historical anthropogenic forcing  
38 in the model. However, in a similar approach with two coarser parameterized convection models, Wehner et  
39 al. (2018) found conflicting human influences on Haiyan's intensity. Patricola and Wehner (2018) found  
40 little evidence of an attributable change in intensity of hurricanes Katrina (2005), Irma (2017), and Maria  
41 (2017) using a regional climate model configured between 3 and 4.5 km resolution. They did, however, find  
42 attributable increases in heavy precipitation totals. These results imply that higher resolution, such as in a  
43 convective permitting 5-km or less mesh model, is required to obtain a robust anthropogenic intensification  
44 of a strong TC by simulating realistic rapid intensification of a TC (Kanada and Wada, 2016; Kanada et al.,  
45 2017a), and that whether the intensification of TCs can be attributed to the recent warming depends on the  
46 case.

47  
48 The dominant factor in the extreme rainfall amounts during Hurricane Harvey's passage onto the U.S. in  
49 2017 was its slow translation speed. But studies published after the event have argued that anthropogenic  
50 climate change contributed to an increase in rain rate, which compounded the extreme local rainfall caused  
51 by the slow translation. Emanuel (2017) used a large set of synthetically-generated storms and concluded  
52 that the occurrence of extreme rainfall as observed in Harvey was substantially enhanced by anthropogenic  
53 changes to the larger-scale ocean and atmosphere characteristics. Trenberth et al. (2018) linked Harvey's  
54 rainfall totals to the anomalously large ocean heat content from the Gulf of Mexico. van Oldenborgh et al.  
55 (2017) and Risser and Wehner (2017) applied extreme value analysis to extreme rainfall records in the

1 Houston, Texas region and both attributed large increases to climate change. Large precipitation increases  
2 during Harvey due to global warming were also found using climate models (van Oldenborgh et al., 2017;  
3 Wang et al., 2018b). Harvey precipitation totals were estimated in these papers to be 3 to 10 times more  
4 probable due to climate change. A best estimate from a regional climate and flood model is that urbanization  
5 increased the risk of the Harvey flooding by a factor of 21 (Zhang et al., 2018c). Anthropogenic effects on  
6 precipitation increases were also predicted in advance from a forecast model for Hurricane Florence in 2018  
7 (Reed et al., 2020).

8  
9 In summary, it is *very likely* that the recent active TC seasons in the North Atlantic, the North Pacific, and  
10 Arabian basins cannot be explained without an anthropogenic influence. The anthropogenic influence on  
11 these changes is principally associated to aerosol forcing, with stronger contributions to the response in the  
12 North Atlantic. It is *more likely than not* that the slowdown of TC translation speed over the USA has  
13 contributions from anthropogenic forcing. It is *likely* that the poleward migration of TCs in the western  
14 North Pacific and the global increase in TC intensity rates cannot be explained entirely by natural variability.  
15 Event attribution studies of specific strong TCs provide limited evidence for anthropogenic effects on TC  
16 intensifications so far, but *high confidence* for increases in TC heavy precipitation. There is *high confidence*  
17 that anthropogenic climate change contributed to extreme rainfall amounts during Hurricane Harvey (2017)  
18 and other intense TCs.

#### 21 11.7.1.5 Projections

22  
23 A summary of studies on TC projections for the late 21st century, particularly studies since AR5, is given by  
24 Knutson et al. (2020), which is an assessment report mandated by the World Meteorological Organization  
25 (WMO). Studies subsequent to Knutson et al. (2020) are generally consistent and the confidence assessments  
26 here closely follow theirs (Cha et al., 2020), although there are some differences due to the different  
27 confidence calibrations between the IPCC and WMO reports.

28  
29 There is not an established theory for the drivers of future changes in the frequency of TCs. Most, but not all,  
30 high-resolution global simulations project significant reductions in the total number of TCs, with the bulk of  
31 the reduction at the weaker end of the intensity spectrum as the climate warms (Knutson et al., 2020). Recent  
32 exceptions based on high-resolution coupled model results are noted in Bhatia et al. (2018) and Vecchi et al.  
33 (2019). Vecchi et al. (2019) showed that the representation of synoptic-scale seeds for TC genesis in their  
34 high-resolution model causes different projections of global TC frequency, and there is evidence for a  
35 decrease in seeds in some projected TC simulations (Sugi et al., 2020). However, other research indicates  
36 that TC seeds are not an independent control on climatological TC frequency, rather the seeds covary with  
37 the large-scale controls on TCs (Patricola et al., 2018). While empirical genesis indices derived from  
38 observations and reanalysis describe well the observed subseasonal and interannual variability of current TC  
39 frequency (Camargo et al., 2007, 2009; Tippett et al., 2011; Menkes et al., 2012), they fail to predict the  
40 decreased TC frequency found in most high-resolution model simulations (Zhang et al., 2010; Camargo,  
41 2013; Wehner et al., 2015), as they generally project an increase as the climate warms. This suggests a  
42 limitation of the use of the empirical genesis indices for projections of TC genesis, in particular due to their  
43 sensitivity to the humidity variable considered in the genesis index for these projections (Camargo et al.,  
44 2014). In a different approach, a statistical-dynamical downscaling framework assuming a constant seeding  
45 rate with warming (Emanuel, 2013, 2021) exhibits increases in TC frequency consistent with genesis indices  
46 based projections, while downscaling with a different model leads to two different scenarios depending on  
47 the humidity variable considered (Lee et al., 2020a). This disparity in the sign of the projected change in  
48 global TC frequency and the difficulty in explaining the mechanisms behind the different signed responses  
49 further emphasizes the lack of process understanding of future changes in tropical cyclogenesis (Walsh et al.,  
50 2015; Hoogewind et al., 2020). Even within a single model, uncertainty in the pattern of future SST changes  
51 leads to large uncertainties (including the sign) in the projected change in TC frequency in individual ocean  
52 basins, although global TCs would appear to be less sensitive (Yoshida et al., 2017; Bacmeister et al., 2018).

53  
54 Changes in SST and atmospheric temperature and moisture play a role in tropical cyclogenesis (Walsh et al.,  
55 2015). Reductions in vertical convective mass flux due to increased tropical stability have been associated

1 with a reduction in cyclogenesis (Held and Zhao, 2011; Sugi et al., 2012). Satoh et al. (2015) further posits  
2 that the robust simulated increase in the number of intense TCs, and hence increased vertical mass flux  
3 associated with intense TCs, must lead to a decrease in overall TC frequency because of this association. The  
4 Genesis Potential Index can be modified to mimic the TC frequency decreases of a model by altering the  
5 treatment of humidity (Camargo et al., 2014), supporting the idea that increased mid-tropospheric saturation  
6 deficit (Emanuel et al., 2008) controls TC frequency, but the approach remains empirical. Other possible  
7 controlling factors, such as a decline in the number of seeds (held constant in Emanuel's downscaling  
8 approach, or dependent on the genesis index formulation in the approach proposed by Lee et al., 2020)  
9 caused by increased atmospheric stability have been proposed, but questioned as an important factor  
10 (Patricola et al., 2018). The resolution of atmospheric models affects the number of seeds, hence TC genesis  
11 frequency (Vecchi et al., 2019; Sugi et al., 2020; Yamada et al., 2021). The diverse and sometimes  
12 inconsistent projected changes in global TC frequency by high-resolution models indicate that better process  
13 understanding and improvement of the models are needed to raise confidence in these changes.  
14

15 Most TC-permitting model simulations (10-60 km or finer grid spacing) are consistent in their projection of  
16 increases in the proportion of intense TCs (Category 4-5), as well as an increase in the intensity of the  
17 strongest TCs defined by maximum wind speed or central pressure fall (Murakami et al., 2012; Tsuboki et  
18 al., 2015; Wehner et al., 2018a; Knutson et al., 2020). The general reduction in the total number of TCs,  
19 which is concentrated in storms weaker than or equal to Category 1, contributes to this increase. The models  
20 are somewhat less consistent in projecting an increase in the frequency of Category 4-5 TCs (Wehner et al.,  
21 2018a). The projected increase in the intensity of the strongest TCs is consistent with theoretical  
22 understanding (e.g., Emanuel, 1987) and observations (e.g., Kossin et al., 2020). For a 2°C global warming,  
23 the median proportion of Category 4-5 TCs increases by 13%, while the median global TC frequency  
24 decreases by 14%, which implies that the median of the global Category 4-5 TC frequency is slightly  
25 reduced by 1% or almost unchanged (Knutson et al., 2020). Murakami et al. (2020) projected a decrease in  
26 TC frequency over the coming century in the North Atlantic due to greenhouse warming, as consistent with  
27 Dunstone et al. (2013), and a reduction in TC frequency almost everywhere in the tropics in response to +1%  
28 CO<sub>2</sub> forcing; exceptions include the central North Pacific (Hawaii region), east of the Philippines in the  
29 North Pacific, and two relatively small regions in the northern Arabian Sea and Bay of Bengal. These  
30 projections can vary substantially between ocean basins, possibly due to differences in regional SST  
31 warming and warming patterns (Sugi et al., 2017; Yoshida et al., 2017; Bacmeister et al., 2018). A summary  
32 of projections of TC characteristics is schematically shown by Figure 11.20.  
33

34 The increase in global TC maximum surface wind speeds is about 5% for a 2°C global warming across a  
35 number of high-resolution multi-decadal studies (Knutson et al., 2020). This indicates the deepening in  
36 global TC minimum surface pressure under the global warming conditions. A regional cloud-permitting  
37 model study shows that the strongest TC in the western North Pacific can be as strong as 857 hPa in  
38 minimum surface pressure with a wind speed of 88 m s<sup>-1</sup> under warming conditions in 2074-2087 (Tsuboki  
39 et al., 2015). TCs are also measured by quantities such as Accumulated Cyclone Energy (ACE) and the  
40 power dissipation index (PDI), which conflate TC intensity, frequency, and duration (Murakami et al., 2014).  
41 Several TC modeling studies (Yamada et al., 2010; Kim et al., 2014a; Knutson et al., 2015) project little  
42 change or decreases in the globally-accumulated value of PDI or ACE, which is due to the decrease in the  
43 total number of TCs.  
44

45 A projected increase in global average TC rainfall rates of about 12% for a 2°C global warming is consistent  
46 with the Clausius-Clapeyron scaling of saturation specific humidity (Knutson et al., 2020). Increases  
47 substantially greater than Clausius-Clapeyron scaling are projected in some regions, which is caused by  
48 increased low-level moisture convergence due to projected TC intensity increases in those regions (Knutson  
49 et al., 2015; Phibbs and Toumi, 2016; Patricola and Wehner, 2018; Liu et al., 2019c). Projections of TC  
50 precipitation using large-ensemble experiments (Kitoh and Endo, 2019) show that the annual maximum 1-  
51 day precipitation total is projected to increase, except for the western North Pacific where there is only a  
52 small change or even a reduction is projected, mainly due to a projected decrease of TC frequency. They also  
53 show that the 10-year return value of extreme Rx1day associated with TCs will greatly increase in a region  
54 extending from Hawaii to the south of Japan. TC tracks and the location of topography relative to TCs  
55 significantly affect precipitation, thus in general, areas on the eastern and southern faces of mountains have

1 more impacts of TC precipitation changes (Hatsuzuka et al., 2020). Projection studies using variable-  
2 resolution models in the North Atlantic (Stansfield et al., 2020) indicate that TC-related precipitation rates  
3 within North Atlantic TCs and the amount of hourly precipitation due to TC are projected to increase by the  
4 end of the century compared to a historical simulation. However, the annual average TC-related Rx5day over  
5 the eastern United States is projected to decrease because of a decrease in landfalling TCs. RCM studies with  
6 around 25–50 km grid spacing are used to study projected changes in TCs. The projected changes of TCs in  
7 Southeast Asia simulated by RCMs are consistent with those of most global climate models, showing a  
8 decrease in TC frequency and an increase in the amount of TC-associated precipitation or an increase in the  
9 frequency of intense TCs (Redmond et al., 2015; Gallo et al., 2019).

10  
11 Projected changes in TC tracks or TC areas of occurrence in the late 21st century vary considerably among  
12 available studies, although there is better agreement in the western North Pacific. Several studies project  
13 either poleward or eastward expansion of TC occurrence over the western North Pacific region, and more TC  
14 occurrence in the central North Pacific (Yamada et al., 2017; Yoshida et al., 2017; Wehner et al., 2018a;  
15 Roberts et al., 2020b). The observed poleward expansion of the latitude of maximum TC intensity in the  
16 western North Pacific is consistently reproduced by the CMIP5 models and downscaled models and these  
17 models show further poleward expansion in the future; the projected mean migration rate of the mean  
18 latitude where TCs reach their lifetime-maximum intensity is  $0.2 \pm 0.1^\circ$  from CMIP5 model results, while it is  
19  $0.13 \pm 0.04^\circ$  from downscaled models in the western North Pacific (Kossin et al., 2014, 2016a). In the North  
20 Atlantic, while the location of TC maximum intensity does not show clear poleward migration  
21 observationally (Kossin et al., 2014), it tends to migrate poleward in projections (Garner et al., 2017). The  
22 poleward migration is less robust among models and observations in the Indian Ocean, eastern North Pacific,  
23 and South Pacific (e.g., Tauvale and Tsuboki, 2019; Ramsay et al. 2018; Cattiaux et al. 2020). There is  
24 presently no clear consensus in projected changes in TC translation speed (Knutson et al., 2020), although  
25 recent studies suggest a slowdown outside of the tropics (Kossin, 2019; Yamaguchi et al., 2020; Zhang et al.,  
26 2020a), but regionally there can even be an acceleration of the storms (Hassanzadeh et al., 2020).

27  
28 The spatial extent, or “size”, of the TC wind-field is an important determinant of storm surge and damage.  
29 No detectable anthropogenic influences on TC size have been identified to date, because TCs in observations  
30 vary in size substantially (Chan and Chan, 2015) and there is no definite theory on what controls TC size,  
31 although this is an area of active research (Chavas and Emanuel, 2014; Chan and Chan, 2018). However,  
32 projections by high-resolution models indicate future broadening of TC wind-fields when compared to TCs  
33 of the same categories (Yamada et al., 2017), while Knutson et al. (2015) simulates a reasonable interbasin  
34 distribution of TC size climatology, but projects no statistically significant change in global average TC size.  
35 A plausible mechanism is that as the tropopause height becomes higher with global warming, the eye wall  
36 areas become wider because the eye walls are inclined outward with height to the tropopause. This effect is  
37 only reproduced in high-resolution convection-permitting models capturing eye walls, and such modeling  
38 studies are not common. Moreover, the projected TC size changes are generally on the order of 10% or less,  
39 and these size changes are still highly variable between basins and studies. Thus, the projected change in  
40 both magnitude and sign of TC size is uncertain.

41  
42 The coastal effects of TCs depend on TC intensity, size, track, and translation speed. Projected increases in  
43 sea level, average TC intensity, and TC rainfall rates each generally act to further elevate future storm surge  
44 and fresh-water flooding (see Section 9.6.4.2). Changes in TC frequency could contribute toward increasing  
45 or decreasing future storm surge risk, depending on the net effects of changes in weaker vs stronger storms.  
46 Several studies (McInnes et al., 2014, 2016; Little et al., 2015; Garner et al., 2017; Timmermans et al., 2017,  
47 2018) have explored future projections of storm surge in the context of anthropogenic climate change with  
48 the influence of both sea level rise and future TC changes. Garner et al. (2017) investigated the near future  
49 changes in the New York City coastal flood hazard, and suggested a small change in storm-surge height  
50 because effects of TC intensification are compensated by the offshore shifts in TC tracks, but concluded that  
51 the overall effect due to the rising sea levels would increase the flood hazard. Future projection studies of  
52 storm surge in East Asia, including China, Japan and Korea, also indicate that storm surge due to TCs  
53 become more severe (Yang et al., 2018b; Mori et al., 2019, 2021; Chen et al., 2020c). For the Pacific islands,  
54 McInnes et al. (2014) found that the future projected increase in storm surge in Fiji is dominated by sea level  
55 rise, and projected TC changes make only a minor contribution. Among various storm surge factors, there is

1 *high confidence* that sea level rise will lead to a higher possibility of extreme coastal water levels in most  
2 regions, with all other factors assumed equal.

3  
4 In the North Atlantic, vertical wind shear, which inhibits TC genesis and intensification, varies in a quasi-  
5 dipole pattern with one center of action in the tropics and another along the U.S. southeast coast (Vimont and  
6 Kossin, 2007). This pattern of variability creates a protective barrier of high shear along the U.S. coast  
7 during periods of heightened TC activity in the tropics (Kossin, 2017), and appears to be a natural part of the  
8 Atlantic ocean-atmosphere climate system (Ting et al., 2019). Greenhouse gas forcing in CMIP5 and the  
9 Community Earth System Model Large Ensemble (CESM-LE; Kay et al., 2015) simulations, however,  
10 erodes the pattern and degrades the natural shear barrier along the U.S. coast. Following the Representative  
11 Concentration Pathway 8.5 (RCP8.5) emission scenario, the magnitude of the erosion of the barrier equals  
12 the amplitude of past natural variability (time of emergence) by the mid-twenty-first century (Ting et al.,  
13 2019). The projected reduction of shear along the U.S. East Coast with warming is consistent among studies  
14 (e.g., Vecchi and Soden, 2007).

15  
16 In summary, average peak TC wind speeds and the proportion of Category 4-5 TCs will *very likely* increase  
17 globally with warming. It is *likely* that the frequency of Category 4-5 TCs will increase in limited regions  
18 over the western North Pacific. It is *very likely* that average TC rain-rates will increase with warming, and  
19 *likely* that the peak rain-rates will increase at greater than the Clausius-Clapeyron scaling rate of 7% per °C  
20 of warming in some regions due to increased low-level moisture convergence caused by regional increases in  
21 TC wind-intensity. It is *likely* that the average location where TCs reach their peak wind-intensity will  
22 migrate poleward in the western North Pacific Ocean as the tropics expand with warming, and that the global  
23 frequency of TCs over all categories will decrease or remain unchanged.

## 24 25 26 **11.7.2 Extratropical storms**

27  
28 This section focuses on extratropical cyclones (ETCs) that are either classified as strong or extreme by using  
29 some measure of their intensity, or by being associated with the occurrence of extremes in variables such as  
30 precipitation or near-surface wind speed (Seneviratne et al., 2012). Since AR5, the high relevance of ETCs  
31 for extreme precipitation events has been well established (Pfahl and Wernli, 2012; Catto and Pfahl, 2013;  
32 Utsumi et al., 2017), with 80% or more of hourly and daily precipitation extremes being associated with  
33 either ETCs or fronts over oceanic mid-latitude regions, and somewhat smaller, but still very large,  
34 proportions of events over mid-latitude land regions (Utsumi et al., 2017). The emphasis in this section is on  
35 individual ETCs that have been identified using some detection and tracking algorithms. Mid-latitude  
36 atmospheric rivers are assessed in Section 8.3.2.8.

### 37 38 39 **11.7.2.1 Observed trends**

40  
41 Chapter 2 (Section 2.3.1.4.3) concluded that there is overall *low confidence* in recent changes in the total  
42 number of ETCs over both hemispheres and that there is *medium confidence* in a poleward shift of the storm  
43 tracks over both hemispheres since the 1980s. Overall, there is also *low confidence* in past-century trends in  
44 the number and intensity of the strongest ETCs due to the large interannual and decadal variability (Feser et  
45 al., 2015; Reboita et al., 2015; Wang et al., 2016; Varino et al., 2018) and due to temporal and spatial  
46 heterogeneities in the number and type of assimilated data in reanalyses, particularly before the satellite era  
47 (Krueger et al., 2013; Tilinina et al., 2013; Befort et al., 2016; Chang and Yau, 2016; Wang et al., 2016).  
48 There is *medium confidence* that the agreement among reanalyses and among detection and tracking  
49 algorithms is higher when considering stronger cyclones (Neu et al., 2013; Pepler et al., 2015; Wang et al.,  
50 2016). Over the Southern Hemisphere, there is *high confidence* that the total number of ETCs with low  
51 central pressures (<980 hPa) has increased between 1979 and 2009, with all eight reanalyses considered by  
52 Wang et al. (2016), showing positive trends and five of them showing statistically significant trends. Similar  
53 results were found by (Reboita et al., 2015) using a different detection and tracking algorithm and a single  
54 reanalysis product. Over the Northern Hemisphere, there is *high agreement* among reanalyses that the  
55 number of cyclones with low central pressures (<970 hPa) has decreased in summer and winter during the

1 period 1979-2010 (Tilinina et al., 2013; Chang et al., 2016). However, changes exhibit substantial decadal  
2 variability and do not show monotonic trends since the 1980s. For example, over the Arctic and North  
3 Atlantic, Tilinina et al. (2013) showed the number of cyclones with very low central pressure (<960 hPa)  
4 increased from 1979 to 1990 and then declined until 2010 in all five reanalyses considered. Over the North  
5 Pacific, the number of cyclones with very low central pressure reached a peak around 2000 and then  
6 decreased until 2010 in the five reanalyses considered (Tilinina et al., 2013).

7  
8 Overall, however, it should be noted that characterising trends in the dynamical intensity of ETCs (e.g., wind  
9 speeds) using the absolute central pressure is problematic because the central pressure depends on the  
10 background mean sea level pressure, which varies seasonally and regionally (e.g., Befort et al., 2016).

#### 11 11.7.2.2 Model evaluation

12  
13  
14  
15 There is *high confidence* that coarse-resolution climate models (e.g., CMIP5 and CMIP6) underestimate the  
16 dynamical intensity of ETCs, including the strongest ETCs, as measured using a variety of metrics, including  
17 mean pressure gradient, mean vorticity and near-surface winds, over most regions (Colle et al., 2013; Zappa  
18 et al., 2013a; Govekar et al., 2014; Di Luca et al., 2016; Trzeciak et al., 2016; Seiler et al., 2018; Priestley et  
19 al., 2020). There is also *high confidence* that most current climate models underestimate the number of  
20 explosive systems (i.e., systems showing a decrease in mean sea level pressure of at least 24 hPa in 24 hours)  
21 over both hemispheres (Seiler and Zwiers, 2016a; Gao et al., 2020; Priestley et al., 2020). There is *high*  
22 *confidence* that the underestimation of the intensity of ETCs is associated with the coarse horizontal  
23 resolution of climate models, with higher horizontal resolution models, including models from HighResMIP  
24 and CORDEX, usually showing better performance (Colle et al., 2013; Zappa et al., 2013a; Di Luca et al.,  
25 2016; Trzeciak et al., 2016; Seiler et al., 2018; Gao et al., 2020; Priestley et al., 2020). The improvement by  
26 higher-resolution models is found even when comparing models and reanalyses after postprocessing data to a  
27 common resolution (Zappa et al., 2013a; Di Luca et al., 2016; Priestley et al., 2020). The systematic bias in  
28 the intensity of ETCs has also been linked to the inability of current climate models to well resolve diabatic  
29 processes, particularly those related to the release of latent heat (Willison et al., 2013; Trzeciak et al., 2016)  
30 and the formation of clouds (Govekar et al., 2014). There is *medium confidence* that climate models simulate  
31 well the spatial distribution of precipitation associated with ETCs over the Northern Hemisphere, together  
32 with some of the main features of the ETC life cycle, including the maximum in precipitation occurring just  
33 before the peak in dynamical intensity (e.g., vorticity) as observed in a reanalysis and observations  
34 (Hawcroft et al., 2018). There is, however, large observational uncertainty in ETC-associated precipitation  
35 (Hawcroft et al., 2018) and limitations in the simulation of frontal precipitation, including too low rainfall  
36 intensity over mid-latitude oceanic areas in both hemispheres (Catto et al., 2015).

#### 37 38 39 11.7.2.3 Detection and attribution, event attribution

40  
41 Chapter 3 (Section 3.3.3.3) concluded that there is *low confidence* in the attribution of observed changes in  
42 the number of ETCs in the Northern Hemisphere and that there is *high confidence* that the poleward shift of  
43 storm tracks in the Southern Hemisphere is linked to human activity, mostly due to emissions of ozone-  
44 depleting substances. Specific studies attributing changes in the most extreme ETCs are not available. The  
45 human influence on individual extreme ETC events has been considered only a few times and there is overall  
46 *low confidence* in the attribution of these changes (NASEM, 2016; Vautard et al., 2019).

#### 47 48 49 11.7.2.4 Projections

50  
51 The frequency of ETCs is expected to change primarily following a poleward shift of the storm tracks as  
52 discussed in Chapters 4 (Section 4.5.1.6, see also Figure 4.31) and 8 (Section 8.4.2.8). There is *medium*  
53 *confidence* that changes in the dynamical intensity (e.g., wind speeds) of ETCs will be small, although  
54 changes in the location of storm tracks can lead to substantial changes in local extreme wind speeds (Zappa  
55 et al., 2013b; Chang, 2014; Li et al., 2014; Seiler and Zwiers, 2016b; Yettella and Kay, 2017; Barcikowska

1 et al., 2018; Kar-Man Chang, 2018). Yettella and Kay (2017) detected and tracked ETCs over both  
2 hemispheres in an ensemble of 30 CESM-LE simulations, differing only in their initial conditions, and found  
3 that changes in mean wind speeds around ETC centres are often negligible between present (1986-2005) and  
4 future (2081-2100) periods. Using 19 CMIP5 models, Zappa et al. (2013b) found an overall reduction in the  
5 number of cyclones associated with low-troposphere (850-hPa) wind speeds larger than  $25 \text{ m s}^{-1}$  over the  
6 North Atlantic and Europe with the number of the 10% strongest cyclones decreasing by about 8% and 6%  
7 in DJF and JJA according to the RCP4.5 scenario (2070-2099 vs. 1976-2005). Over the North Pacific, Chang  
8 (2014) showed that CMIP5 models project a decrease in the frequency of ETCs with the largest central  
9 pressure perturbation (i.e., the depth, strongly related with low-level wind speeds) by the end of the century  
10 according to simulations using the RCP8.5 scenario. Using projections from CMIP5 GCMs under the  
11 RCP8.5 scenario (1981-2000 to 2081-2100), Seiler and Zwiers (2016b) projected a northward shift in the  
12 number of explosive ETCs in the northern Pacific, with fewer and weaker events south, and more frequent  
13 and stronger events north of  $45^\circ\text{N}$ . Using 19 CMIP5 GCMs under the RCP8.5 scenario, (Kar-Man Chang,  
14 2018) found a significant decrease in the number of ETCs associated with extreme wind speeds (2081–2100  
15 vs. 1980–99) over the Northern Hemisphere (average decrease of 17%) and over some smaller regions,  
16 including the Pacific and Atlantic regions.

17  
18 Over the Southern Hemisphere, future changes (RCP8.5 scenario; 1980-1999 to 2081-2100) in extreme  
19 ETCs were studied by Chang (2017) using 26 CMIP5 models and a variety of intensity metrics (850-hPa  
20 vorticity, 850-hPa wind speed, mean sea level pressure and near-surface wind speed). They found that the  
21 number of extreme cyclones is projected to increase by at least 20% and as much as 50%, depending on the  
22 specific metric used to define extreme ETCs. Increases in the number of strong cyclones appear to be robust  
23 across models and for most seasons, although they show strong regional variations with increases occurring  
24 mostly over the southern flank of the storm track, consistent with a shift and intensification of the storm  
25 track. Overall, there is *medium confidence* that projected changes in the dynamical intensity of ETCs depend  
26 on the resolution and formulation (e.g., explicit or implicit representation of convection) of climate models  
27 (Booth et al., 2013; Michaelis et al., 2017; Zhang and Colle, 2017).

28  
29 As reported in AR5 and in Chapter 8 (Section 8.4.2.8), despite small changes in the dynamical intensity of  
30 ETCs, there is *high confidence* that the precipitation associated with ETCs will increase in the future (Zappa  
31 et al., 2013b; Marciano et al., 2015; Pepler et al., 2016; Zhang and Colle, 2017; Michaelis et al., 2017;  
32 Yettella and Kay, 2017; Barcikowska et al., 2018; Zarzycki, 2018; Hawcroft et al., 2018; Kodama et al.,  
33 2019; Bevacqua et al., 2020c; Reboita et al., 2020). There is *high confidence* that increases in precipitation  
34 will follow increases in low-level water vapour (i.e., about 7% per degree of surface warming; Box 11.1) and  
35 will be largest for higher warming levels (Zhang and Colle, 2017). There is *medium confidence* that  
36 precipitation changes will show regional and seasonal differences due to distinct changes in atmospheric  
37 humidity and dynamical conditions (Zappa et al., 2015; Hawcroft et al., 2018), with even decreases in some  
38 specific regions such as the Mediterranean (Zappa et al., 2015; Barcikowska et al., 2018). There is *high*  
39 *confidence* that snowfall associated with wintertime ETCs will decrease in the future, because increases in  
40 tropospheric temperatures lead to a lower proportion of precipitation falling as snow (O’Gorman, 2014;  
41 Rhoades et al., 2018; Zarzycki, 2018). However, there is *medium confidence* that extreme snowfall events  
42 associated with wintertime ETCs will change little in regions where snowfall will be supported in the future  
43 (O’Gorman, 2014; Zarzycki, 2018).

44  
45 In summary, there is *low confidence* in past changes in the dynamical intensity (e.g., maximum wind speeds)  
46 of ETCs and *medium confidence* that in the future these changes will be small, although changes in the  
47 location of storm tracks could lead to substantial changes in local extreme wind speeds. There is *high*  
48 *confidence* that average and maximum ETC precipitation-rates will increase with warming, with the  
49 magnitude of the increases associated with increases in atmospheric water vapour. There is *medium*  
50 *confidence* that projected changes in the intensity of ETCs, including wind speeds and precipitation, depend  
51 on the resolution and formulation of climate models.

### 52 53 54 **11.7.3 Severe convective storms**



1 Severe convective storms are convective systems that are associated with extreme phenomena such as  
2 tornadoes, hail, heavy precipitation (rain or snow), strong winds, and lightning. The assessment of changes in  
3 severe convective storms in SREX (Chapter 3, Seneviratne et al., 2012) and AR5 (Chapter 12, Collins et al.,  
4 2013) is limited and focused mainly on tornadoes and hail storms. SREX assessed that there is *low*  
5 *confidence* in observed trends in tornadoes and hail because of data inhomogeneities and inadequacies in  
6 monitoring systems. Subsequent literature assessed in the Climate Science Special Report (Kossin et al.,  
7 2017) led to the assessment of the observed tornado activity over the 2000s in the United States with a  
8 decrease in the number of days per year with tornadoes and an increase in the number of tornadoes on these  
9 days (*medium confidence*). However, there is *low confidence* in past trends for hail and severe thunderstorm  
10 winds. Climate models consistently project environmental changes that would support an increase in the  
11 frequency and intensity of severe thunderstorms that combine tornadoes, hail, and winds (*high confidence*),  
12 but there is *low confidence* in the details of the projected increase. Regional aspects of severe convective  
13 storms and details of the assessment of tornadoes and hail are also assessed in Chapter 12 (Section 12.3.3.2  
14 for tornadoes; Section 12.3.4.5 for hail; Section 12.4.5.3 for Europe, Section 12.4.6.3 for North America, and  
15 Section 12.7.2 for regional gaps and uncertainties).

### 16 17 18 11.7.3.1 Mechanisms and drivers

19  
20 Severe convective storms are sometimes embedded in synoptic-scale weather systems, such as TCs, ETCs,  
21 and fronts (Kunkel et al., 2013). They are also generated as individual events as mesoscale convective  
22 systems (MCSs) and mesoscale convective complexes (MCCs) (a special type of a large, organized and long-  
23 lived MCS), without being clearly embedded within larger-scale weather systems. In addition to the general  
24 vigorousness of precipitation, hail, and winds associated with MCSs, characteristics of MCSs are viewed in  
25 new perspectives in recent years, probably because of both the development of dense mesoscale observing  
26 networks and advances in high-resolution mesoscale modelling (Sections 11.7.3.2 and 11.7.3.3). The  
27 horizontal scale of MCSs is discussed with their organization of the convective structure and it is examined  
28 with a concept of "convective aggregation" in recent years (Holloway et al., 2017). MCSs sometimes take a  
29 linear shape and stay almost stationary with successive production of cumulonimbus on the upstream side  
30 (back-building type convection), and cause heavy rainfall (Schumacher and Johnson, 2005). Many of the  
31 recent severe rainfall events in Japan are associated with band-shaped precipitation systems (Kunii et al.,  
32 2016; Oizumi et al., 2018; Tsuguti et al., 2018; Kato, 2020), suggesting common characteristics of severe  
33 precipitation, at least in East Asia. The convective modes of severe storms in the United States can be  
34 classified into rotating or linear modes and preferable environmental conditions for these modes, such as  
35 vertical shear, have been identified (Trapp et al., 2005; Smith et al., 2013; Allen, 2018). Cloud microphysics  
36 characteristics of MCSs were examined and the roles of warm rain processes on extreme precipitation were  
37 emphasized recently (Sohn et al., 2013; Hamada et al., 2015; Hamada and Takayabu, 2018). Idealized  
38 studies also suggest the importance of ice and mixed-phase processes of cloud microphysics on extreme  
39 precipitation (Sandvik et al., 2018; Bao and Sherwood, 2019). However, it is unknown whether the types of  
40 MCSs are changing in recent periods or observed ubiquitously all over the world.

41  
42 Severe convective storms occur under conditions preferable for deep convection, that is, conditionally  
43 unstable stratification, sufficient moisture both in lower and middle levels of the atmosphere, and a strong  
44 vertical shear. These large-scale environmental conditions are viewed as necessary conditions for the  
45 occurrence of severe convective systems, or the resulting tornadoes and lightning, and the relevance of these  
46 factors strongly depends on the region (e.g., Antonescu et al., 2016a; Allen, 2018; Tochimoto and Niino,  
47 2018). Frequently used metrics are atmospheric static stability, moisture content, convective available  
48 potential energy (CAPE) and convective inhibition (CIN), wind shear or helicity, including storm-relative  
49 environmental helicity (SREH) (Tochimoto and Niino, 2018; Elsner et al., 2019). These metrics, largely  
50 controlled by large-scale atmospheric circulations or synoptic weather systems, such as TCs and ETCs, are  
51 then generally used to examine severe convective systems. In particular, there is *high confidence* that CAPE  
52 in the tropics and the subtropics increases in response to global warming (Singh et al., 2017a), as supported  
53 by theoretical studies (Singh and O’Gorman, 2013; Seeley and Romps, 2015; Romps, 2016; Agard and  
54 Emanuel, 2017). The uncertainty, however, arises from the balance between factors affecting severe storm  
55 occurrence. For example, the warming of mid-tropospheric temperatures leads to an increase in the freezing

1 level, which leads to increased melting of smaller hailstones, while there may be some offset by stronger  
2 updrafts driven by increasing CAPE, which would favour the growth of larger hailstones, leading to less  
3 melting when falling (Allen, 2018; Mahoney, 2020).

4  
5 There are few studies on relations between changes in severe convective storms and those of the large-scale  
6 circulation patterns. Tornado outbreaks in the United States are usually associated with ETCs with their  
7 frontal systems and TCs (Fuhrmann et al., 2014; Tochimoto and Niino, 2016). In early June in East Asia,  
8 associated with the Baiu/Changma/Mei-yu, severe precipitation events are frequently caused by MCSs.  
9 Severe precipitation events are also caused by remote effects of TCs, known as predecessor rain events  
10 (PREs) (Galarneau et al., 2010). Atmospheric rivers and other coherent types of enhanced water vapour flux  
11 also have the potential to induce severe convective systems (Kamae et al., 2017; Ralph et al., 2018; Waliser  
12 and Guan, 2017; see Section 8.3.2.8.1). Combined with the above drivers, topographic effects also enhance  
13 the intensity and duration of severe convective systems and the associated precipitation (Ducrocq et al.,  
14 2008; Piaget et al., 2015). However, the changes in these drivers are not generally significant, so their  
15 relations to severe convective storms are unclear.

16  
17 In summary, severe convective storms are sometimes embedded in synoptic-scale weather systems, such as  
18 TCs, ETCs, and fronts, and modulated by large-scale atmospheric circulation patterns. The occurrence of  
19 severe convective storms and the associated severe events, including tornadoes, hail, and lightning, is  
20 affected by environmental conditions of the atmosphere, such as CAPE and vertical shear. The uncertainty,  
21 however, arises from the balance between these environmental factors affecting severe storm occurrence.

#### 22 23 24 *11.7.3.2 Observed trends*

25  
26 Observed trends in severe convective storms or MCSs are not well documented, but the climatology of  
27 MCSs has been analysed in specific regions (North America, South America, Europe, Asia; regional aspects  
28 of convective storms are separately assessed in Chapter 12). As the definition of severe convective storms  
29 varies depending on the literature, it is not straightforward to make a synthesizing view of observed trends in  
30 severe convective storms in different regions. However, analysis using satellite observations provides a  
31 global view of MCSs (Kossin et al., 2017). The global distribution of thunderstorms is captured (Zipser et  
32 al., 2006; Liu and Zipser, 2015) by using the satellite precipitation measurements by the Tropical Rainfall  
33 Measuring Mission (TRMM) and Global Precipitation Mission (GPM) (Hou et al., 2014). The climatological  
34 characteristics of MCSs are provided by satellite analyses in South America (Durkee and Mote, 2010;  
35 Rasmussen and Houze, 2011; Rehbein et al., 2018) and those of MCC in the Maritime Continent by  
36 Trismidianto and Satyawardhana (2018). Analysis of the environmental conditions favourable for severe  
37 convective events indirectly indicates the climatology and trends of severe convective events (Allen et al.,  
38 2018; Taszarek et al., 2018, 2019), though favourable conditions depend on the location, such as the  
39 difference for tornadoes associated with ETCs between the United States and Japan (Tochimoto and Niino,  
40 2018).

41  
42 Observed trends in severe convective storms are highly regionally dependent. In the United States, it is  
43 indicated that there is no significant increase in convective storms, and hail and severe thunderstorms  
44 (Kossin et al., 2017; Kunkel et al., 2013). There is an upward trend in the frequency and intensity of extreme  
45 precipitation events in the United States (*high confidence*) (Kunkel et al., 2013; Easterling et al., 2017), and  
46 MCSs have increased in occurrence and precipitation amounts since 1979 (*limited evidence*) (Feng et al.,  
47 2016). Significant interannual variability of hailstone occurrences is found in the Southern Great Plains of  
48 the United States (Jeong et al., 2020). The mean annual number of tornadoes has remained relatively  
49 constant, but their variability of occurrence has increased since the 1970s, particularly over the 2000s, with a  
50 decrease in the number of days per year, but an increase in the number of tornadoes on these days (Brooks et  
51 al., 2014; Elsner et al., 2015, 2019; Kossin et al., 2017; Allen, 2018). There has been a shift in the  
52 distribution of tornadoes, with increases in tornado occurrence in the mid-south of the US and decreases over  
53 the High Plains (Gensini and Brooks, 2018). Trends in MCSs are relatively more visible for particular  
54 aspects of MCSs, such as lengthening of active seasons and dependency on duration. MCSs have increased  
55 in occurrence and precipitation amounts since 1979 (Easterling et al., 2017). Feng et al. (2016) analysed that

1 the observed increases in springtime total and extreme rainfall in the central United States are dominated by  
2 MCSs, with increased frequency and intensity of long-lasting MCSs.

3  
4 Studies on trends in severe convective storms and their ingredients outside of the United States are limited.  
5 Westra et al. (2014) found that there is an increase in the intensity of short-duration convective events  
6 (minutes to hours) over many regions of the world, except eastern China. In Europe, a climatology of  
7 tornadoes shows an increase in detected tornadoes between 1800 to 2014, but this trend might be affected by  
8 the density of observations (Antonescu et al., 2016b, 2016a). An increase in the trend in extreme daily  
9 rainfall is found in southeastern France, where MCSs play a key role in this type of event (Blanchet et al.,  
10 2018; Ribes et al., 2019). Trend analysis of the mean annual number of days with thunderstorms since 1979  
11 in Europe indicates an increase over the Alps and central, southeastern, and eastern Europe, with a decrease  
12 over the southwest (Taszarek et al., 2019). In the Sahelian region, Taylor et al. (2017) analysed MCSs using  
13 satellite observations since 1982 and showed an increase in the frequency of extreme storms. In Bangladesh,  
14 the annual number of propagating MCSs decreased significantly during 1998-2015 based on TRMM  
15 precipitation data (Habib et al., 2019). Prein and Holland (2018) estimated the hail hazard from large-scale  
16 environmental conditions using a statistical approach and showed increasing trends in the United States,  
17 Europe, and Australia. However, trends in hail on regional scales are difficult to validate because of an  
18 insufficient length of observations and inhomogeneous records (Allen, 2018). The high spatial variability of  
19 hail suggests it is reasonable that there would be local signals of both positive and negative trends and the  
20 trends that are occurring in hail globally are uncertain. In China, the total number of days that have either a  
21 thunderstorm or hail have decreased by about 50% from 1961 to 2010, and the reduction in these severe  
22 weather occurrences correlates strongly with the weakening of the East Asian summer monsoon (Zhang et  
23 al., 2017b). More regional aspects of severe convective storms are detailed in Chapter 12.

24  
25 In summary, because the definition of severe convective storms varies depending on the literature and the  
26 region, it is not straightforward to make a synthesizing view of observed trends in severe convective storms  
27 in different regions. In particular, observational trends in tornadoes, hail, and lightning associated with  
28 severe convective storms are not robustly detected due to insufficient coverage of the long-term  
29 observations. There is *medium confidence* that the mean annual number of tornadoes in the United States has  
30 remained relatively constant, but their variability of occurrence has increased since the 1970s, particularly  
31 over the 2000s, with a decrease in the number of days per year and an increase in the number of tornadoes on  
32 these days (*high confidence*). Detected tornadoes have also increased in Europe, but the trend depends on the  
33 density of observations.

### 34 35 36 11.7.3.3 Model evaluation

37  
38 The explicit representation of severe convective storms requires non-hydrostatic models with horizontal grid  
39 spacings below 5 km, denoted as convection-permitting models or storm-resolving models (Section 10.3.1).  
40 Convection-permitting models are becoming available to run over a wide domain, such as a continental scale  
41 or even over the global area, and show realistic climatological characteristics of MCSs (Prein et al., 2015;  
42 Guichard and Couvreur, 2017; Satoh et al., 2019). Such high-resolution simulations are computationally too  
43 expensive to perform at the larger domain and for long periods and alternative methods by using an RCM  
44 with dynamical downscaling are generally used (Section 10.3.1). Convection-permitting models are used as  
45 the flagship project of CORDEX to particularly study projections of thunderstorms (Section 10.3.3).  
46 Simulations of North American MCSs by a convection-permitting model conducted by Prein et al. (2017a)  
47 were able to capture the main characteristics of the observed MCSs, such as their size, precipitation rate,  
48 propagation speed, and lifetime. Cloud-permitting model simulations in Europe also showed sub-daily  
49 precipitation realistically (Ban et al., 2014; Kendon et al., 2014). Evaluation of precipitation conducted using  
50 convection-permitting simulations around Japan showed that finer resolution improves intense precipitation  
51 (Murata et al., 2017). MCSs over Africa simulated using convection-permitting models showed better  
52 extreme rainfall (Kendon et al., 2019) and diurnal cycles and convective rainfall over land than the coarser-  
53 resolution RCMs or GCMs (Stratton et al., 2018; Crook et al., 2019).

54  
55 The other modeling approach is the analysis of the environmental conditions that control characteristics of

1 severe convective storms using the typical climate model results in CMIP5/6 (Allen, 2018). Severe  
2 convective storms are generally formed in environments with large CAPE and tornadic storms are, in  
3 particular, formed with a combination of large CAPE and strong vertical wind shear. As the processes  
4 associated with severe convective storms occur over a wide range of spatial and temporal scales, some of  
5 which are poorly understood and are inadequately sampled by observational networks, the model calibration  
6 approaches are in general difficult and insufficiently validated. Therefore, model simulations and their  
7 interpretations should be done with much caution.

8  
9 In summary, there are typically two kinds of modeling approaches for studying changes in severe convective  
10 storms. One is to use convection-permitting models in wider regions or the global domain in time-sliced  
11 downscaling methods to directly simulate severe convective storms. The other is the analysis of the  
12 environmental conditions that control characteristics of severe convective storms by using coarse-resolution  
13 GCMs. Even in finer-resolution convection-permitting models, it is difficult to directly simulate tornadoes,  
14 hail storms, and lightning, so modeling studies of these changes are limited.

#### 15 16 17 *11.7.3.4 Detection and attribution, event attribution*

18  
19 It is extremely difficult to detect differences in time and space of severe convective storms (Kunkel et al.,  
20 2013). Although some ingredients that are favourable for severe thunderstorms have increased over the  
21 years, others have not; thus, overall, changes in the frequency of environments favourable for severe  
22 thunderstorms have not been statistically significant. Event attribution studies on severe convective events  
23 have now been undertaken for some cases. For the case of the July 2018 heavy rainfall event in Japan (BOX  
24 11.3), Kawase et al. (2019) took a storyline approach to show that the rainfall during this event in Japan was  
25 increased by approximately 7% due to the recent rapid warming around Japan. For the case of the December  
26 2015 extreme rainfall event in Chennai, India, the extremity of the event was equally caused by the warming  
27 trend in the Bay of Bengal SSTs and the strong El Niño conditions (van Oldenborgh et al., 2016; Boyaj et al.,  
28 2018). For hailstorms, such as those that caused disasters in the United States in 2018, detection of the role  
29 of climate change in changing hail storms is more difficult, because hail storms are not, in general, directly  
30 simulated by convection-permitting models and not adequately represented by the environmental parameters  
31 of coarse-resolution GCMs (Mahoney, 2020).

32  
33 In summary, it is extremely difficult to detect and attribute changes in severe convective storms, except for  
34 case study approaches by event attribution. There is *limited evidence* that extreme precipitation associated  
35 with severe convective storms has increased in some cases.

#### 36 37 38 *11.7.3.5 Projections*

39  
40 Future projections of severe convective storms are usually studied either by analysing the environmental  
41 conditions simulated by climate models or by a time slice approach with higher-resolution convection-  
42 permitting models by comparing simulations downscaled with climate model results under historical  
43 conditions and those under hypothesized future conditions (Kendon et al., 2017; Allen, 2018). Up to now,  
44 individual studies using convection-permitting models gave projections of extreme events associated with  
45 severe convective storms in local regions, and it is not generally possible to obtain global or general views of  
46 projected changes of severe convective storms. Prein et al. (2017b) investigated future projections of North  
47 American MCS simulations and showed an increase in MCS frequency and an increase in total MCS  
48 precipitation volume by the combined effect of increases in maximum precipitation rates associated with  
49 MCSs and increases in their size. Rasmussen et al. (2017) investigated future changes in the diurnal cycle of  
50 precipitation by capturing organized and propagating convection and showed that weak to moderate  
51 convection will decrease and strong convection will increase in frequency in the future. Ban et al. (2015)  
52 found the day-long and hour-long precipitation events in summer intensify in the European region covering  
53 the Alps. Kendon et al. (2019) showed future increases in extreme 3-hourly precipitation in Africa. Murata et  
54 al. (2015) investigated future projections of precipitation around Japan and showed a decrease in monthly  
55 mean precipitation in the eastern Japan Sea region in December, suggesting convective clouds become

1 shallower in the future in the winter over the Japan Sea.

2  
3 The other approach is the projection of the environmental conditions that control characteristics of severe  
4 convective storms by analysing climate model results. There is *high confidence* that CAPE, particularly  
5 summertime mean CAPE and high percentiles of the CAPE in the tropics and subtropics, increases in  
6 response to global warming in an ensemble of climate models including those of CMIP5, mainly from  
7 increased low-level specific humidity (Sobel and Camargo, 2011; Singh et al., 2017a; Chen et al., 2020b).  
8 CIN becomes stronger over most land areas under global warming, resulting mainly from reduced low-level  
9 relative humidity over land (Chen et al., 2020b). However, there are large differences within the CMIP5  
10 ensemble for environmental conditions, which contribute to some degree of uncertainty (Allen, 2018).  
11 Because the relation between simulated environments in models and the occurrence of severe convective  
12 storms are in general insufficiently validated, the confidence level of the projection of severe convective  
13 storms with the approach of the environmental conditions is generally *low*.

14  
15 In the United States, projected changes in the environmental conditions show an increase in CAPE and no  
16 changes or decreases in the vertical wind shear, suggesting favourable conditions for an increase in severe  
17 convective storms in the future, but the interpretation of how tornadoes or hail will change is an open  
18 question because of the strong dependence on shear (Brooks, 2013). Diffenbaugh et al. (2013) showed robust  
19 increases in the occurrence of the favourable environments for severe convective storms with increased  
20 CAPE and stronger low-level wind shear in response to future global warming. A downscaling approach  
21 showed that the variability of the occurrence of severe convective storms increases in spring in late 21st  
22 century simulations (Gensini and Mote, 2015). Future changes in hail occurrence in the United States  
23 examined through convection-permitting dynamical downscaling suggested that the hail season may begin  
24 earlier in the year and exhibit more interannual variability with increases in the frequency of large hail in  
25 broad areas over the United States (Trapp et al., 2019). There is *medium confidence* that the frequency and  
26 variability of the favourable environments for severe convective storms will increase in spring, and *low*  
27 *confidence* for summer and autumn (Diffenbaugh et al., 2013; Gensini and Mote, 2015; Hoogewind et al.,  
28 2017). The occurrence of hail events in Colorado in the United States was examined by comparing both  
29 present-day and projected future climates using high-resolution model simulations capable of resolving  
30 hailstorms (Mahoney et al., 2012), which showed hail is almost eliminated at the surface in the future in  
31 most of the simulations, despite more intense future storms and significantly larger amounts of hail generated  
32 in-cloud.

33  
34 Future changes in severe convection environments show enhancement of instability with less robust changes  
35 in the frequency of strong vertical wind shear in Europe (Púčik et al. 2017) and in Japan (Muramatsu et al.  
36 2016). In Japan, the frequency of conditions favourable for strong tornadoes increases in spring and partly in  
37 summer.

38  
39 In summary, the average and maximum rain rates associated with severe convective storms increase in a  
40 warming world in some regions including the USA (*high confidence*). There is *high confidence* from climate  
41 models that CAPE increases in response to global warming in the tropics and subtropics, suggesting more  
42 favourable environments for severe convective storms. The frequency of springtime severe convective  
43 storms is projected to increase in the USA leading to a lengthening of the severe convective storm season  
44 (*medium confidence*), evidence in other regions is limited. There is significant uncertainty about projected  
45 regional changes in tornadoes, hail, and lightning due to limited analysis of simulations using convection-  
46 permitting models (*high confidence*).

#### 47 48 49 **11.7.4 Extreme winds**

50  
51 Extreme winds are defined here in terms of the strongest near-surface wind speeds that are generally  
52 associated with extreme storms, such as TCs, ETCs, and severe convective storms. In previous IPCC reports,  
53 near-surface wind speed (including extremes), has not been assessed as a variable in its own right, but rather  
54 in the context of other extreme atmospheric or oceanic phenomena. The exception was the SREX report  
55 (Seneviratne et al., 2012), which specifically examined past changes and projections of mean and extreme

1 near-surface wind speeds. A strong decline in extreme winds compared to mean winds was reported for the  
2 continental northern mid-latitudes. Due to the small number of studies and uncertainties in terrestrial-based  
3 surface wind measurements, the findings were assigned *low confidence* in the SREX. AR5 reported a  
4 weakening of mean and maximum winds from the 1960s or 1970s to the early 2000s in the tropics and mid-  
5 latitudes and increases in high latitudes, but with *low confidence* in changes in the observed surface winds  
6 over land (Hartmann et al., 2013). Observed trends in mean wind speed over land and the ocean are assessed  
7 in Section 2.3.1.4.4. Aspects of climate impact-drivers for winds are addressed in Section 12.3.3 and 12.5.2.3  
8 and their regional changes are assessed in Section 12.4.

9  
10 Observationally, although not specifically addressing extreme wind speed changes, negative surface wind  
11 speed trends (stilling) were found in the tropics and mid-latitudes of both hemispheres of  $-0.014 \text{ m s}^{-1} \text{ year}^{-1}$ ,  
12 while positive trends were reported at high latitudes poleward of 70 degrees, based on a review of 148  
13 studies (McVicar et al., 2012a). An earlier study attributed the stilling to both changes in atmospheric  
14 circulation and an increase in surface roughness due to an overall increase in vegetation cover (Vautard et  
15 al., 2010). Since then, a number of additional studies have mostly confirmed these general negative mean-  
16 wind trends based on anemometer data for Spain (Azorin-Molina et al., 2017), Turkey, (Dadaser-Celik and  
17 Cengiz, 2014), the Netherlands, (Wever, 2012), Saudi Arabia, (Rehman, 2013), Romania, (Marin et al.,  
18 2014), and China (Chen et al., 2013). Lin et al. (2013) note that wind speed variability over China is greater  
19 at high elevation locations compared to those closer to mean sea level. Hande et al. (2012), using radiosonde  
20 data, found an increase in surface wind speed on Macquarie Island.

21  
22 A number of new studies have examined surface wind speeds over the ocean based on ship-based  
23 measurements, satellite altimeters, and Special Sensor Microwave/Imagers (SSM/I) (Tokinaga and Xie,  
24 2011; Zieger et al., 2014). It has been noted that wind speed trends tend to be stronger in altimeter  
25 measurements, although the spatial patterns of change are qualitatively similar in both instruments (Zieger et  
26 al., 2014). Liu et al. (2016) found positive trends in surface wind speeds over the Arctic Ocean in 20 years of  
27 satellite observations. Small positive trends in mean wind speed were found in 33 years of satellite data,  
28 together with larger trends in the 90th percentile values over global oceans (Ribal and Young, 2019). These  
29 results were consistent with an earlier study that found a positive trend in 1-in-100 year wind speeds (Young  
30 et al., 2012). A positive change in mean wind speeds was found for the Arabian Sea and the Bay of Bengal  
31 (Shanas and Kumar, 2015) and Zheng et al. (2017) found that positive wind speed trends over the ocean  
32 were larger during winter seasons than summer seasons.

33  
34 Changes in extreme winds are associated with changes in the characteristics (locations, frequencies, and  
35 intensities) of extreme storms, including TCs, ETCs, and severe convective storms. For TCs, as assessed in  
36 Section 11.7.1.5, it is projected that the average peak TC wind speeds will increase globally with warming,  
37 while the global frequency of TCs over all categories will decrease or remain unchanged; the average  
38 location where TCs reach their peak wind-intensity will migrate poleward in the western North Pacific  
39 Ocean as the tropics expand with warming. Frequency, intensities, and geographical distributions of extreme  
40 wind events associated with TCs will change according to these TC changes. For ETCs, by the end of the  
41 century, CMIP5 models show the number of ETCs associated with extreme winds will significantly decrease  
42 in the mid- and high latitudes of the Northern Hemisphere in winter, with the projected decrease being larger  
43 over the Atlantic (Kar-Man Chang, 2018), while it will significantly increase irrespective of the season in the  
44 Southern Hemisphere (Chang, 2017)(Section 11.7.2.4). Over the ocean in the subtropics, a large ensemble of  
45 60-km global model simulations indicated that extreme winds associated with storm surges will intensify  
46 over  $15\text{--}35^\circ\text{N}$  in the Northern Hemisphere (Mori et al., 2019). On the other hand, extreme surface wind  
47 speeds will mostly decrease due to decreases in the number and intensity of TCs over most tropical areas of  
48 the Southern Hemisphere (Mori et al., 2019). The projected changes in the frequency of extreme winds are  
49 associated with the future changes in TCs and ETCs.

50  
51 Extreme cyclonic windstorms that share some characteristics with both TCs and ETCs occur regularly over  
52 the Mediterranean Sea and are often referred to as “medicanes” (Ragone et al., 2018; Miglietta and Rotunno,  
53 2019; Ragone et al., 2018; Miglietta and Rotunno, 2019; Zhang et al., 2020e). Medicanes pose substantial  
54 threats to regional islands and coastal zones. A growing body of literature consistently found that the  
55 frequency of medicanes decreases under warming, while the strongest medicanes become stronger

1 (González-Alemán et al., 2019; Tous et al., 2016; Romero and Emanuel, 2017; Romera et al., 2017;  
2 Cavicchia et al., 2014; Romero and Emanuel, 2013; Gaertner et al., 2007). This is also consistent with  
3 expected global changes in TCs under warming (11.7.1). Based on the consistency of these studies, it is  
4 *likely* that medicanes will decrease in frequency, while the strongest medicanes become stronger under  
5 warming scenario projections (*medium confidence*).  
6

7 In summary, the observed intensity of extreme winds is becoming less severe in the lower to mid-latitudes,  
8 while becoming more severe in higher latitudes poleward of 60 degrees (*low confidence*). Projected changes  
9 in the frequency and intensity of extreme winds are associated with projected changes in the frequency and  
10 intensity of TCs and ETCs (*medium confidence*).  
11

## 12 13 **11.8 Compound events**

14  
15 The IPCC SREX (SREX Ch3) first defined compound events as “(1) two or more extreme events occurring  
16 simultaneously or successively, (2) combinations of extreme events with underlying conditions that amplify  
17 the impact of the events, or (3) combinations of events that are not themselves extremes but lead to an  
18 extreme event or impact when combined”. Further definitions of compound events have emerged since the  
19 SREX. Zscheischler et al. (2018) defined compound events broadly as “the combination of multiple drivers  
20 and/or hazards that contributes to societal or environmental risk”. This definition is used in the present  
21 assessment, because of its clear focus on the risk framework established by the IPCC, and also highlighting  
22 that compound events may not necessarily result from dependent drivers. Compound events have been  
23 classified into preconditioned events, where a weather-driven or climate-driven precondition aggravates the  
24 impacts of a hazard; multivariate events, where multiple drivers and/or hazards lead to an impact; temporally  
25 compounding events, where a succession of hazards leads to an impact; and spatially compounding events,  
26 where hazards in multiple connected locations cause an aggregated impact (Zscheischler et al., 2020).  
27 Drivers include processes, variables, and phenomena in the climate and weather domain that may span over  
28 multiple spatial and temporal scales. Hazards (such as floods, heat waves, wildfires) are usually the  
29 immediate physical precursors to negative impacts, but can occasionally have positive outcomes (Flach et  
30 al., 2018).  
31

### 32 33 **11.8.1 Overview**

34  
35 The combination of two or more – not necessarily extreme – weather or climate events that occur i) at the  
36 same time, ii) in close succession, or iii) concurrently in different regions, can lead to extreme impacts that  
37 are much larger than the sum of the impacts due to the occurrence of individual extremes alone. This is  
38 because multiple stressors can exceed the coping capacity of a system more quickly. The contributing events  
39 can be of similar types (clustered multiple events) or of different types (Zscheischler et al., 2020). Many  
40 major weather- and climate-related catastrophes are inherently of a compound nature (Zscheischler et al.,  
41 2018). This has been highlighted for a broad range of hazards, such as droughts, heat waves, wildfires,  
42 coastal extremes, and floods (Westra et al., 2016; AghaKouchak et al., 2020; Ridder et al., 2020). Co-  
43 occurring extreme precipitation and extreme winds can result in infrastructural damage (Martius et al.,  
44 2016); the compounding of storm surge and precipitation extremes can cause coastal floods (Wahl et al.,  
45 2015); the combination of drought and heat can lead to tree mortality (Allen et al., 2015)(see also Section  
46 11.6); wildfires increase occurrences of hailstorms and lightning (Zhang et al., 2019e). Compound storm  
47 types consisting of co-located cyclone, front and thunderstorm systems have a higher chance of causing  
48 extreme rainfall and extreme winds than individual storm types (Dowdy and Catto, 2017). Extremes may  
49 occur at similar times at different locations (De Luca et al., 2020a,b) but affect the same system, for instance,  
50 spatially-concurrent climate extremes affecting crop yields and food prices (Anderson et al., 2019; Singh et  
51 al., 2018). Studies also show an increasing risk for breadbasket regions to be concurrently affected by  
52 climate extremes with increasing global warming, even between 1.5°C and 2°C of global warming (Gaupp et  
53 al., 2019) (Box 11.2). Concomitant extreme conditions at different locations become more probable as  
54 changes in climate extremes are emerging over an increasing fraction of the land area (Sections 11.2.3,  
55 11.2.4, 11.8.2, 11.8.3; Box 11.4).

1  
2 Finally, impacts may occur because of large multivariate anomalies in the climate drivers, if systems are  
3 adapted to historical multivariate climate variability (Flach et al., 2017). For instance, ecosystems are  
4 typically adapted to the local covariability of temperature and precipitation such that a bivariate anomaly  
5 may have a large impact even though neither temperature nor precipitation may be extreme based on a  
6 univariate assessment (Mahony and Cannon, 2018). Given that almost all systems are affected by weather  
7 and climate phenomena at multiple space-time scales (Raymond et al., 2020), it is natural to consider  
8 extremes in a compound event framework. It should be noted, however, that multi-hazard dependencies can  
9 also decrease risk, for instance when hazards are negatively correlated (Hillier et al., 2020). Despite this  
10 recognition, the literature on past and future changes in compound events has been limited, but is growing.  
11 This section assesses examples of types of compound events in available literature.

12  
13 In summary, compound events include the combination of two or more – not necessarily extreme – weather  
14 or climate events that occur i) at the same time, ii) in close succession, or iii) concurrently in different  
15 regions. The land area affected by concurrent extremes has increased (*high confidence*). Concurrent extreme  
16 events at different locations, but possibly affecting similar sectors (e.g., breadbaskets) in different regions,  
17 will become more frequent with increasing global warming, in particular above +2°C of global warming  
18 (*high confidence*).

### 19 20 21 **11.8.2 Concurrent extremes in coastal and estuarine regions**

22  
23 Coastal and estuarine zones are prone to a number of meteorological extreme events and also to concurrent  
24 extremes. A major climate-impact driver in coastal regions around the world is floods (Chapter 12), and flood  
25 occurrence may be influenced by the dependence between storm surge, extreme rainfall, river flow, but also  
26 by sea level rise, waves and tides, as well as groundwater for estuaries. Floods with multiple drivers are often  
27 referred to as “compound floods” (Wahl et al., 2015; Moftakhari et al., 2017; Bevacqua et al., 2020b).

28  
29 At US coasts, the probability of co-occurring storm surge and heavy precipitation is higher for the  
30 Atlantic/Gulf coast relative to the Pacific coast (Wahl et al., 2015). Furthermore, six studied locations on the  
31 US coast with long overlapping time series show an increase in the dependence between heavy precipitation  
32 and storm surge over the last century, leading to more frequent co-occurring storm surge and heavy  
33 precipitation events at the present day (Wahl et al., 2015). Storm surge and extreme rainfall are also  
34 dependent in most locations on the Australian coasts (Zheng et al., 2013) and in Europe along the Dutch  
35 coasts (Ridder et al., 2018), along the Mediterranean Sea, the Atlantic coast and the North Sea (Bevacqua et  
36 al., 2019). The probability of flood occurrence can be assessed via the dependence between storm surge and  
37 river flow (Bevacqua et al., 2020a, 2020b). For instance, the occurrence of a North Sea storm surge in close  
38 succession with an extreme Rhine or Meuse river discharge is much more probable due to their dependence,  
39 compared to if both events would be independent (Kew et al., 2013; Klerk et al., 2015). Significant  
40 dependence between high sea levels and high river discharge are found for more than half of the available  
41 station observations, which are mostly located around the coasts of North America, Europe, Australia, and  
42 Japan (Ward et al., 2018). Combining global river discharge with a global storm surge model, hotspots of  
43 compound flooding have been discovered that are not well covered by observations, including Madagascar,  
44 Northern Morocco, Vietnam, and Taiwan (Couasnon et al., 2020). In the Dutch Noorderzijlvest area, there is  
45 more than a two-fold increase in the frequency of exceeding the highest warning level compared to the case  
46 if storm surge and heavy precipitation were independent (van den Hurk et al., 2015). In other regions and  
47 seasons, the dependence can be insignificant (Wu et al., 2018b) and there can be significant seasonal and  
48 regional differences in the storm surge-heavy precipitation relationship. Assessments of flood probabilities  
49 are often not based on actual flood measurements and instead are estimated from its main drivers including  
50 astronomical tides, storm surge, heavy precipitation, and high streamflow. Such single driver analyses might  
51 underestimate flood probabilities if multiple correlated drivers contribute to flood occurrence (e.g., van den  
52 Hurk et al., 2015).

53  
54 Many coastal areas are also prone to the occurrence of compound precipitation and wind extremes, which  
55 can cause damage, including to infrastructure and natural environments. A high percentage of co-occurring



1 wind and precipitation extremes are found in coastal regions and in areas with frequent tropical cyclones.  
2 Finally, the combination of extreme wave height and duration is also shown to influence coastal erosion  
3 processes (Corbella and Stretch, 2012).

4  
5 Aspects of concurrent extremes in coastal and estuarine environments have increased in frequency and/or  
6 magnitude over the last century in some regions. These include an increase in the dependence between heavy  
7 precipitation and storm surge over the last century, leading to more frequent co-occurring storm surge and  
8 heavy precipitation events in the present day along US coastlines (Wahl et al., 2015). In Europe, the  
9 probability of compound flooding occurrence increases most strongly along the Atlantic coast and the North  
10 Sea under strong warming. This increase is mostly driven by an intensification of precipitation extremes and  
11 aggravated flooding probability due to sea level rise (Bevacqua et al., 2019). At the global scale and under a  
12 high emissions scenario, the concurrence probability of meteorological conditions driving compound  
13 flooding would increase by more than 25% on average along coastlines worldwide by 2100, compared to the  
14 present (Bevacqua et al., 2020b). Sea level extremes and their physical impacts in the coastal zone arise from  
15 a complex set of atmospheric, oceanic, and terrestrial processes that interact on a range of spatial and  
16 temporal scales and will be modified by a changing climate, including sea level rise (McInnes et al., 2016).  
17 Interactions between sea level rise and storm surges (Little et al., 2015), and sea level and fluvial flooding  
18 (Moftakhari et al., 2017) are projected to lead to more frequent and more intense compound coastal flooding  
19 events as sea levels continue to rise.

20  
21 In summary, there is *medium confidence* that over the last century the probability of compound flooding has  
22 increased in some locations, including along the US coastline. There is *medium confidence* that the  
23 occurrence and magnitude of compound flooding in coastal regions will increase in the future due to both sea  
24 level rise and increases in heavy precipitation.

### 25 26 27 **11.8.3 Concurrent droughts and heat waves**

28  
29 Concurrent droughts and heat waves have a number of negative impacts on human society and natural  
30 ecosystems. Studies since SREX and AR5 show several occurrences of observed combinations of drought  
31 and heat waves in various regions.

32  
33 Over most land regions, temperature and precipitation are strongly negatively correlated during summer  
34 (Zscheischler and Seneviratne, 2017), mostly due to land-atmosphere feedbacks (Sections 11.1.6, 11.3.2),  
35 but also because synoptic-scale weather systems favourable for extreme heat are also unfavourable for rain  
36 (Berg et al., 2015). This leads to a strong correlation between droughts and heat waves (Zscheischler and  
37 Seneviratne, 2017). Drought events characterized by low precipitation and extreme high temperatures have  
38 occurred, for example, in California (AghaKouchak et al., 2014), inland eastern Australia (King et al., 2014),  
39 and large parts of Europe (Orth et al., 2016b). The 2018 growing season was both record-breaking dry and  
40 hot in Germany (Zscheischler and Fischer, 2020).

41  
42 The probability of co-occurring meteorological droughts and heat waves has increased in the observational  
43 period in many regions and will continue to do so under unabated warming (Herrera-Estrada and Sheffield,  
44 2017; Zscheischler and Seneviratne, 2017; Hao et al., 2018; Sarhadi et al., 2018; Alizadeh et al., 2020; Wu et  
45 al., 2021). Overall, projections of increases in co-occurring drought and heat waves are reported in northern  
46 Eurasia (Schubert et al., 2014), Europe ; Sedlmeier et al., 2018), southeast Australia (Kirono et al., 2017),  
47 multiple regions of the United States (Differbaugh et al., 2015; Herrera-Estrada and Sheffield 2017),  
48 northwest China (Li et al., 2019c; Kong et al., 2020) and India (Sharma and Mujumdar, 2017). The dominant  
49 signal is related to the increase in heat wave occurrence, which has been attributed to anthropogenic forcing  
50 (11.3.4). This means that even if drought occurrence is unaffected, compound hot and dry events will be  
51 more frequent (Sarhadi et al., 2018; Yu and Zhai, 2020).

52  
53 Drought and heat waves are also associated with fire weather, related through high temperatures, low soil  
54 moisture, and low humidity. Fire weather refers to weather conditions conducive to triggering and sustaining  
55 wildfires, which generally include temperature, soil moisture, humidity, and wind (Chapter 12). Concurrent

1 hot and dry conditions amplify conditions that promote wildfires (Schubert et al., 2014; Littell et al., 2016;  
2 Hope et al., 2019, Dowdy, 2018). Burnt area extent in western US forests (Abatzoglou and Williams, 2016)  
3 and particularly in California (Williams et al., 2019) has been linked to anthropogenic climate change via a  
4 significant increase in vapour pressure deficit, a primary driver of wildfires. A study of the western US  
5 examined the correlation between historical water-balance deficits and annual area burned, across a range of  
6 vegetation types from temperate rainforest to desert (McKenzie and Littell, 2017). The relationship between  
7 temperature and dryness, and wildfire, varied with ecosystem type, and the fire-climate relationship was both  
8 nonstationary and vegetation-dependent. In many fire-prone regions, such as the Mediterranean and China's  
9 Daxing'anling region, projections for increased severity of future drought and heat waves may lead to an  
10 increased frequency of wildfires relative to observed (Ruffault et al., 2018; Tian et al., 2017). Observations  
11 show a long-term trend towards more dangerous weather conditions for bushfires in many regions of  
12 Australia, which is attributable at least in part to anthropogenic climate change (Dowdy, 2018). There is  
13 emerging evidence that recent regional surges in wildland fires are being driven by changing weather  
14 extremes (SRCCL Ch2, Cross-Chapter Box 3; Jia et al., 2019). Between 1979 and 2013, the global burnable  
15 area affected by long fire-weather seasons doubled, and the mean length of the fire-weather season increased  
16 by 19% (Jolly et al., 2015). However, at the global scale, the total burned area has been decreasing between  
17 1998 and 2015 due to human activities mostly related to changes in land use (Andela et al., 2017). Given the  
18 projected *high confidence* increase in compound hot and dry conditions, there is *high confidence* that fire  
19 weather conditions will become more frequent at higher levels of global warming in some regions. This  
20 assessment is also consistent with assessments of Chapter 12 for regional projected changes in fire weather.  
21 The SRCCL Ch2 assessed with *high confidence* that future climate variability is expected to enhance the risk  
22 and severity of wildfires in many biomes such as tropical rainforests.

23  
24 In summary, there is *high confidence* that concurrent heat waves and droughts have increased in frequency  
25 over the last century at the global scale due to human influence. There is *medium confidence* that weather  
26 conditions that promote wildfires (fire weather) have become more probable in southern Europe, northern  
27 Eurasia, the US, and Australia over the last century. There is *high confidence* that compound hot and dry  
28 conditions become more probable in nearly all land regions as global mean temperature increases. There is  
29 *high confidence* that fire weather conditions will become more frequent at higher levels of global warming in  
30 some regions.

31  
32  
33 [START BOX 11.4 HERE]

34  
35  
36 **BOX 11.4: Case study: Global-scale concurrent climate anomalies at the example of the 2015-2016  
37 extreme El Niño and the 2018 boreal spring/summer extremes**

38  
39 Occurrence of concurrent or near-concurrent extremes in different parts of a region, or in different locations  
40 around the world challenges adaptation and risk management capacity. This can occur as a result of natural  
41 climate variability, as climates in different parts of the world are inter-connected through teleconnections. In  
42 addition, in a warming climate, the probability of having several locations being affected simultaneously by  
43 e.g. hot extremes and heat waves increases strongly as a function of global warming, with detectable changes  
44 even for changes as small as +0.5°C of additional global warming (Sections 11.2.5 and 11.3, Cross-chapter  
45 Box 11.1). Recent articles have highlighted the risks associated with concurrent extremes over large spatial  
46 scales (e.g. Lehner and Stocker, 2015; Boers et al., 2019; Gaupp et al., 2019). There is evidence that such  
47 global-scale extremes associated with hot temperature extremes are increasing in occurrence (Sippel et al.,  
48 2015; Vogel et al., 2019). Hereafter, the focus is on two recent global-scale events that featured concurrent  
49 extremes in several regions across the world. The first focuses on concurrent extremes driven by variability  
50 in tropical Pacific SSTs associated with the 2015-2016 extreme El Niño, while the second is a case study of  
51 the impacts of global warming combined with abnormal atmospheric circulation patterns in the 2018 boreal  
52 spring/summer.

53  
54  
55 [START BOX 11.4, FIGURE 1 HERE]

1 **Box 11.4, Figure 1:** Analysis of the percentage of land area affected by temperature extremes larger than two (orange)  
2 or three (blue) standard deviations in June-July-August (JJA) between 30°N and 80°N using a  
3 normalization. The more appropriate estimate is the corrected normalization. These panels show  
4 for both estimates a substantial increase in the overall land area affected by very high hot extremes  
5 since 1990 onward. Adapted from Sippel et al. (2015)  
6

7 **[END BOX 11.4, FIGURE 1 HERE]**  
8  
9

### 10 **The extreme El Niño in 2015-2016**

11  
12 El Niño-Southern Oscillation (ENSO) is one of the phenomena that have the ability to bring multitudes of  
13 extremes in different parts of the world, especially in the extreme cases of El Niño (Annex VI.4).  
14 Additionally, the background climate warming associated with greenhouse gas forcing can significantly  
15 exacerbate extremes in parts of the world even under normal El Niño conditions. The 2015-2016 El Niño  
16 event was one of the three extreme El Niño events since 1980s since the availability of satellite rainfall  
17 observations. According to some measures, it was the strongest El Niño over the past 145 years (Barnard et  
18 al., 2017). The 2015-2016 warmth was unprecedented at the central equatorial Pacific (Niño4: 5°N–5°S,  
19 150°E–150°W) and this exceptional warmth was *unlikely* to have occurred entirely naturally, appearing to  
20 reflect an anthropogenically forced trend (Newman et al., 2018)). In particular, its signal was seen in very  
21 high monthly Global Mean Surface Temperature (GMST) values in late 2015 and early 2016, contributing to  
22 the highest record of GMST in 2016 (Section 2.3.1.1). Both the ENSO amplitude and the frequency of high-  
23 magnitude events since 1950 is higher than over the pre-industrial period (*medium confidence*; Section  
24 2.4.2), suggesting that global extremes similar to those associated with the 2015-2016 El Niño would occur  
25 more frequently under further increases in global warming. Hereafter, the 2015-2016 El Niño event is  
26 referred to as “the 2015-2016 extreme El Niño” (Annex VI.4.1). A brief summary of extreme events that  
27 happened in 2015-2016 is provided in Section 6.2.2, 6.5.1.1 of the Special Report on the Ocean and  
28 Cryosphere in a Changing Climate (SROCC’s). We provide some highlights illustrating extremes that  
29 occurred in different parts of the world during the 2015-2016 extreme El Niño in BOX 11.4-Table 1, as well  
30 as a short summary hereafter.  
31

32 Several regions were strongly affected by droughts in 2015, including Indonesia, Australia, the Amazon  
33 region, Ethiopia, Southern Africa, and Europe. As a result, global measurements of land water anomalies  
34 were particularly low in that year (Humphrey et al., 2018). In 2015, Indonesia experienced a severe drought  
35 and forest fire causing pronounced impact on economy, ecology and human health due to haze crisis (Field  
36 et al., 2016; Huijnen et al., 2016; Patra et al., 2017; Hartmann et al., 2018). The northern part of Australia  
37 experienced high temperatures and low precipitation between late 2015 and early 2016, and the extensive  
38 mangrove trees were damaged along the Gulf of Carpentaria in northern Australia (Duke et al., 2017). The  
39 Amazon region experienced the most intense droughts of this century in 2015-2016. This drought was more  
40 severe than the previous major droughts that occurred in the Amazon in 2005 and 2010 (Lewis et al., 2011;  
41 Erfanian et al., 2017; Panisset et al., 2018). The 2015-2016 Amazon drought impacted the entirety of South  
42 America north of 20°S during the austral spring and summer (Erfanian et al., 2017). It also increased forest  
43 fire incidence by 36% compared to the preceding 12 years (Aragão et al., 2018) and as a consequence,  
44 increased the biomass burning outbreaks and the carbon monoxide (CO) concentration in the area, affecting  
45 air quality (Ribeiro et al., 2018). This out-of-season drought affected the water availability for human  
46 consumption and agricultural irrigation and it also left rivers with very low water levels, without conditions  
47 of ship transportation, due to large sandbanks, preventing the arrival of food, medicines, and fuels (INMET,  
48 2017). Eastern African countries were impacted by drought in 2015. It was found that the drought in  
49 Ethiopia, which was the worst in several decades, was associated with the 2015-2016 extreme El Niño that  
50 developed early in the year (Blunden and Arndt, 2016; Philip et al., 2018b). It was suggested that  
51 anthropogenic warming contributed to the 2015 Ethiopian and southern African droughts by increasing SSTs  
52 and local air temperatures (Funk et al., 2016, 2018b; Yuan et al., 2018a). It has also been suggested that the  
53 2015-2016 extreme El Niño affected circulation patterns in Europe during the 2015-2016 winter (Geng et al.,  
54 2017; Scaife et al., 2017).  
55

1 It was identified that 2015 was a year of a particularly high CO<sub>2</sub> growth rate, possibly related to some of the  
2 mentioned droughts, in particular in Indonesia and the Amazon region, leading to higher CO<sub>2</sub> release in  
3 combination with less CO<sub>2</sub> uptake from land areas (Humphrey et al., 2018). The impact of the 2015-2016  
4 extreme El Niño on vegetation systems via drought was also shown from satellite data (Kogan and Guo,  
5 2017). Overall, tropical forests were a carbon source to the atmosphere during the 2015–2016 El Niño–  
6 related drought, with some estimates suggesting that up to 2.3 PgC were released (Brando et al., 2019).  
7

8 The 2015-2016 extreme El Niño has induced extreme precipitation in some regions. Severe rainfall events  
9 were observed in Chennai city in India in December 2015 and Yangtze river region in China in June-July  
10 2016, and it was shown that these rainfall events are partly attributed to the 2015-2016 extreme El Niño (van  
11 Oldenborgh et al., 2016; Boyaj et al., 2018; Sun and Miao, 2018; Yuan et al., 2018b; Zhou et al., 2018).  
12

13 In 2015, the activity of tropical cyclones was notably high in the North Pacific (Blunden and Arndt, 2016).  
14 Over the western North Pacific, the number of category 4 and 5 Tropical Cyclones (TCs) was 13, which is  
15 more than twice its typical annual value of 6.3 (Zhang et al., 2016a). Similarly, a record-breaking number of  
16 TCs was observed in the eastern North Pacific, particularly in the western part of that domain (Collins et al.,  
17 2016; Murakami et al., 2017a). These extraordinary TC activities were related to the average SST anomaly  
18 during that year, which were associated with the 2015-2016 extreme El Niño and the positive phase of the  
19 Pacific Meridional Mode (PMM) (Murakami et al., 2017a; Hong et al., 2018; Yamada et al., 2019).  
20 However, it has been suggested that the intense TC activities in both the western and the eastern North  
21 Pacific in 2015 were not only due to the El Niño, but also to a contribution of anthropogenic forcing  
22 (Murakami et al., 2017a; Yang et al., 2018d). The impact of the Indian Ocean SST also was suggested to  
23 contribute to the extreme TC activity in the western North Pacific in 2015 (Zhan et al., 2018). In contrast, in  
24 Australia, it was the least active TC season since satellite records began in 1969-70 (Blunden and Arndt,  
25 2017).  
26  
27

#### 28 [START BOX 11.4, TABLE 1 HERE]

29 **Box 11.4, Table 1:** List of events related to the 2015-2016 Extreme El Niño in the literature.  
30  
31

Region	Period	Events	References
Indonesia	July 2015 to June 2016	droughts, forest fire	(Field et al., 2016; Huijnen et al., 2016; Patra et al., 2017; Hartmann et al., 2018)
Northern Australia	Between late 2015 and early 2016	high temperature and drought	(Duke et al., 2017)
Amazon	September 2015 to May 2016	droughts, forest fire	(Jiménez-Muñoz et al., 2016; Erfanian et al., 2017; Aragão et al., 2018; Panisset et al., 2018; Ribeiro et al., 2018)
The entirety of South America north of 20°S	Austral spring and 2015-2016 summer	droughts	(Erfanian et al., 2017)
Ethiopia	February-September 2015	droughts	(Blunden and Arndt, 2016; Philip et al., 2018b)
Southern Africa	November 2015–April 2016	droughts	(Funk et al., 2016, 2018a; Blamey et al., 2018; Yuan et al., 2018a)
Europe	Boreal 2015-2016 winter	effects on of circulation patterns	(Geng et al., 2017; Scaife et al., 2017)
India	May 2016	high temperature	(van Oldenborgh et al., 2018)
India	December 2015	extreme rainfall	(van Oldenborgh et al., 2016; Boyaj et al., 2018)
China	June-July 2016	extreme rainfall	(Sun and Miao, 2018; Yuan et al., 2018b; Zhou et al., 2018)
Western North Pacific	Boreal summer 2015	the large number (13) of category 4 and 5 tropical cyclones	(Blunden and Arndt, 2016; Mueller et al., 2016a; Zhang et al., 2016b; Hong et al., 2018; Yamada et al., 2019)
Eastern North Pacific	Boreal summer 2015	a record-breaking number of tropical cyclones	(Collins et al., 2016; Murakami et al., 2017a)
Global	2015-2016 El Niño	high CO <sub>2</sub> release to the atmosphere associated with	(Humphrey et al., 2018; Brando et al., 2019)

		droughts and fires in several affected regions	
--	--	--	--

1  
2  
3 **[END BOX 11.4, TABLE 1 HERE]**  
4  
5

6 **Global-scale temperature extremes and concurrent precipitation extremes in boreal 2018 spring and summer**  
7

8  
9 In the 2018 boreal spring-summer season (May-August), wide areas of the mid-latitudes in the Northern Hemisphere experienced heat extremes and in part enhanced drought (Kornhuber et al., 2019; Vogel et al., 2019; Box 11.3, Figure 2). The reported impacts included the following (Vogel et al., 2019): 90 deaths from heat strokes in Quebec (Canada), 1469 deaths from heat strokes in Japan (Shimpo et al., 2019a), 48 heat-related deaths in South Korea (Min et al., 2020), heat warning affecting 90,000 students in the USA, fires in numerous countries (Canada (British Columbia), USA (California), Lapland, Latvia), crop losses in the UK, Germany and Switzerland (Vogel et al., 2019) and overall in central and northern Europe (leading to yield reductions of up to 50% for the main crops; Toreti et al., 2019), fish deaths in Switzerland, and melting of roads in the Netherlands and the UK, among others. In addition to the numerous hot and dry extremes, an extremely heavy rainfall event occurred over wide areas of Japan from 28 June to 8 July 2018 (Tsuguti et al., 2018), which was followed by a heat wave (Shimpo et al., 2019b). The heavy precipitation event caused more than 230 deaths in Japan, and was named as “the Heavy Rain Event of July 2018”.

21  
22 The heavy precipitation event was characterized by unusually widespread and persistent rainfall and locally anomalous total precipitation led by band-shaped precipitation systems, which are frequently associated with heavy precipitation events in East Asia (Kato, 2020; Section 11.7.3). The extreme rainfall in Japan was caused by anomalous moisture transport with a combination of abnormal jet condition (Takemi and Unuma, 2019; Takemura et al., 2019; Tsuji et al., 2019; Yokoyama et al., 2020), which can be viewed as an atmospheric river (Yatagai et al., 2019; Sections 8.2.2.8, 11.7.2) caused by intensified inflow velocity and high SST around Japan (Kawase et al., 2019; Sekizawa et al., 2019).

29  
30 This precipitation event and the subsequent heat wave are related to abnormal condition of the jet and North Pacific Subtropical High in this month (Shimpo et al., 2019a; Ren et al., 2020), which caused extreme conditions from Europe, Eurasia, and North America (Kornhuber et al., 2019; Box 11.4, Figure 2). A role of Atlantic SST anomaly on the meandering jets and the subtropical high have been suggested (Liu et al., 2019a). These dynamic and thermodynamic components generally have substantial influence on extreme rainfall in East Asia (Oh et al., 2018), but it is under investigation whether these factors were due to anthropogenic forcing.

37  
38  
39 **[START BOX 11.4, FIGURE 2 HERE]**

40 **Box 11.4, Figure 2:** Meteorological conditions in July 2018. The color shading shows the monthly mean near-surface air temperature anomaly with respect to 1981 to 2010. Contour lines indicate the geopotential height in m, highlighted are the isolines on 12'000 m and 12'300 m, which indicate the approximate positions of the polar-front jet and subtropical jet, respectively. The light blue-green ellipse shows the approximate extent of the strong precipitation event that occurred at the beginning of July in the region of Japan and Korea. All data is from the global ECMWF Reanalysis v5 (ERA5, Hersbach et al., 2020).

47 **[END BOX 11.4, FIGURE 2 HERE]**  
48  
49

50 Regarding the hot extremes that occurred across the Northern Hemisphere in the 2018 boreal May-July time period, Vogel et al. (2019) found that the event was unprecedented in terms of the total area affected by hot extremes (on average about 22% of populated and agricultural areas in the Northern Hemisphere) for that

1 period, but was consistent with a +1°C climate which was the estimated global mean temperature anomaly  
2 around that time (for 2017; SR1.5). This study also found that events similar to the 2018 May-July  
3 temperature extremes would approximately occur 2 out of 3 years under +1.5°C of global warming, and  
4 every year under +2°C of global warming. Imada et al. (2019) also suggests that the mean annual occurrence  
5 of extremely hot days in Japan will be expected to increase by 1.8 times under a global warming level of 2°C  
6 above pre-industrial levels. Kawase et al. (2019) showed that the extreme rainfall in Japan during this event  
7 was increased by approximately 7% due to recent rapid warming around Japan. Hence, it is *virtually certain*  
8 that these 2018 concurrent events would not have occurred without human-induced global warming.  
9 Concurrent events of this type are also projected to happen more frequently under higher levels of global  
10 warming. On the other hand, there is currently *low confidence* in projected changes in the frequency or  
11 strength of the anomalous circulation patterns leading to concurrent extremes (e.g. Cross-Chapter Box 10.1).  
12

13 The case studies presented in this Box illustrate the current state of knowledge regarding the contribution of  
14 human-induced climate change to recent concurrent extremes in the global domain. Recent years have seen a  
15 more frequent occurrence of such events. The heat wave in Europe in the 2019 boreal summer and its  
16 coverage in the global domain is an additional example (Vautard et al., 2020a). However, there are still very  
17 few studies investigating which types of concurrent extreme events could occur under increasing global  
18 warming. It has been noted that such events could also be of particular risk for concurrent impacts in the  
19 world's breadbaskets (Zampieri et al., 2017; Kornhuber et al., 2020).  
20

21 In summary, the 2015-2016 extreme El Niño and the 2018 boreal spring/summer extremes were two  
22 examples of recent concurrent extremes. The El Niño event in 2015-2016 was one of the three extreme El  
23 Niño events since 1980s and there are many extreme events concurrently observed in this period including  
24 droughts, heavy precipitation, and more frequent intense tropical cyclones. Both the ENSO amplitude and  
25 the frequency of high-magnitude events since 1950 is higher than over the pre-industrial period (*medium*  
26 *confidence*), suggesting that global extremes similar to those associated with the 2015-2016 El Niño would  
27 occur more frequently under further increases in global warming. The 2018 boreal spring/summer extremes  
28 were characterized by heat extremes and enhanced droughts in wide areas of the mid-latitudes in the  
29 Northern Hemisphere and extremely heavy rainfall in East Asia. These concurrent events were generally  
30 related to abnormal condition of the jet and North Pacific Subtropical High, but also amplified by  
31 background global warming. It is *virtually certain* that these 2018 concurrent extreme events would not have  
32 occurred without human-induced global warming. Recent years have seen a more frequent occurrence of  
33 such concurrent events. However, it is still unknown which types of concurrent extreme events could occur  
34 under increasing global warming.  
35

36  
37 **[END BOX 11.4 HERE]**  
38  
39

## 40 11.9 Regional information on extremes

41  
42 This section complements the assessments of changes in temperature extremes (Section 11.3), heavy  
43 precipitation (Section 11.4), and droughts (Section 11.6), by providing additional regional details. Owing to  
44 the large number of regions and space limitations, the regional assessment for each of the AR6 reference  
45 regions (see Section 1.5.2.2 for a description) is presented here in a set of tables. The tables are organized  
46 according to types of extremes (temperature, heavy precipitation, droughts) for Africa (Tables 11.4-11.6),  
47 Asia (Table 11.7-11.9), Australasia (Tables 11.10-11.12), Central and South America (Tables 11.13-11.15),  
48 Europe (Tables 11.16-11.18), and North America (Tables 11.19-11.21). Each table contains regional  
49 assessments for observed changes, the human contribution to the observed changes, and projections of  
50 changes in these extremes at 1.5°C, 2°C and 4°C of global warming. Expanded versions of the tables with  
51 full evidence and rationale for assessments are provided in the Chapter Appendix (Tables 11.A.4-11.A.21).  
52  
53

### 54 11.9.1 Overview

55 Sections 11.9.2, 11.9.3., and 11.9.4 provide brief summaries of the underlying evidence used to derive the  
56 **Do Not Cite, Quote or Distribute**

1 regional assessments for temperature extremes, heavy precipitation events, and droughts, respectively. The  
2 assessments take into account evidence from studies based on global datasets (global studies), as well as  
3 regional studies. Global studies include analyses for all continents and AR6 regions with sufficient data  
4 coverage, and provide an important basis for cross-region consistency, as the same data and methods are  
5 used for all regions. However, individual regional studies may include additional information that is missed  
6 in global studies and thus provide an important regional calibration for the assessment.

7  
8 The assessments are presented using the calibrated confidence and likelihood language (Box 1.1). *Low*  
9 *confidence* is assessed when there is *limited evidence*, either because of a lack of available data in the region  
10 and/or a lack of relevant studies. *Low confidence* is also assessed when there is a lack of agreement on the  
11 evidence of a change, which may be due to large variability or inconsistent changes depending on the  
12 considered subregions, time frame, models, assessed metrics, or studies. In cases when the evidence is  
13 strongly contradictory, for example with substantial regional changes of opposite sign, “mixed signal” is  
14 indicated. With an assessment of *low confidence*, the direction of change is not indicated in the tables. A  
15 direction of change (increase or decrease) is provided with an assessment of *medium confidence*, *high*  
16 *confidence*, *likely*, or higher likelihood levels. Likelihood assessments are only provided in the case of *high*  
17 *confidence*. In some cases, there may be confidence in a small or no change.

18  
19 For projections, changes are assessed at three global warming levels (GWLs, CC-Box 11.1): 1.5°C, 2°C and  
20 4°C. Literature based both on GWL projections and on scenario-based projections is used for the  
21 assessments. In the case of literature on scenario-based projections, a mapping between scenarios/time  
22 frames and GWLs was performed as documented in CC-Box 11.1. Projections of changes in temperature and  
23 precipitation extremes are assessed relative to two different baselines: the recent past (1995-2014) and pre-  
24 industrial (1850-1900). With smaller changes relative to the variability, in particular because droughts  
25 happen on longer timescales compared to extremes of daily temperature and precipitation, it is more difficult  
26 to distinguish changes in drought relative to the recent past. As such, changes in droughts are assessed  
27 relative to the pre-industrial baseline, unless indicated otherwise.

### 28 29 30 **11.9.2 Temperature extremes**

31  
32 Tables 11.4, 11.7, 11.10, 11.13, 11.16, and 11.19 include assessments for past chn temperature extremes and  
33 their attribution, as well as future projections. The evidence is mostly drawn from changes in metrics based  
34 on daily maximum and minimum temperatures, similar to those used in Section 11.3. The regional  
35 assessments start from global studies that used consistent analyses for all regions globally with sufficient  
36 data. This includes Dunn et al. (2020) for observed changes and Li et al. (2020) and the Chapter 11  
37 Supplementary Material (11.SM) for projections with the CMIP6 multi-model ensemble. Evidence from  
38 regional studies, and those based on the CMIP5 multi-model ensemble or CORDEX simulations, are then  
39 used to refine the confidence assessments. For attribution, Seong et al. (2020) provide a consistent analysis  
40 for AR6 regions and Wang et al. (2017) for SREX regions. Additional regional studies, including event  
41 attribution analyses (Section 11.2), are used when available. In some regions that were not analysed in Seong  
42 et al. (2020) and with no known event attribution studies, *medium confidence* of a human contribution is  
43 assessed when there is strong evidence of changes from observations that are in the direction of model  
44 projected changes for the future, the magnitude of projected changes increases with global warming, and  
45 there is no other evidence to the contrary. Understanding of how temperature extremes change with the mean  
46 temperature and overwhelming evidence of a human contribution to the observed larger-scale changes in the  
47 mean temperature and temperature extremes further support this assessment.

### 48 49 50 **11.9.3 Heavy precipitation**

1 Tables 11.5, 11.8, 11.11, 11.14, 11.17, and 11.20 include assessments for past changes in heavy precipitation  
2 events and their attribution, as well as future projections. The evidence is mostly drawn from changes in  
3 metrics based on one-day or five-day precipitation amounts, as addressed in Section 11.4. Similar to  
4 temperature extremes, the assessment of changes in heavy precipitation uses global studies, including Dunn  
5 et al. (2020) and Sun et al. (2020) for observed changes, and Li et al. (2020) and the Chapter 11  
6 Supplementary Material (11.SM) for projected changes using the CMIP6 multi-model ensemble. For  
7 attribution, Paik et al. (2020) provided continental analyses where data coverage was sufficient, but no  
8 attribution studies based on global data are available for the regional scale. For each region, regional studies,  
9 and studies based on the CMIP5 multi-model ensemble or CORDEX simulations, are also considered in the  
10 assessments for past changes, attribution, and projections.

11

12

### 13 **11.9.4 Droughts**

14

15 Tables 11.6, 11.9, 11.12, 11.15, 11.18, and 11.21 provide regional tables on past, attributed and projected  
16 changes in droughts. The assessment is subdivided in three drought categories corresponding to four drought  
17 types: i) meteorological droughts, ii) agricultural and ecological droughts, and iii) hydrological droughts (see  
18 Section 11.6). A list of metrics and global studies used for the assessments is provided below. The evidence  
19 from global studies is complemented in each continent with evidence from regional studies. An overview of  
20 studies considered for the assessments in projections is provided in Table 11.3.

21

22 Meteorological droughts are assessed based on observed and projected changes in precipitation-only metrics  
23 such as the Standardized Precipitation Index (SPI) and Consecutive Dry Days (CDD). Observed changes are  
24 assessed based on two global studies, Dunn et al. (2020) for CDD and Spinoni et al. (2019) for SPI. For  
25 projections, evidence for changes at 1.5°C and 2°C of global warming is drawn from Xu et al. (2019) and  
26 Touma et al. (2015) (based on RCP8.5 for 2010-2054 compared to 1961-2005) for SPI (CMIP5) and the  
27 Chapter 11 Supplementary Material (11.SM) for CDD (CMIP6). For projections at 4°C of global warming,  
28 evidence is drawn from several sources, including Touma et al. (2015) and Spinoni et al. (2020) for SPI  
29 (from CMIP5 and CORDEX, respectively), and the Chapter 11 Supplementary Material (11.SM) for CDD  
30 (CMIP6). No global-scale studies are available for the attribution of meteorological drought, and thus this  
31 assessment is based on regional detection and attribution or event attribution studies.

32

33 Agricultural and ecological droughts are assessed based on observed and projected changes in total column  
34 soil moisture, complemented by evidence on changes in surface soil moisture, water-balance (precipitation  
35 minus evapotranspiration (ET)) and metrics driven by precipitation and atmospheric evaporative demand  
36 (AED) such as the SPEI and PDSI (Section 11.6). In the case of the latter, only studies including estimates  
37 based on the Penman-Monteith equation (SPEI-PM and PDSI-PM) are considered because of biases  
38 associated with temperature-only approaches (Section 11.6). In arid regions in which AED-based metrics can  
39 increase strongly in projections, more weight is given to soil moisture projections. For observed changes,  
40 evidence is drawn from several sources: Padrón et al. (2020) for changes in precipitation minus ET, as well  
41 as soil moisture from the multi-model Land Surface Snow and Soil Moisture Model Intercomparison Project  
42 within CMIP6 (LS3MIP, Van Den Hurk et al., 2016; Chapter 11 Supplementary Material (11.SM)); Greve et  
43 al. (2014) for changes in precipitation minus ET, and precipitation minus AED; Spinoni et al. (2019) for  
44 changes in SPEI-PM; and Dai and Zhao (2017) for changes in PDSI-PM. For projections at 1.5°C of global  
45 warming, evidence is drawn from Xu et al. (2019) based on CMIP5 and the Chapter 11 Supplementary  
46 Material (11.SM) based on CMIP6 for changes in total column and surface soil moisture, and from Naumann  
47 et al. (2018) for changes in SPEI-PM, based on EC-Earth simulations driven with SSTs from seven CMIP5  
48 ESMs. For projections at 2°C of global warming, evidence is drawn from Xu et al. (2019) based on CMIP5,  
49 and Cook et al. (2020) (SSP1-2.6, 2071-2100 compared to pre-industrial) and the Chapter 11 Supplementary  
50 Material (11.SM) based on CMIP6, for changes in total column and surface soil moisture; evidence is also  
51 drawn from Naumann et al. (2018) for changes in SPEI-PM. For projections at 4°C of global warming,  
52 evidence is mostly drawn from Cook et al. (2020) (SSP3-7.0, 2071-2100) and the Chapter 11 Supplementary  
53 Material (11.SM) based on CMIP6 for changes in total column and surface soil moisture, and from Vicente-  
54 Serrano et al. (2020) for changes in SPEI-PM based on CMIP5. No global-scale studies with regional-scale



1 information are available for the attribution of agricultural and ecological droughts, and thus this assessment  
 2 is based on regional detection and attribution or event attribution studies.

3  
 4 Hydrological droughts are assessed based on observed and projected changes in low flows, complemented  
 5 by information on changes in mean runoff. For observed changes, evidence is drawn from three studies (Dai  
 6 and Zhao, 2017; Gudmundsson et al., 2019, 2021). For projected changes at 1.5°C of global warming,  
 7 evidence is drawn from Touma et al. (2015) based on analyses of the Standardized Runoff Index (SRI)  
 8 (CMIP5, based on 2010-2054 compared to 1961-2005), complemented with regional studies when available.  
 9 For projected changes at 2°C of global warming, evidence is also drawn from Cook et al. (2020) for changes  
 10 in runoff in CMIP6 (Scenario SSP1-2.6, 2071-2100), and from Zhai et al. (2020) for changes in low flows  
 11 based on simulations with a single model. For projected changes at 4°C of global warming, evidence is  
 12 drawn from Touma et al. (2015) based on CMIP5 analyses of SRI, Cook et al. (2020) for changes in surface  
 13 and total runoff based on CMIP6, and Giuntoli et al. (2015) for changes in low flows based on the Inter-  
 14 Sectoral Impact Model Intercomparison Project (ISI-MIP) based on six Global Hydrological Models  
 15 (GHMs) and five GCMs, including an analysis of inter-model signal-to-noise ratio. One global-scale study  
 16 with regional-scale information is available for the attribution of hydrological droughts (Gudmundsson et al.,  
 17 2021), but only in a few AR6 regions. This information was complemented with evidence from regional  
 18 detection and attribution, and event attribution studies when available.

19  
 20  
 21 **[START TABLE 11.3 HERE]**

22  
 23 **Table 11.3:** Global analyses considered for the assessments of drought projections. “MET” refers to meteorological  
 24 droughts, “AGR/ECOL” to agricultural and ecological droughts, and “HYDR” to hydrological droughts  
 25

Reference	Model data	Index	Drought type	Projection horizon(s)	Baseline
Chapter 11 Suppl. Material (11.SM)	CMIP6	CDD, Soil moisture (total, surface)	MET	1.5°C, 2°C, 4°C	1850-1900
Cook et al. (2020)	CMIP6	Soil moisture (total, surface), Runoff (total, surface)	AGR/ECOL, HYDR	2071-2011, SSP1-2.6 (~2°C, CC-Box 11.1; Chapter 4, Table 4.2) 2071-2011, SSP3-7.3 (~4°C, CC-Box 11.1; Chapter 4, Table 4.2)	1850-1900
Xu et al. (2019)	CMIP5	SPI, Soil moisture (total, surface)	MET, AGR/ECOL	1.5°C, 2°C	1971-2000
Touma et al. (2015)	CMIP5	SPI, SRI	MET, HYDR	2010-2054, RCP8.5 (~1.5°C, CC-Box 11.1, 11.SM.1) 2055-2099, RCP8.5 (~3.5°C, CC-Box 11.1, 11.SM.1)	1961-2005
Spinoni et al. (2020)	CORDEX (CMIP5 driving GCMs, RCMs)	SPI	MET	2071-2100, RCP4.5 (~2.5°C, CC-Box 11.1, 11.SM.1) 2071-2100, RCP8.5 (~4.5°C, CC-Box 11.1, 11.SM.1)	1981-2010
Naumann et al. (2018)	1 GCM (EC-EARTH3-HR v3.1) driven with SST fields from 7 CMIP5 GCMs	SPEI-PM	AGR/ECOL	1.5°C, 2°C, (3°C)	0.6°C
Vicente-Serrano et al. (2020)	CMIP5	SPEI-PM	AGR/ECOL	2070-2100, RCP8.5 (~4.5°C, CC-Box 11.1, 11.SM.1)	1970-2000
Giuntoli et al. (2015)	ISI-MIP (6 GHMs and 5 CMIP5 GCMs)	Low-flows days	HYDR	2066-2099, RCP8.5 (~4°C, CC-Box 11.1, 11.SM.1)	1972-2005
Zhai et al. (2020)	1 GHM (VIC) driven by 4 CMIP5 GCMs	Extreme low runoff	HYDR	1.5°C, 2°C	2006-2015

26  
 27  
 28 **[END TABLE 11.3 HERE]**  
 29  
 30

## 1 Frequently Asked Questions

2

### 3 **FAQ 11.1: How do changes in climate extremes compare with changes in climate averages?**

4 *Human-caused climate change alters the frequency and intensity of climate variables (e.g., surface*  
5 *temperature) and phenomena (e.g., tropical cyclones) in a variety of ways. We now know that the ways in*  
6 *which average and extreme conditions have changed (and will continue to change) depend on the variable*  
7 *and the phenomenon being considered. Changes in local surface temperature extremes follow closely the*  
8 *corresponding changes in local average surface temperatures. On the contrary, changes in precipitation*  
9 *extremes (heavy precipitation) generally do not follow those in average precipitation and can even move in*  
10 *the opposite direction (e.g., with average precipitation decreasing but extreme precipitation increasing).*

11 Climate change will manifest very differently depending on which region, which season and which variable  
12 we are interested in. For example, over some parts of the Arctic, temperatures will warm at rates about 3-4  
13 times higher during winter compared to summer months. And in summer, most of northern Europe will  
14 experience larger temperature increases than most places in Southeast South America and Australasia, with  
15 differences that can be larger than 1°C depending on the level of global warming. In general, differences  
16 across regions and seasons arise because the underlying physical processes differ drastically across regions  
17 and seasons.

18 Climate change will also manifest differently for different weather regimes and can lead to contrasting  
19 changes in average and extreme conditions. Observations of the recent past and climate model projections  
20 show that, in most places, changes in daily temperatures are dominated by a general warming in which both  
21 the climatological average and extreme values are shifted towards higher temperatures, making warm  
22 extremes more frequent and cold extremes less frequent. The top panels in FAQ 11.1, Figure 1 show  
23 projected changes in surface temperature for long-term average conditions (left) and for extreme hot days  
24 (right) during the warm season (summer in mid- to high-latitudes). Projected increases in long-term average  
25 temperature differ substantially in different places, varying from less than 3°C in some places in central  
26 South Asia and southern South America to over 7°C in some places in North America, north Africa and the  
27 Middle East. Changes in extreme hot days follow changes in average conditions quite closely, although in  
28 some places the warming rates for extremes can be intensified (e.g., southern Europe and the Amazon basin)  
29 or weakened (e.g., northern Asia and Greenland) compared to average values.

30 Recent observations and global and regional climate model projections point to changes in precipitation  
31 extremes (including both rainfall and snowfall extremes) differing drastically from those in average  
32 precipitation. The bottom panels in FAQ 11.1, Figure 1 show projected changes in the long-term average  
33 precipitation (left) and in heavy precipitation (right). Averaged precipitation changes show striking regional  
34 differences, with substantial drying in places such as southern Europe and northern South America and  
35 wetting in places such as Middle East and southern South America. Changes in extreme heavy precipitation  
36 are much more uniform, with systematic increases over nearly all land regions. The physical reasons behind  
37 the different response of averaged and extreme precipitation are now well understood. The intensification of  
38 extreme precipitation is driven by the increase in atmospheric water vapour (about 7% per 1°C of warming  
39 near the surface), although this is modulated by various dynamical changes. In contrast, changes in average  
40 precipitation are driven not only by moisture increases but also by slower processes that constrain future  
41 changes to on be only about 2–3% per 1°C of warming near the surface.

42 In summary, the specific relationship between changes in average and extreme conditions strongly depends  
43 on the variable or phenomenon being considered. At the local scale, average and extreme surface  
44 temperature changes are strongly related, while average and extreme precipitation changes are often weakly  
45 related. For both variables, the changes in average and extreme conditions vary strongly across different  
46 places due to the effect of local and regional processes.

47

### 48 **[START FAQ 11.1, FIGURE 1 HERE]**

**FAQ 11.1, Figure 1:** Global maps of future changes in surface temperature (top panels) and precipitation (bottom panels) for long-term average (left) and extreme conditions (right). All changes were estimated using the CMIP6 ensemble mean for a scenario with a global warming of 4°C relative to 1850-

1900 temperatures. Average surface temperatures refer to the warmest three-month season (summer in mid- to high-latitudes) and extreme temperature refer to the hottest day in a year. Precipitation changes, which can include both rainfall and snowfall changes, are normalized by 1850-1900 values and shown in percentage; extreme precipitation refers to the largest daily rainfall in a year.

1 **[END FAQ 11.1, FIGURE 1 HERE]**

2

3 **[END FAQ 11.1 HERE]**

4

1 **[START FAQ 11.2 HERE]**

2  
3 **FAQ 11.2: Will unprecedented extremes occur as a result of human-induced climate change?**

4  
5 *Climate change has already increased the magnitude and frequency of extreme hot events and decreased the*  
6 *magnitude and frequency of extreme cold events, and, in some regions, intensified extreme precipitation*  
7 *events. As the climate moves away from its past and current states, we will experience extreme events that*  
8 *are unprecedented, either in magnitude, frequency, timing or location. The frequency of these unprecedented*  
9 *extreme events will increase with increasing global warming. Additionally, the combined occurrence of*  
10 *multiple unprecedented extremes may result in large and unprecedented impacts.*

11  
12 Human-induced climate change has already affected many aspects of the climate system. In addition to the  
13 increase in global surface temperature, many types of weather and climate extremes have changed. In most  
14 regions, the frequency and intensity of hot extremes have increased and those of cold extremes have  
15 decreased. The frequency and intensity of heavy precipitation events have increased at a global scale and  
16 over a majority of land regions. Although extreme events such as land and marine heatwaves, heavy  
17 precipitation, drought, tropical cyclones, and associated wildfires and coastal flooding have occurred in the  
18 past and will continue to occur in the future, they often come with different magnitudes or frequencies in a  
19 warmer world. For example, future heatwaves will last longer and have higher temperatures, and future  
20 extreme precipitation events will be more intense in several regions. Certain extremes, such as extreme cold,  
21 will be less intense and less frequent with increasing warming.

22  
23 Unprecedented extremes – that is, events not experienced in the past – will occur in the future in five  
24 different ways (FAQ 11.2, Figure 1). First, events that are considered to be extreme in the current climate  
25 will occur in the future with unprecedented magnitudes. Second, future extreme events will also occur with  
26 unprecedented frequency. Third, certain types of extremes may occur in regions that have not previously  
27 encountered those types of events. For example, as the sea level rises, coastal flooding may occur in new  
28 locations, and wildfires are already occurring in areas, such as parts of the Arctic, where the probability of  
29 such events was previously low. Fourth, extreme events may also be unprecedented in their timing. For  
30 example, extremely hot temperatures may occur either earlier or later in the year than they have in the past.

31  
32 Finally, compound events, where multiple extreme events of either different or similar types occur  
33 simultaneously and/or in succession, may be more probable or severe in the future. These compound events  
34 can often impact ecosystems and societies more strongly than when such events occur in isolation. For  
35 example, a drought along with extreme heat will increase the risk of wildfires and agriculture damages or  
36 losses. As individual extreme events become more severe as a result of climate change, the combined  
37 occurrence of these events will create unprecedented compound events. This could exacerbate the intensity  
38 and associated impacts of these extreme events.

39  
40 Unprecedented extremes have already occurred in recent years, relative to the 20th century climate. Some  
41 recent extreme hot events would have had very little chance of occurring without human influence on the  
42 climate (see FAQ 11.3). In the future, unprecedented extremes will occur as the climate continues to warm.  
43 Those extremes will happen with larger magnitudes and at higher frequencies than previously experienced.  
44 Extreme events may also appear in new locations, at new times of the year, or as unprecedented compound  
45 events. Moreover, unprecedented events will become more frequent with higher levels of warming, for  
46 example at 3°C of global warming compared to 2°C of global warming.

47  
48 **[START FAQ 11.2, FIGURE 1 HERE]**

49  
50 **FAQ 11.2, Figure 1:** New types of unprecedented extremes that will occur as a result of climate change.

51  
52 **[END FAQ 11.2, FIGURE 1 HERE]**

53  
54 **[END FAQ 11.2 HERE]**

1 **[START FAQ 11.3 HERE]**

2  
3 **FAQ 11.3: Did climate change cause that recent extreme event in my country?**

4  
5 *While it is difficult to identify the exact causes of a particular extreme event, the relatively new science of*  
6 *event attribution is able to quantify the role of climate change in altering the probability and magnitude of*  
7 *some types of weather and climate extremes. There is strong evidence that characteristics of many individual*  
8 *extreme events have already changed because of human-driven changes to the climate system. Some types of*  
9 *highly impactful extreme weather events have occurred more often and have become more severe due to*  
10 *these human influences. As the climate continues to warm, the observed changes in the probability and/or*  
11 *magnitude of some extreme weather events will continue as the human influences on these events increase.*

12  
13 It is common to question whether human-caused climate change caused a major weather- and climate-related  
14 disaster. When extreme weather and climate events do occur, both exposure and vulnerability play an  
15 important role in determining the magnitude and impacts of the resulting disaster. As such, it is difficult to  
16 attribute a specific disaster directly to climate change. However, the relatively new science of event  
17 attribution enables scientists to attribute aspects of specific extreme weather and climate events to certain  
18 causes. Scientists cannot answer directly whether a particular event was caused by climate change, as  
19 extremes do occur naturally and any specific weather and climate event is the result of a complex mix of  
20 human and natural factors. Instead, scientists quantify the relative importance of human and natural  
21 influences on the magnitude and/or probability of specific extreme weather events. Such information is  
22 important for disaster risk reduction planning, because improved knowledge about changes in the probability  
23 and magnitude of relevant extreme events enables better quantification of disaster risks.

24  
25 On a case-by-case basis, scientists can now quantify the contribution of human influences to the magnitude  
26 and probability of many extreme events. This is done by estimating and comparing the probability or  
27 magnitude of the same type of event between the current climate – including the increases in greenhouse gas  
28 concentrations and other human influences – and an alternate world where the atmospheric greenhouse gases  
29 remained at pre-industrial levels. FAQ 11.3 Figure 1 illustrates this approach using differences in  
30 temperature and probability between the two scenarios as an example. Both the pre-industrial (blue) and  
31 current (red) climates experience hot extremes, but with different probabilities and magnitudes. Hot extremes  
32 of a given temperature have a higher probability of occurrence in the warmer current climate than in the  
33 cooler pre-industrial climate. Additionally, an extreme hot event of a particular probability will be warmer in  
34 the current climate than in the pre-industrial climate. Climate model simulations are often used to estimate  
35 the occurrence of a specific event in both climates. The change in the magnitude and/or probability of the  
36 extreme event in the current climate compared to the pre-industrial climate is attributed to the difference  
37 between the two scenarios, which is the human influence.

38  
39 Attributable increases in probability and magnitude have been identified consistently for many hot extremes.  
40 Attributable increases have also been found for some extreme precipitation events, including hurricane  
41 rainfall events, but these results can vary among events. In some cases, large natural variations in the climate  
42 system prevent attributing changes in the probability or magnitude of a specific extreme to human influence.  
43 Additionally, attribution of certain classes of extreme weather (e.g., tornadoes) is beyond current modelling  
44 and theoretical capabilities. As the climate continues to warm, larger changes in probability and magnitude  
45 are expected, and as a result it will be possible to attribute future temperature and precipitation extremes in  
46 many locations to human influences. Attributable changes may emerge for other types of extremes as the  
47 warming signal increases.

48  
49 In conclusion, human-caused global warming has resulted in changes in a wide variety of recent extreme  
50 weather events. Strong increases in probability and magnitude, attributable to human influence, have been  
51 found for many heat waves and hot extremes around the world.

52  
53  
54 **[START FAQ11.3 FIGURE 1 HERE]**

1 **FAQ 11.3, Figure 1:** Changes in climate result in changes in the magnitude and probability of extremes. Example of  
2 how temperature extremes differ between a climate with pre-industrial greenhouse gases (shown  
3 in blue) and the current climate (shown in orange) for a representative region. The horizontal  
4 axis shows the range of extreme temperatures, while the vertical axis shows the annual chance of  
5 each temperature event's occurrence. Moving towards the right indicates increasingly hotter  
6 extremes that are more rare (less probable). For hot extremes, an extreme event of a particular  
7 temperature in the pre-industrial climate would be more probable (vertical arrow) in the current  
8 climate. An event of a certain probability in the pre-industrial climate would be warmer  
9 (horizontal arrow) in the current climate. While the climate under greenhouse gases at the pre-  
10 industrial level experiences a range of hot extremes, such events are hotter and more frequent in  
11 the current climate.

12

13 **[END FAQ11.3 FIGURE 1 HERE]**

14

15 **[END FAQ11.3 HERE]**

16

17

1 **Large tables**

2 Color scale for tables for changes in temperature extremes and heavy precipitation

	Fact	Virtually certain	Extremely likely	Very likely	Likely	High confidence	Medium confidence	Low confidence
Increasing hot extremes, decreasing cold extremes								
Decreasing hot extremes, increasing cold extremes								
Inconsistent sign								

3  
4 Color scale for tables for changes in droughts

	Fact	Virtually certain	Extremely likely	Very likely	Likely	High confidence	Medium confidence	Low confidence
Increasing drought								
Decreasing drought								
Inconsistent sign								

5  
6 **[START TABLE 11.4 HERE]**

7  
8 **Table 11.4:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in Africa,  
9 subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details

All Africa	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
	Insufficient data for the continent, but there is <i>high confidence</i> of an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes in all subregions with sufficient data	Limited evidence for the continent, but there is <i>medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes for all subregions with sufficient data	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level ((Li et al., 2020)
	<i>Medium confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014))

		frequency of cold extremes.	Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	<i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)	<i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Mediterranean (MED) <sup>2</sup>	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Peña-Angulo et al., 2020; El Kenawy et al., 2013; Acero et al., 2014; Fioravanti et al., 2016; Ruml et al., 2017; Türkeş and Erlat, 2018; Donat et al., 2013, 2014, 2016; Filahi et al., 2016; Driouech et al., 2021; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Sippel and Otto, 2014; Wilcox et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Zollo et al., 2016; Weber et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Tomozeiu et al., 2014; Abaurrea et al., 2018; Nastos and Kapsomenakis, 2015; Cardell et al., 2020; Zollo et al., 2016; Weber et al., 2018; Coppola et al., 2021a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Nastos and Kapsomenakis, 2015; Tomozeiu et al., 2014; Cardell et al., 2020; Zollo et al., 2016; Giorgi et al., 2014; Driouech et al., 2020; Coppola et al., 2021a; Engelbrecht et al., 2015)

<sup>2</sup> This region includes both northern Africa and southern Europe  
Do Not Cite, Quote or Distribute



	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Sahara (SAH)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Donat et al., 2014a; Moron et al., 2016; Dunn et al., 2020)	<b>Strong evidence of changes from observations</b> that are in the direction of model projected changes for the future. The magnitude of projected changes increases with global warming.	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020 ) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium</i> confidence in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with pre-industrial)

			<i>Very likely</i> (compared with pre-industrial).	<i>Extremely likely</i> (compared with pre-industrial)	with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Western Africa (WAF)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Barry et al., 2018; Chaney et al., 2014; Dunn et al., 2020; Mouhamed et al., 2013; Perkins-Kirkpatrick and Lewis, 2020)	Strong evidence of changes from observations that are in the direction of model projected changes for the future. The magnitude of projected changes increases with global warming.	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Northern Eastern Africa (NEAF)	Increases in the intensity and frequency of hot extremes and decreases in the intensity and frequency of cold extremes (Perkins-Kirkpatrick and Lewis, 2020; Chaney et al., 2014; Gebrechorkos et al.,	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than

	2018; Omondi et al., 2014; Dunn et al., 2020)	extremes (Otto et al., 2015; Philip et al., 2020; Marthews et al., 2015; Kew et al., 2021; Funk et al., 2015)	the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Medium confidence</i> in the increase in the intensity and frequency of hot extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Central Africa (CAF)	Insufficient data to assess trends (Dunn et al., 2020)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of

			frequency of hot extremes (Weber et al., 2018)	hot extremes (Weber et al., 2018; Coppola et al., 2021a)	hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Low confidence</i>	<i>Low confidence</i>	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
South Eastern Africa (SEAF)	Increases in the intensity and frequency of hot extremes and decreases in the intensity and frequency of cold extremes (Perkins-Kirkpatrick and Lewis, 2020; Gebrechorkos et al., 2018; Omondi et al., 2014; Chaney et al., 2014)	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Otto et al., 2015; Philip et al., 2020; Marthews et al., 2015; Kew et al., 2021; Funk et al., 2015)	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)</p>
	<i>Medium confidence</i> in the increase in the intensity and frequency of hot extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>

			Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Western Southern Africa (WSAF)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Russo et al., 2016; Perkins-Kirkpatrick and Lewis, 2020; Kruger and Nxumalo, 2017; Mbokodo et al., 2020; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Eastern Southern Africa (ESAF)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and	Robust evidence of a human contribution to the observed increase in the intensity and frequency of	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity

	frequency of cold extremes (Dunn et al., 2020; Russo et al., 2016; Perkins-Kirkpatrick and Lewis, 2020; Kruger and Nxumalo, 2017; Mbokodo et al., 2020)	hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017)	and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Likely increase</i> in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Madagascar (MDG)	Increases in the intensity and frequency of hot extremes and decreases in the intensity and frequency of cold extremes (Vincent et al., 2011; Donat et al., 2013)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).

			Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018)	Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Weber et al., 2018; Coppola et al., 2021a)	Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in the intensity and frequency of hot extremes (Coppola et al., 2021a; Engelbrecht et al., 2015; Giorgi et al., 2014)
	<i>Medium confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Low confidence</i>	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

1

2 [END TABLE 11.4 HERE]

3

4 [START TABLE 11.5 HERE]

5

6 **Table 11.5:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in Africa, subdivided  
7 by AR6 regions. See Sections 11.9.1 and 11.9.3 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Africa	Insufficient data to assess trends	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation:

Do Not Cite, Quote or Distribute

11-130

Total pages: 345

					<i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Mediterranean (MED) <sup>3</sup>	Lack of agreement on the evidence of trends (Sun et al., 2020; Casanueva et al., 2014; de Lima et al., 2015; Gajić-Čapka et al., 2015; Ribes et al., 2019; Peña-Angulo et al., 2020; Rajczak and Schär, 2017; Jacob et al., 2018; Coppola et al., 2021a; Donat et al., 2014; Mathbout et al., 2018; Dunn et al., 2020)	Limited evidence (Añel et al., 2014; U.S. Department of Agriculture Economic Research Service, 2016)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Cardell et al., 2020; Zollo et al., 2016; Samuels et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1 day and Rx5 day events compared to the 1°C warming level (Li et al., 2020a) and more than 0% in annual Rx1 day and Rx5 day and less than -2% in annual Rx30 day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020; Zollo et al., 2016; Samuels et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1 day and Rx5 day events compared to the 1°C warming level (Li et al., 2020a) and more than 2% in annual Rx1 day and Rx5 day and less than -2% in annual Rx30 day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020; Trambly and Somot, 2018; Zollo et al., 2016; Samuels et al., 2018; Monjo et al., 2016; Rajczak et al., 2013; Coppola et al., 2021b; Driouech et al., 2020)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)
Sahara (SAH)	Insufficient data to assess trends (Sun et al., 2020; Dunn et al., 2020)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1 day and Rx5 day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1 day and Rx5 day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1 day,	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 30% in the 50-year Rx1 day and Rx5 day events

<sup>3</sup> This region includes both northern Africa and southern Europe  
Do Not Cite, Quote or Distribute



			annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	Rx5day, and Rx30day compared to pre-industrial (Annex).	compared to the 1°C warming level (Li et al., 2020a) and more than 40% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Western Africa (WAF)	Insufficient data and a lack of agreement on the evidence of trends (Mouhamed et al., 2013; Chaney et al., 2014; Sanogo et al., 2015; Zittis, 2017; Barry et al., 2018; Sun et al., 2020; Dunn et al., 2020)	Limited evidence (Parker et al., 2017)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Nikulin et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Nikulin et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 25% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 30% in annual Rx1day and Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Giorgi et al., 2014; Dosio et al., 2019; Akinsanola and Zhou, 2018; Coppola et al., 2021b)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
North Eastern Africa (NEAF)	Insufficient data to assess trends (Sun et al., 2020; Dunn et al., 2020)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020;

			50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).	events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	Annex). Median increase of more than 25% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 35% in annual Rx1day and Rx5day and 30% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Central Africa (CAF)	Insufficient data to assess trends (Sun et al., 2020; Dunn et al., 2020)	Limited evidence (Otto et al., 2013)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Nikulin et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Nikulin et al., 2018; Déqué et al., 2017; Coppola et al., 2021b)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 30% in annual Rx1day and Rx5day and 20% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Diedhiou et al. 2018; Fotso-Nguemo et al. 2018; Sonkoué et al. 2019; Coppola et al., 2021b)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

South Eastern Africa (SEAF)	Insufficient data to assess trends (Sun et al., 2020; Dunn et al., 2020)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 25% in annual Rx1day and Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
West Southern Africa (WSAF)	Intensification of heavy precipitation (Sun et al., 2020; Donat et al., 2013)	Limited evidence	CMIP6 models project inconsistent changes in the region (Li et al., 2020, Annex)	CMIP6 models project inconsistent changes in the region (Li et al., 2020, Annex)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Pinto et al., 2016; Dosio et al., 2019; Giorgi et al., 2014; Coppola et al., 2021b)
	<i>Medium confidence</i> in the intensification of heavy precipitation.	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014))

					<i>Likely</i> (compared with pre-industrial)
East Southern Africa (ESAF)	Intensification of heavy precipitation (Sun et al., 2020; Donat et al., 2013)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Pinto et al., 2016; Dosio et al., 2019; Giorgi et al., 2014; Coppola et al., 2021b)
	<i>Medium confidence</i> in the intensification of heavy precipitation.	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
Madagascar (MDG)	Insufficient data to assess trends and trends in available data are not significant (Sun et al., 2020; Dunn et al., 2020; Donat et al., 2013; Vincent et al., 2011)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Weber et al., 2018)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Weber et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).

					Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Weber et al., 2018)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)

1  
2  
3  
4  
5  
6  
7  
8  
9

[END TABLE 11.5 HERE]

[START TABLE 11.6 HERE]

**Table 11.6:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET), agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in Africa, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.4 for details.

Region and drought type		Observed trends	Human contribution	Projections		
				1.5 °C	2 °C	4 °C
MED <sup>4</sup>	MET	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED
	AGR ECOL	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED
	HYDR	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED	ENTRY IDENTICAL TO EU-MED
Sahara (SAH)	MET	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed signals</b> (seasonally and geographically varying) and non-robust changes (Cook et al., 2020). Slightly reduced drying based on CDD (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signals</b> (seasonally and geographically varying) and non-robust changes (Cook et al., 2020). Slightly reduced drying based on CDD (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signals</b> (seasonally and geographically varying) and non-robust changes (Cook et al., 2020). Reduced drying based on CDD (Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence</b> and inconsistent signals in CMIP6 (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Limited evidence and inconsistent signals</b> in CMIP6 (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Limited evidence and inconsistent signals</b> in CMIP6. (Cook et al., 2020; Vicente-Serrano et al.,

<sup>4</sup> This region includes both northern Africa and southern Europe

						2020a)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Inconsistent trends</b> (Touma et al., 2015; Cook et al., 2020)
Western Africa (WAF)	MET	<b>Medium confidence:</b> Increased drying based on CDD and SPI (Chaney et al., 2014; Barry et al., 2018; Spinoni et al., 2019; Dunn et al., 2020)	<b>Low confidence: Mixed signals</b> (Lawal et al., 2016; Knutson and Zeng, 2018).  Drying attributable in fraction of region to climate change over 1901-2010 and 1951-2010 time frames, but trend reversal from 1981-2010 (Knutson and Zeng, 2018)  No evidence that late onset of 2015 wet season in Nigeria was due to human contribution (Lawal et al., 2016)	<b>Low confidence: Mixed signal.</b> Mean increase of CDD over larger part of Guinea Coast in 25 CORDEX AFR runs, 1.5°C minus 1971-2000 (Klutse et al., 2016); slight increase in SPI-based meteorological drought frequency and magnitude in the Niger and Volta river basin in CORDEX simulations (Oguntunde et al., 2020); but inconsistent changes in CDD in CMIP6 GCMs (Diedhiou et al., 2018)(Chapter 11 Supplementary Material (11.SM)), as well as in mean precipitation in CMIP6 GCMs (Cook et al., 2020)	<b>Low confidence: Mixed signal.</b> Mean increase of CDD over larger part of Guinea Coast in 25 CORDEX AFR runs, 1.5°C minus 1971-2000 (Klutse et al., 2016); slight increase in SPI-based meteorological drought frequency and magnitude in the Niger and Volta river basin in CORDEX simulations (Oguntunde et al., 2020); but inconsistent changes in CDD in CMIP6 GCMs (Diedhiou et al., 2018)(Chapter 11 Supplementary Material (11.SM)), as well as in mean precipitation in CMIP6 GCMs (Cook et al., 2020)	<b>Medium confidence: Increase</b> in meteorological droughts, mostly on seasonal time scale. Seasonal CDD increases in the region for MAM and JJA (Dosio et al., 2019), increase in SPI-based meteorological drought frequency and magnitude in Niger and Volta river basins (Oguntunde et al., 2020); and slight increase in SPI-based meteorological drought for overall region (Spinoni et al., 2020). Mixed signal in annual CDD (Akisanola and Zhou, 2018; Dosio et al., 2019)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Medium confidence:</b> Increased drying based on water-balance estimates and SPEI-PM, with stronger signals for SPEI-PM (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent signals</b> (geographical and inter-model variations) in soil moisture and SPEI-PM (Naumann et al., 2018; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent signals</b> (geographical and inter-model variations) in soil moisture and SPEI-PM (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Mixed signal.</b> Inconsistent changes depending on subregion, indices, and season (Naumann et al., 2018; Cook et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)). Most projections show a drying in Western half of domain.
	HYDR	<b>Medium confidence:</b> Decrease in streamflow (Dai and Zhao, 2017; Trambly et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Inconsistent signal</b> (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Inconsistent projections and/or non-robust changes</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
North Eastern Africa (NEAF)	MET	<b>Low confidence: Mixed signals.</b> Increasing drought in part of the region, in particular in recent two decades; but decreasing drought trends in other part of domain (NOTE: wetting trend in Horn of Africa in Spinoni et al. 2019)(Funk	<b>Low confidence: Limited evidence</b> on attribution of long-term trends.  Robust evidence that recent meteorological drought events (in 2016 and 2017) are not attributable to anthropogenic climate	<b>Low confidence: Inconsistent trends.</b> Inconsistent and weak signals in SPI (Nguvava et al., 2019; Xu et al., 2019a), with high spatial variation (Nguvava et al., 2019); inconsistent signals in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent trends.</b> Inconsistent changes in CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Nguvava et al., 2019; Xu et al., 2019a); but tendency towards increase in mean precipitation (Cook et al., 2020).	<b>Medium confidence: Decrease</b> in meteorological drought (Sillmann et al., 2013b; Dosio et al., 2019; Cook et al., 2020; Spinoni et al., 2020)  Sillmann et al. (2013), (2081-2100)/1981-2000, rcp8.5, CMIP3-CMIP5

		et al., 2015a; Nicholson, 2017; Spinoni et al., 2019) “No trends in observations in Ethiopia”; “large variability” (Philip et al., 2018b)	change (Lott et al., 2013; Marthews et al., 2015; Uhe et al., 2017; Funk et al., 2018b; Otto et al., 2018a; Philip et al., 2018b; Kew et al., 2021)	Nguvava et al. (2019): projections at 1.5°C GWL in Cordex AFR data, compared to 1971-2000: non significant changes in SPI-12-based meteorological drought frequency and intensity.	Nguvava et al. (2019): projections at 2°C GWL in Cordex AFR data, compared to 1971-2000: non significant changes in SPI-12-based meteorological drought frequency and intensity.	Decrease of CDD  Dosio et al. (2019), (2070-2099/1981-2010), rcp 8.5, 23 RCM: Decrease in CDD
	AGR ECOL	<b>Low confidence: Inconsistent trends</b> (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b> because of lack of studies	<b>Low confidence: Inconsistent trends</b> (Naumann et al., 2018; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends</b> , but tendency to wetting (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Decrease</b> in soil moisture-based drought (Cook et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence on attribution of long-term trends</b> (Fenta et al., 2017)	<b>Low confidence: Limited evidence. One study shows lack of signal</b> (Touma et al., 2015)	<b>Low confidence: Limited evidence</b> due to lack of studies;inconsistent trends (Touma et al., 2015; Cook et al., 2020)	<b>Medium confidence: Decrease</b> in hydrological drought compared to pre-industrial conditions and recent past (Giuntoli et al., 2015; Cook et al., 2020) but some inconsistent signals (Touma et al., 2015)
Central Africa (CAF)	MET	<b>Medium confidence Decrease</b> in SPI (Spinoni et al., 2019) and mean rainfall. (Aguilar et al., 2009; Hua et al., 2016; Dai and Zhao, 2017)	<b>Low confidence: Inconsistent</b> signal in observations vs models for 1951-2010 trends (Knutson and Zeng, 2018); no signal in single-model based study (Otto et al., 2013)	<b>Low confidence; Mixed signal.</b> Drying tendency (increasing CDD) in CORDEX AFR simulations compared to 1971-2000 (Mba et al., 2018); but tendency towards less drying (CDD decrease) in CMIP6 GCMs (Chapter 11 Supplementary Material (11.SM)), consistent with increase in precipitation at higher warming levels (Cook et al., 2020). Inconsistent signals in SPI in CMIP5 GCMs (Xu et al., 2019a)	<b>Low confidence; Mixed signal. Robust drying tendency</b> (increasing CDD) in CORDEX AFR simulations compared to 1971-2000 (Mba et al., 2018); but inconsistent signal in CMIP6 GCMs (with tendency towards CDD decrease (Chapter 11 Supplementary Material (11.SM))); consistent with projected increase in mean precipitation (Cook et al., 2020)); inconsistent signals in CDD in CMIP5 GCMs (Sonkoué et al., 2019). Decrease frequency of SPI-based droughts in CMIP5 (Xu et al., 2019a).	<b>Low confidence: Mixed signal,</b> depending on multi-model experiment and considered index (Fotso-Nguemo et al., 2018; Dosio et al., 2019; Sonkoué et al., 2019; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Increase in mean precipitation in CMIP6 GCMs (Cook et al., 2020). Increase in CDD (increase in meteorological drought) in CORDEX AFR simulations (Dosio et al., 2019; Fotso-Nguemo et al., 2019) but inconsistent CDD signals in CMIP6 (with tendency towards CDD decrease; Chapter 11 Supplementary Material (11.SM)) and CMIP5 GCMs (Sonkoué et al., 2019). Increase in SPI (less drying) in CMIP5 GCMs (Spinoni et al., 2020).
	AGR ECOL	<b>Medium confidence Decrease</b> in water-balance availability or SPEI, but some regional variability and index dependency of trends (Greve et al., 2014;	<b>Low confidence: Limited evidence</b> due to lack of studies	<b>Low confidence: Inconsistent signals.</b> Slight tendency towards soil moisture wetting in CMIP5 (Xu et al., 2019a) and CMIP6 (Chapter 11 Supplementary Material	<b>Low confidence: Inconsistent signals.</b> Inconsistent trends in duration vs frequency of soil moisture-based drought events in CMIP5 (Xu et al., 2019a); slight mean soil moisture wetting in CMIP6	<b>Low confidence. Inconsistent signals.</b> Tendency towards wetting in CMIP6 soil moisture (Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM));

		Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)		(11.SM)); and slight increase (less drying in SPEI-PM (Naumann et al., 2018)	(Chapter 11 Supplementary Material (11.SM)); slight wetting of SPEI-PM based events (Naumann et al., 2018).	inconsistent signals in SPEI-PM (Vicente-Serrano et al., 2020a)
	HYDR	<b>Low confidence: Limited evidence.</b> Decrease in streamflow from 1950-2012 in southern part of domain (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence and inconsistent trends</b> in mean runoff in two studies (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Inconsistent</b> projections and/or non-robust changes (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
South Eastern Africa (SEAF)	MET	<b>Low confidence: Inconsistent trends</b> in SPI (Spinoni et al., 2019) but occurrence of strong drought events in recent years (Funk et al., 2015a; Nicholson, 2017)	<b>Low confidence: Limited evidence</b> on attribution of long-term trends.  Robust evidence that recent drought events are not attributable to anthropogenic climate change (Uhe et al., 2017; Funk et al., 2018b)	<b>Low confidence: Inconsistent changes</b> (Osima et al., 2018; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and lack of signal (Nangombe et al., 2018)  Xu et al. (2019): Inconsistent or weak trends in SPI  Osima et al. (2018): Cordex AFR data, CTL 1971-2000, RCP8.5, consistent increase of CDD over southern part  Chapter 11 Supplementary Material (11.SM): Inconsistent changes in CDD	<b>Low confidence: Inconsistent changes</b> (Osima et al., 2018; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and lack of signal (Nangombe et al., 2018)  Xu et al. (2019): Inconsistent or weak trends in SPI  Osima et al. (2018): Cordex AFR data, CTL 1971-2000, RCP8.5, Robust increase of CDD over southern part.  Chapter 11 Supplementary Material (11.SM): inconsistent changes in CDD	<b>Low confidence: Inconsistent trends</b> between studies and subregions (Sillmann et al., 2013b; Dosio et al., 2019; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).  Inconsistent or no changes in SPI (Vicente-Serrano et al., 2020a)  Sillmann et al. (2013), (2081-2100)/1981-2000, rcp8.5, CMIP3-CMIP5: Decrease of CDD  Dosio et al. (2019), (2070-2099/1981-2010), rcp 8.5, 23 RCM: Decrease in CDD  Inconsistent trends in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM))
	AGR ECOL	<b>Low confidence: Inconsistent trends</b> (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b> due to lack of studies	<b>Low confidence: Inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence; Inconsistent trends</b> (Xu et al., 2019a; Cook et al., 2020) (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends</b> (Cook et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Inconsistent trends</b> (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence; Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence;</b> inconsistent trends in runoff in two studies (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Inconsistent trends.</b> Increase in runoff in a study based on CMIP6 (Cook et al., 2020) but inconsistent or non-robust trends in studies based on ISIMIP and CMIP5 ensembles (Giuntoli et al., 2015; Touma et al., 2015)



Western Southern Africa (WSAF)	MET	<p><b>Low confidence: Inconsistent trends</b> (Spinoni et al., 2019; Dunn et al., 2020)</p> <p>Dunn et al. (2020): Conflicting trends in CDD depending on time frame</p>	<p><b>Low confidence: Limited evidence and inconsistent observed trends.</b></p> <p>But recent meteorological drought attributable to anthropogenic climate change (Bellprat et al., 2015)</p> <p>Recent meteorological drought (2015/2016 drought in southern Africa) attributable to anthropogenic climate change (Otto et al., 2018b; Funk et al., 2018a; Yuan et al., 2018; Pascale et al., 2020)</p>	<p><b>Medium confidence: Increase.</b> Increases in dryness (CDD) (Maúre et al., 2018)(Chapter 11 Supplementary Material (11.SM)) both compared to pre-industrial climate and recent past. Increase in CDD for changes of +0.5°C in global warming based on CMIP5 for overall SREX/AR5 South Africa region (Wartenburger et al., 2017), but only weak shift in mean precipitation in large-ensemble single-model experiment for +0.5°C of global warming (Nangombe et al., 2018). Slight but weaker increase in SPI compared to CDD (Abiodun et al., 2019; Xu et al., 2019a; Naik and Abiodun, 2020)</p> <p>Maúre et al. (2018): 25 Cordex AFR run, CTL 1971-2000, RCP8.5, -Increase of CDD</p> <p>NB: Weaker signals in SPI (Xu et al., 2019a)</p> <p>Cordex AFR data, CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890) (Abiodun et al., 2019; Naik and Abiodun, 2020)</p> <p>Non-significant increase in SPI-based drought frequency and intensity</p>	<p><b>High confidence: Increases</b> in dryness (CDD, DF, NDD) (Maúre et al., 2018; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM)); slight but weaker increase in SPI (Abiodun et al., 2019; Naik and Abiodun, 2019; Xu et al., 2019)</p> <p>Maúre et al. (2018): 25 Cordex AFR run ,CTL 1971-2000, RCP8.5, -Increase of CDD</p> <p>Coppola et al. (2021b), (2041-2060)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6 Increase in DF (drought frequency) and NDD (number of dry days )</p> <p>NB: Weaker signals in SPI (Xu et al., 2019a)</p> <p>Cordex AFR data, CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890) ) (Abiodun et al., 2019; Naik and Abiodun, 2020): Non-significant increase in SPI-based drought frequency and intensity.</p>	<p><b>Likely: Increase</b> (CDD amd SPI) (Sillmann et al., 2013b; Giorgi et al., 2014; Touma et al., 2015; Pinto et al., 2016; Abiodun et al., 2019; Dosio et al., 2019; Naik and Abiodun, 2020; Spinoni et al., 2020; Coppola et al., 2021b)</p> <p>Using Cordex , CTL :1981-2010,RCP 8.5 2071-2100 (Spinoni et al., 2020)</p> <p>Robust increase of drought frequency and severity (SPI-12 )</p> <p>Based on Giorgi et al., 2014, 5GCM/IRCM, CTL: 1976-2005, rcp 8.5, 2071-2100: Increase of CDD</p> <p>Sillmann et al. (2013), (2081-2100)/1981-2000, rcp8.5, CMIP3-CMIP5 Increase of CDD</p> <p>Coppola et al. (2021b), (2080-2099)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6</p> <p>Increase in DF (drought frequency) and NDD (number of dry days )</p> <p>Dosio et al. (2019) (2070-2099/1981-2010), rcp 8.5, 23 RCM: Increase in CDD</p> <p>Pinto et al. (2016): (2069-2098/1976-2005), rcp 8.5,4 GCM/2RCM: Increase in CDD.</p>
	AGR ECOL	<p><b>Medium confidence: Drought increase</b> based on water-balance estimates and SPEI (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020)</p>	<p><b>Low confidence: Limited evidence:</b></p> <p>Given small number of studies based on soil moisture (Yuan et al., 2018a) and atmospheric drought indices (Nangombe et al., 2020)</p>	<p><b>Medium confidence; Drought increase.</b> Decrease in SM both compared to recent past (Xu et al., 2019) and pre-industrial (Chapter 11 Supplementary Material (11.SM)) baselines; but conflicting changes of drought magnitude based on SPEI-PM compared to 0.6°C baseline (Naumann et al., 2018)</p>	<p><b>High confidence: Drought increase.</b> Decrease in SM (Xu et al., 2019) (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020); but conflicting changes of drought magnitude based on SPEI-PM (Naumann et al., 2018)</p>	<p><b>Likely: Drought increase.</b> Decrease in SM (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020) and SPEI-PM (Vicente-Serrano et al., 2020a)</p>

	HYDR	<b>Low confidence: Limited evidence.</b> Decrease in runoff in larger AR5 “Southern Africa” region, but weaker signal depending on time frame (Gudmundsson et al., 2019, 2021); non significant drying tendency (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal ( Touma et al., 2015)	<b>Medium confidence; Increased drying</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020a)	<b>Medium confidence: Increased drying</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Eastern Southern Africa (ESAF)	MET	<b>Medium confidence: Dominant increase in meteorological drought</b> in SPI and CDD (Spinoni et al., 2019; Dunn et al., 2020)	<b>Low confidence: Limited evidence</b> on attribution of long-term trends.  <i>Medium confidence</i> that human-influence has contributed to stronger recent meteorological drought.(Bellprat et al., 2015; Funk et al., 2018a; Yuan et al., 2018a)	<b>Medium confidence: Increases</b> in meteorological drought based on CDD (Maúre et al., 2018)(Chapter 11 Supplementary Material (11.SM)) both compared to pre-industrial climate and recent past. Non-significant increase in SPI-based drought (Abiodun et al., 2017); lack of signal in SPI compared to recent past (1970-2000) (Xu et al., 2019a). Increase in CDD for changes of +0.5°C in global warming based on CMIP5 for overall SREX/AR5 South Africa region (Wartenburger et al., 2017), but only weak shift in mean precipitation in large-ensemble single-model experiment for +0.5°C of global warming (Nangombe et al., 2018).  Maúre et al. (2018): 25 Cordex AFR run ,CTL 1971-2000, RCP8.5, -Increase of CDD  Cordex AFR data,CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890) (Abiodun et al., 2019) SPI non-significant drought frequency & intensity increase	<b>High confidence: Increase</b> in meteorological drought based on (CDD,DF,NDD) (Maúre et al., 2018; Coppola et al., 2021b)(Chapter 11 Supplementary Material (11.SM)) and SPI (Abiodun et al., 2019; Xu et al., 2019a) both compared to recent past and pre-industrial period.  Maúre et al. (2018): 25 Cordex AFR run ,CTL 1971-2000, RCP8.5: Increase of CDD  (Coppola et al., 2021b), (2041-2060)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6: Increase in DF (drought frequency) and NDD (number of dry days )  Abiodun et al. (2019): Cordex AFR data,CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890): increase in SPI-based meteorological drought frequency and intensity.  Xu et al. (2019): Drying in SPI at 2°C compared to 1970-2000 conditions.	<b>Likely: Increase</b> in meteorological drought (CDD amd SPI) (Sillmann et al., 2013b; Giorgi et al., 2014; Touma et al., 2015; Pinto et al., 2016; Dosio et al., 2019; Spinoni et al., 2020; Coppola et al., 2021b)(Chapter 11 Supplementary Material (11.SM)) Using Cordex, CTL :1981-2010,RCP 8.5, 2071-2100 (Spinoni et al., 2020) Robust increase of drought frequency and severity (SPI-12,SPEI-12)  Based on Giorgi et al. (2014), 5GCM/1RCM, CTL: 1976-2005, rcp 8.5, 2071-2100: Increase of CDD  Sillmann et al. (2013), (2081-2100)/1981-2000, rcp8.5, CMIP3-CMIP5 Increase of CDD  (Coppola et al., 2021b), (2080-2099)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6 Increase in DF (drought frequency) and NDD (number of dry days )  Dosio et al. (2019), (2070-2099/1981-2010), rcp 8.5, 23 RCM Increase in CDD  Pinto et al. (2016): (2069-2098/1976-2005), rcp 8.5,4 GCM/2RCM: Increase in CDD

	AGR ECOL	<b>Medium confidence Increase</b> , based on water-balances estimates, PDSI and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b> (Yuan et al., 2018a)	<b>Medium confidence: Increase</b> in drought. Decrease in SM both compared to recent past (Xu et al., 2019) and pre-industrial (Chapter 11 Supplementary Material (11.SM)) baselines; but inconsistent changes of drought magnitude based on SPEI-PM compared to +0.6°C baseline (Naumann et al., 2018)	<b>Medium confidence: Increase</b> in drought; decrease in SM (Xu et al., 2019a; Cook et al., 2020) (Chapter 11 Supplementary Material (11.SM)); but inconsistent changes of in drought magnitude based on SPEI-PM (Naumann et al., 2018)	<b>High confidence: Increase</b> in drought; decrease in SM (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020) and SPEI-PM (Vicente-Serrano et al., 2020a)
	HYDR	<b>Low confidence: Limited evidence.</b> Decrease in runoff in larger AR5 “Southern Africa” region, but weaker signal depending on time frame (Gudmundsson et al., 2019, 2021); non significant drying tendency (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Medium confidence; Increased drying</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020a).	<b>Medium confidence: Increased drying</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Mada-gascar (MDG)	MET	<b>Low confidence: Inconsistent trends</b> (Vincent et al., 2011; Spinoni et al., 2019)	<b>Low confidence: Limited evidence</b>	<b>Medium confidence: Increase</b> in meteorological drought based on SPI compared to recent past (Abiodun et al., 2019; Xu et al., 2019a) and CDD compared to pre-industrial baseline (Chapter 11 Supplementary Material (11.SM)).  Abiodun et al. (2019): Cordex AFR data, CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890) SPI (drought frequency & intensity increase)	<b>High confidence: Increase</b> in meteorological drought based on several metrics, including SPI (Abiodun et al., 2019; Xu et al., 2019a), CDD (Chapter 11 Supplementary Material (11.SM)), and DF (drought frequency) and NDD (number of dry days) (Coppola et al., 2021b)  (Coppola et al., 2021b), (2041-2060)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6 Increase in DF (drought frequency) and NDD (number of dry days )  Abiodun et al. (2019): Cordex AFR data, CTL 1971-2000, RCP8.5, pre-industrial reference period (1861-1890): Increase in SPI-based drought frequency and intensity.	<b>Likely: Increase</b> in meteorological drought based on CDD and SPI (Sillmann et al., 2013b; Giorgi et al., 2014; Touma et al., 2015; Pinto et al., 2016; Dosio et al., 2019; Spinoni et al., 2020; Coppola et al., 2021b)  Sillmann et al. (2013), (2081-2100)/1981-2000, rcp8.5, CMIP3-CMIP5 Increase of CDD  Spinoni et al. (2020): Using CORDEX, CTL:1981-2010, RCP 8.5, 2071-2100 Robust increase of drought frequency and severity (SPI-12)  (Coppola et al., 2021b), (2080-2099)/1995-2014, rcp 8.5, CMIP5-CORDEX-CMIP6 Increase in DF (drought frequency) and NDD (number of dry days )  Dosio et al. (2019), (2070-2099/1981-2010), rcp 8.5, 23 RCM: Increase in CDD

	AGR ECOL	<b>Low confidence: Inconsistent trends</b> based on water-balance estimates, PDSI and SPEI (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent</b> or weak trends (Xu et al., 2019) (Chapter 11 Supplementary Material (11.SM)) (Naumann et al., 2018)	<b>Medium confidence: Increase</b> in drought. Decrease in SM (Chapter 11 Supplementary Material (11.SM)); (Cook et al., 2020) and in SPEI-PM (Naumann et al., 2018)	<b>High confidence: Increase</b> in drought. Robust decrease in SM (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020) and SPEI-PM (Vicente-Serrano et al., 2020a)
	HYDR	<b>Low confidence: Limited evidence.</b> Inconsistent trends in one study (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Inconsistent</b> trends. Inconsistent trends (Cook et al., 2020) or weak drying (Touma et al., 2015; Zhai et al., 2020b)	<b>Medium confidence: Increase</b> in drought based on two studies based on CMIP5 (Giuntoli et al., 2015; Touma et al., 2015), but some inconsistent trends in CMIP6 mean runoff trends (Cook et al., 2020)

1  
2  
3  
4  
5  
6  
7  
8

[END TABLE 11.6 HERE]

[START TABLE 11.7 HERE]

**Table 11.7:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in Asia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Asia	Most subregions show a <i>very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Hu et al., 2020; Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>very likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes:	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and

			<i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)	frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Russian Arctic (RAR)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Donat et al., 2016a; Sui et al., 2017; Dunn et al., 2020)	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wang et al., 2017c)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Arabian Peninsula (ARP)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and	Strong evidence of changes from observations that are in the direction of model projected changes for the	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity

	<p>frequency of cold extremes (Dunn et al., 2020; Almazroui et al., 2014; Barlow et al., 2016; Donat et al., 2014; Nazrul Islam et al., 2015; Rahimi and Hejabi, 2018; Donat et al., 2014; Rahimi et al., 2018)</p>	<p>future. The magnitude of projected changes increases with global warming.</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Almazroui, 2019b)</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Almazroui, 2019b)</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Almazroui, 2019b)</p>
	<p><i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes</p>	<p><i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
West Central Asia (WCA)	<p>Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Hu et al., 2016; Jiang et al., 2013; Dunn et al., 2020)</p>	<p>Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Dong et al., 2018; Kim et al., 2019)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 6°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from</p>

			CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)	CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)	CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
West Siberia (WSB)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Degeffie et al., 2014; Salnikov et al., 2015; Donat et al., 2016a; Zhang et al., 2019c, 2019b; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wang et al., 2017; Seong et al., 2020; Dong et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the	<i>High confidence</i> in a human contribution to the observed increase in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared

	intensity and frequency of cold extremes	frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Siberia (ESB)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dashkhuu et al., 2015; Donat et al., 2016a; Zhang et al., 2019c; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wang et al., 2017; Seong et al., 2020; Dong et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014))



					<i>Virtually certain</i> (compared with pre-industrial)
Russian Far East (RFE)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Donat et al., 2016; Dunn et al., 2020; Zhang et al., 2019b)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Dong et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Xu et al. 2017; Han et al. 2018; Khlebnikova et al. 2019).
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Asia (EAS)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Lin et al., 2017; Lu et al., 2016, 2018; Wang et al.,	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020;	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than

	2013a; Yin et al., 2017; Zhou et al., 2016; Dunn et al., 2020)	Wang et al., 2017; Imada et al., 2014, 2019; Kim et al., 2018; Lu et al., 2016, 2018; Takahashi et al., 2016; Ye and Li, 2017; Zhou et al., 2016)	the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Guo et al., 2018; Imada et al., 2019; Li et al., 2018c; Seo et al., 2014; Sui et al., 2018; Wang et al., 2017a, 2017c; Xu et al., 2016a; Zhou et al., 2014; Shi et al., 2018; Sun et al., 2019a)	1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Guo et al., 2018; Imada et al., 2019; Li et al., 2018c; Seo et al., 2014; Sui et al., 2018; Wang et al., 2017a, 2017c; Xu et al., 2016a; Zhou et al., 2014; Shi et al., 2018; Sun et al., 2019a)	4°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Guo et al., 2018; Imada et al., 2019; Li et al., 2018c; Seo et al., 2014; Sui et al., 2018; Wang et al., 2017a, 2017c; Xu et al., 2016a; Zhou et al., 2014; Shi et al., 2018; Sun et al., 2019a)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Central Asia (ECA)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Dong et al., 2018; Kim et al., 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in

			<p>compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)</p>	<p>annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)</p>	<p>annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018)</p>
	<p><i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes</p>	<p><i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
Tibetan Plateau (TIB)	<p>Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Donat et al., 2016a; Hu et al., 2016; Sun et al., 2017; Yin et al., 2019; Zhang et al., 2019; Dunn et al., 2020)</p>	<p>Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Yin et al., 2019)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Zhou et al., 2014; Singh and Goyal, 2016; Zhang et al.,</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Zhou et al., 2014; Singh and Goyal, 2016; Zhang et al.,</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Zhou et al., 2014; Singh and Goyal, 2016; Zhang et al.,</p>

			2016a; Xu et al., 2017; Han et al., 2018; Li et al., 2018a)	2016a; Xu et al., 2017; Han et al., 2018; Li et al., 2018a)	2016a; Xu et al., 2017; Han et al., 2018; Li et al., 2018a)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
South Asia (SAS)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Chakraborty et al., 2018; Dimri, 2019; Donat et al., 2016; Dunn et al., 2020; Roy, 2019; Sheikh et al., 2015; Rohini et al., 2016; Zahid and Rasul, 2012)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Wehner et al., 2016; Kumar, 2017; van Oldenborgh et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Ali et al., 2019; Han et al., 2018; Kharin et al., 2018; Sillmann et al., 2013; Xu et al., 2017; Murari et al., 2015; Nasim et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Ali et al., 2019; Han et al., 2018; Kharin et al., 2018; Sillmann et al., 2013; Xu et al., 2017; Murari et al., 2015; Nasim et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Ali et al., 2019; Han et al., 2018; Kharin et al., 2018; Sillmann et al., 2013; Xu et al., 2017; Murari et al., 2015; Nasim et al., 2018)
	<i>High confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014))

	and frequency of cold extremes	and frequency of cold extremes.	industrial) Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	with pre-industrial) Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	<i>Virtually certain</i> (compared with pre-industrial) Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Southeast Asia (SEA)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Donat et al., 2016a; Supari et al., 2017; Cheong et al., 2018; Zhang et al., 2019c; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; King et al., 2016; Min et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Han et al., 2018; Kharin et al., 2018; Xu et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Kharin et al., 2018; Xu et al., 2017).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex)  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Kharin et al., 2018; Xu et al., 2017)..
	<i>High confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

					with pre-industrial)
--	--	--	--	--	----------------------

1  
2  
3  
4  
5  
6  
7

[END TABLE 11.7 HERE]

[START TABLE 11.8 HERE]

**Table 11.8:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in Asia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.3 for details

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Asia	Significant intensification of heavy precipitation (Sun et al., 2020)	Robust evidence of a human contribution to the observed intensification of heavy precipitation	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)
	<i>Likely</i> intensification of heavy precipitation	Human influence <i>likely</i> contributed to the observed intensification of heavy precipitation	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Russian Arctic (RAR)	Insufficient data and a lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 25% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 25% in annual Rx1day and Rx5day and 20% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et

			al., 2013b; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	al., 2013b; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	al., 2013b; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Arabian Peninsula (ARP)	Insufficient data and a lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020; Atif et al., 2020; Donat et al., 2014; Rahimi and Fatemi, 2019)	Limited evidence		CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 40% in annual Rx1day and Rx5day and 45% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
West Central Asia (WCA)	Intensification of heavy precipitation (Sun et al., 2020; Hu et al., 2016; Zhang et al., 2017).	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of

			heavy precipitation (Han et al., 2018)	heavy precipitation (Han et al., 2018)	heavy precipitation (Han et al., 2018)
	<i>Medium confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
West Siberia (WSB)	Significant intensification of heavy precipitation (Sun et al., 2020; Zhang et al., 2017)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)
	<i>High confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Siberia (ESB)	Intensification of heavy precipitation (Knutson and Zeng, 2018; Sun et al., 2020; Dunn et al., 2020)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and



			Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)	Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018; Khlebnikova et al., 2019b)
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Russian Far East (RFE)	Intensification of heavy precipitation (Sun et al., 2020)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 25% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 25% in annual Rx1day and Rx5day and 20% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Kharin et al., 2018)
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Asia (EAS)	Intensification of heavy precipitation (Sun et al.,	Disagreement among studies (Chen and Sun, 2017; Li et	CMIP6 models project an increase in the intensity and	CMIP6 models project a robust increase in the intensity and	CMIP6 models project a robust increase in the intensity and

	2020; Dunn et al., 2020; Baek et al., 2017; Nayak et al., 2017; Ye and Li, 2017; Zhou et al., 2016)	al., 2017; Burke et al., 2016; Zhou et al., 2013; Ma et al., 2017)	frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Ahn et al., 2016; Guo et al., 2018; Hatsuzuka et al., 2020; Kawase et al., 2019; Kim et al., 2018; Kusunoki, 2018; Kusunoki and Mizuta, 2013; Li et al., 2018a; Nayak and Dairaku, 2016; Ohba and Sugimoto, 2020, 2019; Seo et al., 2014; Wang et al., 2017a, 2017b; Zhou et al., 2014; Li et al., 2018b)	frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Ahn et al., 2016; Guo et al., 2018; Hatsuzuka et al., 2020; Kawase et al., 2019; Kim et al., 2018; Kusunoki, 2018; Kusunoki and Mizuta, 2013; Li et al., 2018a; Nayak and Dairaku, 2016; Ohba and Sugimoto, 2020, 2019; Seo et al., 2014; Wang et al., 2017a, 2017b; Zhou et al., 2014; Li et al., 2018b)	frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Ahn et al., 2016; Guo et al., 2018; Hatsuzuka et al., 2020; Kawase et al., 2019; Kim et al., 2018; Kusunoki, 2018; Kusunoki and Mizuta, 2013; Li et al., 2018a; Nayak and Dairaku, 2016; Ohba and Sugimoto, 2020, 2019; Seo et al., 2014; Wang et al., 2017a, 2017b; Zhou et al., 2014; Li et al., 2018b)
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
East Central Asia (ECA)	Intensification of heavy precipitation (Sun et al., 2020)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 25% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).

	<i>Medium confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Tibetan Plateau (TIB)	Intensification of heavy precipitation (Sun et al., 2020; Jiang et al., 2013; Hu et al., 2016; Ge et al., 2017; Zhan et al., 2017; Liu et al., 2019)	Limited evidence	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Zhou et al., 2014; Zhang et al., 2015c; Gao et al., 2018; Han et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Zhou et al., 2014; Zhang et al., 2015c; Gao et al., 2018; Han et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 25% in annual Rx1day and Rx5day and 20% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Zhou et al., 2014; Zhang et al., 2015c; Gao et al., 2018; Han et al., 2018)
	<i>Medium confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
South Asia (SAS)	Significant intensification of heavy precipitation (Kim et al., 2019; Malik et al., 2016; Pai et al., 2015; Rohini et al., 2016; Roxy et al., 2017; Sheikh et al., 2015; Singh et al., 2014; Dunn et al., 2020; Hussain and Lee, 2013; Kim et al., 2019; Malik et al., 2016)	Disagreement among studies (Mukherjee et al., 2018a) (Singh et al., 2014a; van Oldenborgh et al., 2016)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 25% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 30% in annual Rx1day and Rx5day and 25% in annual Rx30day compared to pre-industrial (Annex).

			Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Mukherjee et al., 2018a; Ali et al., 2019b; Rai et al., 2019)	Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Mukherjee et al., 2018a; Ali et al., 2019b; Rai et al., 2019)	Additional evidence from CMIP5 simulations for an increase in the intensity of heavy precipitation (Sillmann et al., 2013b; Xu et al., 2017; Han et al., 2018; Mukherjee et al., 2018a; Ali et al., 2019b; Rai et al., 2019)
	<i>High confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
Southeast Asia (SEA)	Intensification of heavy precipitation (Sun et al., 2020; Cheong et al., 2018; Li et al., 2018c; Siswanto et al., 2015; Supari et al., 2017; Villafuerte and Matsumoto, 2015)	Evidence of a human contribution for some events (Otto et al., 2018a), but cannot be generalized	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 0% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Xu et al., 2017; Han et al., 2018; Tangang et al., 2018; Trinh-Tuan et al., 2019; Basconcillo et al., 2016; Ge et al., 2017; Han et al., 2018; Marzin et al., 2015; Tangang et al., 2018; Trinh-Tuan et al., 2019; Xu et al., 2017)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Xu et al., 2017; Han et al., 2018; Tangang et al., 2018; Trinh-Tuan et al., 2019; Basconcillo et al., 2016; Ge et al., 2017; Han et al., 2018; Marzin et al., 2015; Tangang et al., 2018; Trinh-Tuan et al., 2019; Xu et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Xu et al., 2017; Han et al., 2018; Tangang et al., 2018; Trinh-Tuan et al., 2019; Basconcillo et al., 2016; Ge et al., 2017; Han et al., 2018; Marzin et al., 2015; Tangang et al., 2018; Trinh-Tuan et al., 2019; Xu et al., 2017)
	<i>Medium confidence in the intensification of heavy precipitation</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014))

			past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	2014)) <i>Likely</i> (compared with pre-industrial)	<i>Extremely likely</i> (compared with pre-industrial)
--	--	--	--	--	--

1  
2  
3  
4  
5  
6  
7  
8

[END TABLE 11.8 HERE]

[START TABLE 11.9 HERE]

**Table 11.9:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET), agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in Asia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.4 for details

Region/ Drought type		Observed trends	Human contribution	Projections		
				+1.5 °C	+2 °C	+4 °C
Russian Arctic (RAR)	MET	<b>Low confidence: Limited evidence.</b> Tendency towards decrease in CDD (Dunn et al., 2020). Lack of data in (Spinoni et al., 2019).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> Slight decrease in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Limited evidence,</b> but some evidence of decrease in dry spell duration (Khlebnikova et al., 2019b)(Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Decrease in drought severity</b> based on SPI (Touma et al., 2015; Spinoni et al., 2020) and CDD (Chapter 11 Supplementary Material (11.SM)).
	AGR, ECOL	<b>Low confidence: Inconsistent trends</b> (Greve et al., 2014; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent changes</b> in soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent changes</b> in soil moisture, variations across subregions (Xu et al., 2019a) (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends.</b> Inconsistent trends across models and subregions for surface and total soil moisture (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)); Slight drying in PDSI (Dai et al., 2018); inconsistent trends or wetting in SPEI-PM in CMIP5 (Cook et al., 2014b; Vicente-Serrano et al., 2020a).

	HYDR	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Inconsistent changes.</b> Increasing runoff in CMIP6 (Cook et al., 2020) , inconsistent signal in SRI depending on subregion in CMIP5(Touma et al., 2015), or lack of signal (Zhai et al., 2020b) in available studies.  (Cook et al., 2020): Increasing runoff in one study based on CMIP6 GCMs  (Zhai et al., 2020b): Lack of signal in one study based on single hydrological model driven by HAPPI-MIP GCM simulations  Touma et al. (2015): Inconsistent signal in SRI depending on subregion (CMIP5 GCMs)	<b>Low confidence: Mixed signals</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Arabian Peninsula (ARP)	MET	<b>Low confidence: Inconsistent or no signal</b> (Almazroui, 2019a; Almazroui and Islam, 2019).  (Dunn et al., 2020): Wetting based on CDD in part of domain, but missing data in large fraction of region.  (Spinoni et al., 2019): Missing data in this region.	<b>Low confidence: Limited evidence</b> (Barlow and Hoell, 2015; Barlow et al., 2016)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Limited evidence and inconsistent trends</b> (Touma et al., 2015; Tabari and Willems, 2018)(Chapter 11 Supplementary Material (11.SM)).  (Touma et al., 2015): Inconsistent projections in CMIP5 (Tabari and Willems, 2018): Dominant lack of signal  Chapter 11 Supplementary Material (11.SM)): decreasing dryness based on CDD
	AGR, ECOL	<b>Low confidence: Limited evidence.</b> Drying in fraction of region in one study, but missing data in rest of region (Greve et al., 2014).  (Greve et al., 2014) : Drying in part of region, but missing data in large fraction of region.  (Padrón et al., 2020) : Missing data.  (Spinoni et al., 2019) : Missing data.	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))  (Naumann et al., 2018): Missing data	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Mixed signal</b> between different metrics, including total and surface soil moisture (Chapter 11 Supplementary Material (11.SM))(Rajsekhar and Gorelick, 2017; Dai et al., 2018; Lu et al., 2019; Cook et al., 2020), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).

	HYDR	<b>Low confidence: Limited evidence</b> Drying in one study in northern part of region but missing data in rest of region (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal ( Touma et al., 2015).	<b>Low confidence: Limited evidence and inconsistent trends</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Inconsistent trends</b> between models and studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
West Central Asia (WCA)	MET	<b>Low confidence: Inconsistent trends</b> between subregions, based both on CDD and SPI (Spinoni et al., 2019; Dunn et al., 2020; Sharafati et al., 2020; Yao et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> Inconsistent or weak trends in available analyses (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent, weak and/or non-significant trends</b> in SPI and CDD (Xu et al., 2019a; Spinoni et al., 2020; Yao et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signals</b> between models and between regions (Touma et al., 2015; Han et al., 2018; Tabari and Willems, 2018; Spinoni et al., 2020; Yao et al., 2020)(Chapter 11 Supplementary Material (11.SM))
	AGR, ECOL	<b>Medium confidence: Increase in drought severity.</b> Dominant signal shows drying for soil moisture, water-balance (precipitation-evapotranspiration), PDSI-PM and SPEI-PM, but with some differences between subregions and studies (Greve et al., 2014; Dai and Zhao, 2017; Li et al., 2017c; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence.</b> One study by Li et al. (2017) concluded that anthropogenic forcing has increased AED and contributed to drought severity over the last decades.	<b>Low confidence: Mixed signals</b> in changes in drought severity, depending on model and index (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Weak signals and inconsistent trends between models for total and surface soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)), but increased drying based on SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signals</b> in changes in drought severity, depending on model and index (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020) (Chapter 11 Supplementary Material (11.SM)).  Weak signals and inconsistent trends between models and subregions for total and surface soil moisture (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), but increased drying based on SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increased drying</b> in several metrics, but substantial intermodel spread and lack of signal for total soil moisture (Dai et al., 2018; Cook et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).  Increase in drought severity based on surface soil moisture (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020). (Chapter 11 Supplementary Material (11.SM)): only median, not 83.5%ile), PDSI (Dai et al., 2018), and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a); but increase in median response and substantial intermodel spread for total soil moisture (Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Medium confidence: Increase of hydrological drought severity</b> (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020); but large intermodel spread (only 2/3 of models showing signal) (Touma et al., 2015) and weak signal-to-noise ratio in eastern half of domain (Giuntoli et al., 2015).

Western Siberia (WSB)	MET	<p><b>Medium confidence: Decrease in dryness</b> based on SPI and CDD, but some inconsistent trends in part of domain (Zhang et al., 2017a, 2019b; Khlebnikova et al., 2019b; Spinoni et al., 2019; Dunn et al., 2020).</p> <p>Khlebnikova et al. (2019): In part mixed signals within domain (Dunn et al., 2020): Mostly decreasing trend, including significant changes. (Spinoni et al., 2019): Mostly decreasing trends</p>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent evidence</b> in CMIP5 (Xu et al., 2019a) and CMIP6 projections (Chapter 11 Supplementary Material (11.SM)).	<p><b>Low confidence: Inconsistent trends</b> (Chapter 11 Supplementary Material (11.SM)) or slight decrease in drought (Khlebnikova et al., 2019b; Xu et al., 2019a; Spinoni et al., 2020).</p> <p>(Khlebnikova et al., 2019b): Mostly decrease in CDD in a regional climate model driven by several CMIP5 models (RCP8.5, 2050-2059 relative to 1990-1999)</p> <p>Chapter 11 Supplementary Material (11.SM): Tendency towards decrease but lack of model agreement.</p>	<p><b>Low confidence: Inconsistent trends</b> , but slight decrease in some studies (Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).</p> <p>Spinoni et al. (2020): Slight decrease Touma et al. (2015): Tendency towards decrease but partly lack of model agreement.</p> <p>Chapter 11 Supplementary Material (11.SM): Lack of model agreement</p>
	AGR, ECOL	<p><b>Low confidence: Inconsistent trends</b> according to subregions or indices based on soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Li et al., 2017c; Spinoni et al., 2019; Padrón et al., 2020).</p>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> among different metrics and models. Inconsistent soil moisture projections in CMIP5 (Xu et al., 2019a) and CMIP6 (Chapter 11 Supplementary Material (11.SM)), and decrease in drought severity based on SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<p><b>Low confidence: Inconsistent trends</b> among different metrics. No signal with total soil moisture (Chapter 11 Supplementary Material (11.SM)) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020), and wetting trend with surface soil moisture (Xu et al., 2019a).</p>	<p><b>Low confidence: Mixed signals</b> between different models and metrics, including total and surface soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), surface soil moisture in CMIP5 (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).</p> <p>Difference in signal in CMIP6 vs CMIP5: CMIP6 models show drying in soil moisture, while CMIP5 models show wetting (Cook et al., 2020)</p>



	HYDR	<b>Low confidence: Limited evidence.</b> One study suggests increasing weak (wetting) trend in runoff (Dai and Zhao, 2017). Some increase in runoff at stations from 1951-1990 and 1961-2000 (Gudmundsson et al., 2019)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows drying (Touma et al., 2015).	<b>Low confidence: Inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)  (Cook et al., 2020): Inconsistent trends including large seasonal variations  (Zhai et al., 2020b): Inconsistent trends in study with single hydrological model driven with HAPPI-MIP GCM simulations  (Touma et al., 2015): Increase in the frequency of hydrological droughts based on SRI in CMIP5	<b>Low confidence: Inconsistent trends.</b> Mixed signal among studies and low signal to noise ratio (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Eastern Siberia (ESB)	MET	<b>Medium confidence: Decrease</b> in the duration and frequency of meteorological droughts (Khlebnikova et al., 2019b; Spinoni et al., 2019; Dunn et al., 2020).  (Khlebnikova et al., 2019b): Decrease in fraction of dry days and decrease in mean CDD, but inconsistent trends for maximum CDD, for 1991-2015 compared to 1966-1990  (Dunn et al., 2020): Significant CDD decrease  (Spinoni et al., 2019): Mostly decrease in SPI, but partly mixed signals and inconsistent trends	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> Tendency towards decrease in SPI in CMIP5 (Xu et al., 2019a) and CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Decrease</b> in frequency and severity of meteorological droughts (Khlebnikova et al., 2019b; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  (Khlebnikova et al., 2019b): Projections with a regional climate model driven with several CMIP5 GCMs (RCP8.5, 2050-2059 compared with 1990-1999): Mostly decrease in CDD but increase in part of domain, in particular in the south	<b>Medium confidence: Decrease</b> in meteorological drought severity (Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).

	AGR, ECOL	<b>Low confidence: Inconsistent trends</b> depending on subregion and index based on soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed signal</b> in changes in drought severity depending on models and metrics.  Inconsistent trends in soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)), , but wetting tendency for SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> in changes in drought severity depending on models and metrics.  Inconsistent trends in soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) , but wetting tendency for SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> in drought changes depending on models and metrics, including total and surface soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), surface soil moisture in CMIP5 (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).  Difference in signal in CMIP6 vs CMIP5: CMIP6 models show drying in soil moisture, while CMIP5 models show wetting (Cook et al., 2020)
	HYDR	<b>Low confidence: Limited evidence.</b> One study suggests increasing (wetting) trend in runoff (Dai and Zhao, 2017). Some increase in runoff at stations from 1951-1990 and 1961-2000 (Gudmundsson et al., 2019)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)  (Cook et al., 2020): Inconsistent trends including large seasonal variations  (Zhai et al., 2020b): Inconsistent trends in one study based on single hydrological model driven by HAPPI-MIP GCM simulations  (Touma et al., 2015): Mixed signal.	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Russian Far East (RFE)	MET	<b>Low confidence: Mixed signals</b> between subregions and studies (Knutson and Zeng, 2018; Khlebnikova et al., 2019b; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence.</b> One study, Wilcox et al. in (Herring et al., 2015), but mostly inconclusive.	<b>Low confidence: Limited evidence.</b> Weak decrease in available analyses (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Decrease</b> (Khlebnikova et al., 2019b; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)). (Khlebnikova et al., 2019b): Regional climate model driven by several CMIP5 models (RCP8.5, 2050-2059 relative to 1990-1999): Mostly decrease in CDD but also increase in part of region (Kamchatka Peninsual).	<b>Medium confidence: Decrease</b> in drought severity (Touma et al., 2015; Han et al., 2018; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).

	AGR, ECOL	<b>Low confidence: Inconsistent trends</b> depending on subregion based on soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> depending on model and index.  Inconsistent trends in total and surface soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM)), but wetting trends from CMIP5-based surface soil moisture (Xu et al., 2019a) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).  (Naumann et al., 2018) : EC-Earth driven by SSTs from several CMIP5 models.	<b>Low confidence: Inconsistent trends</b> depending on model and index.  Inconsistent trends in CMIP6 total and surface soil moisture (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), but wetting trends from CMIP5-based surface soil moisture (Xu et al., 2019a) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signals</b> between different models and metrics, including CMIP6 total and surface soil moisture (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), and CMIP5-based surface soil moisture (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).  Difference in signal in CMIP6 vs CMIP5: CMIP6 models show drying in soil moisture, while CMIP5 models show wetting (Cook et al., 2020).
	HYDR	<b>Low confidence: Limited evidence.</b> One study suggests decreasing (drying) trend in runoff (Dai and Zhao, 2017).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Inconsistent trends.</b> Available studies show inconsistent signal with high seasonal variations (Cook et al., 2020) or weak signal (Touma et al., 2015; Zhai et al., 2020b).	<b>Low confidence: Inconsistent signal</b> among studies and metrics, with generally weak drying trend in summer season (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
East Asia (EAS)	MET	<b>Low confidence:</b> Lack of signal and mixed trends between subregions (Spinoni et al., 2019; Zhang et al., 2019a; Dunn et al., 2020; Li et al., 2020b). Drying trends in Southwestern China (Qin et al., 2015a) and Northern China (Qin et al., 2015b), but not for overall China (Li et al., 2020b).	<b>Low confidence: Limited evidence</b> (Qin et al., 2015a; Herring et al., 2019).	<b>Low confidence: Limited evidence.</b> Inconsistent subregional trends (Xu et al., 2019a) or drying tendency (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent trends</b> depending on model, region or index (Guo et al., 2018; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  (Spinoni et al., 2020): Tendency towards decreased in drought severity based on SPI.  (Huang et al., 2018a): Important subregional differences in SPI projections in a single GCM  Chapter 11 Supplementary Material (11.SM): Tendency towards drying based on CDD (increasing CDD), but inconsistent trends depending on model.  (Xu et al., 2019a): Inconsistent subregional trends based on SPI.	<b>Low confidence: Inconsistent trends</b> between different models and important spatial variability (Zhou et al., 2014; Touma et al., 2015; Kusunoki, 2018a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  (Zhou et al., 2014): Tendency towards wetting in the north and drying in the south based on CDD.  (Kusunoki, 2018a): Increasing CDD (drying trend) over Japan based on one GCM.

AGR, ECOL	<p><b>Medium confidence: Increase</b> in drying, especially since ca. 1990; but wetting tendency beforehand and partly inconsistent subregional trends. Large-scale studies based on observed soil moisture, modelled soil moisture or water balance driven by meteorological observations, and SPEI-PM, show drying in northern part of domain (northern China, Russian part of domain, Japan) as well as in Southwest China (east of Tibetan Plateau), but there are some inconsistent trends in part of region or some studies, as well as for different time frames (Greve et al., 2014; Chen and Sun, 2015b; Cheng et al., 2015; Qiu et al., 2016; Dai and Zhao, 2017; Jia et al., 2018; Spinoni et al., 2019; Li et al., 2020b; Padrón et al., 2020). Identified trends are also confirmed by regional studies (Liu et al., 2015; Qin et al., 2015b; Liang et al., 2020; Wang et al., 2020). Most of the drying trend took place since 1990, with wetting trend beforehand (Chen and Sun, 2015b; Wu et al., 2020b).</p>	<p><b>Low confidence: Limited evidence.</b></p> <p>Zhang et al. (2020) concluded that anthropogenic forcing contributed to 2018 drought, principally as consequence of enhanced AED.</p> <p>One study suggests that soil moisture drought conditions in northern China have been intensified by agriculture (Liu et al., 2015).</p>	<p><b>Low confidence: Inconsistent trends</b> depending on model, subregion and index (Huang et al., 2018a; Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)) .</p> <p>(Huang et al., 2018a): Inconsistent projections in a study with a single GCM for the time frame 2016-2050 (for different scenarios) compared to 1960-2005, i.e corresponding to 1.5°C projections compared to recent past.</p>	<p><b>Low confidence: Mixed signals</b> depending on model, subregion and index (Gao et al., 2017b; Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)).</p> <p>(Gao et al., 2017b): Study for very small region (Loess Plateau).</p>	<p><b>Medium confidence: Increasing dryness</b> as dominant signal in projections and over larger part of domain, but also inconsistent signal for some indices and part of the domain (Cook et al., 2014b, 2020; Cheng et al., 2015; Dai et al., 2018; Naumann et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).</p>
--------------	---	---	---	--	---

	<p>HYDR</p>	<p><b>Medium confidence: Increase</b> in hydrological drought in the region, in particular in northern China; inconsistent trends in part of the region (Liu et al., 2015; Dai and Zhao, 2017; Zhang et al., 2018b).</p> <p>Drying in large part of domain, in particular in northern China (Zhao and Dai, 2017)</p> <p>Increase of hydrological droughts in the Yangtze river (Zhang et al., 2018b).</p>	<p><b>Low confidence: Limited evidence and mixed signals.</b> Available evidence suggests that a combination of change in climatic drivers (precipitation, Epot) and human drivers (agriculture, water management) are responsible for trends (Liu et al., 2015; Zhang et al., 2018b).</p> <p>Increasing hydrological droughts trends in the Yangtze river are dominantly driven by precipitation, but increases in potential evaporation and human activities also play a role (Zhang et al., 2018b). Drought conditions in northern China (soil moisture and runoff) have been intensified by agriculture (Liu et al., 2015).</p>	<p><b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).</p>	<p><b>Low confidence: Limited evidence and inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b).</p>	<p><b>Low confidence: Inconsistent</b> trend between models and studies, and generally low signal-to-noise ratio (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)</p> <p>(Touma et al., 2015; Cook et al., 2020): Generally inconsistent trends between models, with low model agreement.</p> <p>(Giuntoli et al., 2015): Trend towards drying but generally low signal-to-noise ratio except in small subregion.</p>
--	-------------	---	---	--	--	---

Eastern Central Asia (ECA)	MET	<b>Low confidence: Inconsistent trends</b> between subregions, with overall tendency to decrease (Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence. Limited evidence;</b> slight decrease in meteorological drought in available analyses (Xu et al., 2013) (Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Decrease</b> in drought severity, with weakly inconsistent changes for some indices (Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM))  (Spinoni et al., 2019): Strong decrease in drought for SPI-based metrics in RCP4.5 compared to 1981-2010  (Xu et al., 2019a): Decrease in frequency of SPI-based events but slight increase or inconsistent changes in duration of SPI-based events.  Chapter 11 Supplementary Material (11.SM): substantial decrease in CDD	<b>Medium confidence: Decrease</b> in drought severity (Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR, ECOL	<b>Medium confidence: Increase</b> in drying, but some conflicting trends between drought metrics and sub-regions (Greve et al., 2014; Cheng et al., 2015; Dai and Zhao, 2017; Li et al., 2017c; Spinoni et al., 2019; Padrón et al., 2020; Zhang et al., 2020c).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed signal</b> in changes in drought severity, lack of signal based in total column soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> in changes in drought severity. Inconsistent trends in total and surface soil moisture, with stronger tendency to wetting, (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), and drying based on the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed trends</b> between different models and drought metrics (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited evidence.</b> Mostly inconsistent trends in one study (Dai and Zhao, 2017).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Limited evidence and inconsistent trends</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Mixed trends.</b> Model disagreement and inconsistent changes among studies, seasons and metrics, with overall low signal-to-noise ratio (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020).

Tibetan Plateau (TIB)	MET	<b>Low confidence:</b> Inconsistent trends (Jiang et al., 2013; Donat et al., 2016a; Hu et al., 2016; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> Weak or inconsistent trends in available analyses (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends,</b> but tendency towards wetting (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM))  (Spinoni et al., 2019): No data in the region (Cook et al., 2020): Only analysis of mean precipitation but tendency towards wetting in all seasons in the region (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)): Weak trends but tendency towards wetting.	<b>Low confidence: Inconsistent trends</b> between models, but tendency towards wetting and decrease in drought (Zhou et al., 2014; Touma et al., 2015)(Chapter 11 Supplementary Material (11.SM)).  (Zhou et al., 2014): Decrease of CDD is projected but there is large uncertainty
	AGR, ECOL	<b>Low confidence: Inconsistent trends.</b> Spatially varying trends, with slight tendency to overall wetting(Cheng et al., 2015; Dai and Zhao, 2017; Jia et al., 2018; Zhang et al., 2018a; Li et al., 2020c; Wang et al., 2020).  (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020): Missing data in most of region.	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> between models, indices and subregions (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent trends</b> between models, indices and subregions (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent trends</b> between models, indices and subregions (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).
	HYDR	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Limited evidence and inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Inconsistent trends</b> between models and studies, and low signal-to-nois ratio (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)

South Asia (SAS)	MET	<b>Medium confidence: Increase in meteorological drought.</b> Subregional differences but drying is dominant (Mishra et al., 2014b; Malik et al., 2016; Guhathakurta et al., 2017; Spinoni et al., 2019; Dunn et al., 2020) (see also Section 10.6.3)	<b>Low confidence: Limited evidence</b> (Fadnavis et al., 2019)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Inconsistent trends</b> , with light tendency to decreased drying (Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends</b> depending on model and subregion, with light tendency to decreases in meteorological drought in CMIP5 and CMIP6 (Mishra et al., 2014b; Touma et al., 2015; Salvi and Ghosh, 2016; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)); light increased drying in NDD in CORDEX-CORE (Coppola et al., 2021b). Overall poor climate model performance for South Asia monsoon in CMIP5 and CORDEX (Mishra et al., 2014a; Saha et al., 2014; Sabecrali et al., 2015; Singh et al., 2017b). See also Section 10.6.3 for assessment for changes in Indian summer monsoon rainfall.
	AGR, ECOL	<b>Low confidence: Lack of signal and inconsistent trends</b> depending on subregion based on soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Mishra et al., 2014b; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020) and decrease of the drying effect of the atmospheric evaporative demand (Jhajharia et al., 2015).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends in drought between models and subregions</b> (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends in drought between models, subregions and studies</b> , but slight dominant tendency towards wetting(Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020)(CMIP6.ANNEX-CH11)	<b>Medium confidence: Decreased drying trend</b> (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2014b, 2020; Mishra et al., 2014b; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Limited evidence.</b> Inconsistent trends or limited data in available studies (Zhao and Dai, 2017; Gudmundsson et al., 2019, 2021).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015).	<b>Low confidence: Limited evidence.</b> Lack of signal in CMIP5 (Touma et al., 2015). Decrease in dryness in CMIP6 (Cook et al., 2020); mostly inconsistent trends in HAPPI-MIP driven simulations with one hydrological model (Zhai et al., 2020b).	<b>Low confidence: Inconsistent trends</b> between models and studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)



<p>Southeast Asia (SEA)</p>	<p>MET</p>	<p><b>Low confidence: Inconsistent trends</b> between subregions (Spinoni et al., 2019; Dunn et al., 2020).</p>	<p><b>Low confidence: Limited evidence</b> (Mcbride et al., 2015; King et al., 2016b) although the the equatorial Asia drought of 2015 has been attributed to anthropogenic warming effects (Shiogama et al., 2020).</p>	<p><b>Low confidence: Limited evidence</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))</p>	<p><b>Low confidence: Inconsistent trends</b> between models, subregions and studies (Tangang et al., 2018; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)) but with overall drying in CMIP6 and CORDEX simulations (Tangang et al., 2018; Cook et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM)).</p> <p>(Tangang et al., 2018): Projected drying based on CDD in CORDEX simulations for Indonesia</p> <p>(Xu et al., 2019a): Inconsistent trends across region based on SPI, but with slight drying over Indonesia</p> <p>(Spinoni et al., 2020): Wetting trend based on SPI</p> <p>(Chapter 11 Supplementary Material (11.SM)): Drying trend based on CDD</p>	<p><b>Medium confidence: Increase</b> in drying in CMIP6 and CORDEX simulations (Cook et al., 2020; Supari et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM)). but inconsistent trends or wetting in CMIP5-based projections(Touma et al., 2015; Cook et al., 2020; Spinoni et al., 2020; Supari et al., 2020)(</p> <p>(Supari et al., 2020): Strong drying trend based on CDD in CORDEX simulations for Indonesia</p> <p>(Coppola et al., 2021b): Drying based on number of dry days (NDD) in CORDEX-CORE projects</p> <p>(Cook et al., 2020): Decreasing trend in mean precipitation which is only found in CMIP6 and not in CMIP5.</p> <p>Chapter 11 Supplementary Material (11.SM): Strong projected drying trend based on CDD in CMIP6 projections</p> <p>(Touma et al., 2015): Inconsistent trends in SPI in CMIP5 projections</p> <p>(Spinoni et al., 2020): Wetting trend based on SPI in CMIP5 projections.</p> <p>(Cai et al., 2014a, 2015, 2018): An increasing frequency of precipitation deficits is projected as a consequence of an increasing frequency of extreme El Niño.</p>
-----------------------------	------------	---	--	--	--	---

AGR, ECOL	<b>Low confidence: Inconsistent trends</b> depending on subregion and index based on soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> depending on model, subregion, index or study (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent trends</b> depending on model, subregion, index or study (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signal</b> depending on model and metric. Drying tendency based on CMIP6 soil moisture projections (Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), inconsistent trends in CMIP5 surface soil moisture (Dai et al., 2018; Lu et al., 2019), but wetting trends with PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a) in studies driven with CMIP5 data. ,  (Cook et al., 2020): Drying trend in SEA in CMIP6), but not in CMIP5.
HYDR	<b>Low confidence: Limited evidence.</b> Regionally inconsistent trends in one study (Dai and Zhao, 2017).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows decrease in hydrological drought (Touma et al., 2015).	<b>Low confidence: Limited evidence and inconsistent trends</b> in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence:</b> Inconsistent trend between models and studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)

1  
2  
3  
4  
5  
6  
7  
8

[END TABLE 11.9 HERE]

[START TABLE 11.10 HERE]

**Table 11.10:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in Australasia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Australasia	Significant increases in the intensity and frequency of hot extremes and decreases in the intensity and frequency of cold extremes (CSIRO and BOM, 2015; Jakob and Walland, 2016; Alexander and Arblaster, 2017)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Hu et al., 2020; Wang et al., 2017).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)  Additional evidence from CMIP5 simulations for an	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)  Additional evidence from CMIP5 simulations for an	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)  Additional evidence from CMIP5 simulations for an

			increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>very likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Northern Australia (NAU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Perkins and Alexander, 2013; Wang et al., 2013c; CSIRO and BOM, 2015; Donat et al., 2016a; Alexander and Arblaster, 2017; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wang et al., 2017; Hu et al., 2020; Seong et al., 2020; Knutson et al., 2014; Lewis and Karoly, 2014; Perkins et al., 2014; Arblaster et al., 2014; Hope et al., 2015, 2016; Perkins and Gibson, 2015)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)
	<i>High confidence</i> in the increase in the intensity and	<i>High confidence</i> in a human contribution to the observed	Increase in the intensity and frequency of hot extremes:	Increase in the intensity and frequency of hot extremes:	Increase in the intensity and frequency of hot extremes:

	frequency of hot extremes and <i>likely</i> decrease in the intensity and frequency of cold extremes	increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	<i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	<i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Central Australia (CAU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Perkins and Alexander, 2013; Wang et al., 2013c; CSIRO and BOM, 2015; Donat et al., 2016a; Alexander and Arblaster, 2017; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes ((Wang et al., 2017c; Hu et al., 2020; Seong et al., 2020; Knutson et al., 2014; Lewis and Karoly, 2014; Perkins et al., 2014; Arblaster et al., 2014; Hope et al., 2015, 2016; Perkins and Gibson, 2015; King et al., 2014)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with pre-industrial)

			<i>Very likely</i> (compared with pre-industrial)	<i>Extremely likely</i> (compared with pre-industrial)	with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Eastern Australia (EAU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Perkins and Alexander, 2013; Wang et al., 2013c; CSIRO and BOM, 2015; Donat et al., 2016a; Alexander and Arblaster, 2017; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes ((Wang et al., 2017c; Hu et al., 2020; Seong et al., 2020; Knutson et al., 2014; Lewis and Karoly, 2014; Perkins et al., 2014; Arblaster et al., 2014; Hope et al., 2015, 2016; Perkins and Gibson, 2015; King et al., 2015)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Southern Australia (SAU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity

	<p>frequency of cold extremes (Perkins and Alexander, 2013; Wang et al., 2013c; Dittus et al., 2014; CSIRO and BOM, 2015; Crimp et al., 2016; Donat et al., 2016a; Alexander and Arblaster, 2017; Dunn et al., 2020)</p>	<p>and decrease in the intensity and frequency of cold extremes (Wang et al., 2017c; Hu et al., 2020; Seong et al., 2020; Black and Karoly, 2016; Knutson et al., 2014; Lewis and Karoly, 2014; Perkins et al., 2014; Arblaster et al., 2014; Hope et al., 2015, 2016; Perkins and Gibson, 2015)</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)</p>	<p>and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Alexander and Arblaster, 2017; Herold et al., 2018; Evans et al., 2020; Grose et al., 2020)</p>
	<p><i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes</p>	<p>Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
New Zealand (NZ)	<p>Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Caloiero, 2017; Dunn et al., 2020; Ministry for the Environment &amp; Stats NZ, 2020; Harrington, 2020)</p>	<p>Limited evidence (Seong et al., 2020; Wang et al., 2017)</p>	<p>CMIP6 models project an increase in the intensity and frequency of TXx events and a decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3°C in annual TXx and TNn compared to pre-industrial</p>

	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Low confidence</i>	Increase in the intensity and frequency of hot extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	(Annex). Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	(Annex). Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
--	--	-----------------------	---	---	---

[END TABLE 11.10 HERE]

[START TABLE 11.11 HERE]

**Table 11.11:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in Australasia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.3 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Australasia	Limited evidence (Jakob and Walland, 2016; Guerreiro et al., 2018b; Dey et al., 2019b; Dunn et al., 2020; Sun et al., 2020)	Limited evidence	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
Northern Australia (NAU)	Intensification of heavy	Limited evidence (Dey et al.,	CMIP6 models project	CMIP6 models project an	CMIP6 models project a robust

	precipitation (Donat et al., 2016a; Alexander and Arblaster, 2017; Evans et al., 2017; Dey et al., 2019b; Dunn et al., 2020; Sun et al., 2020)	2019a)	inconsistent changes in the region (Li et al., 2020a)	increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).	increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
Central Australia (CAU)	Limited evidence (Donat et al., 2016a; Alexander and Arblaster, 2017; Evans et al., 2017; Dey et al., 2019b; Dunn et al., 2020; Sun et al., 2020).	Limited evidence	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
Eastern Australia (EAU)	Lack of agreement on the evidence of trends (Donat et al., 2016a; Alexander and Arblaster, 2017; Evans et al.,	Limited evidence	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020;



	2017; Dey et al., 2019b; Dunn et al., 2020; Sun et al., 2020)				Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)
Southern Australia (SAU)	Limited evidence (Donat et al., 2016a; Alexander and Arblaster, 2017; Evans et al., 2017; Dey et al., 2019b; Dunn et al., 2020; Sun et al., 2020)	Limited evidence	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)
New Zealand (NZ)	Lack of agreement on the evidence of trends (Donat et al., 2016a; Dunn et al., 2020; MfE and Stats NZ, 2020)	Limited evidence (Rosier et al., 2016)	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and

					more than 15% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)

1  
2  
3  
4  
5  
6  
7  
8  
9

[END TABLE 11.11 HERE]

[START TABLE 11.12 HERE]

**Table 11.12:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET), agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in Australasia, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.4 for details.

Region and drought type		Observed trends	Human contribution	Projections		
				+1.5 °C	+2 °C	+4 °C
Northern Australia (NAU)	MET	<b>Medium confidence:</b> Decrease in the frequency and intensity of meteorological droughts (Gallant et al., 2013; Delworth and Zeng, 2014; Alexander and Arblaster, 2017; Knutson and Zeng, 2018; Dey et al., 2019a; Dunn et al., 2020)	<b>Low confidence</b> in attribution (Delworth and Zeng, 2014; Knutson and Zeng, 2018; Dey et al., 2019a).	<b>Low confidence: Increases or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Model disagreement in SPI projections (Spinoni et al., 2020)  Increase in CDD-based drought in CMIP5, but generally not significant (Alexander and Arblaster, 2017)  Slight increase in CDD-based drought in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Increases or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Large intermodel spread in changes in SPI in CMIP5 projections (Kirono et al., 2020)  Model disagreement in SPI projections (Spinoni et al., 2020)  Increase in CDD-based drought in CMIP5, but generally not significant (Alexander and Arblaster, 2017)  Slight increase in CDD-based	<b>Low confidence: Increases or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Grose et al., 2020; Kirono et al., 2020; Spinoni et al., 2020; Ukkola et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Large intermodel spread in changes in SPI in CMIP5 projectons, but slight drying for median (Kirono et al., 2020)  Model disagreement in SPI projections (Spinoni et al., 2020)  Increase in CDD-based drought in CMIP5, but generally not significant (Alexander and Arblaster, 2017)

					drought CMIP6 (Chapter 11 Supplementary Material (11.SM))	Increase in CDD-based drought in CMIP6 (Grose et al., 2020)(Chapter 11 Supplementary Material (11.SM))  Inconsistent trends in mean precipitation in CORDEX RCMs, but drying trend on annual scale at northern tip of region (Evans et al., 2020)
	AGR ECOL	<b>Medium confidence: Decrease</b> in agricultural and ecological drought  Decrease in frequency (but not intensity) of soil moisture-based droughts (Gallant et al., 2013). Inconsistent signals in changes in water-balance (Greve et al., 2014; Padrón et al., 2019). Decrease in agricultural and ecological drought based on SPEI-PM from 1950-2009 (Beguería et al., 2014; Spinoni et al., 2019) and PDSI_PM (Dai and Zhao, 2017)	<b>Low confidence Limited evidence</b>  Lack of studies although (Lewis et al., 2019b) supported an anthropogenic attribution of 2018 drought associated with more extreme temperatures that exacerbated AED and ET, and depleting soil moisture.	<b>Low confidence: Increase or non-robust</b> (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020).(Chapter 11 Supplementary Material (11.SM))  Cook et al. (2020): non-robust changes in surface and column soil moisture in both summer and winter half years (CMIP6 projections)	<b>Low confidence: Increase or non-robust</b> (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020).(Chapter 11 Supplementary Material (11.SM))  Cook et al. (2020): non-robust changes in surface and column soil moisture in both summer and winter half years (CMIP6 projections)  Kirono et al. (2020): Standardized soil moisture index based on surface soil moisture: drying trend for median in CMIP5 but large intermodal spread	<b>Low confidence: Increase or non-robust</b> , with higher increases in SPEI-PM but non-robust changes in CMIP6 soil moisture (Naumann et al., 2018; Cook et al., 2020; Kirono et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).  Cook et al. (2020): non-robust changes in surface and column soil moisture in both summer and winter half years (CMIP6 projections)  Kirono et al. (2020): Standardized soil moisture index based on surface soil moisture : drying trend for median in CMIP5, but larger inter-model spread
	HYDR	<b>Low confidence</b> because of <b>lack of data and studies</b>	<b>Low confidence Limited evidence</b> because of lack of data and studies	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence</b> and generally <b>non-robust</b> change in two studies (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Non-robust changes or high model disagreement</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Central Australia (CAU)	MET	<b>Medium confidence: decrease</b> in the frequency/intensity of droughts (Gallant et al., 2013; Beguería et al., 2014; Delworth and Zeng, 2014; Greve et al., 2014; Alexander and Arblaster, 2017; Knutson and Zeng, 2018).	<b>Low confidence</b> in attribution (Delworth and Zeng, 2014; Knutson and Zeng, 2018).	<b>Low confidence: Inconsistent or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Tendency to increasing SPI-based drought in CMIP6, but to decreasing SPI-based drought in CORDEX (Spinoni et al., 2020)	<b>Low confidence: Inconsistent or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Tendency to increasing SPI-based drought in CMIP6, but to decreasing SPI-based drought in CORDEX (Spinoni et al., 2020)  Kirono et al. (2020): CMIP6 models	<b>Low confidence: Inconsistent or non-robust</b> changes in meteorological droughts (Alexander and Arblaster, 2017; Grose et al., 2020; Kirono et al., 2020; Spinoni et al., 2020; Ukkola et al., 2020) (Chapter 11 Supplementary Material (11.SM)).  Tendency to increasing SPI-based drought in CMIP6, but to decreasing SPI-based drought in CORDEX (Spinoni et al., 2020)

					project increased in SPI in much of region for 2006-2100 under RCP8.5	Kirono et al. (2020); CMIP6 models project increased in SPI in much of region for 2006-2100 under RCP8.5
	AGR ECOL	<b>Low confidence: Inconsistent</b> changes in frequency/intensity of droughts (Gallant et al., 2013; Beguería et al., 2014; Delworth and Zeng, 2014; Greve et al., 2014; Dai and Zhao, 2017; Knutson and Zeng, 2018; Padrón et al., 2019; Spinoni et al., 2019)	<b>Low confidence</b> because of <b>lack of studies</b>	<b>Low confidence: Inconsistent changes</b> both in soil moisture and SPEI-PM (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent changes</b> both in soil moisture and SPEI-PM (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Increased drying</b> for some metrics or part of domain for soil moisture and SPEI-PM with stronger changes for SPEI-PM (Naumann et al., 2018; Cook et al., 2020; Kirono et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence</b> because of <b>lack of data and studies</b>	<b>Low confidence Limited evidence</b> , because of lack of studies	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence</b> and generally <b>non-robust</b> change in two studies (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Non-robust changes or high model disagreement</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Eastern Australia (EAU)	MET	<b>Low confidence: Inconsistent trends</b> (Gallant et al., 2013; Delworth and Zeng, 2014; Alexander and Arblaster, 2017; Knutson and Zeng, 2018; Spinoni et al., 2019)  Gallant et al. (2013): Inconsistent trends, wetting on average in MDB Delworth and Zeng (2014): no trend Knutson and Zeng (2018): no trend Alexander and Arblaster (2017); Dunn et al. (2020): no trends in CDD Spinoni et al. (2019): Inconsistent trends, some increased severity in part of the region	<b>Low confidence</b> in attribution (Delworth and Zeng, 2014; King et al., 2014; Knutson and Zeng, 2018)	<b>Low confidence: Increase</b> in meteorological droughts based on CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Kirono et al., 2020), but weak signals and lack of other studies at this GWL.	<b>Medium confidence: Increases</b> in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)) .	<b>Medium confidence:</b> Increases in meteorological droughts (Alexander and Arblaster, 2017; Grose et al., 2020; Kirono et al., 2020; Spinoni et al., 2020; Ukkola et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Low confidence: Inconsistent trends</b> (Gallant et al., 2013; Beguería et al., 2014; Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence</b> because of lack of studies although enhanced AED driven by extreme temperatures increased the severity of the 2019 drought (van Oldenborgh et al., 2021)	<b>Low confidence: Inconsistent changes</b> in soil moisture and SPEI-PM, but tendency to <b>increase</b> (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Increase</b> in drought based on soil moisture and SPEI-PM, but partly inconsistent changes for some studies (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>High confidence: Increased drying</b> for some metrics or part of domain for soil moisture and SPEI-PM with stronger changes for SPEI-PM (Naumann et al., 2018; Cook et al., 2020; Kirono et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))

	HYDR	<b>Low confidence: Limited evidence</b> because of <b>lack of data and studies</b> (Zhang et al., 2016d)	<b>Low confidence: Limited evidence</b> , because of lack of studies	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Lack of studies</b> and generally <b>non-robust</b> change in two studies (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Non-robust changes or high model disagreement</b> (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
Southern Australia (SAU)	MET	<b>Low confidence: Mixed signal</b> depending on subregion, index and season (Gallant et al., 2013; Delworth and Zeng, 2014; Alexander and Arblaster, 2017; Spinoni et al., 2019; Dunn et al., 2020; Rauniyar and Power, 2020)(Dai and Zhao, 2017).  Gallant et al. (2013): Wetting in eastern part, drying in eastern part  Rauniyar and Power (2020): Recovery from Millenium drought  Delworth and Zeng (2014): Only drying in the western part, not in the eastern part  Alexander and Arblaster (2017); Dunn et al. (2020): Overall decreasing CDD trends  Spinoni et al. (2019): Decreasing droughts in most of domain	<b>Low confidence: Mixed signal</b> in observations.  Increase in the frequency/intensity of meteorological droughts can be attributed to anthropogenic forcing (greenhouse gases, ozone and aerosols) (Delworth and Zeng, 2014; Karoly et al., 2016; Knutson and Zeng, 2018) (Cai et al., 2014b).	<b>Medium confidence: Increase</b> overall in meteorological droughts based on CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Kirono et al., 2020); but weak signals and lack of other studies at this GWL.	<b>Medium confidence: Increases</b> in meteorological droughts (Alexander and Arblaster, 2017; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence:</b> Increases in meteorological droughts (Alexander and Arblaster, 2017; Grose et al., 2020; Kirono et al., 2020; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Medium confidence: Increase.</b> Dominant increasing drying signal but some inconsistent trends depending on subregion and index; strongest drying trend in Western SAU. (Gallant et al., 2013; Beguería et al., 2014; Greve et al., 2018; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence,</b> Enhanced AED driven by extreme temperatures increased the severity of the 2019 drought (van Oldenborgh et al., 2021)	<b>Medium confidence: Increase</b> in soil moisture and SPEI-PM, but partly inconsistent changes for some studies (Naumann et al., 2018; Xu et al., 2019a; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Increase</b> in drought based on soil moisture and SPEI-PM, but partly inconsistent changes for some studies (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Kirono et al., 2020)(Chapter 11 Supplementary Material (11.SM))	<b>High confidence: Increased drying</b> for some metrics or part of domain for soil moisture and SPEI-PM with stronger changes for SPEI-PM (Naumann et al., 2018; Cook et al., 2020; Kirono et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM))

	HYDR	<b>Medium confidence: Increasing drying signal in the southeast and particularly the southwest. Some dependence on time frame</b> in available studies (Gudmundsson et al., 2019, 2021)(Zhang et al., 2016d)	<b>Low confidence : Limited evidence</b> because of lack of studies (Cai and Cowan, 2008)	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Medium confidence: Increase in drought,</b> but some Inconsistent and non-robust change including subregional/seasonal differences (Touma et al., 2015; Zheng et al., 2019; Cook et al., 2020)	<b>Medium confidence: Increase in drought,</b> but some inconsistent changes depending on season or study (Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
New Zealand (NZ)	MET	<b>Low confidence: Inconsistent</b> changes (Caloiero, 2015; Spinoni et al., 2015; Knutson and Zeng, 2018)	<b>Low confidence</b> in attribution of trends (Harrington et al., 2014, 2016; Knutson and Zeng, 2018).	<b>Low confidence: Lack of studies and lack of signal</b> for CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent</b> changes, but increase in Northern Island (MfE, 2018; MfE and Stats NZ, 2020; Spinoni et al., 2020).(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent</b> changes, but increase in Northern Island (MfE, 2018; MfE and Stats NZ, 2020; Spinoni et al., 2020).(Chapter 11 Supplementary Material (11.SM))
	AGR ECOL	<b>Low confidence: Inconsistent</b> trends. Increase in drying in part of the country based on soil moisture and SPEI-PM (Beguería et al., 2014; Spinoni et al., 2019; MfE and Stats NZ, 2020); decrease in PDSI-PM (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b> because of lack of studies	<b>Low confidence: Lack of studies and lack of signal</b> for soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Inconsistent</b> changes, but increase in Northern Island (MfE, 2018; MfE and Stats NZ, 2020; Spinoni et al., 2020).	<b>Low confidence: Inconsistent</b> changes, but increase in Northern Island (MfE, 2018; MfE and Stats NZ, 2020; Spinoni et al., 2020).
	HYDR	<b>Low confidence: Lack of data and studies</b>	<b>Low confidence: Lack of studies</b>	<b>Low confidence: Lack of studies</b>	<b>Low confidence: Lack of studies</b>	<b>Low confidence: Lack of studies</b>

1  
2  
3  
4  
5  
6  
7  
8

[END TABLE 11.12 HERE]

[START TABLE 11.13 HERE]

**Table 11.13:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in Central and South America, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Central and South America	Most subregions show a <i>likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Hu et al., 2020; Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)

			Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a)	Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a)	Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a)
	<i>High confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
South Central America (SCA)	Increases in the intensity and frequency of hot extremes and decreases in the intensity and frequency of cold extremes (Dunn et al. 2020; Aguilar et al. 2005)	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wang et al. 2017, Seong et al. 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Imbach et al., 2018; Angeles-Malaspina et al., 2018; Chou et al., 2014)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Coppola et al., 2021b; Angeles-Malaspina et al., 2018; Chou et al., 2014)
	<i>Medium confidence</i> in the	<i>Medium confidence</i> in a	Increase in the intensity and	Increase in the intensity and	Increase in the intensity and

	increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Caribbean (CAR)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Angeles-Malaspina, González-Cruz & Ramírez-Beltran; 2018; McLean et al., 2015; Dunn et al., 2020)	Strong evidence of changes from observations that are in the direction of model projected changes for the future. The magnitude of projected changes increases with global warming.	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Annex). Median increase of more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Angeles-Malaspina, González-Cruz & Ramírez-Beltran; 2018; Chou et al., 2014)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than XC in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than XC in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Coppola et al., 2021b; Angeles-Malaspina, González-Cruz & Ramírez-Beltran; 2018; Chou et al., 2014; Hall et al., 2013)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)



			frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Northwestern South America (NWS)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Northern South America (NSA)	Significant increases in the intensity and frequency of hot	Evidence of a human contribution to the observed	CMIP6 models project a robust increase in the intensity and	CMIP6 models project a robust increase in the intensity and	CMIP6 models project a robust increase in the intensity and

	<p>extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020), Avila-Diaz et al.; 2020; Geirinhas et al., 2018; Dunn et al., 2020)</p>	<p>increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)</p>	<p>frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).</p>	<p>frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).</p>	<p>frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014)</p>
	<p><i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes</p>	<p><i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
South American Monsoon (SAM)	<p>Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020; Avila-Diaz et al., 2020; Geirinhas et al., 2018; Dunn et al., 2020)</p>	<p>Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in</p>

			TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Northeastern South America (NES)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020), Avila-Diaz et al., 2020; Geirinhas et al., 2018; Dunn et al., 2020)	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes

			(Chou et al., 2014a).	(Chou et al., 2014a).	cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Southwestern South America (SWS)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020; Olmo et al., 2020; Dunn et al., 2020)	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared

		cold extremes.	Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	with pre-industrial) Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Southeastern South America (SES)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Dereczynski et al., 2020; Avila-Diaz et al., 2020; Geirinhas et al., 2018; Rusticucci et al., 2017; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014).
	<i>High confidence</i> in the increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

Southern South America (SSA)	Inconsistent trends and insufficient data (Dereczynski et al., 2020; Ceccherini et al., 2016; (1980-2014) Dunn et al., 2020)		CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Chou et al., 2014a).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 2.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (López-Franca et al., 2016; Coppola et al., 2021b; Chou et al., 2014).
	<i>Low confidence</i>	<i>Low confidence</i>	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial).	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

1  
2  
3  
4  
5  
6

[END TABLE 11.13 HERE]

[START TABLE 11.14 HERE]

1 **Table 11.14:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in Central and South  
2 America, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.3 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Central and South America	Insufficient data to assess trends	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 0% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
South Central America (SCA)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Stephenson et al., 2014)	Limited evidence	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Imbach et al., 2018; Chou et al., 2014).	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014).	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014; Coppola et al., 2021b; Kusunoki et al., 2019; Nakaegawa et al., 2013)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Low confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)
Caribbean (CAR)	Insufficient data and a lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020; McLean et al., 2015; Stephenson et al., 2014)	Evidence of a human contribution for some events (Patricola and Wehner, 2018), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Coppola et al., 2021b; Chou et al., 2014; Nakaegawa et al.,

	<i>Low confidence</i>	<i>Low confidence.</i>	<i>Low confidence</i>	<i>Low confidence</i>	2013; Hall et al., 2013) <i>Low confidence</i>
Northwestern South America (NWS)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020)	Disagreement among studies (Li et al., 2019; Otto et al., 2018a)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)
	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>
Northern South America (NSA)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020; Avila-Diaz et al., 2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1 day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1 day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)
South American Monsoon (SAM)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020; Avila-Diaz et al., 2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1 day and Rx5day and 2% in annual Rx30day compared	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1 day and Rx5day and 4% in annual Rx30day compared



				to pre-industrial (Annex). Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.	to pre-industrial (Annex). Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)
Northeastern South America (NES)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020; Avila-Diaz et al., 2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Conflicting projections by the CMIP6 multi-model ensemble and limited RCM simulations; more weight is given to the CMIP6 results.
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)
Southwestern South America (SWS)	Insufficient data coverage and trends in available data are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020; Olmo et al., 2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)

	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>	<i>Low confidence</i>
Southeastern South America (SES)	Significant intensification of heavy precipitation Dunn et al., 2020; Dereczynski et al., 2020; Olmo et al., 2020; Avila-Diaz et al. (2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)
	<i>High confidence</i> inintensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)
Southern South America (SSA)	Insufficient data coverage and trends are generally not significant (Sun et al., 2020; Dunn et al., 2020; Dereczynski et al., 2020)	Evidence of a human contribution for some events (Li et al., 2019d), but cannot be generalized	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Chou et al., 2014)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 2% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Chou et al., 2014a)

	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
--	-----------------------	-----------------------	---	--	---

1  
2 [END TABLE 11.14 HERE]

3  
4  
5 [START TABLE 11.15 HERE]

6  
7 **Table 11.15:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET),  
8 agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in Central and South America, subdivided by AR6 regions. See Sections 11.9.1  
9 and 11.9.4 for details.

Region		Observed trends	Human contribution	Projections		
				+1.5 °C	+2 °C	+4 °C
South Central America (SCA)	MET	<b>Low confidence: Mixed signal.</b> Dominant decrease in drought duration but mixed trends between subregions (Aguilar et al., 2005; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b> Available evidence suggests increase in drought severity (Chapter 11 Supplementary Material (11.SM) (Chou et al., 2014a; Imbach et al., 2018))  (Chou et al., 2014a): RCM simulations with Eta model driven with 2 different GCMs.	<b>Medium confidence: Increase</b> in drought severity (Chou et al., 2014a; Imbach et al., 2018; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM))  (Chou et al., 2014a): RCM simulations with Eta model driven with 2 different GCMs.	<b>High confidence: Increase</b> in drought severity (Nakaegawa et al., 2013; Chou et al., 2014a; Touma et al., 2015; Corrales-Suastegui et al., 2019; Kusunoki et al., 2019; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM) ).  (Chou et al., 2014a): RCM simulations with Eta model driven with 2 different GCMs.
	AGR ECOL	<b>Low confidence: Mixed signal.</b> Mixed trends in different subregions and in different drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Mixed signal</b> in drought trends. Inconsistent drying trend (but stronger tendency towards drying) based on total column soil moisture (Imbach et al., 2018; Xu et al., 2019a)( <b>Chapter 11 Supplementary Material (11.SM)</b> ) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase</b> in drought based on total and surface soil moisture (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM) and on SPEI-PM (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020).	<b>High confidence: Increase</b> in drought severity with different metrics and high agreement between studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Insufficient evidence</b> (Dai and Zhao, 2017; Gudmundsson et al., 2021).	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b> One study shows inconsistent changes (Touma et al.,	<b>Low confidence: Limited evidence.</b> Inconsistent changes (Touma et al., 2015) or drying in	<b>Medium confidence: Increase</b> in drought severity (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et

				2015)	part of region (Cook et al., 2020)	al., 2015; Cook et al., 2020)
Caribbean (CAR)	MET	<b>Low confidence: Mixed signal.</b> Mixed trends between subregions, but some evidence of increases in drought duration (Stephenson et al., 2014; McLean et al., 2015; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Increase</b> in drought duration (Chou et al., 2014a); inconsistent changes in CDD (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Limited evidence and inconsistent changes.</b> One study suggests increase in drought duration (Chou et al., 2014a), but CMIP6 projections show inconsistent changes in CDD (Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Increase</b> in drought duration (Chapter 11 Supplementary Material (11.SM)) (Nakaegawa et al., 2013; Chou et al., 2014a; Stennett-Brown et al., 2017; Coppola et al., 2021b)
	AGR ECOL	<b>Low confidence: Mixed signal.</b> Mixed trends between subregions with PDSI-PM and SPEI-PM (Dai and Zhao, 2017; Spinoni et al., 2019).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> in total column and surface soil moisture (Chapter 11 Supplementary Material (11.SM) , and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase</b> , but including mixed signal in changes of drought severity, with inconsistent trends in total soil moisture, (Chapter 11 Supplementary Material (11.SM) , and drying trend based on SPEI-PM (Naumann et al., 2018; Gu et al., 2020). See also Chapter 12.	<b>Medium confidence: Increase.</b> Drying trend with surface soil moisture (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a). Total soil moisture shows weak (Chapter 11 Supplementary Material (11.SM) or no signal (Cook et al., 2020)
	HYDR	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b>
North-western South America (NWS)	MET	<b>Low confidence: Mixed signal.</b> Mixed trends between subregions (Skansi et al., 2013; Spinoni et al., 2019; Dereczynski et al., 2020; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Inconsistent trends</b> (Chapter 11 Supplementary Material (11.SM) (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a)	<b>Low confidence: Mixed signal</b> between different studies and models (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a; Spinoni et al., 2020) (Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Increase.</b> Dominant signal is positive CDD trend (increasing dryness; Chapter 11 Supplementary Material (11.SM)); also some mixed signals between different studies (Chou et al., 2014a; Duffy et al., 2015; Touma et al., 2015; Spinoni et al., 2020; Coppola et al., 2021b)
	AGR ECOL	<b>Low confidence: Mixed trends</b> between subregions and drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed trends</b> based on different metrics, including decrease in total column soil moisture, (Chapter 11 Supplementary Material (11.SM) , weak drying with surface soil moisture (Xu et al., 2019a) and wetting based on the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> in changes in drought severity with drying in total column soil moisture, (Chapter 11 Supplementary Material (11.SM) , lack of signal in the surface soil moisture (Xu et al., 2019a) and wetting trends with SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed trend</b> between different drought metrics (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a) (Chapter 11 Supplementary Material (11.SM)).
	HYDR	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b>	<b>Low confidence: Limited evidence.</b> One study shows inconsistent changes (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Inconsistent changes (Touma et al., 2015; Cook et al., 2020)	<b>Low confidence: Lack of signal</b> (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)

Northern South America (NSA)	MET	<b>Low confidence: Mixed trends</b> between subregions, but some evidence of increased drought duration (Skansi et al., 2013; Marengo and Espinoza, 2016; Spinoni et al., 2019; Avila-Diaz et al., 2020; Dereczynski et al., 2020; Dunn et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Medium confidence:</b> Available evidence suggests drying (Chapter 11 Supplementary Material (11.SM) (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a).	<b>Medium confidence: Increase</b> in drought severity (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a; Spinoni et al., 2020) (Chapter 11 Supplementary Material (11.SM).	<b>High confidence: Increase</b> in drought severity (Chou et al., 2014a; Duffy et al., 2015; Touma et al., 2015; Marengo and Espinoza, 2016; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM).
	AGR ECOL	<b>Low confidence: Mixed trends</b> between subregions and different drought metrics, including soil moisture, PDSI-PM and SPEI-PM, but some evidence of decrease in drought severity (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Medium confidence: Increase</b> in drying. Tendency towards increase in drought severity in total and surface soil moisture (Chapter 11 Supplementary Material (11.SM) (Xu et al., 2019a) inconsistent trends in studies based on the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase.</b> Tendency towards increase in drought severity in total soil moisture (Chapter 11 Supplementary Material (11.SM) , surface soil moisture (Xu et al., 2019a) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>High confidence: Increase</b> in drought severity with different metrics and high agreement between studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited evidence.</b> Available evidence suggests lack of signal (Marengo and Espinoza, 2016; Gudmundsson et al., 2021)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows mixed trends (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Tendency to drying in two studies (Touma et al., 2015; Cook et al., 2020)	<b>High confidence: Increase</b> in drought severity (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
South American Monsoon (SAM)	MET	<b>Medium confidence: Increase</b> in the frequency and severity of meteorological droughts based on SPI and CDD (Spinoni et al., 2019; Avila-Diaz et al., 2020; Dereczynski et al., 2020).	<b>Low confidence: Limited evidence</b> and recent droughts as in 2010 were not attributed to anthropogenic climate change (Shiogama et al., 2013).	<b>Medium confidence: Increase</b> meteorological droughts (Chapter 11 Supplementary Material (11.SM) (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a).. Drying trends in CDD in CMIP6 and SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a) but divergent trends in an RCM driven by two GCMs (Chou et al., 2014a)	<b>Medium confidence: Increase</b> in meteorological droughts (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a) (Chapter 11 Supplementary Material (11.SM). Drying trend in CDD in CMIP6 and SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a) but divergent trends in an RCM driven by two GCMs (Chou et al., 2014a) and weak trends in CMIP5-based SPI projections (Spinoni et al., 2020).	<b>High confidence: Increase</b> in drought severity (Chou et al., 2014a; Touma et al., 2015; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM).
	AGR ECOL	<b>Low confidence: Mixed trends</b> depending on subregions and drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Medium confidence: Increase</b> in agricultural and ecological droughts based on total column and surface soil moisture, (Chapter 11 Supplementary Material (11.SM) (Xu et al., 2019a), and inconsistent signal	<b>High confidence: Increase</b> in drought severity with different metrics (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020) (Chapter 11 Supplementary Material (11.SM).	<b>High confidence: Increase</b> in drought severity with different metrics and high agreement between studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al.,

				with SPEI-PM (Naumann et al., 2018; Gu et al., 2020).		2020a).
	HYDR	<b>Low confidence: Limited evidence.</b> Available evidence suggests lack of signal (Gudmundsson et al., 2021)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows mixed signal (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Mixed signal (Touma et al., 2015) or tendency to drying (Cook et al., 2020)	<b>High confidence: Increase</b> in drought severity (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020).
North-eastern South America (NES)	MET	<b>High confidence: Increase</b> in drought duration (Marengo et al., 2017; Brito et al., 2018; Spinoni et al., 2019; Avila-Diaz et al., 2020; Dereczynski et al., 2020; Dunn et al., 2020)	<b>Low confidence:</b> Low confidence in human influence on meteorological drought in the region (Otto et al., 2015b; Martins et al., 2018).	<b>Medium confidence: Increase</b> of CDD (Chapter 11 Supplementary Material (11.SM)(Chou et al., 2014a) and SPI (Xu et al. 2019, Touma et al. 2015). Increase in CDD for change of +0.5°C in global warming based on CMIP5 (Wartenburger et al., 2017)(SR15, Ch3)	<b>Medium confidence: Increase</b> in drought severity (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a; Spinoni et al., 2020) (Chapter 11 Supplementary Material (11.SM).	<b>Medium confidence: Increase</b> in drought severity (Chou et al., 2014a; Touma et al., 2015; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM).
	AGR ECOL	<b>Medium confidence: Increase</b> in drought severity based on different drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Lack of signal</b> based on different metrics, including total and surface column soil moisture, (Chapter 11 Supplementary Material (11.SM) (Xu et al., 2019a), and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase.</b> Dominant increase in drying with some inconsistencies between different drought metrics and models (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020) (Chapter 11 Supplementary Material (11.SM)	<b>Medium confidence: Increase</b> in drought severity with different metrics and high agreement between different studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited evidence.</b> One study shows an increase in drought severity (Gudmundsson et al., 2021)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows a weak drying (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Weak drying (Touma et al., 2015) or inconsistent trends (Cook et al., 2020)	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020).
South-western South America (SWS)	MET	<b>Medium confidence: Increase</b> in drought duration and severity (Skansi et al., 2013; Garreaud et al., 2017, 2020; Saurral et al., 2017; Boisier et al., 2018; Dereczynski et al., 2020; Dunn et al., 2020)	<b>Medium confidence</b> that human-induced climate change has contributed to long-term trends and Central Chile drought between 2010 and 2018 (Boisier et al., 2016; Garreaud et al., 2020)	<b>Low confidence: Inconsistent trends</b> Increase in meteorological drought based on CDD in CMIP6 GCMs (Chapter 11 Supplementary Material (11.SM), but inconsistent trends in SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a) and substantial model spread in Eta-RCM driven with two GCMs (Chou et al., 2014a).	<b>Low confidence: Mixed trends</b> between studies and models. Increase in meteorological drought based on CDD in CMIP6 GCMs (Chapter 11 supplementary Material (11.SM) , but inconsistent trends in SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a) and substantial model spread in Eta-RCM driven with two GCMs (Chou et al., 2014a).	<b>Medium confidence: Increase</b> in drought severity (Chou et al., 2014a; Touma et al., 2015; Spinoni et al., 2020).
	AGR ECOL	<b>Low confidence: Mixed trends</b> according to subregions and different drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed trends</b> based on different metrics, including decrease in total column and surface soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM) , weak drying in total and surface soil moisture in CMIP5 (Xu et al., 2019a), and weak signal based on	<b>Medium confidence: Increase</b> in drought severity based on total and surface soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM) and CMIP5 (Xu et al., 2019a), and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>High confidence: Increase</b> in drought severity with different metrics and high agreement between studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).

				the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).		
	HYDR	<b>Low confidence: Limited evidence.</b> General lack of signal in one study (Gudmundsson et al., 2021) but streamflow decrease in subregions in another study (Boisier et al. (2018))	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows drying ( Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Strong drying in (Cook et al., 2020); weak drying in (Touma et al., 2015)	<b>High confidence: Increase</b> in drought severity (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020).
South-eastern South America (SES)	MET	<b>Low confidence: Mixed signals</b> in observed trends depending on subregion (Saurral et al., 2017; Knutson and Zeng, 2018; Spinoni et al., 2019; Dereczynski et al., 2020; Dunn et al., 2020)	<b>Low confidence: Limited evidence.</b> Wetting trend in models and observations in part of region in one study (Knutson and Zeng, 2018).	<b>Low confidence: Inconsistent trends.</b> Weak drying trend based on CDDCMIP6 (Chapter 11 Supplementary Material (11.SM) , inconsistent trend between models based on SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a) and lack of signal in study with one RCM driven by two GCMs (Chou et al., 2014a).	<b>Low confidence: Mixed signals</b> between studies and models (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a; Spinoni et al., 2020) (Chapter 11 Supplementary Material (11.SM) .	<b>Low confidence: Mixed signals</b> between studies and models (Chou et al., 2014a; Touma et al., 2015; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM).
	AGR ECOL	<b>Low confidence: Mixed trends</b> according to subregions and different drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed trends</b> based on different metrics, including lack of signal in total column soil moisture, (Chapter 11 Supplementary Material (11.SM) , weak drying with surface soil moisture (Xu et al., 2019a) and wetting based on the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> in changes in drought severity with different metrics, (Chapter 11 Supplementary Material (11.SM), (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020).	<b>Low confidence: Mixed signals</b> Inconsistent trends or lack of signal in total and surface soil moisture(Chapter 11 Supplementary Material (11.SM) (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020); decreasing drought severity in PDSI and SPEI-PM (Cook et al., 2014b; Dai et al., 2018; Vicente-Serrano et al., 2020a).
	HYDR	<b>Medium confidence: Decrease.</b> Reduction of hydrological droughts (Dai and Zhao, 2017; Rivera and Penalba, 2018)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows mixed signal (Touma et al., 2015)	<b>Low confidence: Limited evidence.</b> Mixed signal (Touma et al., 2015) or wetting (Cook et al., 2020)	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020).
Southern South America (SSA)	MET	<b>Medium confidence: Increase</b> in the frequency of droughts (Skansi et al., 2013; Spinoni et al., 2019; Dereczynski et al., 2020; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Lack of signal</b> (Chapter 11 Supplementary Material (11.SM) (Chou et al., 2014a).	<b>Medium confidence:</b> Increase in drought severity (Chou et al., 2014a; Touma et al., 2015; Xu et al., 2019a; Spinoni et al., 2020) (Chapter 11 Supplementary Material (11.SM).	<b>Medium confidence:</b> Increase in drought severity (Chou et al., 2014a; Touma et al., 2015; Spinoni et al., 2020; Coppola et al., 2021b) (Chapter 11 Supplementary Material (11.SM)
	AGR ECOL	<b>Low confidence: Mixed trends</b> depending on subregions and drought metrics, including soil moisture, PDSI-PM and SPEI-PM (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Medium confidence: Increase</b> in drought severity considering total column soil moisture, (Chapter 11 Supplementary Material (11.SM) , and surface soil moisture (Xu et al., 2019a) and weak drying with the SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>High confidence: Increase</b> in drought severity (Naumann et al., 2018; Xu et al., 2019a; Gu et al., 2020) (Chapter 11 Supplementary Material (11.SM).	<b>High confidence: Increase</b> in drought severity with different metrics and high agreement between studies (Chapter 11 Supplementary Material (11.SM) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited</b>	<b>Low confidence:</b>	<b>Low confidence: Limited</b>	<b>Low confidence: Limited</b>	<b>High confidence:</b> Increase in

		evidence and lack of signal (Gudmundsson et al., 2021)	Limited evidence	evidence. One study shows drying (Touma et al., 2015)	evidence. Drying (Touma et al., 2015; Cook et al., 2020) or inconsistent trend (Zhai et al., 2020b).	drought severity (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
--	--	--	------------------	---	--	---

[END TABLE 11.15 HERE]

[START TABLE 11.16 HERE]

**Table 11.16:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in Europe, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Europe	All subregions show a <i>very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Hu et al., 2020; Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>very likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Greenland/Iceland (GIC)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes	Strong evidence of changes from observations that are in the direction of model projected changes for the future. The magnitude of	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events

1  
2  
3  
4  
5  
6  
7  
8



	(Peña-Angulo et al., 2020; Mermild et al., 2014; Sui et al., 2017; Dunn et al., 2020)	projected changes increases with global warming.	et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wehner et al., 2018; Cardell et al., 2020).	(Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wehner et al., 2018; Cardell et al., 2020).	(Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardell et al., 2020; Sillmann et al., 2013).
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes.	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Mediterranean (MED) <sup>5</sup>	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Peña-Angulo et al., 2020; El Kenawy et al., 2013; for Spain, Acero et al., 2014; Fioravanti et al., 2016; Ruml et al., 2017; Türkeş and Erlat, 2018; Donat et al., 2013,	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020); (Wang et al., 2017c); (Sippel and Otto, 2014); (Wilcox et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in

<sup>5</sup> This region includes both northern Africa and southern Europe  
Do Not Cite, Quote or Distribute

	2014, 2016; Filahi et al., 2016; Driouech et al., 2020; Dunn et al.2020)		compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Zollo et al., 2016); Weber et al., 2018)	annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Tomozeiu et al., 2014; Abaurrea et al., 2018; Nastos and Kapsomenakis, 2015; Cardell et al., 2020; Zollo et al., 2016; Weber et al., 2018; Coppola et al., 2021a)	annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardoso et al., 2019; Nastos and Kapsomenakis, 2015; Tomozeiu et al., 2014; Cardell et al., 2020; Zollo et al., 2016; Giorgi et al., 2014; Driouech et al., 2020; Coppola et al., 2021a; Engelbrecht et al., 2015)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Western and Central Europe (WCE)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Christidis et al., 2015; Scherrer et al., 2016; Shevchenko et al., 2014; Twardosz and Kossowska-Cezak, 2013; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Sippel et al., 2017, 2018; Dong et al., 2014, 2016; Sippel et al., 2016; Christidis et al., 2015; Cattiaux and Ribes, 2018; Leach et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 5.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 6°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from

			CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes (Lau and Nath, 2014; Lhotka et al., 2018)	CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes (Russo et al., 2015; Lau and Nath, 2014; Lhotka et al., 2018)	CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes (Lau and Nath, 2014; Lhotka et al., 2018)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Eastern Europe (EEU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Peña-Angulo et al., 2020; Zhang et al., 2019b; Donat et al., 2016; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Sippel and Otto, 2014; Leach et al., 2020; Hauser et al., 2016)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wehner et al., 2018; Cardell et al., 2020).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wehner et al., 2018; Cardell et al., 2020; Khlebnikova et al., 2019).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Cardell et al., 2020; Khlebnikova et al., 2019; Sillmann et al., 2013)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014))	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-

	intensity and frequency of cold extremes	and decrease in the intensity and frequency of cold extremes	<p><i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Likely</i> (compared with the recent past (1995-2014))  <i>Very likely</i> (compared with pre-industrial)</p>	<p><i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Very likely</i> (compared with the recent past (1995-2014))  <i>Extremely likely</i> (compared with pre-industrial)</p>	<p>2014))  <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Virtually certain</i> (compared with the recent past (1995-2014))  <i>Virtually certain</i> (compared with pre-industrial)</p>
Northern Europe (NEU)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Matthes et al., 2015; Vikhamar-Schuler et al., 2016; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020; Wang et al., 2017; Otto et al., 2012; Massey et al., 2012; Christiansen et al., 2018; King et al., 2015; Roth et al., 2018)	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Jacob et al., 2018; Laliberté et al., 2015; Sigmond et al., 2018; Dosio and Fischer, 2018; Forzieri et al., 2016)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Jacob et al., 2018; Laliberté et al., 2015; Sigmond et al., 2018; Dosio and Fischer, 2018; Forzieri et al., 2016)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Jacob et al., 2018; Laliberté et al., 2015; Sigmond et al., 2018; Dosio and Fischer, 2018; Forzieri et al., 2016)</p>
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<p>Increase in the intensity and frequency of hot extremes:  <i>Likely</i> (compared with the recent past (1995-2014))  <i>Very likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Likely</i> (compared with the recent past (1995-2014))</p>	<p>Increase in the intensity and frequency of hot extremes:  <i>Very likely</i> (compared with the recent past (1995-2014))  <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Very likely</i> (compared with the recent past (1995-2014))</p>	<p>Increase in the intensity and frequency of hot extremes:  <i>Virtually certain</i> (compared with the recent past (1995-2014))  <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes:  <i>Virtually certain</i> (compared with pre-industrial)</p>

			<i>Very likely</i> (compared with pre-industrial)	<i>Extremely likely</i> (compared with pre-industrial)	with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
--	--	--	---	--	--

1  
2 **[END TABLE 11.16 HERE]**

3  
4  
5 **[START TABLE 11.17 HERE]**

6  
7 **Table 11.17:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in Europe, subdivided  
8 by AR6 regions. See Sections 11.9.1 and 11.9.3 for details.

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All Europe	Significant intensification of heavy precipitation (Sun et al., 2020)	Robust evidence of a human contribution to the observed intensification of heavy precipitation (Paik et al., 2020)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 0% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)
	<i>Likely</i> intensification of heavy precipitation	Human influence <i>likely</i> contributed to the observed intensification of heavy precipitation	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
Greenland/Iceland (GIC)	Intensification of heavy precipitation (Peña-Angulo et al., 2020)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 30% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 30% in annual Rx1day and Rx5day and 35% in annual Rx30day compared to pre-industrial (Annex).

			Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020)	Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020)	Additional evidence from CMIP5 and CORDEX simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020)
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
Mediterranean (MED) <sup>6</sup>	Lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020; Casanueva et al., 2014; de Lima et al., 2015; Gajić-Čapka et al., 2015; Ribes et al., 2019; Peña-Angulo et al., 2020; Jacob et al., 2018; Rajczak and Schär, 2017; Coppola et al., 2021a; Donat et al., 2014; Mathbout et al., 2018)	Limited evidence (Añel et al., 2014; U.S. Department of Agriculture Economic Research Service, 2016)	CMIP6 models, CMIP5 models, and RCMs project inconsistent changes in the region (Li et al., 2020; Cardell et al., 2020; Zollo et al., 2016; Samuels et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 0% in annual Rx1day and Rx5day and less than -2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020; Zollo et al., 2016; Samuels et al., 2018)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 8% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 2% in annual Rx1day and Rx5day and less than -2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Cardell et al., 2020; Trambly and Somot, 2018; Zollo et al., 2016; Samuels et al., 2018; Monjo et al., 2016; Rajczak et al., 2013; Coppola et al., 2021b; Driouech et al., 2020)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014))	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014))	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014))

<sup>6</sup> This region includes both northern Africa and southern Europe  
Do Not Cite, Quote or Distribute

			<i>Medium confidence</i> (compared with pre-industrial)	<i>High confidence</i> (compared with pre-industrial)	<i>High confidence</i> (compared with pre-industrial)
Western and Central Europe (WCE)	Intensification of heavy precipitation (Sun et al., 2020; Casanueva et al., 2014; Croitoru et al., 2013; Fischer et al., 2015; Roth et al., 2014; Willems, 2013).	Disagreement among studies (Wilcox et al., 2018; Philip et al., 2018; Schaller et al., 2014, Vautard et al., 2015)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 0% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Rajczak and Schär, 2017; Donnelly et al., 2017)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Rajczak and Schär, 2017; Donnelly et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Rajczak and Schär, 2017; Madsen et al., 2014)
	<i>Medium confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
Eastern Europe (EEU)	Significant intensification of heavy precipitation (Sun et al., 2020; Dunn et al., 2020; Ashabokov et al., 2017)	Limited evidence	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and CORDEX simulations for an increase in	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5/CMIP3 and CORDEX simulations for an increase in

			the intensity of heavy precipitation (Cardell et al., 2020)	the intensity of heavy precipitation (Cardell et al., 2020)	the intensity of heavy precipitation (Cardell et al., 2020; Rajczak et al., 2013)
	<i>High confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
Northern Europe (NEU)	Significant intensification of heavy precipitation (Sun et al., 2020; Dunn et al., 2020)	Robust evidence of a human contribution to the observed intensification of heavy precipitation in winter (Schaller et al., 2016; Vautard et al., 2016; Otto et al., 2018b), but not in summer (Schaller et al., 2014; Otto et al., 2015c; Wilcox et al., 2018)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 0% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Donnelly et al., 2017)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Donnelly et al., 2017; Ramos et al., 2016; Romero and Emanuel, 2017)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Madsen et al., 2014; Ramos et al., 2016; Romero and Emanuel, 2017; Donnelly et al., 2017)
	<i>High confidence</i> in the intensification of heavy precipitation  <i>High confidence</i> in the changes in flood seasonality  <i>High confidence</i> in the increase in extreme snow-melt events	<i>High confidence</i> in a human contribution to the observed intensification of heavy precipitation in winter.	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)

1  
2  
3

[END TABLE 11.17 HERE]

Do Not Cite, Quote or Distribute

11-211

Total pages: 345



## [START TABLE 11.18 HERE]

**Table 11.18:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET), agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in Europe, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.4 for details.

Region and drought types		Observed trends	Detection and attribution; event attribution	Projections		
				+1.5 °C	+2 °C	+4 °C
Greenland/ Iceland (GIC)	MET	<b>Low confidence: Limited evidence, given limited number of studies and limited data</b> (Walsh et al., 2020; Dunn et al., 2020)	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence: Limited evidence</b> given limited number of studies (Walsh et al., 2020); tendency to decrease in meteorological drought based on CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Touma et al., 2015)	<b>Low confidence: Limited evidence</b> given limited number of studies (Walsh et al., 2020); tendency to decrease in meteorological drought based on CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Touma et al., 2015) ; also consistent with mixed index combining SPI and SPEI in Iceland (Spinoni et al., 2018b)  Based on (Spinoni et al., 2018b) in Iceland [11 EUROCORDEX RCPs 4.5 AND 8.5] Based on the Standardized Precipitation Index: Decrease of drought frequency.	<b>Low confidence: Limited evidence</b> given limited number of studies (Walsh et al., 2020) ; tendency to decrease in meteorological drought based on CDD (Chapter 11 Supplementary Material (11.SM)) and SPI (Touma et al., 2015); also consistent with mixed index combining SPI and SPEI in Iceland (Spinoni et al., 2018b)
	AGR ECOL	<b>Low confidence: Limited evidence, given limited number of studies and limited data</b> (Walsh et al., 2020).	<b>Low confidence: Limited evidence</b> because of lack of studies	<b>Low confidence: Limited evidence</b> because of lack of studies (Walsh et al., 2020) and <b>inconsistent</b> changes in soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Limited evidence</b> because of lack of studies (Walsh et al., 2020) and <b>inconsistent</b> changes in soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Limited evidence</b> because of lack of studies (Walsh et al., 2020) and <b>inconsistent</b> changes in soil moisture in CMIP6 (Chapter 11 Supplementary Material (11.SM))
	HYDR	<b>Low confidence: Limited evidence</b> given <b>limited number of studies and limited data</b> (Walsh et al., 2020)	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>

Mediterranean (MED) <sup>7</sup>	MET	<b>Low confidence: Mixed signals.</b> Observed land precipitation trends show pronounced variability within the region, with magnitude and sign of trend in the past century depending on time period (Donat et al., 2014a; Stagge et al., 2017; Zittis, 2017; Mathbout et al., 2018a). There is <i>low confidence</i> in an increase of drought frequency and severity based on SPI (Spinoni et al., 2015; Gudmundsson and Seneviratne, 2016; Peña-Angulo et al., 2020; Vicente-Serrano et al., 2021; MedECC, 2020; Driouech et al., 2021)	<b>Low confidence: Mixed signals.</b> There are mixed signals within the region and <i>low confidence</i> in human influence on meteorological drought over MED (Kelley et al., 2015; Gudmundsson and Seneviratne, 2016; Knutson and Zeng, 2018; Wilcox et al., 2018)	<b>Medium confidence: Increase.</b> With <i>medium confidence</i> both CMIP5 and CMIP6 show a decline in winter and summer total precipitation and increase in number of CDD (percentage precipitation change per degree of local warming is <i>with high confidence</i> larger in JJA than DJF) (Interactive Atlas, Cardell et al., 2020; Li et al., 2020)(Chapter 11 Supplementary Material (11.SM)). Also weak increase in meteorological drought based on SPI (Touma et al., 2015; Xu et al., 2019a).	<b>Medium confidence: Increase.</b> With <i>medium confidence</i> both CMIP5 and CMIP6 show a decline in winter and summer total precipitation and increase in number of CDD (percentage precipitation change per degree of local warming is <i>with high confidence</i> larger in JJA than DJF) (Interactive Atlas, Cardell et al., 2020; Li et al., 2020)(Chapter 11 Supplementary Material (11.SM)). Also weak increase in meteorological drought based on SPI (Touma et al., 2015; Xu et al., 2019a).	<b>High confidence: Increase.</b> With <i>high confidence</i> both CMIP5 and CMIP6 (and EURO-CORDEX) show a decline in winter and summer total precipitation and increase in number of CDD. Drought intensity and frequency increase with <i>high confidence</i> , particularly in the southern Mediterranean (Samuels et al., 2018; Cardell et al., 2020; Cook et al., 2020; Li et al., 2020a; Spinoni et al., 2020; Coppola et al., 2021a)( Chapter 11 Supplementary Material (11.SM); Interactive Atlas) (Driouech et al., 2020)
	AGR ECOL	<b>Medium confidence: Increase.</b> Increases in probability and intensity of agricultural and ecological droughts based on soil moisture and water-balance deficits, but weakens signals in some studies (Greve et al., 2014; Hanel et al., 2018; García-Herrera et al., 2019; Moravec et al., 2019; Padrón et al., 2020; Markonis et al., 2021). Also increases based on analyses using the Standardized Precipitation Evapotranspiration Index (SPEI) and the Palmer Drought Severity Index (PDSI). Increase of drought severity in South Europe (Stagge et al., 2017; Spinoni et al., 2019; Dai and Zhao, 2017), the Iberian Peninsula (Vicente-Serrano et al., 2014; González-Hidalgo et al., 2018).  (Markonis et al., 2021): Increase in duration of agricultural droughts based on soil moisture deficits from 1901-2015.	<b>Medium confidence: of attribution of increasing trend in ecological and agricultural drought,</b> based on soil moisture and water-balance metrics (Mariotti et al., 2015; García-Herrera et al., 2019; Marvel et al., 2019; Padrón et al., 2020)  García-Herrera et al. (2019): Attribution of the 2016/2017 drought in southwestern Europe to climate change based on NCEP trends in soil moisture for weather analogues to 2016/2017 event.  Mariotti et al. (2015): Attributable trend to CC: Decrease in soil moisture in summer that agrees with CMIP5 models.	<b>Medium confidence: Drought increase for pre-industrial and recent past baselines.</b>  <i>Recent past baseline:</i>  Decreasing soil water availability during drought events compared to 1971-2000, even when accounting for adaptation to mean conditions (Samaniego et al., 2018).  Increasing drought duration and frequency compared to 1971-2000 (Xu et al., 2019a)  Increasing drought magnitude based on SPEI-PM compared to +0.6°C baseline, using simulations within single ESM driven with sea surface temperature and sea ice conditions of 7 ESMS (Naumann et al., 2018)	<b>High confidence: Drought increase for pre-industrial and recent past baselines.</b>  <i>Recent past baseline:</i>  Decreasing soil water availability during drought events compared to 1971-2000, even when accounting for adaptation to mean conditions; about twice larger signal compared to response at +1.5°C (Samaniego et al., 2018).  Increasing drought duration and frequency compared to 1971-2000, with about twice larger signal compared to response at +1.5°C (Xu et al., 2019a)  Increasing drought magnitude based on SPEI-PM compared to +0.6°C baseline, using simulations within single ESM driven with sea surface temperature and sea ice conditions of 7 ESMS (Naumann et al., 2018)	<b>Very likely: Drought increase for pre-industrial and recent past baselines.</b>  <i>Recent past baseline:</i>  Based on projections at +3°C: Large decreasing soil water availability during drought events compared to 1971-2000, even when accounting for adaptation to mean conditions; more than three times larger signal compared to response at +1.5°C (Samaniego et al., 2018).  Based on projections at +3°C: About five-fold increase in drought magnitude based on SPEI-PM compared to +0.6°C baseline, using simulations within single ESM driven with sea surface temperature and sea ice conditions of 7 ESMS (Naumann et al., 2018)  <i>Pre-industrial baseline:</i>

<sup>7</sup> This region includes both northern Africa and southern Europe

		<p>(García-Herrera et al., 2019): Increase in soil moisture anomalies for weather analogues to 2016/2017 drought events in 1985-2018 vs 1948-1984.</p> <p>(Padrón et al., 2020): Weak signals in water-balance (precipitation-evapotranspiration) deficits in the dry season (1985-2014)-(1902-1950)</p> <p>(Greve et al., 2014): Increase in water-balance (precipitation-evapotranspiration) deficits on annual scale, (1985-2005) - (1948-1968)</p> <p>(Hanel et al., 2018): Significant decrease in soil moisture in Southern Europe from 1766-2015 from hydrological model driven with reconstructed meteorological data.</p>	<p>However: no emergence yet in soil moisture or P-E at grid cell scale (see CC-Box A.1. on Uncertainty).</p> <p>Padrón et al. (2020): Increasing drying trend in P-E during dry season over land areas, including in Mediterranean region (but attribution done at global scale, not regional scale)</p> <p>Marvel et al. (2019): Attributable drying trend in larger continental region with tree-ring data including strong signal in Mediterranean from 1900-1950 and currently increasing again after masking from aerosols.</p>	<p><i>Pre-industrial baseline:</i></p> <p>Decrease in soil moisture during drought events in CMIP6 models at +1.5°C vs pre-industrial baseline (Chapter 11 Supplementary Material (11.SM))</p>	<p><i>Pre-industrial baseline:</i></p> <p>Decreases of surface and total soil moisture, in both AMJJAS and ONDJFM half years (Cook et al., 2020)</p> <p>Decrease in soil moisture during drought events in CMIP6 models at +2°C vs pre-industrial baseline (Chapter 11 Supplementary Material (11.SM))</p>	<p>Strong decreases of surface and total soil moisture, in both spring-summer (AMJJAS) and fall-winter (ONDJFM) half years, with about twice larger response compared to +2°C (Cook et al., 2020)</p> <p>Very large decrease in soil moisture during drought events in CMIP6 models at +4°C vs pre-industrial baseline (Chapter 11 Supplementary Material (11.SM))</p>
<p>HYDR</p>		<p><b>High confidence:</b> Increase in frequency and severity of hydrological droughts, particularly in northern part of the domain (Lorenzo-Lacruz et al., 2013; Dai and Zhao, 2017; Gudmundsson et al., 2017, 2019, 2021) (Section 8.3.1.6).</p>	<p><b>Medium confidence:</b> Increase. Model-based assessment shows with <i>medium confidence</i> a human fingerprint on increased hydrological drought, related to rising temperature and atmospheric demand (Gudmundsson et al., 2017, 2021) and recent events. There is <i>medium confidence</i> that change in land use and terrestrial water management contribute to trends in hydrological drought (Teuling et al., 2019; Vicente-Serrano et al., 2019)</p>	<p><b>Medium confidence:</b> Increase in hydrological drought for both pre-industrial and recent past baseline</p> <p><i>Recent past baseline:</i></p> <p>Forzieri et al. (2014): 20 yr deficit volumes are projected to increase by 50% by the 2020s compared to 1961-1990 (based on simulations with LISFLOOD model driven by 12 RCM simulations with different GCM-RCM pairs; CMIP3 GCMs, A1B scenario). Frequency of hydrological droughts is projected to increase (Touma et al., 2015).</p>	<p><b>High confidence:</b> Increase.</p> <p><i>Recent past baseline:</i></p> <p>Forzieri et al. (2014) [LISFLOOD simulations driven by 12 RCM-GCM pairs using CMIP3 GCMs]: Strong increase in the 20-yr return level minimum flow and deficit volumes in 2050 in A1B scenario compared to 1961-1990.</p> <p>Roudier et al. (2016) [11 RCMs]: Increase in the severity of the low flows at +2°C compared to 1971-2000 conditions in the Iberian Peninsula, Southern France and Greece.</p> <p>Schewe et al. (2014). Decrease between 30-50% of the annual runoff compared to 1980-2010.</p>	<p><b>Very likely:</b> Increase</p> <p><i>Recent past baseline:</i></p> <p>Forzieri et al. (2014) [LISFLOOD simulations driven by 12 RCM-GCM pairs using CMIP5 GCMs]: Strong increase in the 20-yr return level minimum flow and deficit volumes in 2080 in A1B scenario compared to 1961-1990</p> <p>Prudhomme et al. (2014) [5 CMIP5 models driving 7 global impact models. RCP8.5, 2070-2099] Strong increase (40-60%) of dry days compared to 1976-2005</p> <p>Giuntoli et al. (2015) (5 CMIP5 models driving 6 global hydrology models): 50-60% increase in frequency of days under low flow in 2066-2099 compared to 1972-2005. Strong signal to noise</p>

					<p>Touma et al. (2015): Increase in the frequency of hydrological droughts relative to 1961-2005.</p> <p><i>Pre-industrial baseline</i></p> <p>Cook et al. (2020) [13 CMIP6 models and SSP1-2.6] Decrease in surface and total runoff, in both spring-summer (AMJJAS) and fall-winter (ONDJFM) half years, with strongest decreases for total runoff in spring-summer half -year-</p>	<p>ratio in terms of model agreement, strongest hot spot globally.</p> <p><i>Pre-industrial baseline</i></p> <p>Cook et al. (2020) [13 CMIP6 models and SSP3-7.0. Very strong decrease (40-60%) of total runoff in spring-summer half-year in southern Europe. Also strong decreases (&gt;20%) for total runoff in fall-winter half-year, and for surface runoff in both half years (AMJJAS, ONDJFM).</p>
Western and Central Europe (WCE)	MET	<p><b>Low confidence:</b> Limited evidence in change in severity. <b>Small and non-significant</b> changes and some dependency on season and location. Small and non significant changes in the frequency of dry spells (Zolina et al., 2013), CDD (Dunn et al., 2020), and in drought severity (SPI) (Orlowsky and Seneviratne, 2013; Stagge et al., 2017; Caloiero et al., 2018; Spinoni et al., 2019); but wet days decrease in summer (Gobiet et al., 2014).</p>	<p><b>Low confidence: No signal or varying signal</b> depending on considered index (Gudmundsson and Seneviratne, 2016; Hauser et al., 2017)</p>	<p><b>Low confidence: Inconsistent</b> signal in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)) and in SPI in CMIP5 (Orlowsky and Seneviratne, 2013; Touma et al., 2015; Xu et al., 2019a).</p>	<p><b>Low confidence: Inconsistent</b> signal, but with weak tendency to drying in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)) and SPI in CMIP5 (Orlowsky and Seneviratne, 2013; Touma et al., 2015; Xu et al., 2019a)</p>	<p><b>Medium confidence: Increase</b> based on CDD (Chapter 11 Supplementary Material (11.SM)). Also partial drying based on CMIP5 SPI, but strong geographical gradients and trends in part not significant (Orlowsky and Seneviratne, 2013; Touma et al., 2015; Vicente-Serrano et al., 2020a). Summer decrease in wet day projected in Switzerland (Fischer et al., 2015).</p>
	AGR ECOL	<p><b>Medium confidence: Increase.</b> Dominant signal shows an increase in available studies based on soil moisture models and SPEI-PM (Greve et al., 2014; Trnka et al., 2015b; Hanel et al., 2018; Moravec et al., 2019; Spinoni et al., 2019; Padrón et al., 2020; Markonis et al., 2021), despite some conflicting trends in some subregions (Spinoni et al., 2019; Padrón et al., 2020).</p>	<p><b>Low confidence: Limited evidence</b> due to limited number of studies; one study suggests attribution of the 2017 drought event to climate change due to decreasing trends in soil moisture (García-Herrera et al. 2019)</p>	<p><b>Low confidence: Inconsistent</b> signal in CMIP6 (Chapter 11 Supplementary Material (11.SM)) or <b>weak</b> (Xu et al., 2019a) or insignificant signal (Samaniego et al., 2018), mostly in summer season. A bit stronger signal based on SPEI-PM projections (Naumann et al., 2018)</p>	<p><b>Medium confidence: Increase</b> of drought frequency and severity based on some AGR and ECOL drought metrics, for surface soil moisture and SPEI-PM (Chapter 11 Supplementary Material (11.SM))(Naumann et al., 2018; Samaniego et al., 2018; Xu et al., 2019a), mostly for summer season, but inconsistent trends for CMIP6 total soil moisture (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020)</p>	<p><b>Medium confidence: Increase</b> of drought frequency and severity based on some AGR and ECOL drought metrics, for CMIP6 surface soil moisture, root-zone soil moisture in hydrological models, and SPEI-PM (Chapter 11 Supplementary Material (11.SM))(Naumann et al., 2018; Samaniego et al., 2018; Xu et al., 2019a; Cook et al., 2020), mostly in summer season, but inconsistent trends for CMIP6 total soil moisture (Chapter 11 Supplementary Material (11.SM)) despite projected drying in substantial fraction of domain, in particular over France (Cook et al., 2020)</p>
	HYDR	<p><b>Low confidence: Weak or insignificant</b> trends (Stahl et al., 2010; Bard et al., 2015; Caillouet et al., 2017; Moravec et al., 2019; Vicente-Serrano et al., 2019;</p>	<p><b>Low confidence: Limited evidence</b> because of <b>lack of studies.</b></p>	<p><b>Low confidence: No or weak changes;</b> CORDEX simulations: no change in most of domain, slight wetting over the Alps (Forzieri et al., 2014; Touma et al., 2015; Marx et al.,</p>	<p><b>Medium confidence: Increase</b> in drying, mostly in western part of domain: summer season surface runoff compared to pre-industrial (Cook et al., 2020); annual discharge in substantial part of domain (Schewe et al.,</p>	<p><b>Medium confidence: Increase</b> based on several lines of evidence: Tendency towards drying but geographical variations (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)</p>

		Gudmundsson et al., 2021)		2018)	2014); increase in duration and magnitude of low flows over France, decrease in eastern part of domain (Touma et al., 2015; Roudier et al., 2016); CORDEX simulations. drying in western and southeastern parts of domain, but wetting over the Alps (Forzieri et al., 2014; Marx et al., 2018)	
Eastern Europe (EEU)	MET	<b>Low confidence: Inconsistent or insignificant changes.</b> Inconsistent or insignificant changes in CDD (Khlebnikova et al., 2019b; Dunn et al., 2020). No change or insignificant changes in SPI (Stagge et al., 2017; Caloiero et al., 2018; Spinoni et al., 2019)	<b>Low confidence: Limited evidence</b> because of <b>lack of studies.</b>	<b>Low confidence: Inconsistent changes.</b> Inconsistent changes in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)) and inSPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a).	<b>Low confidence: Inconsistent changes.</b> Inconsistent changes in CDD in CMIP6 (Chapter 11 Supplementary Material (11.SM)) and in SPI in CMIP5 (Touma et al., 2015; Xu et al., 2019a)	<b>Low confidence: Inconsistent changes.</b> Inconsistent CDD changes in CMIP6 (Chapter 11 Supplementary Material (11.SM)) and weak decrease in drying or inconsistent changes in SPI projections (Touma et al., 2015; Spinoni et al., 2020; Vicente-Serrano et al., 2020a)
	AGR ECOL	<b>Low confidence: Inconsistent or weak changes</b> (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence</b> based on different metrics: <b>Inconsistent</b> trends in both CMIP6 surface and total soil moisture (Chapter 11 Supplementary Material (11.SM)); weak trends in CMIP5 soil moisture (Xu et al., 2019a) or SPEI-PM (Naumann et al., 2018) projections	<b>Low confidence</b> based on different metrics : <b>Inconsistent</b> trends in both CMIP6 surface and total soil moisture (Chapter 11 Supplementary Material (11.SM)); weak trends in CMIP5 soil moisture (Xu et al., 2019a) or SPEI-PM (Naumann et al., 2018) projections	<b>Low confidence: Inconsistent</b> trends based on different metrics: Slight wetting or inconsistent trends in total soil moisture (Chapter 11 Supplementary Material (11.SM)); (Cook et al., 2020); slight drying in surface soil moisture (Chapter 11 Supplementary Material (11.SM)); (Cook et al., 2020). Increasing drying of measures based on evaporative demand (Naumann et al., 2018)
	HYDR	<b>Low confidence: No enough data and limited studies</b> (Gudmundsson et al., 2021)	<b>Low confidence: Limited evidence</b> because of <b>lack of studies</b>	<b>Low confidence: Limited evidence. One study shows lack of signal</b> (Touma et al., 2015)	<b>Low confidence: Inconsistent</b> changes. Some studies with increases in drought/decrease in runoff: (Forzieri et al., 2014) [11 RCMs forced with CMIP5 models and the LISFLOOD model] : Decrease in the 20 yr return level minimum flow and deficit volumes; (Cook et al., 2020); decrease in summer surface runoff in CMIP6 models. Some studies with no change in HYDR drought or runoff: (Touma et al., 2015; Roudier et al., 2016) [11 RCMs]: No substantial changes in the severity of the low flows; (Schewe et al., 2014): No substantial changes in the annual runoff.	<b>Medium confidence: Weak increase.</b> (Forzieri et al., 2014) [11 RCMs forced with CMIP5 models and the LISFLOOD model] : Decrease in the 20 yr return level minimum flow and deficit. (Cook et al., 2020) [13 CMIP6 models and SSP3-7.0. Moderate decrease (20%) of total runoff in eastern Europe during the warm season.. (Prudhomme et al., 2014) [5 CMIP5 models and 7 global impact models. RCP8.5] Small increase (10%) of dry days. (Giuntoli et al., 2015): Weak increase in probability of low flow but low signal to noise ratio.

Northern Europe (NEU)	MET	<i>Medium confidence:</i> Decrease in intensity and frequency; but dependence on considered index, time frame and region, including negligible trends over shorter periods or some subregions (Orlowsky and Seneviratne, 2013; Stagge et al., 2017; Spinoni et al., 2019; Dunn et al., 2020)	<i>Medium confidence:</i> <b>Human contribution to decrease</b> (Gudmundsson and Seneviratne, 2016).	<i>Medium confidence:</i> Decrease of drought frequency and severity based on SPI indices (Touma et al., 2015; Xu et al., 2019a), but unclear sign in CDD (Chapter 11 Supplementary Material (11.SM)).	<i>Medium confidence:</i> Decrease of drought frequency and severity based on SPI indices (Touma et al., 2015; Xu et al., 2019a), but unclear sign in CDD (Chapter 11 Supplementary Material (11.SM)).	<i>Medium confidence:</i> Decrease of drought frequency and severity based on SPI indices (Touma et al., 2015; Spinoni et al., 2020; Vicente-Serrano et al., 2020a) but unclear sign and drying tendency in CDD (Chapter 11 Supplementary Material (11.SM)). Same assessment for pre-industrial and recent past baselines.
	AGR ECOL	<i>Low confidence:</i> Overall <b>weak</b> signals and signs depend on considered season and index (Greve et al., 2014; Spinoni et al., 2019; Padrón et al., 2020; Markonis et al., 2021)	<i>Low confidence:</i> <b>Limited evidence</b> because of <b>lack of studies</b>	<i>Low confidence:</i> <b>Inconsistent</b> signal in CMIP6 total soil moisture at +1.5°C compared to pre-industrial baseline (Chapter 11 Supplementary Material (11.SM)). Overall inconsistency of signals between studies for different indices (e.g. total soil moisture, surface soil moisture, SPEI-PM) independently of global warming level (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020), but some spatial variations in trends and stronger signals in summer (Samaniego et al., 2018). Same assessment for pre-industrial and recent past baseline.	<i>Low confidence:</i> <b>Inconsistent</b> signal in CMIP6 total soil moisture at +2°C compared to pre-industrial baseline (Chapter 11 Supplementary Material (11.SM)). Overall inconsistency of signals between studies for different indices (e.g. total soil moisture, surface soil moisture, SPEI-PM) independently of global warming level (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020); but some spatial variations in trends and stronger signals in summer and over Scandinavia compared to UK (Samaniego et al., 2018). Same assessment for pre-industrial and recent past baseline.	<i>Low confidence:</i> <b>Inconsistent</b> signal in CMIP6 total soil moisture at +4°C compared to pre-industrial baseline (Chapter 11 Supplementary Material (11.SM)). Overall inconsistency of signals between studies for different indices (e.g. total soil moisture, surface soil moisture, SPEI-PM) independently of global warming level (Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Vicente-Serrano et al., 2020a), but some spatial variations in trends and stronger signals in summer and over Scandinavia compared to UK (Samaniego et al., 2018). Same assessment for pre-industrial and recent past baseline.
	HYDR	<i>Medium confidence:</i> Decrease in hydrological drought for overall region, but trends are weak, can be of different sign in sub-regions, and are dependent on time frame (Harrigan et al., 2018; Kay et al., 2018; Barker et al., 2019; Gudmundsson et al., 2019, 2021; Vicente-Serrano et al., 2019)	<i>Low confidence:</i> <b>Limited evidence</b> because of <b>lack of studies</b>	<i>Low confidence:</i> <b>Weak and inconsistent signals.</b> Slight increase in Scandinavia, slight decrease or no change in the UK (Forzieri et al., 2014; Touma et al., 2015; Marx et al., 2018)	<i>Low confidence:</i> <b>Inconsistent</b> changes, generally with drying in ESMs (CMIP5, CMIP6) and wetting in CORDEX (Forzieri et al., 2014; Touma et al., 2015; Roudier et al., 2016; Dai et al., 2018; Marx et al., 2018; Cook et al., 2020).  Cook et al. (2020): Weak increase in hydrological drought (decrease in runoff) in summer in Scandinavia.	<i>Medium confidence:</i> Weak <b>increase</b> in hydrological drought in summer but low signal-to-noise ratio (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)

					<p>Roudier et al. (2016): Decrease in magnitude and duration of low-flows (wetting trend)</p> <p>Dai et al. (2018): CMIP5, RCP4.5, (2070-2099)-(1970-1999): slight drying trend, but lack of model agreement.</p> <p>Forzieri et al. (2014), for (2050 compared to 1961-1990 baseline), CORDEX simulations: Decrease in magnitude of low-flow in Scandinavia no change in UK,</p> <p>Marx et al. (2018), CORDEX simulations: slight wetting in Scandinavia</p>	
--	--	--	--	--	--	--

1  
2  
3  
4  
5  
6

[START TABLE 11.19 HERE]

**Table 11.19:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for temperature extremes in North America, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.2 for details

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All North America	Most subregions show a <i>likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Seong et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Human influence <i>very likely</i> contributed to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<p>Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Extremely likely</i> (compared with the recent past (1995-</p>	<p>Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p> <p>Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-</p>

			with pre-industrial)	2014)) <i>Virtually certain</i> (compared with pre-industrial)	2014)) <i>Virtually certain</i> (compared with pre-industrial)
North Central America (NCA)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (García-Cueto et al., 2019; Martínez-Austria and Bandala, 2017; Montero-Martínez et al., 2018; Dunn et al., 2020)	Strong evidence of changes from observations that are in the direction of model projected changes for the future. The magnitude of projected changes increases with global warming.	CMIP6 models project an increase in the intensity and frequency of TXx events and a decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Kharin et al., 2013; Sillmann et al., 2013b; Alexandru, 2018; Wehner et al., 2018b)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Kharin et al., 2013; Sillmann et al., 2013b; Alexandru, 2018; Wehner et al., 2018b)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 3.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 4.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Kharin et al., 2013; Sillmann et al., 2013b; Alexandru, 2018; Wehner et al., 2018b)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
W. North America (WNA)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and	Evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a decrease in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a decrease in	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust



	frequency of cold extremes (Vose et al., 2017; Dunn et al., 2020)	and decrease in the intensity and frequency of cold extremes (Seager et al., 2015; Angélil et al., 2017)	the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Palipane and Grotjahn, 2018; Wehner et al., 2018b)	the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Palipane and Grotjahn, 2018; Wehner et al., 2018b)	decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Palipane and Grotjahn, 2018; Wehner et al., 2018b)
	<i>Likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>Medium confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
C. North America (CNA)	Weak and inconsistent trends (Dunn et al., 2020)	Evidence of a human contribution for some events but cannot be generalized	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 3°C in annual TXx and TNn compared to pre-	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn

			industrial (Annex). Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Wehner et al., 2018b)	industrial (Annex). Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Wehner et al., 2018b)	compared to pre-industrial (Annex). Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al., 2017; Wehner et al., 2018b)
	<i>Low confidence</i>	<i>Low confidence</i>	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
E. North America (ENA)	Weak and inconsistent trends (Dunn et al., 2020)	Evidence of a human contribution for some events, but cannot be generalized	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al.,	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al.,	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Vose et al.,

			2017; Wehner et al., 2018b; Zhang et al., 2019d).	2017; Wehner et al., 2018b; Zhang et al., 2019d).	2017; Wehner et al., 2018b; Zhang et al., 2019d).
	<i>Low confidence</i>	<i>Low confidence</i>	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
N. E. North America (NEN)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Vincent et al., 2018; Zhang et al., 2019c; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wan et al., 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Li et al., 2018d; Zhang et al., 2019d)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Li et al., 2018d; Zhang et al., 2019d)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Li et al., 2018d; Zhang et al., 2019d)
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

			frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
N. W. North America (NWN)	Significant increases in the intensity and frequency of hot extremes and significant decreases in the intensity and frequency of cold extremes (Vincent et al., 2018; Zhang et al., 2019c; Dunn et al., 2020)	Robust evidence of a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Wan et al., 2019)	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 0.5°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 1.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Bennett and Walsh, 2015; Li et al., 2018d; Zhang et al., 2019d).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 1°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 2.5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Bennett and Walsh, 2015; Li et al., 2018d; Zhang et al., 2019d).	CMIP6 models project a robust increase in the intensity and frequency of TXx events and a robust decrease in the intensity and frequency of TNn events (Li et al., 2020; Annex). Median increase of more than 4°C in the 50-year TXx and TNn events compared to the 1°C warming level (Li et al., 2020) and more than 5°C in annual TXx and TNn compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes (Bennett and Walsh, 2015; Li et al., 2018d; Zhang et al., 2019d).
	<i>Very likely</i> increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	<i>High confidence</i> in a human contribution to the observed increase in the intensity and frequency of hot extremes and decrease in the intensity and frequency of cold extremes	Increase in the intensity and frequency of hot extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Increase in the intensity and frequency of hot extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)  Decrease in the intensity and frequency of cold extremes: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)

1 **[END TABLE 11.19 HERE]**2  
3  
4 **[START TABLE 11.20 HERE]**5  
6 **Table 11.20:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for heavy precipitation in North America,  
7 subdivided by AR6 regions. See Sections 11.9.1 and 11.9.3 for details

Region	Observed trends	Detection and attribution; event attribution	Projections		
			1.5 °C	2 °C	4 °C
All North America	Significant intensification of heavy precipitation (Sun et al., 2020; Dunn et al., 2020)	Robust evidence of a human contribution to the observed intensification of heavy precipitation (Kirchmeier-Young and Zhang, 2020; Paik et al., 2020)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020a). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a)
	<i>Likely</i> intensification of heavy precipitation	Human influence <i>likely</i> contributed to the observed intensification of heavy precipitation	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Virtually certain</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)
North Central America (NCA)	Trends are generally not significant (Sun et al., 2020; Dunn et al., 2020; Donat et al., 2016; García-Cueto et al., 2019)	Disagreement among studies (Eden et al., 2016; Pall et al., 2017; Hoerling et al., 2014)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 2% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 0% in annual Rx30day compared to pre-industrial (Annex).	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)

			with pre-industrial)	industrial)	
W. North America (WNA)	Lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020; Easterling et al. 2017; Wu 2015)	Evidence of a human contribution for some events (Easterling et al., 2017; Kirchmeier-Young and Zhang, 2020), but cannot be generalized	CMIP6 models project inconsistent changes in the region (Li et al., 2020a)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Easterling et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Easterling et al., 2017)
	<i>Low confidence</i>	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Low confidence</i> (compared with the recent past (1995-2014)) <i>Medium confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)
C. North America (CNA)	Significant intensification of heavy precipitation (Dunn et al., 2020; Easterling et al., 2017; Wu, 2015; Emanuel, 2017; Risser and Wehner, 2017; Trenberth et al., 2018; van Oldenborgh et al., 2017; Wang et al., 2018).	Evidence of a human contribution to the observed intensification of heavy precipitation (Easterling et al., 2017; Kirchmeier-Young and Zhang, 2020; Emanuel, 2017; Risser and Wehner, 2017; Trenberth et al., 2018; van Oldenborgh et al., 2017; Wang et al., 2018)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 4% in annual Rx1day and Rx5day and 2% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 10% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity

			(Easterling et al., 2017)	(Easterling et al., 2017)	of heavy precipitation (Easterling et al., 2017; Knutson et al., 2015; Kossin et al., 2017)
	<i>High confidence</i> in the intensification of heavy precipitation	<i>Medium confidence</i> in a human contribution to the intensification of heavy precipitation.	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
E. North America (ENA)	Significant intensification of heavy precipitation (Sun et al., 2020; Dunn et al., 2020; Easterling et al., 2017; Wu, 2015; Emanuel, 2017; Risser and Wehner, 2017; Trenberth et al., 2018; van Oldenborgh et al., 2017; Wang et al., 2018), but a lack of a significant trend over Canada (Shephard et al., 2014; Mekis et al., 2015; Vincent et al., 2018)	Evidence of a human contribution for some events (Easterling et al., 2017; Teufel et al., 2019; Kirchmeier-Young and Zhang, 2020), but cannot be generalized	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1 day and Rx5day and 4% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019; Easterling et al., 2017)	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 4% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1 day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019; Easterling et al., 2017)	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 15% in the 50-year Rx1 day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 15% in annual Rx1 day and Rx5day and 10% in annual Rx30day compared to pre-industrial (Annex).  Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019; Easterling et al., 2017; Knutson et al., 2015; Kossin et al., 2017)
	<i>High confidence</i> in the intensification of heavy precipitation	<i>Low confidence</i>	Intensification of heavy precipitation: <i>Medium confidence</i> (compared with the recent past (1995-2014)) <i>High confidence</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)	Intensification of heavy precipitation: <i>Very likely</i> (compared with the recent past (1995-2014)) <i>Extremely likely</i> (compared with pre-industrial)
N. E. North America (NEN)	Limited evidence (Shephard et al., 2014; Mekis et al., 2015; Vincent et al., 2018)	Evidence of a human contribution for some events (Szeto et al., 2015), but cannot be generalized	CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in	CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in

			<p>Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 8% in annual Rx1day and Rx5day and 6% in annual Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019d)</p>	<p>the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day and Rx5day and 8% in annual Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019d)</p>	<p>the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day and Rx5day and 15% in annual Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Zhang et al., 2019d)</p>
	<i>Low confidence</i>	<i>Low confidence</i>	<p>Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014)) <i>Likely</i> (compared with pre-industrial)</p>	<p>Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with pre-industrial)</p>	<p>Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014)) <i>Virtually certain</i> (compared with pre-industrial)</p>
N. W. North America (NWN)	Lack of agreement on the evidence of trends (Sun et al., 2020; Dunn et al., 2020; Mekis et al., 2015; Shephard et al., 2014; Vincent et al., 2018)	Evidence of a human contribution for some events (Teufel et al., 2017; Kirchmeier-Young and Zhang, 2020), but cannot be generalized	<p>CMIP6 models project an increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 2% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 6% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Bennett and Walsh, 2015; Zhang et al., 2019d)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 6% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 10% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Bennett and Walsh, 2015; Zhang et al., 2019d)</p>	<p>CMIP6 models project a robust increase in the intensity and frequency of heavy precipitation (Li et al., 2020; Annex). Median increase of more than 20% in the 50-year Rx1day and Rx5day events compared to the 1°C warming level (Li et al., 2020a) and more than 20% in annual Rx1day, Rx5day, and Rx30day compared to pre-industrial (Annex).</p> <p>Additional evidence from CMIP5 and RCM simulations for an increase in the intensity of heavy precipitation (Bennett and Walsh, 2015; Zhang et al., 2019d)</p>
	<i>Low confidence</i>	<i>Low confidence</i>	<p>Intensification of heavy precipitation: <i>High confidence</i> (compared with the recent past (1995-2014))</p>	<p>Intensification of heavy precipitation: <i>Likely</i> (compared with the recent past (1995-2014)) <i>Very likely</i> (compared with</p>	<p>Intensification of heavy precipitation: <i>Extremely likely</i> (compared with the recent past (1995-2014))</p>



			<i>Likely</i> (compared with pre-industrial)	pre-industrial)	<i>Virtually certain</i> (compared with pre-industrial)
--	--	--	--	-----------------	---

[END TABLE 11.20 HERE]

[START TABLE 11.21 HERE]

**Table 11.21:** Observed trends, human contribution to observed trends, and projected changes at 1.5°C, 2°C and 4°C of global warming for meteorological droughts (MET), agricultural and ecological droughts (AGR/ECOL), and hydrological droughts (HYDR) in North America, subdivided by AR6 regions. See Sections 11.9.1 and 11.9.4 for details.

Region and drought types		Observed trends	Human contribution	Projections		
				+1.5 °C	+2 °C	+4 °C
North Central America (NCA)	MET	<b>Low confidence: Inconsistent</b> changes in the duration and frequency of droughts, (Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: No signal</b> in precipitation (Funk et al., 2014; Swain et al., 2014; Wang and Schubert, 2014)	<b>Low confidence: Limited evidence.</b> Evidence suggests tendency towards drying (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Increase</b> in drought duration(Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).  Xu et al. (2019): Strong drying signal for meteorological drought duration using SPI at 2°C compared to recent past.  Spinoni et al. (2020): for RCP4.5 compared to recent past: SPI-based drying trends in CORDEX GCMs, but inconsistent signals in CORDEX RCMs.	<b>High confidence: Increase</b> in meteorological drought severity in the majority of models (Sillmann et al., 2013b; Touma et al., 2015; Escalante-Sandoval and Nuñez-Garcia, 2017; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Low evidence:</b> No signal in the duration and severity of droughts based on soil moisture, PDSI and SPEI and conflicting trend depending of the subregion (Greve et al., 2014; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low evidence: Mixed signal</b> between the different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM)), surface soil moisture (Xu et al., 2019a) and a weak drying by SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase</b> of drought severity. This is consistent between the different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM)), surface soil moisture (Xu et al., 2019a) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Likely: Increase</b> of drought severity. This is consistent between the different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM)), surface soil moisture (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows inconsistent trends(Touma et al., 2015)	<b>Low confidence: Limited evidence .</b> Inconsistent trends in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020), but slight stronger

1  
2  
3  
4  
5  
6  
7  
8  
9  
10

						tendency towards drying.
W. North America (WNA)	MET	<b>Low confidence: Inconsistent trends</b> depending on subregion (Swain and Hayhoe, 2015; Wehner et al., 2017; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence and inconsistent trends</b> depending on models and seasons (Swain and Hayhoe, 2015; Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM))	<b>Low confidence: Limited evidence and inconsistent trends</b> depending on models and seasons (Swain and Hayhoe, 2015; Xu et al., 2019a; Spinoni et al., 2020).	<b>Low confidence: Mixed signal</b> among models, seasons, and studies (Swain and Hayhoe, 2015; Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)), with tendency towards drying in the spring and wetting in summer (Swain and Hayhoe, 2015).
	AGR ECOL	<b>Medium confidence: Increase.</b> Dominant increase but some inconsistent trends based on soil moisture, water-balance estimates, PDSI and SPEI, but some inconsistent trends depending study, index and the subregion (Greve et al., 2014; Griffin and Anchukaitis, 2014; Williams et al., 2015, 2020; Ahmadalipour and Moradkhani, 2017; Dai and Zhao, 2017; Spinoni et al., 2019; Padrón et al., 2020)	<b>Medium confidence: Human contribution to observed trend.</b>  Williams et al. (2020) concluded human-induced climate change contributed to the strong soil moisture deficits recorded in the last two decades in western North America through VPD (and AED) increases associated with higher air temperatures and lower air humidity. Williams et al. (2015) and Griffin and Anchukaitis (2014) concluded that increased AED has had an increased contribution to drought severity over the last decades, and played a dominant role in the intensification of the 2012-2014 drought in California	<b>Low evidence: Inconsistent signal</b> between models, with weak tendency to increased drying in total and surface soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and the SPEI-PM (Naumann et al., 2018; Gu et al., 2020). Weak soil moisture drying projection for California (Louise et al., 2018)	<b>Medium confidence: Increase</b> of drought severity. There are differences depending on metrics and models, with weak median drying and substantial intermodel spread for total soil moisture (Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)) and larger drying for surface soil moisture (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020). Stronger soil moisture drying in southern part of domain (Cook et al., 2020).	<b>Medium confidence: Increase</b> of drought severity. There are differences depending on metrics and models, with weak drying in total column soil moisture (Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), and substantial drying with surface soil moisture (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Mixed signal</b> between different time frames and subregions (Gudmundsson et al., 2019, 2021; Poshtiri and Pal, 2016; Dudley et al., 2020). Strong spatial variability in the recent trends of low flows in the region (Poshtiri and Pal,	<b>Low confidence: Mixed signal</b> for overall region in observations. But evidence that temperature increase has been the main driver of increased hydrological drought in California and in the Colorado basin	<b>Low confidence: Limited evidence.</b> One study shows drying (Touma et al., 2015)	<b>Medium confidence: Increase</b> in hydrological drought (more intense low flows, less runoff and more frequent hydrological droughts) (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b) Particularly strong evidence of increasing hydrological droughts in regions dependent on snow pack	<b>Medium confidence: Increase</b> in hydrological droughts (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020) Particularly strong evidence of increasing hydrological droughts in regions dependent on snow pack reservoirs (Wehner et al., 2017; Ackerly et al., 2018; Rhoades et al.,

		2016) but dominant increase of hydrological drought in California and in the Colorado basin (Xiao et al., 2018b; Milly and Dunne, 2020).	(Milly and Dunne, 2020; Shukla et al., 2015; Xiao et al., 2018; Udall and Overpeck, 2017).		reservoirs (Wehner et al., 2017; Ackerly et al., 2018; Rhoades et al., 2018)	2018)
C. North America (CNA)	MET	<b>Medium confidence:</b> Decrease in the duration and frequency of meteorological droughts, (Wehner et al., 2017; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b> (Rupp et al., 2013; Easterling et al., 2017)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signal</b> among different models (Sillmann et al., 2013b; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Mixed signal</b> among different models (Sillmann et al., 2013b; Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)); drying trend in spring and summer (Swain and Hayhoe, 2015).
	AGR ECOL	<b>Low confidence: Mixed signal</b> based on soil moisture, water-balance estimates, PDSI and SPEI and conflicting trend depending of the subregion (Greve et al., 2014; Dai and Zhao, 2017; Seager et al., 2019; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence.</b> Human influence on surface soil moisture deficits due to increased evapotranspiration caused by higher temperatures. (Easterling et al., 2017)	<b>Medium confidence: Increase</b> in drought. Dominant signal shows drought increase based on total and surface soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Medium confidence: Increase</b> in drought severity or frequency. Changes are consistent between different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), surface soil moisture (Xu et al., 2019a) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>High confidence: Increase</b> of drought severity. Changes are consistent between different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020), surface soil moisture (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020), PDSI (Dai et al., 2018), and SPEI-PM (Cook et al., 2014b; Feng et al., 2017; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Mixed signal.</b> No signal in changes (Gudmundsson et al., 2021; Mo and Lettenmaier, 2018; Dudley et al., 2020). Poshtiri and Pal (2016) show strong spatial variability in the recent trends of low flows although there is an increase of hydrological droughts in the Missouri (Martin et al., 2020; Woodhouse and Wise, 2020) and in the Colorado basins (Xiao et al., 2018b; Milly and Dunne, 2020) Wetting trend in (Dai and Zhao, 2017)	<b>Low confidence: Inconsistent trends in observations.</b> Two studies suggest that emperature increase has been the main driver of increased hydrological drought in the Missouri basin (Martin et al., 2020; Woodhouse and Wise, 2020).	<b>Low confidence: Limited evidence.</b> One study shows drying (Touma et al., 2015)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b).	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
E. North America (ENA)	MET	<b>Low confidence: Inconsistent</b> trends depending on the region (Wehner et al., 2017; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b> (Easterling et al., 2017)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Xu et al., 2019a) (Chapter 11 Supplementary Material (11.SM)).	<b>Low confidence: Limited evidence</b> (Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Increase</b> in drought severity in the majority of models, but weaker or inconsistent trends in part of region (Sillmann et al., 2013b; Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR	<b>Low confidence: Mixed</b>	<b>Low confidence:</b>	<b>Low confidence:</b>	<b>Low confidence: Inconsistent</b>	<b>Medium confidence: Increase of</b>

	ECOL	<b>signal.</b> Inconsistent trends depending on metric, subregion, time frame and studies, based on soil moisture, water-balance estimates, PDSI, and SPEI (Greve et al., 2014; Dai and Zhao, 2017; Park Williams et al., 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Limited evidence.</b> Human influence on surface soil moisture deficits due to increased evapotranspiration caused by higher temperatures. (Easterling et al., 2017)	<b>Inconsistent</b> trends between models, metrics and studies based on total and surface soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	trends between models, metrics and studies based on total and surface soil moisture (Xu et al., 2019a; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), and SPEI-PM (Naumann et al., 2018; Gu et al., 2020), but with stronger tendency towards drying.	drought severity. Consistent signal between different drought metrics including total column soil moisture, (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020), surface soil moisture (Dai et al., 2018; Lu et al., 2019), PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a).
	HYDR	<b>Low confidence: Limited evidence.</b> Decrease in low flows from 1971-2020, but not since 1950 (Gudmundsson et al., 2019, 2021). Poshtiri and Pal, (2016) and Dudley et al., (2020) show strong spatial variability in the recent trends of low flows in the region.	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal ( Touma et al., 2015)	<b>Low confidence: Limited evidence and inconsistent trends</b> (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Mixed signal</b> among models and studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020)
N. E. North America (NEN)	MET	<b>Low confidence: No or limited signal</b> in duration and frequency of droughts (Bonsal et al., 2019; Dunn et al., 2020)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> Available evidence suggest decrease in meteorological drought (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Decrease</b> in meteorological drought (Sillmann et al., 2013b; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Decrease</b> in meteorological drought (Touma et al., 2015; Spinoni et al., 2020; Vicente-Serrano et al., 2020a)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Low confidence: Mixed signal</b> between different drought metrics and strong spatial differences (Greve et al., 2014; Dai and Zhao, 2017; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Mixed signal</b> between different models and metrics. Substantial intermodal variations and weak drying trend in soil moisture(Xu et al., 2019a)( Chapter 11 Supplementary Material (11.SM)) and slight decrease in drought severity in SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> between different models and drought metrics. Substantial intermodel spread for total column soil moisture, with overall weak or no change (Chapter 11 Supplementary Material (11.SM))(Cook et al., 2020), slight drying in surface soil moisture (Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)) and tendency to wetting trend in SPEI-PM (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> between models and different drought metrics, including total column soil moisture, which shows inconsistent changes (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2020), surface soil moisture, which suggest drying (Dai et al., 2018; Lu et al., 2019; Cook et al., 2020)(Chapter 11 Supplementary Material (11.SM)), and PDSI (Dai et al., 2018) and SPEI-PM (Cook et al., 2014b; Vicente-Serrano et al., 2020a), which show tendency fo wetting trend.
	HYDR	<b>Low confidence: Limited evidence.</b> Inconsistent trends in one study (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows inconsistent signals (Touma et al., 2015)	<b>Low confidence: Inconsistent trends and limited evidence.</b> Available studies suggest inconsistent trends in low flow (Zhai et al., 2020b) and the	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020). Some evidence

					SRI (Touma et al., 2015), and seasonally inconsistent trends in runoff, with decrease in summer and increase in winter (Cook et al., 2020).	( <i>medium confidence</i> ) for strong seasonality of trends, with decrease in summer and increase in winter (Giuntoli et al., 2015; Cook et al., 2020).
N. W. North America (NWN)	MET	<b>Low confidence: Mixed signal</b> with conflicting trends depending on the region (Bonsal et al., 2019; Spinoni et al., 2019; Dunn et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited and inconsistent evidence.</b> Some evidence points to decrease in meteorological drought severity or intensity based on SPI (Xu et al., 2019a) and CDD (Chapter 11 Supplementary Material (11.SM))	<b>Medium confidence: Decrease</b> in meteorological drought severity or intensity (Sillmann et al., 2013b; Xu et al., 2019a; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).	<b>Medium confidence: Decrease</b> in meteorological drought severity in the majority of models (Sillmann et al., 2013b; Swain and Hayhoe, 2015; Touma et al., 2015; Spinoni et al., 2020)(Chapter 11 Supplementary Material (11.SM)).
	AGR ECOL	<b>Low confidence: No signal or inconsistent signals</b> in the duration and severity of droughts based on soil moisture, PDSI and SPEI and conflicting trend depending of the subregion (Greve et al., 2014; Dai and Zhao, 2017; Park Williams et al., 2017; Spinoni et al., 2019; Padrón et al., 2020).	<b>Low confidence: Limited evidence</b>	<b>Low evidence: Mixed signal</b> in changes in drought severity. Inconsistent changes between models in CMIP6 and CMIP5 total and surface soil moisture(Xu et al., 2019a)(Chapter 11 Supplementary Material (11.SM)); SPEI-PM also suggests inconsistent changes drought severity (Naumann et al., 2018; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> between different models, drought metrics and studies, including total and surfacemoisture, as well as SPEI-PM(Chapter 11 Supplementary Material (11.SM))(Naumann et al., 2018; Xu et al., 2019a; Cook et al., 2020; Gu et al., 2020).	<b>Low confidence: Mixed signal</b> between different models and drought metrics, including total and surface soil moisture, PDSI and SPEI-PM (Chapter 11 Supplementary Material (11.SM)) (Cook et al., 2014b, 2020; Dai et al., 2018; Lu et al., 2019; Vicente-Serrano et al., 2020a), with slight larger tendency towards wetting.
	HYDR	<b>Low confidence: Limited evidence.</b> Regionally inconsistent trends in one study (Dai and Zhao, 2017)	<b>Low confidence: Limited evidence</b>	<b>Low confidence: Limited evidence.</b> One study shows lack of signal (Touma et al., 2015)	<b>Low confidence: Limited evidence and inconsistent</b> signals in available studies (Touma et al., 2015; Cook et al., 2020; Zhai et al., 2020b)	<b>Low confidence: Mixed signal</b> among studies (Prudhomme et al., 2014; Giuntoli et al., 2015; Touma et al., 2015; Cook et al., 2020), but slight stronger tendency towards wetting.

1  
2 [END TABLE 11.21 HERE]  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13

1 **Acknowledgements**

2

3 Nate McDowell, Alexis Berg, Jamie Hannaford, Jack Scheff, Lena Tallaksen, Tim Brodribb, Peter Stott,

4 Peter Thorne, Francis Zwiers.

5

1 **References**

- 2
- 3 Abatzoglou, J. T., and Williams, A. P. (2016). Impact of anthropogenic climate change on wildfire across western US  
4 forests. *Proc. Natl. Acad. Sci.*, 201607171. doi:10.1073/pnas.1607171113.
- 5 Abaurrea, J., Asín, J., and Cebrián, A. C. (2018). Modelling the occurrence of heat waves in maximum and minimum  
6 temperatures over Spain and projections for the period 2031-60. *Glob. Planet. Change* 161, 244–260.  
7 doi:10.1016/j.gloplacha.2017.11.015.
- 8 Abiodun, B. J., Adegoke, J., Abatan, A. A., Ibe, C. A., Egbebiyi, T. S., Engelbrecht, F., et al. (2017). Potential impacts  
9 of climate change on extreme precipitation over four African coastal cities. *Clim. Change* 143.  
10 doi:10.1007/s10584-017-2001-5.
- 11 Abiodun, B. J., Makhanya, N., Petja, B., Abatan, A. A., and Oguntunde, P. G. (2019). Future projection of droughts  
12 over major river basins in Southern Africa at specific global warming levels. *Theor. Appl. Climatol.* 137, 1785–  
13 1799. doi:10.1007/s00704-018-2693-0.
- 14 Acar Deniz, Z., and Gönençgil, B. (2015). Trends of summer daily maximum temperature extremes in Turkey. *Phys.*  
15 *Geogr.* 36, 268–281. doi:10.1080/02723646.2015.1045285.
- 16 Acero, F. J., García, J. A., Gallego, M. C., Parey, S., and Dacunha-Castelle, D. (2014). Trends in summer extreme  
17 temperatures over the Iberian Peninsula using nonurban station data. *J. Geophys. Res. Atmos.* 119, 39–53.  
18 doi:10.1002/2013JD020590.
- 19 Ackerly, D., Jones, A., Stacey, M., and Riordan, B. (2018). “San Francisco Bay Area Summary Report,” in *California’s*  
20 *Fourth Climate Change Assessment* SUM-CCCA4-2018-005. (Berkeley, CA, USA: University of California  
21 Berkeley). Available at: <https://www.climateassessment.ca.gov/regions/>.
- 22 Adnan, M., Rehman, N., and Shahbir, J. (2016). Predicting the Frequency and Intensity of Climate Extremes by  
23 Regression Models. *J. Climatol. Weather Forecast.* 04. doi:10.4172/2332-2594.1000185.
- 24 Aerenson, T., Tebaldi, C., Sanderson, B., and Lamarque, J.-F. (2018). Changes in a suite of indicators of extreme  
25 temperature and precipitation under 1.5 and 2 degrees warming. *Environ. Res. Lett.* 13, 035009.  
26 doi:10.1088/1748-9326/aaafd6.
- 27 Aerts, J. C. J. H., Botzen, W. J., Clarke, K. C., Cutter, S. L., Hall, J. W., Merz, B., et al. (2018). Integrating human  
28 behaviour dynamics into flood disaster risk assessment. *Nat. Clim. Chang.* 8, 193–199. doi:10.1038/s41558-018-  
29 0085-1.
- 30 Agard, V., and Emanuel, K. (2017). Clausius-Clapeyron scaling of peak CAPE in continental convective storm  
31 environments. *J. Atmos. Sci.* 74, 3043–3054. doi:10.1175/JAS-D-16-0352.1.
- 32 Agel, L., and Barlow, M. (2020). How well do CMIP6 historical runs match observed northeast U.S. precipitation and  
33 extreme precipitation–related circulation? *J. Clim.* 33, 9835–9848. doi:10.1175/JCLI-D-19-1025.1.
- 34 AghaKouchak, A. (2014). A baseline probabilistic drought forecasting framework using standardized soil moisture  
35 index: Application to the 2012 United States drought. *Hydrol. Earth Syst. Sci.* 18, 2485–2492. doi:10.5194/hess-  
36 18-2485-2014.
- 37 AghaKouchak, A., Cheng, L., Mazdiyasni, O., and Farahmand, A. (2014). Global warming and changes in risk of  
38 concurrent climate extremes: Insights from the 2014 California drought. *Geophys. Res. Lett.* 41, 8847–8852.  
39 doi:10.1002/2014GL062308.
- 40 AghaKouchak, A., Chiang, F., Huning, L. S., Love, C. A., Mallakpour, I., Mazdiyasni, O., et al. (2020). Climate  
41 Extremes and Compound Hazards in a Warming World. *Annu. Rev. Earth Planet. Sci.* 48, 519–548.  
42 doi:10.1146/annurev-earth-071719-055228.
- 43 Aguilar, E., Barry, A. A., Brunet, M., Ekang, L., Fernandes, A., Massoukina, M., et al. (2009). Changes in temperature  
44 and precipitation extremes in western central Africa, Guinea Conakry, and Zimbabwe, 1955-2006. *J. Geophys.*  
45 *Res. Atmos.* 114. doi:10.1029/2008JD011010.
- 46 Aguilar, E., Peterson, T. C., Obando, P. R., Frutos, R., Retana, J. A., Solera, M., et al. (2005). Changes in precipitation  
47 and temperature extremes in Central America and northern South America, 1961–2003. *J. Geophys. Res.* 110,  
48 D23107. doi:10.1029/2005JD006119.
- 49 Ahlswede, B., and Thomas, R. Q. (2017). Community Earth System Model Simulations Reveal the Relative Importance  
50 of Afforestation and Forest Management to Surface Temperature in Eastern North America. *Forests* 8, 499.  
51 doi:10.3390/f8120499.
- 52 Ahmadalipour, A., and Moradkhani, H. (2017). Analyzing the uncertainty of ensemble-based gridded observations in  
53 land surface simulations and drought assessment. *J. Hydrol.* 555, 557–568. doi:10.1016/j.jhydrol.2017.10.059.
- 54 Ahmed, M., Anchukaitis, K. J., Asrat, A., Borgaonkar, H. P., Braidia, M., Buckley, B. M., et al. (2013). Continental-  
55 scale temperature variability during the past two millennia. *Nat. Geosci.* 6, 339–346. doi:10.1038/ngeo1797.
- 56 Ahn, J.-B., Jo, S., Suh, M.-S., Cha, D.-H., Lee, D.-K., Hong, S.-Y., et al. (2016). Changes of precipitation extremes  
57 over South Korea projected by the 5 RCMs under RCP scenarios. *Asia-Pacific J. Atmos. Sci.* 52, 223–236.  
58 doi:10.1007/s13143-016-0021-0.
- 59 Aich, V., Liersch, S., Vetter, T., Fournet, S., Andersson, J. C. M., Calmanti, S., et al. (2016). Flood projections within  
60 the Niger River Basin under future land use and climate change. *Sci. Total Environ.* 562, 666–677.

- 1 doi:10.1016/j.scitotenv.2016.04.021.
- 2 Akinsanola, A. A., Kooperman, G. J., Pendergrass, A. G., Hannah, W. M., and Reed, K. A. (2020). Seasonal  
3 representation of extreme precipitation indices over the United States in CMIP6 present-day simulations. *Environ.*  
4 *Res. Lett.* 15, 094003. doi:10.1088/1748-9326/ab92c1.
- 5 Akinsanola, A. A., and Zhou, W. (2018). Projections of West African summer monsoon rainfall extremes from two  
6 CORDEX models. *Clim. Dyn.* doi:10.1007/s00382-018-4238-8.
- 7 Akinyemi, F. O., and Abiodun, B. J. (2019). Potential impacts of global warming levels 1.5 °C and above on climate  
8 extremes in Botswana. *Clim. Change* 154, 387–400. doi:10.1007/s10584-019-02446-1.
- 9 Albergel, C., Dorigo, W., Reichle, R. H., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., et al. (2013). Skill and Global  
10 Trend Analysis of Soil Moisture from Reanalyses and Microwave Remote Sensing. *J. Hydrometeorol.* 14, 1259–  
11 1277. doi:10.1175/JHM-D-12-0161.1.
- 12 Alexander, L. V. (2016). Global observed long-term changes in temperature and precipitation extremes: A review of  
13 progress and limitations in IPCC assessments and beyond. *Weather Clim. Extrem.* 11, 4–16.  
14 doi:10.1016/J.WACE.2015.10.007.
- 15 Alexander, L. V., and Arblaster, J. M. (2017). Historical and projected trends in temperature and precipitation extremes  
16 in Australia in observations and CMIP5. *Weather Clim. Extrem.* 15, 34–56.
- 17 Alexander, L. V., Fowler, H. J., Bador, M., Behrangi, A., Donat, M. G., Dunn, R., et al. (2019). On the use of indices to  
18 study extreme precipitation on sub-daily and daily timescales. *Environ. Res. Lett.* 14, 125008. doi:10.1088/1748-  
19 9326/ab51b6.
- 20 Alexandru, A. (2018). Consideration of land-use and land-cover changes in the projection of climate extremes over  
21 North America by the end of the twenty-first century. *Clim. Dyn.* 50, 1949–1973. doi:10.1007/s00382-017-3730-  
22 x.
- 23 Alfieri, L., Bisselink, B., Dottori, F., Naumann, G., de Roo, A., Salamon, P., et al. (2017). Global projections of river  
24 flood risk in a warmer world. *Earth's Futur.* 5, 171–182. doi:10.1002/2016EF000485.
- 25 Ali, H., Fowler, H. J., Lenderink, G., Lewis, E., and Pritchard, D. (2021). Consistent large-scale response of hourly  
26 extreme precipitation to temperature variation over land. *Geophys. Res. Lett.* doi:10.1029/2020GL090317.
- 27 Ali, H., and Mishra, V. (2018). Contributions of Dynamic and Thermodynamic Scaling in Subdaily Precipitation  
28 Extremes in India. *Geophys. Res. Lett.* 45, 2352–2361. doi:10.1002/2018GL077065.
- 29 Ali, H., Modi, P., and Mishra, V. (2019a). Increased flood risk in Indian sub-continent under the warming climate.  
30 *Weather Clim. Extrem.* 25, 100212. doi:10.1016/j.wace.2019.100212.
- 31 Ali, S., Eum, H.-I., Cho, J., Dan, L., Khan, F., Dairaku, K., et al. (2019b). Assessment of climate extremes in future  
32 projections downscaled by multiple statistical downscaling methods over Pakistan. *Atmos. Res.* 222, 114–133.  
33 doi:10.1016/j.atmosres.2019.02.009.
- 34 Alizadeh, M. R., Adamowski, J., Nikoo, M. R., AghaKouchak, A., Dennison, P., and Sadegh, M. (2020). A century of  
35 observations reveals increasing likelihood of continental-scale compound dry-hot extremes. *Sci. Adv.* 6, eaaz4571.  
36 doi:10.1126/sciadv.aaz4571.
- 37 Alkama, R., and Cescatti, A. (2016). Biophysical climate impacts of recent changes in global forest cover. *Science* (80-  
38 ). 351, 600–604. doi:10.1126/science.aac8083.
- 39 Allan, R. P., Barlow, M., Byrne, M. P., Cherchi, A., Douville, H., Fowler, H. J., et al. (2020). Advances in  
40 understanding large-scale responses of the water cycle to climate change. *Ann. N. Y. Acad. Sci.* 1472, 49–75.  
41 doi:10.1111/nyas.14337.
- 42 Allen, C. D., Breshears, D. D., and McDowell, N. G. (2015). On underestimation of global vulnerability to tree  
43 mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6, art129. doi:10.1890/ES15-  
44 00203.1.
- 45 Allen, J. T. (2018). “Climate Change and Severe Thunderstorms,” in *Oxford Research Encyclopedia of Climate Science*  
46 (Oxford, UK: Oxford University Press), 1–65. doi:10.1093/acrefore/9780190228620.013.62.
- 47 Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., et al. (2018). “Framing and  
48 Context,” in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above*  
49 *pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the*  
50 *global response to the threat of climate change*, eds. V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J.  
51 Skea, P. R. Shukla, et al. (In Press), 49–92. Available at: <https://www.ipcc.ch/sr15/chapter/chapter-1>.
- 52 Almazroui, M. (2019a). Assessment of meteorological droughts over Saudi Arabia using surface rainfall observations  
53 during the period 1978–2017. *Arab. J. Geosci.* 12. doi:10.1007/s12517-019-4866-2.
- 54 Almazroui, M. (2019b). Temperature Changes over the CORDEX-MENA Domain in the 21st Century Using CMIP5  
55 Data Downscaled with RegCM4: A Focus on the Arabian Peninsula. *Adv. Meteorol.* 2019.  
56 doi:10.1155/2019/5395676.
- 57 Almazroui, M., and Islam, M. N. (2019). Coupled Model Inter-comparison Project Database to Calculate Drought  
58 Indices for Saudi Arabia: A Preliminary Assessment. *Earth Syst. Environ.* 3, 419–428. doi:10.1007/s41748-019-  
59 00126-9.
- 60 Almazroui, M., Islam, M. N., Dambul, R., and Jones, P. D. (2014). Trends of temperature extremes in Saudi Arabia. *Int.*  
61 *J. Climatol.* 34, 808–826. doi:10.1002/joc.3722.



- 1 Almazroui, M., and Saeed, S. (2020). Contribution of extreme daily precipitation to total rainfall over the Arabian  
2 Peninsula. *Atmos. Res.* 231, 104672. doi:10.1016/j.atmosres.2019.104672.
- 3 Almeida, C. T., Oliveira-Júnior, J. F., Delgado, R. C., Cubo, P., and Ramos, M. C. (2017). Spatiotemporal rainfall and  
4 temperature trends throughout the Brazilian Legal Amazon, 1973–2013. *Int. J. Climatol.* 37, 2013–2026.  
5 doi:10.1002/joc.4831.
- 6 AlSarmi, S. H., and Washington, R. (2014). Changes in climate extremes in the Arabian Peninsula: analysis of daily  
7 data. *Int. J. Climatol.* 34, 1329–1345. doi:10.1002/joc.3772.
- 8 Althoff, D., Rodrigues, L. N., and da Silva, D. D. (2020). Impacts of climate change on the evaporation and availability  
9 of water in small reservoirs in the Brazilian savannah. *Clim. Change* 159, 215–232. doi:10.1007/s10584-020-  
10 02656-y.
- 11 Altman, J., Ukhvatkina, O. N., Omelko, A. M., Macek, M., Plener, T., Pejcha, V., et al. (2018). Poleward migration of  
12 the destructive effects of tropical cyclones during the 20th century. *Proc. Natl. Acad. Sci.* 115, 11543 LP–11548.  
13 doi:10.1073/pnas.1808979115.
- 14 Amann, B., Szidat, S., and Grosjean, M. (2015). A millennial-long record of warm season precipitation and flood  
15 frequency for the North-western Alps inferred from varved lake sediments: implications for the future. *Quat. Sci.*  
16 *Rev.* 115, 89–100. doi:10.1016/j.quascirev.2015.03.002.
- 17 Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., et al. (2017). A human-driven  
18 decline in global burned area. *Science (80-. )*. 356, 1356–1362. doi:10.1126/science.aal4108.
- 19 Anderegg, W. R. L., Berry, J. A., Smith, D. D., Sperry, J. S., Anderegg, L. D. L., and Field, C. B. (2012). The roles of  
20 hydraulic and carbon stress in a widespread climate-induced forest die-off. *Proc. Natl. Acad. Sci.* 109, 233–237.  
21 doi:10.1073/pnas.1107891109.
- 22 Anderegg, W. R. L., Kane, J. M., and Anderegg, L. D. L. (2013). Consequences of widespread tree mortality triggered  
23 by drought and temperature stress. *Nat. Clim. Chang.* 3, 30–36. doi:10.1038/nclimate1635.
- 24 Anderegg, W. R. L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A. F. A., Choat, B., et al. (2016). Meta-analysis reveals  
25 that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. *Proc. Natl.*  
26 *Acad. Sci. U. S. A.* 113, 5024–5029. doi:10.1073/pnas.1525678113.
- 27 Anderegg, W. R. L., Trugman, A. T., Badgley, G., Konings, A. G., and Shaw, J. (2020). Divergent forest sensitivity to  
28 repeated extreme droughts. *Nat. Clim. Chang.* doi:10.1038/s41558-020-00919-1.
- 29 Anderson-Teixeira, K. J., Snyder, P. K., Twine, T. E., Cuadra, S. V., Costa, M. H., and DeLucia, E. H. (2012). Climate-  
30 regulation services of natural and agricultural ecoregions of the Americas. *Nat. Clim. Chang.* 2, 177–181.  
31 doi:10.1038/nclimate1346.
- 32 Anderson, R. G., Canadell, J. G., Randerson, J. T., Jackson, R. B., Hungate, B. A., Baldocchi, D. D., et al. (2011).  
33 Biophysical considerations in forestry for climate protection. *Front. Ecol. Environ.* 9, 174–182.  
34 doi:10.1890/090179.
- 35 Anderson, W. B., Seager, R., Baethgen, W., Cane, M., and You, L. (2019). Synchronous crop failures and climate-  
36 forced production variability. *Sci. Adv.* 5, eaaw1976. doi:10.1126/sciadv.aaw1976.
- 37 Añel, J. A., López-Moreno, J. I., Otto, F. E. L., Vicente-Serrano, S. M., Schaller, N., Massey, N., et al. (2014a). The  
38 extreme snow accumulation in the Western Spanish Pyrenees during winter and spring 2013 [in “Explaining  
39 Extreme Events of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S73–S74. doi:10.1175/1520-  
40 0477-95.9.S1.1.
- 41 Añel, J. A., López-Moreno, J. I., Otto, F. E. L., Vicente Serrano, S. M., Schaller, Nathalie; Massey, N., Buisán, S. T., et  
42 al. (2014b). The extreme snow accumulation in the western Spanish Pyrenees during winter and spring 2013 [in  
43 “Explaining Extreme Events of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S73–S76.  
44 doi:10.1175/1520-0477-95.9.S1.1.
- 45 Angeles-Malaspina, M., González-Cruz, J. E., and Ramírez-Beltrán, N. (2018). Projections of Heat Waves Events in the  
46 Intra-Americas Region Using Multimodel Ensemble. *Adv. Meteorol.* 2018, 1–16. doi:10.1155/2018/7827984.
- 47 Angéilil, O., Perkins-Kirkpatrick, S., Alexander, L. V., Stone, D., Donat, M. G., Wehner, M., et al. (2016). Comparing  
48 regional precipitation and temperature extremes in climate model and reanalysis products. *Weather Clim. Extrem.*  
49 13, 35–43. doi:10.1016/j.wace.2016.07.001.
- 50 Angéilil, O., Stone, D. A., Tadross, M., Tummon, F., Wehner, M., and Knutti, R. (2014). Attribution of extreme weather  
51 to anthropogenic greenhouse gas emissions: Sensitivity to spatial and temporal scales. *Geophys. Res. Lett.* 41,  
52 2150–2155. doi:10.1002/2014GL059234.
- 53 Angéilil, O., Stone, D., Perkins-Kirkpatrick, S., Alexander, L. V., Wehner, M., Shiogama, H., et al. (2018). On the  
54 nonlinearity of spatial scales in extreme weather attribution statements. *Clim. Dyn.* 50, 2739–2752.  
55 doi:10.1007/s00382-017-3768-9.
- 56 Angéilil, O., Stone, D., Wehner, M., Paciorek, C. J., Krishnan, H., and Collins, W. (2017). An Independent Assessment  
57 of Anthropogenic Attribution Statements for Recent Extreme Temperature and Rainfall Events. *J. Clim.* 30, 5–16.  
58 doi:10.1175/JCLI-D-16-0077.1.
- 59 Antonescu, B., Schultz, D. M., Holzer, A., and Groenemeijer, P. (2016a). Tornadoes in Europe: An Underestimated  
60 Threat. *Bull. Am. Meteorol. Soc.* 98, 713–728. doi:10.1175/BAMS-D-16-0171.1.
- 61 Antonescu, B., Schultz, D. M., Lomas, F., and Kühne, T. (2016b). Tornadoes in Europe: Synthesis of the Observational

- 1 Datasets. *Mon. Weather Rev.* 144, 2445–2480. doi:10.1175/MWR-D-15-0298.1.
- 2 Apurv, T., Sivapalan, M., and Cai, X. (2017). Understanding the Role of Climate Characteristics in Drought  
3 Propagation. *Water Resour. Res.* 53, 9304–9329. doi:10.1002/2017WR021445.
- 4 Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., et al. (2018). 21st  
5 Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nat. Commun.* 9,  
6 536. doi:10.1038/s41467-017-02771-y.
- 7 Arblaster, J. M., Lim, E.-P., Hendon, H. H., Trewin, B., Wheeler, M. C., Liu, G., et al. (2014). Understanding  
8 Australia's hottest September on record [in "Explaining Extreme Events of 2013 from a Climate Perspective"].  
9 *Bull. Am. Meteorol. Soc.* 95, S37–S41. doi:10.1175/1520-0477-95.9.S1.1.
- 10 Archer, D. R., and Fowler, H. J. (2018). Characterising flash flood response to intense rainfall and impacts using  
11 historical information and gauged data in Britain. *J. Flood Risk Manag.* 11, S121–S133. doi:10.1111/jfr3.12187.
- 12 Archer, D. R., Parkin, G., and Fowler, H. J. (2016). Assessing long term flash flooding frequency using historical  
13 information. *Hydrol. Res.* 48, 1–16. doi:10.2166/nh.2016.031.
- 14 Archfield, S. A., Hirsch, R. M., Viglione, A., and Blöschl, G. (2016). Fragmented patterns of flood change across the  
15 United States. *Geophys. Res. Lett.* 43, 10,232–10,239. doi:10.1002/2016GL070590.
- 16 Argüeso, D., Di Luca, A., and Evans, J. P. (2016). Precipitation over urban areas in the western Maritime Continent  
17 using a convection-permitting model. *Clim. Dyn.* 47, 1143–1159. doi:10.1007/s00382-015-2893-6.
- 18 Argüeso, D., Hidalgo-Muñoz, J. M., Ga´ Miz-Fortis, S. R., Esteban-Parra, M. J., and Castro-Díez, Y. (2012). Evaluation  
19 of WRF Mean and Extreme Precipitation over Spain: Present Climate (1970–99). *J. Clim.* 25, 4883–4895.  
20 doi:10.1175/JCLI-D-11-00276.1.
- 21 Armstrong, W. H., Collins, M. J., and Snyder, N. P. (2014). Hydroclimatic flood trends in the northeastern United  
22 States and linkages with large-scale atmospheric circulation patterns. *Hydrol. Sci. J.* 59, 1636–1655.  
23 doi:10.1080/02626667.2013.862339.
- 24 Arnell, N. W., and Gosling, S. N. (2016). The impacts of climate change on river flood risk at the global scale. *Clim.*  
25 *Change* 134, 387–401. doi:10.1007/s10584-014-1084-5.
- 26 Arnone, E., Pumo, D., Viola, F., Noto, L. V., and La Loggia, G. (2013). Rainfall statistics changes in Sicily. *Hydrol.*  
27 *Earth Syst. Sci.* 17, 2449–2458. doi:10.5194/hess-17-2449-2013.
- 28 Ashabokov, B. A., Tashilova, A. A., Kesheva, L. A., and Taubekova, Z. A. (2017). Trends in precipitation parameters  
29 in the climate zones of southern Russia (1961–2011). *Russ. Meteorol. Hydrol.* 42, 150–158.  
30 doi:10.3103/S1068373917030025.
- 31 Asmerom, Y., Polyak, V. J., Rasmussen, J. B. T., Burns, S. J., and Lachniet, M. (2013). Multidecadal to multicentury  
32 scale collapses of Northern Hemisphere monsoons over the past millennium. *Proc. Natl. Acad. Sci. U. S. A.* 110,  
33 9651–9656. doi:10.1073/pnas.1214870110.
- 34 Atif, R. M., Almazroui, M., Saeed, S., Abid, M. A., Islam, M. N., and Ismail, M. (2020). Extreme precipitation events  
35 over Saudi Arabia during the wet season and their associated teleconnections. *Atmos. Res.* 231, 104655.  
36 doi:10.1016/j.atmosres.2019.104655.
- 37 Ault, T. R. (2020). On the essentials of drought in a changing climate. *Science (80-. )*. 368, 256–260.  
38 doi:10.1126/science.aaz5492.
- 39 Ault, T. R., Cole, J. E., Overpeck, J. T., Pederson, G. T., and Meko, D. M. (2014). Assessing the Risk of Persistent  
40 Drought Using Climate Model Simulations and Paleoclimate Data. *J. Clim.* 27, 7529–7549. doi:10.1175/JCLI-D-  
41 12-00282.1.
- 42 Avila-Diaz, A., Benezoli, V., Justino, F., Torres, R., and Wilson, A. (2020). Assessing current and future trends of  
43 climate extremes across Brazil based on reanalyses and earth system model projections. *Clim. Dyn.* 55, 1403–  
44 1426. doi:10.1007/s00382-020-05333-z.
- 45 Ávila, A., Justino, F., Wilson, A., Bromwich, D., and Amorim, M. (2016). Recent precipitation trends, flash floods and  
46 landslides in southern Brazil. *Environ. Res. Lett.* 11, 114029. doi:10.1088/1748-9326/11/11/114029.
- 47 Avila, F. B., Dong, S., Menang, K. P., Rajczak, J., Renom, M., Donat, M. G., et al. (2015). Systematic investigation of  
48 gridding-related scaling effects on annual statistics of daily temperature and precipitation maxima: A case study  
49 for south-east Australia. *Weather Clim. Extrem.* 9, 6–16. doi:10.1016/j.wace.2015.06.003.
- 50 Azorin-Molina, C., Vicente-Serrano, S. M., McVicar, T. R., Revuelto, J., Jerez, S., and López-Moreno, J. I. (2017).  
51 Assessing the impact of measurement time interval when calculating wind speed means and trends under the  
52 stilling phenomenon. *Int. J. Climatol.* 37, 480–492. doi:10.1002/joc.4720.
- 53 Azorin-Molina, C., Vicente-Serrano, S. M., Sanchez-Lorenzo, A., McVicar, T. R., Morán-Tejeda, E., Revuelto, J., et al.  
54 (2015). Atmospheric evaporative demand observations, estimates and driving factors in Spain (1961–2011). *J.*  
55 *Hydrol.* 523, 262–277. doi:10.1016/j.jhydrol.2015.01.046.
- 56 Baburaj, P. P., Abhilash, S., Mohankumar, K., and Sahai, A. K. (2020). On the Epochal Variability in the Frequency of  
57 Cyclones during the Pre-Onset and Onset Phases of the Monsoon over the North Indian Ocean. *Adv. Atmos. Sci.*  
58 37, 634–651. doi:10.1007/s00376-020-9070-5.
- 59 Bacmeister, J. T., Reed, K. A., Hannay, C., Lawrence, P., Bates, S., Truesdale, J. E., et al. (2018). Projected changes in  
60 tropical cyclone activity under future warming scenarios using a high-resolution climate model. *Clim. Change*  
61 146, 547–560. doi:10.1007/s10584-016-1750-x.

- 1 Bador, M., Boé, J., Terray, L., Alexander, L. V., Baker, A., Bellucci, A., et al. (2020). Impact of Higher Spatial  
2 Atmospheric Resolution on Precipitation Extremes Over Land in Global Climate Models. *J. Geophys. Res. Atmos.*  
3 125. doi:10.1029/2019JD032184.
- 4 Bador, M., Terray, L., and Boé, J. (2016). Detection of anthropogenic influence on the evolution of record-breaking  
5 temperatures over Europe. *Clim. Dyn.* 46, 2717–2735. doi:10.1007/s00382-015-2725-8.
- 6 Baek, H.-J., Kim, M.-K., and Kwon, W.-T. (2017). Observed short- and long-term changes in summer precipitation  
7 over South Korea and their links to large-scale circulation anomalies. *Int. J. Climatol.* 37, 972–986.  
8 doi:10.1002/joc.4753.
- 9 Baek, S. H., Steiger, N. J., Smerdon, J. E., and Seager, R. (2019). Oceanic Drivers of Widespread Summer Droughts in  
10 the United States Over the Common Era. *Geophys. Res. Lett.* 46, 8271–8280. doi:10.1029/2019GL082838.
- 11 Bai, P., Liu, X., Liang, K., and Liu, C. (2016). Investigation of changes in the annual maximum flood in the Yellow  
12 River basin, China. *Quat. Int.* 392, 168–177. doi:10.1016/j.quaint.2015.04.053.
- 13 Balaguru, K., Foltz, G. R., and Leung, L. R. (2018). Increasing Magnitude of Hurricane Rapid Intensification in the  
14 Central and Eastern Tropical Atlantic. *Geophys. Res. Lett.* 45, 4238–4247. doi:10.1029/2018GL077597.
- 15 Balaji, M., Chakraborty, A., and Mandal, M. (2018). Changes in tropical cyclone activity in north Indian Ocean during  
16 satellite era (1981–2014). *Int. J. Climatol.* 38, 2819–2837. doi:10.1002/joc.5463.
- 17 Ballesteros Cánovas, J. A., Trappmann, D., Shekhar, M., Bhattacharyya, A., and Stoffel, M. (2017). Regional flood-  
18 frequency reconstruction for Kullu district, Western Indian Himalayas. *J. Hydrol.* 546, 140–149.  
19 doi:10.1016/j.jhydrol.2016.12.059.
- 20 Balsamo, G., Albergel, C., Beljaars, A., Boussetta, S., Brun, E., Cloke, H., et al. (2015). ERA-Interim/Land: a global  
21 land surface reanalysis data set. *Hydrol. Earth Syst. Sci.* 19, 389–407. doi:10.5194/hess-19-389-2015.
- 22 Ban, N., Schmidli, J., and Schär, C. (2014). Evaluation of the new convective-resolving regional climate modeling  
23 approach in decade-long simulations. *J. Geophys. Res. Atmos.* 119, 7889–7907. doi:10.1002/2014JD021478.
- 24 Ban, N., Schmidli, J., and Schär, C. (2015). Heavy precipitation in a changing climate: Does short-term summer  
25 precipitation increase faster? *Geophys. Res. Lett.* 42, 1165–1172. doi:10.1002/2014GL062588.
- 26 Bandyopadhyay, N., Bhuiyan, C., and Saha, A. K. (2016). Heat waves, temperature extremes and their impacts on  
27 monsoon rainfall and meteorological drought in Gujarat, India. *Nat. Hazards* 82, 367–388. doi:10.1007/s11069-  
28 016-2205-4.
- 29 Bao, J., and Sherwood, S. C. (2019). The Role of Convective Self-Aggregation in Extreme Instantaneous Versus Daily  
30 Precipitation. *J. Adv. Model. Earth Syst.* 11, 19–33. doi:10.1029/2018MS001503.
- 31 Bao, J., Sherwood, S. C., Alexander, L. V., and Evans, J. P. (2017). Future increases in extreme precipitation exceed  
32 observed scaling rates. *Nat. Clim. Chang.* 7, 128–132. doi:10.1038/nclimate3201.
- 33 Barbero, R., Fowler, H. J., Lenderink, G., and Blenkinsop, S. (2017). Is the intensification of precipitation extremes  
34 with global warming better detected at hourly than daily resolutions? *Geophys. Res. Lett.* 44, 974–983.  
35 doi:10.1002/2016GL071917.
- 36 Barcikowska, M. J., Weaver, S. J., Feser, F., Russo, S., Schenk, F., Stone, D. A., et al. (2018). Euro-Atlantic winter  
37 storminess and precipitation extremes under 1.5° C vs. 2° C warming scenarios. *Earth Syst. Dyn.* 9. Available at:  
38 <https://doi.org/10.5194/esd-9-679-2018>.
- 39 Bard, A., Renard, B., Lang, M., Giuntoli, I., Korck, J., Koboltschnig, G., et al. (2015). Trends in the hydrologic regime  
40 of Alpine rivers. *J. Hydrol.* 529, 1823–1837. doi:10.1016/j.jhydrol.2015.07.052.
- 41 Barella-Ortiz, A., and Quintana Seguí, P. (2019). Evaluation of drought representation and propagation in regional  
42 climate model simulations across Spain. *Hydrol. Earth Syst. Sci.* 23, 5111–5131. doi:10.5194/hess-23-5111-2019.
- 43 Barichivich, J., Gloor, E., Peylin, P., Brienen, R. J. W., Schöngart, J., Espinoza, J. C., et al. (2018). Recent  
44 intensification of Amazon flooding extremes driven by strengthened Walker circulation. *Sci. Adv.* 4.  
45 doi:10.1126/sciadv.aat8785.
- 46 Barker, L. J., Hannaford, J., Chiveron, A., and Svensson, C. (2016). From meteorological to hydrological drought  
47 using standardised indicators. *Hydrol. Earth Syst. Sci.* 20, 2483–2505. doi:10.5194/hess-20-2483-2016.
- 48 Barker, L. J., Hannaford, J., Parry, S., Smith, K. A., Tanguy, M., and Prudhomme, C. (2019). Historic hydrological  
49 droughts 1891–2015: Systematic characterisation for a diverse set of catchments across the UK. *Hydrol. Earth  
50 Syst. Sci.* 23, 4583–4602. doi:10.5194/hess-23-4583-2019.
- 51 Barkhordarian, A., Saatchi, S. S., Behrangi, A., Loikith, P. C., and Mechoso, C. R. (2019). A Recent Systematic  
52 Increase in Vapor Pressure Deficit over Tropical South America. *Sci. Rep.* 9, 15331. doi:10.1038/s41598-019-  
53 51857-8.
- 54 Barlow, M., and Hoell, A. (2015). Drought in the Middle East and Central–Southwest Asia During Winter 2013/14.  
55 *Bull. Am. Meteorol. Soc.* 96, S71–S76. doi:10.1175/BAMS-D-15-00127.1.
- 56 Barlow, M., Zaitchik, B., Paz, S., Black, E., Evans, J., and Hoell, A. (2016). A review of drought in the Middle East and  
57 southwest Asia. *J. Clim.* 29, 8547–8574. doi:10.1175/JCLI-D-13-00692.1.
- 58 Barnard, P. L., Hoover, D., Hubbard, D. M., Snyder, A., Ludka, B. C., Allan, J., et al. (2017). Extreme oceanographic  
59 forcing and coastal response due to the 2015–2016 El Niño. *Nat. Commun.* 8, 14365. doi:10.1038/ncomms14365.
- 60 Barnhart, T. B., Molotch, N. P., Livneh, B., Harpold, A. A., Knowles, J. F., and Schneider, D. (2016). Snowmelt rate  
61 dictates streamflow. *Geophys. Res. Lett.* 43, 8006–8016. doi:10.1002/2016GL069690.

- 1 Barraqué, B. (2017). The common property issue in flood control through land use in France. *J. Flood Risk Manag.* 10,  
2 182–194. doi:10.1111/jfr3.12092.
- 3 Barriopedro, D., Fischer, E. M., Luterbacher, J., Trigo, R. M., and Garcia-Herrera, R. (2011). The Hot Summer of 2010:  
4 Redrawing the Temperature Record Map of Europe. *Science* (80-. ). 332, 220–224. doi:10.1126/science.1201224.
- 5 Barros, V. R., Boninsegna, J. A., Camilloni, I. A., Chidiak, M., Magrín, G. O., and Rusticucci, M. (2015). Climate  
6 change in Argentina: trends, projections, impacts and adaptation. *Wiley Interdiscip. Rev. Clim. Chang.* 6, 151–  
7 169. doi:10.1002/wcc.316.
- 8 Barry, A. A., Caesar, J., Klein Tank, A. M. G., Aguilar, E., McSweeney, C., Cyrille, A. M., et al. (2018). West Africa  
9 climate extremes and climate change indices. *Int. J. Climatol.* 38, e921–e938. doi:10.1002/joc.5420.
- 10 Bartók, B., Wild, M., Folini, D., Lüthi, D., Kotlarski, S., Schär, C., et al. (2017). Projected changes in surface solar  
11 radiation in CMIP5 global climate models and in EURO-CORDEX regional climate models for Europe. *Clim.*  
12 *Dyn.* 49, 2665–2683. doi:10.1007/s00382-016-3471-2.
- 13 Basara, J. B., Christian, J. I., Wakefield, R. A., Otkin, J. A., Hunt, E. H., and Brown, D. P. (2019). The evolution,  
14 propagation, and spread of flash drought in the Central United States during 2012. *Environ. Res. Lett.* 14, 084025.  
15 doi:10.1088/1748-9326/ab2cc0.
- 16 Basconcello, J., Lucero, A., Solis, A., Sandoval Jr, R., Bautista, E., Koizumi, T., et al. (2016). Statistically downscaled  
17 projected changes in seasonal mean temperature and rainfall in Cagayan Valley, Philippines. *J. Meteorol. Soc.*  
18 *Japan. Ser. II* 94, 151–164.
- 19 Bathiany, S., Dakos, V., Scheffer, M., and Lenton, T. M. (2018). Climate models predict increasing temperature  
20 variability in poor countries. *Sci. Adv.* 4, eaar5809. doi:10.1126/sciadv.aar5809.
- 21 Befort, D. J., Wild, S., Kruschke, T., Ulbrich, U., and Leckebusch, G. C. (2016). Different long-term trends of extra-  
22 tropical cyclones and windstorms in ERA-20C and NOAA-20CR reanalyses. *Atmos. Sci. Lett.* 17, 586–595.  
23 doi:10.1002/asl.694.
- 24 Beguería, S., Vicente-Serrano, S. M., Reig, F., and Latorre, B. (2014). Standardized precipitation evapotranspiration  
25 index (SPEI) revisited: Parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *Int.*  
26 *J. Climatol.* 34. doi:10.1002/joc.3887.
- 27 Bell, S. S., Chand, S. S., Tory, K. J., Dowdy, A. J., Turville, C., and Ye, H. (2019). Projections of southern hemisphere  
28 tropical cyclone track density using CMIP5 models. *Clim. Dyn.* 52, 6065–6079. doi:10.1007/s00382-018-4497-4.
- 29 Bellprat, O., García-Serrano, J., Fučkar, N. S., Massonnet, F., Guemas, V., and Doblas-Reyes, F. J. (2016). The Role of  
30 Arctic Sea Ice and Sea Surface Temperatures on the Cold 2015 February Over North America. *Bull. Am.*  
31 *Meteorol. Soc.* 97, S36–S41. doi:10.1175/BAMS-D-16-0159.1.
- 32 Bellprat, O., Lott, F. C., Gulizia, C., Parker, H. R., Pampuch, L. A., Pinto, I., et al. (2015). Unusual past dry and wet  
33 rainy seasons over Southern Africa and South America from a climate perspective. *Weather Clim. Extrem.* 9, 36–  
34 46. doi:10.1016/J.WACE.2015.07.001.
- 35 Belušić, D., de Vries, H., Dobler, A., Landgren, O., Lind, P., Lindstedt, D., et al. (2020). HCLIM38: a flexible regional  
36 climate model applicable for different climate zones from coarse to convection-permitting scales. *Geosci. Model*  
37 *Dev.* 13, 1311–1333. doi:10.5194/gmd-13-1311-2020.
- 38 Benestad, R. E., Parding, K. M., Erlandsen, H. B., and Mezghani, A. (2019). A simple equation to study changes in  
39 rainfall statistics. *Environ. Res. Lett.* 14, 84017. doi:10.1088/1748-9326/ab2bb2.
- 40 Benestad, R. E., van Oort, B., Justino, F., Stordal, F., Parding, K. M., Mezghani, A., et al. (2018). Downscaling  
41 probability of long heatwaves based on seasonal mean daily maximum temperatures. *Adv. Stat. Climatol.*  
42 *Meteorol. Oceanogr.* 4, 37–52. doi:10.5194/ascmo-4-37-2018.
- 43 Benito, G., Brázdil, R., Herget, J., and Machado, M. J. (2015). Quantitative historical hydrology in Europe. *Hydrol.*  
44 *Earth Syst. Sci.* 19, 3517–3539. doi:10.5194/hess-19-3517-2015.
- 45 Bennett, K. E., and Walsh, J. E. (2015). Spatial and temporal changes in indices of extreme precipitation and  
46 temperature for Alaska. *Int. J. Climatol.* 35, 1434–1452. doi:10.1002/joc.4067.
- 47 Berg, A., Findell, K., Lintner, B., Giannini, A., Seneviratne, S. I., van den Hurk, B., et al. (2016). Land–atmosphere  
48 feedbacks amplify aridity increase over land under global warming. *Nat. Clim. Chang.* 6, 869–874.  
49 doi:10.1038/nclimate3029.
- 50 Berg, A., Lintner, B. R., Findell, K., and Giannini, A. (2017a). Uncertain soil moisture feedbacks in model projections  
51 of Sahel precipitation. *Geophys. Res. Lett.* 44, 6124–6133. doi:10.1002/2017GL073851.
- 52 Berg, A., Lintner, B. R., Findell, K., Seneviratne, S. I., van den Hurk, B., Ducharme, A., et al. (2015). Interannual  
53 Coupling between Summertime Surface Temperature and Precipitation over Land: Processes and Implications for  
54 Climate Change. *J. Clim.* 28, 1308–1328. doi:10.1175/JCLI-D-14-00324.1.
- 55 Berg, A., and Sheffield, J. (2018). Climate Change and Drought: the Soil Moisture Perspective. *Curr. Clim. Chang.*  
56 *Reports* 4, 180–191. doi:10.1007/s40641-018-0095-0.
- 57 Berg, A., Sheffield, J., and Milly, P. C. D. (2017b). Divergent surface and total soil moisture projections under global  
58 warming. *Geophys. Res. Lett.* 44, 236–244. doi:10.1002/2016GL071921.
- 59 Bergaoui, K., Mitchell, D., Otto, F., Allen, M., Zaaboul, R., McDonnell, R., et al. (2015). The Contribution of Human-  
60 Induced Climate Change to the Drought of 2014 in the Southern Levant Region. *Bull. Am. Meteorol. Soc.* 96,  
61 S66–S70. doi:10.1175/BAMS-D-15-00129.1.

- 1 Berghuijs, W. R., Aalbers, E. E., Larsen, J. R., Trancoso, R., and Woods, R. A. (2017). Recent changes in extreme  
2 floods across multiple continents. *Environ. Res. Lett.* 12, 114035. doi:10.1088/1748-9326/aa8847.
- 3 Berghuijs, W. R., Woods, R. A., Hutton, C. J., and Sivapalan, M. (2016). Dominant flood generating mechanisms  
4 across the United States. *Geophys. Res. Lett.* 43, 4382–4390. doi:10.1002/2016GL068070.
- 5 Berthou, S., Kendon, E. J., Rowell, D. P., Roberts, M. J., Tucker, S., and Stratton, R. A. (2019a). Larger Future  
6 Intensification of Rainfall in the West African Sahel in a Convection-Permitting Model. *Geophys. Res. Lett.* 46,  
7 13299–13307. doi:10.1029/2019GL083544.
- 8 Berthou, S., Rowell, D. P., Kendon, E. J., Roberts, M. J., Stratton, R. A., Crook, J. A., et al. (2019b). Improved  
9 climatological precipitation characteristics over West Africa at convection-permitting scales. *Clim. Dyn.* 53,  
10 1991–2011. doi:10.1007/s00382-019-04759-4.
- 11 Bessho, K., Date, K., Hayashi, M., Ikeda, A., Imai, T., Inoue, H., et al. (2016). An Introduction to Himawari-8/9--  
12 Japan's New-Generation Geostationary Meteorological Satellites. *J. Meteorol. Soc. Japan. Ser. II* 94, 151–183.  
13 doi:10.2151/jmsj.2016-009.
- 14 Betts, R. A., Alfieri, L., Bradshaw, C., Caesar, J., Feyen, L., Friedlingstein, P., et al. (2018). Changes in climate  
15 extremes, fresh water availability and vulnerability to food insecurity projected at 1.5°C and 2°C global warming  
16 with a higher-resolution global climate model. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376.  
17 doi:10.1098/rsta.2016.0452.
- 18 Beusch, L., Gudmundsson, L., and Seneviratne, S. I. (2020). Emulating Earth system model temperatures with  
19 MESMER: From global mean temperature trajectories to grid-point-level realizations on land. *Earth Syst. Dyn.*  
20 11, 139–159. doi:10.5194/esd-11-139-2020.
- 21 Bevacqua, E., Maraun, D., Vousdoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., et al. (2019). Higher  
22 probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate  
23 change. *Sci. Adv.* 5, eaaw5531. doi:10.1126/sciadv.aaw5531.
- 24 Bevacqua, E., Vousdoukas, M. I., Shepherd, T. G., and Vrac, M. (2020a). Brief communication: The role of using  
25 precipitation or river discharge data when assessing global coastal compound flooding. *Nat. Hazards Earth Syst.*  
26 *Sci.* 20, 1765–1782. doi:10.5194/nhess-20-1765-2020.
- 27 Bevacqua, E., Vousdoukas, M. I., Zappa, G., Hodges, K., Shepherd, T. G., Maraun, D., et al. (2020b). More  
28 meteorological events that drive compound coastal flooding are projected under climate change. *Commun. Earth*  
29 *Environ.* 1, 47. doi:10.1038/s43247-020-00044-z.
- 30 Bevacqua, E., Zappa, G., and Shepherd, T. G. (2020c). Shorter cyclone clusters modulate changes in European  
31 wintertime precipitation extremes. *Environ. Res. Lett.* 8, 808–812. doi:10.1088/1748-9326/abbde7.
- 32 Bezerra, B. G., Silva, L. L., Santos e Silva, C. M., and de Carvalho, G. G. (2018). Changes of precipitation extremes  
33 indices in São Francisco River Basin, Brazil from 1947 to 2012. *Theor. Appl. Climatol.* 135, 565–576.  
34 doi:10.1007/s00704-018-2396-6.
- 35 Bhatia, K. T., Vecchi, G. A., Knutson, T. R., Murakami, H., Kossin, J., Dixon, K. W., et al. (2019). Recent increases in  
36 tropical cyclone intensification rates. *Nat. Commun.* 10, 1–9. doi:10.1038/s41467-019-08471-z.
- 37 Bhatia, K., Vecchi, G., Murakami, H., Underwood, S., and Kossin, J. (2018). Projected response of tropical cyclone  
38 intensity and intensification in a global climate model. *J. Clim.* 31, 8281–8303. doi:10.1175/JCLI-D-17-0898.1.
- 39 Bindoff, N. L., Stott, P. A., AchutaRao, K., Allen, M. R., Gillett, N., Gutzler, D., et al. (2013). “Detection and  
40 Attribution of Climate Change: from Global to Regional,” in *Climate Change 2013: The Physical Science Basis.*  
41 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
42 *Change*, eds. T. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, et al. (Cambridge, United  
43 Kingdom and New York, NY, USA: Cambridge University Press), 867–952.  
44 doi:10.1017/CBO9781107415324.022.
- 45 Biscarini, C., Francesco, S. D., Ridolfi, E., and Manciola, P. (2016). On the simulation of floods in a narrow bending  
46 valley: The malpasset dam break case study. *Water (Switzerland)* 8. doi:10.3390/w8110545.
- 47 Bitencourt, D. P., Fuentes, M. V., Maia, P. A., and Amorim, F. T. (2016). Frequência, Duração, Abrangência Espacial e  
48 Intensidadedas Ondas de Calor no Brasil. *Rev. Bras. Meteorol.* 31, 506–517. doi:10.1590/0102-  
49 778631231420150077.
- 50 Black, M. T., and Karoly, D. J. (2016). Southern Australia's Warmest October on Record: The Role of ENSO and  
51 Climate Change. *Bull. Am. Meteorol. Soc.* 97, S118–S121. doi:10.1175/BAMS-D-16-0124.1.
- 52 Blamey, R. C., Kolusu, S. R., Mahlalela, P., Todd, M. C., and Reason, C. J. C. (2018). The role of regional circulation  
53 features in regulating El Niño climate impacts over southern Africa: A comparison of the 2015/2016 drought with  
54 previous events. *Int. J. Climatol.* 38, 4276–4295. doi:10.1002/joc.5668.
- 55 Blanchet, J., Molinié, G., and Touati, J. (2018). Spatial analysis of trend in extreme daily rainfall in southern France.  
56 *Clim. Dyn.* 51, 799–812. doi:10.1007/s00382-016-3122-7.
- 57 Bloomfield, J. P., and Marchant, B. P. (2013). Analysis of groundwater drought building on the standardised  
58 precipitation index approach. *Hydrol. Earth Syst. Sci.* 17, 4769–4787. doi:10.5194/hess-17-4769-2013.
- 59 Bloomfield, J. P., Marchant, B. P., and McKenzie, A. A. (2019). Changes in groundwater drought associated with  
60 anthropogenic warming. *Hydrol. Earth Syst. Sci.* 23, 1393–1408. doi:10.5194/hess-23-1393-2019.
- 61 Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., et al. (2017). Changing climate shifts

- 1 timing of European floods. *Science* (80-. ). 357, 588–590. doi:10.1126/science.aan2506.
- 2 Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A. P., Parajka, J., Merz, B., et al. (2019). Changing climate both  
3 increases and decreases European river floods. *Nature* 573, 108–111. doi:10.1038/s41586-019-1495-6.
- 4 Blunden, J., and Arndt, D. S. (2016). State of the Climate in 2015. *Bull. Am. Meteorol. Soc.* 97, Si-S275.  
5 doi:10.1175/2016BAMSStateoftheClimate.1.
- 6 Blunden, J., and Arndt, D. S. (2017). State of the Climate in 2016. *Bull. Am. Meteorol. Soc.* 98, Si-S280.  
7 doi:10.1175/2017BAMSStateoftheClimate.1.
- 8 Boé, J., Somot, S., Corre, L., and Nabat, P. (2020). Large discrepancies in summer climate change over Europe as  
9 projected by global and regional climate models: causes and consequences. *Clim. Dyn.* 54, 2981–3002.  
10 doi:10.1007/s00382-020-05153-1.
- 11 Boers, N., Goswami, B., Rheinwalt, A., Bookhagen, B., Hoskins, B., and Kurths, J. (2019). Complex networks reveal  
12 global pattern of extreme-rainfall teleconnections. *Nature* 566, 373–377. doi:10.1038/s41586-018-0872-x.
- 13 Boisier, J. P., Alvarez-Garretón, C., Cordero, R. R., Damiani, A., Gallardo, L., Garreaud, R. D., et al. (2018).  
14 Anthropogenic drying in central-southern Chile evidenced by long-term observations and climate model  
15 simulations. *Elem Sci Anth* 6, 74. doi:10.1525/elementa.328.
- 16 Boisier, J. P., Rondanelli, R., Garreaud, R. D., and Muñoz, F. (2016). Anthropogenic and natural contributions to the  
17 Southeast Pacific precipitation decline and recent megadrought in central Chile. *Geophys. Res. Lett.* 43, 413–421.  
18 doi:10.1002/2015GL067265.
- 19 Bonan, G. B., Williams, M., Fisher, R. A., and Oleson, K. W. (2014). Modeling stomatal conductance in the earth  
20 system: linking leaf water-use efficiency and water transport along the soil–plant–atmosphere continuum. *Geosci.*  
21 *Model Dev.* 7, 2193–2222. doi:10.5194/gmd-7-2193-2014.
- 22 Bonfils, C. J. W., Santer, B. D., Fyfe, J. C., Marvel, K., Phillips, T. J., and Zimmerman, S. R. H. (2020). Human  
23 influence on joint changes in temperature, rainfall and continental aridity. *Nat. Clim. Chang.* 10, 726–731.  
24 doi:10.1038/s41558-020-0821-1.
- 25 Bonsal, B. R., Aider, R., Gachon, P., and Lapp, S. (2013). An assessment of Canadian prairie drought: Past, present,  
26 and future. *Clim. Dyn.* 41, 501–516. doi:10.1007/s00382-012-1422-0.
- 27 Bonsal, B. R., Peters, D. L., Seglenieks, F., Rivera, A., and Berg, A. (2019). Changes in freshwater availability across  
28 Canada. *Canada's Chang. Clim. Rep.*, 261–342.
- 29 Booth, J. F., Wang, S., and Polvani, L. (2013). Midlatitude storms in a moister world: lessons from idealized baroclinic  
30 life cycle experiments. *Clim. Dyn.* 41, 787–802. doi:10.1007/s00382-012-1472-3.
- 31 Borga, M., Stoffel, M., Marchi, L., Marra, F., and Jakob, M. (2014). Hydrogeomorphic response to extreme rainfall in  
32 headwater systems: Flash floods and debris flows. *J. Hydrol.* 518, 194–205. doi:10.1016/j.jhydrol.2014.05.022.
- 33 Borodina, A., Fischer, E. M., and Knutti, R. (2017a). Models are likely to underestimate increase in heavy rainfall in the  
34 extratropical regions with high rainfall intensity. *Geophys. Res. Lett.* 44, 7401–7409.  
35 doi:10.1002/2017GL074530.
- 36 Borodina, A., Fischer, E. M., and Knutti, R. (2017b). Potential to Constrain Projections of Hot Temperature Extremes.  
37 *J. Clim.* 30, 9949–9964. doi:10.1175/JCLI-D-16-0848.1.
- 38 Bosshard, T., Carambia, M., Goergen, K., Kotlarski, S., Krahe, P., Zappa, M., et al. (2013). Quantifying uncertainty  
39 sources in an ensemble of hydrological climate-impact projections. *Water Resour. Res.* 49, 1523–1536.  
40 doi:10.1029/2011WR011533.
- 41 Boyaj, A., Ashok, K., Ghosh, S., Devanand, A., and Dandu, G. (2018). The Chennai extreme rainfall event in 2015:  
42 The Bay of Bengal connection. *Clim. Dyn.* 50, 2867–2879. doi:10.1007/s00382-017-3778-7.
- 43 Bozkurt, D., Rojas, M., Boisier, J. P., Rondanelli, R., Garreaud, R., and Gallardo, L. (2019). Dynamical downscaling  
44 over the complex terrain of southwest South America: present climate conditions and added value analysis. *Clim.*  
45 *Dyn.* 53, 6745–6767. doi:10.1007/s00382-019-04959-y.
- 46 Bozkurt, D., Rojas, M., Boisier, J. P., and Valdivieso, J. (2018). Projected hydroclimate changes over Andean basins in  
47 central Chile from downscaled CMIP5 models under the low and high emission scenarios. *Clim. Change* 150,  
48 131–147. doi:10.1007/s10584-018-2246-7.
- 49 Brando, P. M., Paolucci, L., Ummenhofer, C. C., Ordway, E. M., Hartmann, H., Cattau, M. E., et al. (2019). Droughts,  
50 Wildfires, and Forest Carbon Cycling: A Pantropical Synthesis. *Annu. Rev. Earth Planet. Sci.* 47, 555–581.  
51 doi:10.1146/annurev-earth-082517-010235.
- 52 Brandon, C. M., Woodruff, J. D., Lane, D. P., and Donnelly, J. P. (2013). Tropical cyclone wind speed constraints from  
53 resultant storm surge deposition: A 2500 year reconstruction of hurricane activity from St. Marks, FL.  
54 *Geochemistry, Geophys. Geosystems* 14, 2993–3008. doi:10.1002/ggge.20217.
- 55 Bregy, J. C., Wallace, D. J., Minzoni, R. T., and Cruz, V. J. (2018). 2500-year paleotempestological record of intense  
56 storms for the northern Gulf of Mexico, United States. *Mar. Geol.* 396, 26–42.  
57 doi:10.1016/j.margeo.2017.09.009.
- 58 Breña-Naranjo, J., Laverde B., M., and Pedrozo-Acuña, A. (2016). Changes in pan evaporation in Mexico from 1961 to  
59 2010. *Int. J. Climatol.* 37. doi:10.1002/joc.4698.
- 60 Breshears, D. D., Adams, H. D., Eamus, D., Mcdowell, N. G., Law, D. J., Will, R. E., et al. (2013). The critical  
61 amplifying role of increasing atmospheric moisture demand on tree mortality and associated regional die-off.

- 1 *Front. Plant Sci.* 4. doi:10.3389/fpls.2013.00266.
- 2 Bright, R. M., Davin, E., O'Halloran, T., Pongratz, J., Zhao, K., and Cescatti, A. (2017). Local temperature response to  
3 land cover and management change driven by non-radiative processes. *Nat. Clim. Chang.* 7, 296–302.  
4 doi:10.1038/nclimate3250.
- 5 Brito, S. S. B., Cunha, A. P. M. A., Cunningham, C. C., Alvalá, R. C., Marengo, J. A., and Carvalho, M. A. (2018).  
6 Frequency, duration and severity of drought in the Semiarid Northeast Brazil region. *Int. J. Climatol.* 38, 517–  
7 529. doi:10.1002/joc.5225.
- 8 Brocca, L., Hasenauer, S., Lacava, T., Melone, F., Moramarco, T., Wagner, W., et al. (2011). Soil moisture estimation  
9 through ASCAT and AMSR-E sensors: An intercomparison and validation study across Europe. *Remote Sens.*  
10 *Environ.* 115, 3390–3408. doi:10.1016/j.rse.2011.08.003.
- 11 Brodrribb, T. J., Powers, J., Cochard, H., and Choat, B. (2020). Hanging by a thread? Forests and drought. *Science* (80-  
12 ). 368, 261–266. doi:10.1126/science.aat7631.
- 13 Brohan, P., Compo, G. P., Brönnimann, S., Allan, R. J., Auchmann, R., Brugnara, Y., et al. (2016). The 1816 ‘year  
14 without a summer’ in an atmospheric reanalysis. *Clim. Past Discuss.*, 1–11. doi:10.5194/cp-2016-78.
- 15 Brooks, H. E. (2013). Severe thunderstorms and climate change. *Atmos. Res.* 123, 129–138.  
16 doi:10.1016/j.atmosres.2012.04.002.
- 17 Brooks, H. E., Carbin, G. W., and Marsh, P. T. (2014). Increased variability of tornado occurrence in the United States.  
18 *Science* (80- ). 346, 349 LP-352. doi:10.1126/science.1257460.
- 19 Brown, S. J. (2020). Future changes in heatwave severity, duration and frequency due to climate change for the most  
20 populous cities. *Weather Clim. Extrem.* 30, 100278. doi:10.1016/j.wace.2020.100278.
- 21 Brunner, L., Hegerl, G. C., and Steiner, A. K. (2017). Connecting atmospheric blocking to European temperature  
22 extremes in spring. *J. Clim.* 30, 585–594. doi:10.1175/JCLI-D-16-0518.1.
- 23 Brunner, L., Schaller, N., Anstey, J., Sillmann, J., and Steiner, A. K. (2018). Dependence of Present and Future  
24 European Temperature Extremes on the Location of Atmospheric Blocking. *Geophys. Res. Lett.* 45, 6311–6320.  
25 doi:10.1029/2018GL077837.
- 26 Brunner, M. I., and Tallaksen, L. M. (2019). Proneness of European Catchments to Multiyear Streamflow Droughts.  
27 *Water Resour. Res.* doi:10.1029/2019WR025903.
- 28 Bucchignani, E., Zollo, A. L., Cattaneo, L., Montesarchio, M., and Mercogliano, P. (2017). Extreme weather events  
29 over China: assessment of COSMO-CLM simulations and future scenarios. *Int. J. Climatol.* 37, 1578–1594.  
30 doi:10.1002/joc.4798.
- 31 Büntgen, U., Tegel, W., Carrer, M., Krusic, P. J., Hayes, M., and Esper, J. (2015). Commentary to Wetter et al. (2014):  
32 Limited tree-ring evidence for a 1540 European ‘Megadrought.’ *Clim. Change* 131, 183–190.  
33 doi:10.1007/s10584-015-1423-1.
- 34 Burdanowitz, J., Buehler, S. A., Bakan, S., and Klepp, C. (2019). The sensitivity of oceanic precipitation to sea surface  
35 temperature. *Atmos. Chem. Phys.* 19, 9241–9252. doi:10.5194/acp-19-9241-2019.
- 36 Burgman, R. J., and Jang, Y. (2015). Simulated U.S. drought response to interannual and decadal pacific SST  
37 variability. *J. Clim.* 28, 4688–4705. doi:10.1175/JCLI-D-14-00247.1.
- 38 Burke, C., Stott, P., Ciavarella, A., and Sun, Y. (2016). Attribution of Extreme Rainfall in Southeast China During May  
39 2015. *Bull. Am. Meteorol. Soc.* 97, S92–S96. doi:10.1175/BAMS-D-16-0144.1.
- 40 Burn, D. H., and Whitfield, P. H. (2016). Changes in floods and flood regimes in Canada. *Can. Water Resour. J.* 41,  
41 139–150. doi:10.1080/07011784.2015.1026844.
- 42 Buttle, J. M., Allen, D. M., Caissie, D., Davison, B., Hayashi, M., Peters, D. L., et al. (2016). Flood processes in  
43 Canada: Regional and special aspects. *Can. Water Resour. J.* 41, 7–30. doi:10.1080/07011784.2015.1131629.
- 44 Byrne, M. P., and O’Gorman, P. A. (2015). The Response of Precipitation Minus Evapotranspiration to Climate  
45 Warming: Why the “Wet-Get-Wetter, Dry-Get-Drier” Scaling Does Not Hold over Land\*. *J. Clim.* 28, 8078–  
46 8092. doi:10.1175/JCLI-D-15-0369.1.
- 47 Byrne, M. P., and O’Gorman, P. A. (2018). Trends in continental temperature and humidity directly linked to ocean  
48 warming. *Proc. Natl. Acad. Sci.* 115, 4863–4868. doi:10.1073/pnas.1722312115.
- 49 Cabré, M. F., Solman, S., and Núñez, M. (2016). Regional climate change scenarios over southern South America for  
50 future climate (2080-2099) using the MM5 Model. Mean, interannual variability and uncertainties. *Atmósfera* 29,  
51 35–60. doi:10.20937/ATM.2016.29.01.04.
- 52 Cai, W., Borlace, S., Lengaigne, M., Van Rensch, P., Collins, M., Vecchi, G., et al. (2014a). Increasing frequency of  
53 extreme El Niño events due to greenhouse warming. *Nat. Clim. Chang.* 4, 111–116. doi:10.1038/nclimate2100.
- 54 Cai, W., and Cowan, T. (2008). Evidence of impacts from rising temperature on inflows to the Murray-Darling Basin.  
55 *Geophys. Res. Lett.* 35, 2–6. doi:10.1029/2008GL033390.
- 56 Cai, W., Purich, A., Cowan, T., Van Rensch, P., and Weller, E. (2014b). Did climate change-induced rainfall trends  
57 contribute to the Australian millennium drought? *J. Clim.* 27, 3145–3168. doi:10.1175/JCLI-D-13-00322.1.
- 58 Cai, W., Santoso, A., Wang, G., Yeh, S. W., An, S. II, Cobb, K. M., et al. (2015). ENSO and greenhouse warming. *Nat.*  
59 *Clim. Chang.* 5, 849–859. doi:10.1038/nclimate2743.
- 60 Cai, W., Wang, G., Gan, B., Wu, L., Santoso, A., Lin, X., et al. (2018). Stabilised frequency of extreme positive Indian  
61 Ocean Dipole under 1.5°C warming. *Nat. Commun.* 9, 4–11. doi:10.1038/s41467-018-03789-6.

- 1 Caillouet, L., Vidal, J.-P., Sauquet, E., Devers, A., and Graff, B. (2017). Ensemble reconstruction of spatio-temporal  
2 extreme low-flow events in France since 1871. *Hydrol. Earth Syst. Sci.* 21, 2923–2951. doi:10.5194/hess-21-  
3 2923-2017.
- 4 Callaghan, J., and Power, S. B. (2011). Variability and decline in the number of severe tropical cyclones making land-  
5 fall over eastern Australia since the late nineteenth century. *Clim. Dyn.* 37, 647–662. doi:10.1007/s00382-010-  
6 0883-2.
- 7 Caloiero, T. (2015). Analysis of rainfall trend in New Zealand. *Environ. Earth Sci.* 73, 6297–6310.  
8 doi:10.1007/s12665-014-3852-y.
- 9 Caloiero, T. (2017). Trend of monthly temperature and daily extreme temperature during 1951–2012 in New Zealand.  
10 *Theor. Appl. Climatol.* 129, 111–127. doi:10.1007/s00704-016-1764-3.
- 11 Caloiero, T., Veltri, S., Caloiero, P., and Frustaci, F. (2018). Drought Analysis in Europe and in the Mediterranean  
12 Basin Using the Standardized Precipitation Index. *Water* 10, 1043. doi:10.3390/w10081043.
- 13 Camargo, S. J. (2013). Global and regional aspects of tropical cyclone activity in the CMIP5 models. *J. Clim.* 26, 9880–  
14 9902. doi:10.1175/JCLI-D-12-00549.1.
- 15 Camargo, S. J., Emanuel, K. A., and Sobel, A. H. (2007). Use of a Genesis Potential Index to Diagnose ENSO Effects  
16 on Tropical Cyclone Genesis. *J. Clim.* 20, 4819–4834. doi:10.1175/JCLI4282.1.
- 17 Camargo, S. J., Giulivi, C. F., Sobel, A. H., Wing, A. A., Kim, D., Moon, Y., et al. (2020). Characteristics of Model  
18 Tropical Cyclone Climatology and the Large-Scale Environment. *J. Clim.* 33, 4463–4487. doi:10.1175/JCLI-D-  
19 19-0500.1.
- 20 Camargo, S. J., Tippett, M. K., Sobel, A. H., Vecchi, G. A., and Zhao, M. (2014). Testing the performance of tropical  
21 cyclone genesis indices in future climates using the HiRAM model. *J. Clim.* 27, 9171–9196. doi:10.1175/JCLI-D-  
22 13-00505.1.
- 23 Camargo, S. J., Wheeler, M. C., and Sobel, A. H. (2009). Diagnosis of the MJO modulation of tropical cyclogenesis  
24 using an empirical index. *J. Atmos. Sci.* 66, 3061–3074. doi:10.1175/2009JAS3101.1.
- 25 Camargo, S. J., and Wing, A. A. (2016). Tropical cyclones in climate models. *Wiley Interdiscip. Rev. Clim. Chang.* 7,  
26 211–237. doi:10.1002/wcc.373.
- 27 Camuffo, D., Bertolin, C., Diodato, N., Cocheo, C., Barriendos, M., Dominguez-Castro, F., et al. (2013). Western  
28 Mediterranean precipitation over the last 300 years from instrumental observations. *Clim. Change* 117, 85–101.  
29 doi:10.1007/s10584-012-0539-9.
- 30 Cannon, A. J., and Innocenti, S. (2019). Projected intensification of sub-daily and daily rainfall extremes in convection-  
31 permitting climate model simulations over North America: implications for future intensity–duration–frequency  
32 curves. *Nat. Hazards Earth Syst. Sci.* 19, 421–440. doi:10.5194/nhess-19-421-2019.
- 33 Cardell, M. F., Amengual, A., Romero, R., and Ramis, C. (2020). Future extremes of temperature and precipitation in  
34 Europe derived from a combination of dynamical and statistical approaches. *Int. J. Climatol.* 40, 4800–4827.  
35 doi:10.1002/joc.6490.
- 36 Cardoso, R. M., Soares, P. M. M., Lima, D. C. A., and Miranda, P. M. A. (2019). Mean and extreme temperatures in a  
37 warming climate: EURO CORDEX and WRF regional climate high-resolution projections for Portugal. *Clim.*  
38 *Dyn.* 52, 129–157. doi:10.1007/s00382-018-4124-4.
- 39 Carril, A., Cavalcanti, I., Menéndez, C., Sörensson, A., López-Franca, N., Rivera, J., et al. (2016). Extreme events in  
40 the La Plata basin: a retrospective analysis of what we have learned during CLARIS-LPB project. *Clim. Res.* 68,  
41 95–116. doi:10.3354/cr01374.
- 42 Carvalho, L. M. V. (2020). Assessing precipitation trends in the Americas with historical data: A review. *WIREs Clim.*  
43 *Chang.* 11, e627. doi:10.1002/wcc.627.
- 44 Casanueva, A., Rodríguez-Puebla, C., Frías, M. D., and González-Reviriego, N. (2014). Variability of extreme  
45 precipitation over Europe and its relationships with teleconnection patterns. *Hydrol. Earth Syst. Sci.* 18, 709–725.  
46 doi:10.5194/hess-18-709-2014.
- 47 Cattiaux, J., and Ribes, A. (2018). Defining Single Extreme Weather Events in a Climate Perspective. *Bull. Am.*  
48 *Meteorol. Soc.* 99, 1557–1568. doi:10.1175/BAMS-D-17-0281.1.
- 49 Catto, J. L., Jakob, C., and Nicholls, N. (2015). Can the CMIP5 models represent winter frontal precipitation? *Geophys.*  
50 *Res. Lett.* doi:10.1002/2015GL066015.
- 51 Catto, J. L., and Pfahl, S. (2013). The importance of fronts for extreme precipitation. *J. Geophys. Res. Atmos.* 118,  
52 10791–10801. doi:10.1002/jgrd.50852.
- 53 Cavalcanti, I. F. A., Carril, A. F., Penalba, O. C., Grimm, A. M., Menéndez, C. G., Sanchez, E., et al. (2015).  
54 Precipitation extremes over La Plata Basin – Review and new results from observations and climate simulations.  
55 *J. Hydrol.* 523, 211–230. doi:10.1016/j.jhydrol.2015.01.028.
- 56 Cavanaugh, N. R., and Shen, S. S. P. (2015). The Effects of Gridding Algorithms on the Statistical Moments and Their  
57 Trends of Daily Surface Air Temperature. *J. Clim.* 28, 9188–9205. doi:10.1175/JCLI-D-14-00668.1.
- 58 Cavicchia, L., von Storch, H., and Gualdi, S. (2014). Mediterranean Tropical-Like Cyclones in Present and Future  
59 Climate. *J. Clim.* 27, 7493–7501. doi:10.1175/JCLI-D-14-00339.1.
- 60 Ceccherini, G., Russo, S., Amezttoy, I., Marchese, A. F., and Carmona-Moreno, C. (2017). Heat waves in Africa 1981–  
61 2015, observations and reanalysis. *Nat. Hazards Earth Syst. Sci.* 17, 115–125. doi:10.5194/nhess-17-115-2017.



- 1 Ceccherini, G., Russo, S., Amezttoy, I., Romero, C. P., and Carmona-Moreno, C. (2016). Magnitude and frequency of  
2 heat and cold waves in recent decades: the case of South America. *Nat. Hazards Earth Syst. Sci.* 16, 821–831.  
3 doi:10.5194/nhess-16-821-2016.
- 4 Cha, E. J., Knutson, T. R., Lee, T.-C., Ying, M., and Nakaegawa, T. (2020). Third assessment on impacts of climate  
5 change on tropical cyclones in the Typhoon Committee Region – Part II: Future projections. *Trop. Cyclone Res.*  
6 *Rev.* 9, 75–86. doi:https://doi.org/10.1016/j.tcr.2020.04.005.
- 7 Chai, R., Sun, S., Chen, H., and Zhou, S. (2018). Changes in reference evapotranspiration over China during 1960–  
8 2012: Attributions and relationships with atmospheric circulation. *Hydrol. Process.* 32, 3032–3048.  
9 doi:10.1002/hyp.13252.
- 10 Chakraborty, D., Sehgal, V. K., Dhakar, R., Varghese, E., Das, D. K., and Ray, M. (2018). Changes in daily maximum  
11 temperature extremes across India over 1951–2014 and their relation with cereal crop productivity. *Stoch.*  
12 *Environ. Res. Risk Assess.* 32. doi:10.1007/s00477-018-1604-3.
- 13 Chan, K. T. F., and Chan, J. C. L. (2015). Global climatology of tropical cyclone size as inferred from QuikSCAT data.  
14 *Int. J. Climatol.* 35, 4843–4848. doi:10.1002/joc.4307.
- 15 Chan, K. T. F., and Chan, J. C. L. (2018). The Outer-Core Wind Structure of Tropical Cyclones. *J. Meteorol. Soc.*  
16 *Japan. Ser. II* 96, 297–315. doi:10.2151/jmsj.2018-042.
- 17 Chan, S. C., Kendon, E. J., Berthou, S., Fosser, G., Lewis, E., and Fowler, H. J. (2020). Europe-wide precipitation  
18 projections at convection permitting scale with the Unified Model. *Clim. Dyn.* 55, 409–428. doi:10.1007/s00382-  
19 020-05192-8.
- 20 Chand, S. S., Dowdy, A. J., Ramsay, H. A., Walsh, K. J. E., Tory, K. J., Power, S. B., et al. (2019). Review of tropical  
21 cyclones in the Australian region: Climatology, variability, predictability, and trends. *WIREs Clim. Chang.* 10,  
22 e602. doi:10.1002/wcc.602.
- 23 Chaney, N. W., Sheffield, J., Villarini, G., and Wood, E. F. (2014). Development of a high-resolution gridded daily  
24 meteorological dataset over sub-Saharan Africa: Spatial analysis of trends in climate extremes. *J. Clim.*  
25 doi:10.1175/JCLI-D-13-00423.1.
- 26 Chang, E. K. M. (2014). Impacts of background field removal on CMIP5 projected changes in Pacific winter cyclone  
27 activity. *J. Geophys. Res. Atmos.* 119, 4626–4639. doi:10.1038/175238c0.
- 28 Chang, E. K. M. (2017). Projected Significant Increase in the Number of Extreme Extratropical Cyclones in the  
29 Southern Hemisphere. *J. Clim.* 30, 4915–4935. doi:10.1175/JCLI-D-16-0553.1.
- 30 Chang, E. K. M., Ma, C. G., Zheng, C., and Yau, A. M. W. (2016). Observed and projected decrease in Northern  
31 Hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature. *Geophys. Res.*  
32 *Lett.* doi:10.1002/2016GL068172.
- 33 Chang, E. K. M., and Yau, A. M. W. (2016). Northern Hemisphere winter storm track trends since 1959 derived from  
34 multiple reanalysis datasets. *Clim. Dyn.* 47, 1435–1454. doi:10.1007/s00382-015-2911-8.
- 35 Chapman, S., Watson, J. E. M., Salazar, A., Thatcher, M., and McAlpine, C. A. (2017). The impact of urbanization and  
36 climate change on urban temperatures: a systematic review. *Landsc. Ecol.* 32, 1921–1935. doi:10.1007/s10980-  
37 017-0561-4.
- 38 Chavas, D. R., and Emanuel, K. (2014). Equilibrium tropical cyclone size in an idealized state of axisymmetric  
39 radiative-convective equilibrium. *J. Atmos. Sci.* 71, 1663–1680. doi:10.1175/JAS-D-13-0155.1.
- 40 Chen, C.-T., and Knutson, T. (2008). On the Verification and Comparison of Extreme Rainfall Indices from Climate  
41 Models. *J. Clim.* 21, 1605–1621. doi:10.1175/2007JCLI1494.1.
- 42 Chen, D., Guo, J., Yao, D., Lin, Y., Zhao, C., Min, M., et al. (2019). Mesoscale Convective Systems in the Asian  
43 Monsoon Region From Advanced Himawari Imager: Algorithms and Preliminary Results. *J. Geophys. Res.*  
44 *Atmos.* 124, 2210–2234. doi:10.1029/2018JD029707.
- 45 Chen, G.-S., Notaro, M., Liu, Z., and Liu, Y. (2012). Simulated Local and Remote Biophysical Effects of Afforestation  
46 over the Southeast United States in Boreal Summer. *J. Clim.* 25, 4511–4522. doi:10.1175/JCLI-D-11-00317.1.
- 47 Chen, H., and Sun, J. (2015a). Assessing model performance of climate extremes in China: an intercomparison between  
48 CMIP5 and CMIP3. *Clim. Change* 129, 197–211. doi:10.1007/s10584-014-1319-5.
- 49 Chen, H., and Sun, J. (2015b). Changes in Drought Characteristics over China Using the Standardized Precipitation  
50 Evapotranspiration Index. *J. Clim.* 28, 5430–5447. doi:10.1175/JCLI-D-14-00707.1.
- 51 Chen, H., and Sun, J. (2017a). Anthropogenic warming has caused hot droughts more frequently in China. *J. Hydrol.*  
52 544, 306–318. doi:10.1016/j.jhydrol.2016.11.044.
- 53 Chen, H., and Sun, J. (2017b). Characterizing present and future drought changes over eastern China. *Int. J. Climatol.*  
54 37, 138–156. doi:10.1002/joc.4987.
- 55 Chen, H., and Sun, J. (2017c). Contribution of human influence to increased daily precipitation extremes over China.  
56 *Geophys. Res. Lett.* doi:10.1002/2016GL072439.
- 57 Chen, H., Sun, J., Lin, W., and Xu, H. (2020a). Comparison of CMIP6 and CMIP5 models in simulating climate  
58 extremes. *Sci. Bull.* 65, 1415–1418. doi:10.1016/j.scib.2020.05.015.
- 59 Chen, J., Dai, A., Zhang, Y., and Rasmussen, K. L. (2020b). Changes in Convective Available Potential Energy and  
60 Convective Inhibition under Global Warming. *J. Clim.* 33, 2025–2050. doi:10.1175/JCLI-D-19-0461.1.
- 61 Chen, J., Wang, Z., Tam, C.-Y., Lau, N.-C., Lau, D.-S. D., and Mok, H.-Y. (2020c). Impacts of climate change on

- 1 tropical cyclones and induced storm surges in the Pearl River Delta region using pseudo-global-warming method.  
2 *Sci. Rep.* 10, 1965. doi:10.1038/s41598-020-58824-8.
- 3 Chen, L., and Dirmeyer, P. A. (2019). Global observed and modelled impacts of irrigation on surface temperature. *Int.*  
4 *J. Climatol.* 39, 2587–2600. doi:10.1002/joc.5973.
- 5 Chen, L., Li, D., and Pryor, S. C. (2013). Wind speed trends over China: Quantifying the magnitude and assessing  
6 causality. *Int. J. Climatol.* 33, 2579–2590. doi:10.1002/joc.3613.
- 7 Chen, R., Wen, Z., and Lu, R. (2016). Evolution of the circulation anomalies and the quasi-biweekly oscillations  
8 associated with extreme heat events in Southern China. *J. Clim.* 29, 6909–6921. doi:10.1175/JCLI-D-16-0160.1.
- 9 Chen, S., Li, Y., Kim, J., and Kim, S. W. (2017). Bayesian change point analysis for extreme daily precipitation. *Int. J.*  
10 *Climatol.* 37, 3123–3137. doi:10.1002/joc.4904.
- 11 Chen, X., Zhao, P., Ouyang, L., Zhu, L., Ni, G., and Schäfer, K. V. R. (2020d). Whole-plant water hydraulic integrity to  
12 predict drought-induced Eucalyptus urophylla mortality under drought stress. *For. Ecol. Manage.* 468.  
13 doi:10.1016/j.foreco.2020.118179.
- 14 Chen, Y., Li, W., Jiang, X., Zhai, P., and Luo, Y. (2021). Detectable Intensification of Hourly and Daily Scale  
15 Precipitation Extremes across Eastern China. *J. Clim.* 34, 1185–1201. doi:10.1175/JCLI-D-20-0462.1.
- 16 Chen, Y., and Zhai, P. (2017). Revisiting summertime hot extremes in China during 1961–2015: Overlooked compound  
17 extremes and significant changes. *Geophys. Res. Lett.* 44, 5096–5103. doi:10.1002/2016GL072281.
- 18 Cheng, L., Hoerling, M., Smith, L., and Eischeid, J. (2018). Diagnosing Human-Induced Dynamic and Thermodynamic  
19 Drivers of Extreme Rainfall. *J. Clim.* 31, 1029–1051. doi:10.1175/JCLI-D-16-0919.1.
- 20 Cheng, S., Guan, X., Huang, J., Ji, F., and Guo, R. (2015). Long-term trend and variability of soil moisture over East  
21 Asia. *J. Geophys. Res. Atmos.* 120, 8658–8670. doi:10.1002/2015JD023206.
- 22 Cheong, W. K., Timbal, B., Golding, N., Sirabaha, S., Kwan, K. F., Cinco, T. A., et al. (2018). Observed and modelled  
23 temperature and precipitation extremes over Southeast Asia from 1972 to 2010. *Int. J. Climatol.* 38, 3013–3027.  
24 doi:10.1002/joc.5479.
- 25 Chevuturi, A., Klingaman, N. P., Turner, A. G., and Hannah, S. (2018). Projected Changes in the Asian-Australian  
26 Monsoon Region in 1.5°C and 2.0°C Global-Warming Scenarios. *Earth's Futur.* 6, 339–358.  
27 doi:10.1002/2017EF000734.
- 28 Cho, C., Li, R., Wang, S.-Y., Yoon, J.-H., and Gillies, R. R. (2016). Anthropogenic footprint of climate change in the  
29 June 2013 northern India flood. *Clim. Dyn.* 46, 797–805. doi:10.1007/s00382-015-2613-2.
- 30 Chou, S. C., Lyra, A., Mourão, C., Dereczynski, C., Pilotto, I., Gomes, J., et al. (2014a). Assessment of Climate Change  
31 over South America under RCP 4.5 and 8.5 Downscaling Scenarios. *Am. J. Clim. Chang.* 03, 512–527.  
32 doi:10.4236/ajcc.2014.35043.
- 33 Chou, S. C., Lyra, A., Mourão, C., Dereczynski, C., Pilotto, I., Gomes, J., et al. (2014b). Evaluation of the Eta  
34 Simulations Nested in Three Global Climate Models. *Am. J. Clim. Chang.* 03, 438–454.  
35 doi:10.4236/ajcc.2014.35039.
- 36 Choy, C., Chong, S., Kong, D., and Cayan, E. O. (2015). A Discussion of the Most Intense Tropical Cyclones in the  
37 Western North Pacific From 1978 to 2013. *Trop. Cyclone Res. Rev.* 4, 1–11.  
38 doi:https://doi.org/10.6057/2015TCRR01.01.
- 39 Christensen, J. H., Kumar Krishna, K., Aldrian, E., An, S.-I., Cavalcanti, I. F. A., Castro, M. de, et al. (2013). “Climate  
40 Phenomena and their Relevance for Future Regional Climate Change,” in *Climate Change 2013: The Physical*  
41 *Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel*  
42 *on Climate Change*, eds. T. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, et al.  
43 (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1217–1308.  
44 doi:10.1017/CBO9781107415324.028.
- 45 Christiansen, B., Alvarez-Castro, C., Christidis, N., Ciavarella, A., Colfescu, I., Cowan, T., et al. (2018). Was the Cold  
46 European Winter of 2009/10 Modified by Anthropogenic Climate Change? An Attribution Study. *J. Clim.* 31,  
47 3387–3410. doi:10.1175/JCLI-D-17-0589.1.
- 48 Christidis, N., Ciavarella, A., and Stott, P. A. (2018). Different Ways of Framing Event Attribution Questions: The  
49 Example of Warm and Wet Winters in the United Kingdom Similar to 2015/16. *J. Clim.* 31, 4827–4845.  
50 doi:10.1175/JCLI-D-17-0464.1.
- 51 Christidis, N., Jones, G. S., and Stott, P. A. (2015). Dramatically increasing chance of extremely hot summers since the  
52 2003 European heatwave. *Nat. Clim. Chang.* 5, 46–50. doi:10.1038/nclimate2468.
- 53 Christidis, N., and Stott, P. A. (2014). Change in the Odds of Warm Years and Seasons Due to Anthropogenic Influence  
54 on the Climate. *J. Clim.* 27, 2607–2621.
- 55 Christidis, N., and Stott, P. A. (2016). Attribution analyses of temperature extremes using a set of 16 indices. *Weather*  
56 *Clim. Extrem.* 14, 24–35. doi:10.1016/j.wace.2016.10.003.
- 57 Christidis, N., Stott, P. A., and Ciavarella, A. (2014). The effect of anthropogenic climate change on the cold spring of  
58 2013 in the United Kingdom. *Bull. Am. Meteorol. Soc.* 95, S15–S18. doi:10.1175/1520-0477-95.9.S1.1.
- 59 Christidis, N., Stott, P. A., Karoly, D. J., and Ciavarella, A. (2013a). An Attribution Study of the Heavy Rainfall Over  
60 Eastern Australia in March 2012 [in “Explaining Extreme Events of 2012 from a Climate Perspective”]. *Bull. Am.*  
61 *Meteorol. Soc.* 94, S58–S61. doi:10.1175/BAMS-D-13-00085.1.

- 1 Christidis, N., Stott, P. A., Scaife, A. A., Arribas, A., Jones, G. S., Copsey, D., et al. (2013b). A New HadGEM3-A-  
2 Based System for Attribution of Weather- and Climate-Related Extreme Events. *J. Clim.* 26, 2756–2783.  
3 doi:10.1175/JCLI-D-12-00169.1.
- 4 Cioffi, F., Lall, U., Rus, E., and Krishnamurthy, C. K. B. (2015). Space-time structure of extreme precipitation in  
5 Europe over the last century. *Int. J. Climatol.* 35, 1749–1760. doi:10.1002/joc.4116.
- 6 Clark, R. T., and Brown, S. J. (2013). Influences of Circulation and Climate Change on European Summer Heat  
7 Extremes. *J. Clim.* 26, 9621–9632. doi:10.1175/JCLI-D-12-00740.1.
- 8 Clark, R. T., Brown, S. J., and Murphy, J. M. (2006). Modeling Northern Hemisphere Summer Heat Extreme Changes  
9 and Their Uncertainties Using a Physics Ensemble of Climate Sensitivity Experiments. *J. Clim.* 19, 4418–4435.  
10 doi:10.1175/JCLI3877.1.
- 11 Clarke, H., and Evans, J. P. (2019). Exploring the future change space for fire weather in southeast Australia. *Theor.*  
12 *Appl. Climatol.* 136, 513–527. doi:10.1007/s00704-018-2507-4.
- 13 Colle, B. A., Zhang, Z., Lombardo, K. A., Chang, E., Liu, P., and Zhang, M. (2013). Historical evaluation and future  
14 prediction of eastern North American and Western Atlantic extratropical cyclones in the CMIP5 models during  
15 the cool season. *J. Clim.* doi:10.1175/JCLI-D-12-00498.1.
- 16 Collins, J. M., Klotzbach, P. J., Maue, R. N., Roache, D. R., Blake, E. S., Paxton, C. H., et al. (2016). The record-  
17 breaking 2015 hurricane season in the eastern North Pacific: An analysis of environmental conditions. *Geophys.*  
18 *Res. Lett.* 43, 9217–9224. doi:10.1002/2016GL070597.
- 19 Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., et al. (2013). “Long-term Climate  
20 Change: Projections, Commitments and Irreversibility,” in *Climate Change 2013: The Physical Science Basis.*  
21 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
22 *Change*, eds. T. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, et al. (Cambridge, United  
23 Kingdom and New York, NY, USA: Cambridge University Press), 1029–1136.  
24 doi:10.1017/CBO9781107415324.024.
- 25 Collins, M., Sutherland, M., Bouwer, L., Cheong, S.-M., and Frolicher, T. (2019). “Extremes, Abrupt Changes and  
26 Managing Risks,” in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, eds. H.-O.  
27 Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, et al. (In Press), 589–656.  
28 Available at: <https://www.ipcc.ch/srocc/chapter/chapter-6>.
- 29 Condon, L. E., Atchley, A. L., and Maxwell, R. M. (2020). Evapotranspiration depletes groundwater under warming  
30 over the contiguous United States. *Nat. Commun.* 11. doi:10.1038/s41467-020-14688-0.
- 31 Contractor, S., Donat, M. G., and Alexander, L. V (2020a). Changes in observed daily precipitation over global land  
32 areas since 1950. *J. Clim.*, 1–53. doi:10.1175/JCLI-D-19-0965.1.
- 33 Contractor, S., Donat, M. G., Alexander, L. V, Ziese, M., Meyer-Christoffer, A., Schneider, U., et al. (2020b). Rainfall  
34 Estimates on a Gridded Network (REGEN) – a global land-based gridded dataset of daily precipitation from 1950  
35 to 2016. *Hydrol. Earth Syst. Sci.* 24, 919–943. doi:10.5194/hess-24-919-2020.
- 36 Cook, B. I., Anchukaitis, K. J., Touchan, R., Meko, D. M., and Cook, E. R. (2016a). Spatiotemporal drought variability  
37 in the Mediterranean over the last 900 years. *J. Geophys. Res. Atmos.* 121, 2060–2074.  
38 doi:10.1002/2015JD023929.
- 39 Cook, B. I., Ault, T. R., and Smerdon, J. E. (2015). Unprecedented 21st century drought risk in the American Southwest  
40 and Central Plains. *Sci. Adv.* 1, e1400082. doi:10.1126/sciadv.1400082.
- 41 Cook, B. I., Cook, E. R., Smerdon, J. E., Seager, R., Williams, A. P., Coats, S., et al. (2016b). North American  
42 megadroughts in the Common Era: Reconstructions and simulations. *Wiley Interdiscip. Rev. Clim. Chang.* 7,  
43 411–432. doi:10.1002/wcc.394.
- 44 Cook, B. I., Mankin, J. S., and Anchukaitis, K. J. (2018). Climate Change and Drought: From Past to Future. *Curr.*  
45 *Clim. Chang. Reports* 4, 164–179. doi:10.1007/s40641-018-0093-2.
- 46 Cook, B. I., Mankin, J. S., Marvel, K., Williams, A. P., Smerdon, J. E., and Anchukaitis, K. J. (2020). Twenty-First  
47 Century Drought Projections in the CMIP6 Forcing Scenarios. *Earth’s Futur.* 8, e2019EF001461.  
48 doi:10.1029/2019EF001461.
- 49 Cook, B. I., Palmer, J. G., Cook, E. R., Turney, C. S. M., Allen, K., Fenwick, P., et al. (2016c). The paleoclimate  
50 context and future trajectory of extreme summer hydroclimate in eastern Australia. *J. Geophys. Res.* 121, 812-  
51 820,838.
- 52 Cook, B. I., Seager, R., Williams, A. P., Puma, M. J., McDermid, S., Kelley, M., et al. (2019). Climate Change  
53 Amplification of Natural Drought Variability: The Historic Mid-Twentieth-Century North American Drought in a  
54 Warmer World. *J. Clim.* 32, 5417–5436. doi:10.1175/JCLI-D-18-0832.1.
- 55 Cook, B. I., Smerdon, J. E., Seager, R., and Coats, S. (2014a). Global warming and 21st century drying. *Clim. Dyn.* 43,  
56 2607–2627. doi:10.1007/s00382-014-2075-y.
- 57 Cook, B. I., Smerdon, J. E., Seager, R., and Coats, S. (2014b). Global warming and 21st century drying. *Clim. Dyn.* 43,  
58 2607–2627. doi:10.1007/s00382-014-2075-y.
- 59 Cook, B. I., Smerdon, J. E., Seager, R., and Cook, E. R. (2014c). Pan-Continental Droughts in North America over the  
60 Last Millennium. *J. Clim.* 27, 383–397. doi:10.1175/JCLI-D-13-00100.1.
- 61 Cook, E. R., Woodhouse, C. A., Mark Eakin, C., Meko, D. M., and Stahle, D. W. (2004). Long-Term Aridity Changes

- 1 in the Western United States. *Science* (80- ). 306, 1015–1018. doi:10.1126/science.1102586.
- 2 Coppola, E., Nogherotto, R., Ciarlo', J. M., Giorgi, F., Meijgaard, E., Kadygrov, N., et al. (2021a). Assessment of the  
3 European Climate Projections as Simulated by the Large EURO-CORDEX Regional and Global Climate Model  
4 Ensemble. *J. Geophys. Res. Atmos.* 126. doi:10.1029/2019JD032356.
- 5 Coppola, E., Raffaele, F., Giorgi, F., Giuliani, G., Xuejie, G., Ciarlo, J. M., et al. (2021b). Climate hazard indices  
6 projections based on CORDEX-CORE, CMIP5 and CMIP6 ensemble. *Clim. Dyn.* doi:10.1007/s00382-021-  
7 05640-z.
- 8 Corbella, S., and Stretch, D. D. (2012). Multivariate return periods of sea storms for coastal erosion risk assessment.  
9 *Nat. Hazards Earth Syst. Sci.* 12, 2699–2708. doi:10.5194/nhess-12-2699-2012.
- 10 Corrales-Suastegui, A., Fuentes-Franco, R., and Pavia, E. G. (2019). The mid-summer drought over Mexico and Central  
11 America in the 21st century. *Int. J. Climatol.* doi:10.1002/joc.6296.
- 12 Couasnon, A., Eilander, D., Muis, S., Veldkamp, T. I. E., Haigh, I. D., Wahl, T., et al. (2020). Measuring compound  
13 flood potential from river discharge and storm surge extremes at the global scale. *Nat. Hazards Earth Syst. Sci.*  
14 20, 489–504. doi:10.5194/nhess-20-489-2020.
- 15 Coumou, D., Di Capua, G., Vavrus, S., Wang, L., and Wang, S. (2018). The influence of Arctic amplification on mid-  
16 latitude summer circulation. *Nat. Commun.* 9, 2959. doi:10.1038/s41467-018-05256-8.
- 17 Coumou, D., Lehmann, J., and Beckmann, J. (2015). The weakening summer circulation in the Northern Hemisphere  
18 mid-latitudes. *Science* (80- ). 348, 324–327. doi:10.1126/science.1261768.
- 19 Cowan, T., Hegerl, G. C., Colfescu, I., Bollasina, M., Purich, A., and Bosch, G. (2016). Factors Contributing to  
20 Record-Breaking Heat Waves over the Great Plains during the 1930s Dust Bowl. *J. Clim.* 30, 2437–2461.  
21 doi:10.1175/JCLI-D-16-0436.1.
- 22 Cowan, T., Purich, A., Perkins, S., Pezza, A., Bosch, G., and Sadler, K. (2014). More frequent, longer, and hotter heat  
23 waves for Australia in the Twenty-First Century. *J. Clim.* 27, 5851–5871. doi:10.1175/JCLI-D-14-00092.1.
- 24 Crimp, S. J., Zheng, B., Khimashia, N., Gobbett, D. L., Chapman, S., Howden, M., et al. (2016). Recent changes in  
25 southern Australian frost occurrence: Implications for wheat production risk. *Crop Pasture Sci.* 67, 801–811.  
26 doi:10.1071/CP16056.
- 27 Croitoru, A.-E., Chitoroiu, B.-C., Ivanova Todorova, V., and Torică, V. (2013). Changes in precipitation extremes on  
28 the Black Sea Western Coast. *Glob. Planet. Change* 102, 10–19. doi:10.1016/j.gloplacha.2013.01.004.
- 29 Croitoru, A.-E., and Piticar, A. (2013). Changes in daily extreme temperatures in the extra-Carpathians regions of  
30 Romania. *Int. J. Climatol.* 33, 1987–2001. doi:10.1002/joc.3567.
- 31 Croitoru, A.-E., Piticar, A., and Burada, D. C. (2016). Changes in precipitation extremes in Romania. *Quat. Int.* 415,  
32 325–335. doi:10.1016/J.QUAINT.2015.07.028.
- 33 Crook, J., Klein, C., Folwell, S., Taylor, C. M., Parker, D. J., Stratton, R., et al. (2019). Assessment of the  
34 Representation of West African Storm Lifecycles in Convection-Permitting Simulations. *Earth Sp. Sci.* 6, 818–  
35 835. doi:10.1029/2018EA000491.
- 36 CSIRO and BOM (2015). “Climate change in Australia: Projections for Australia’s NRM regions,” in *Climate Change*  
37 *in Australia: Information for Australia’s Natural Resource Management Regions* (Australia: Commonwealth  
38 Scientific and Industrial Research Organisation (CSIRO) and Bureau of Meteorology (BOM)), 216. Available at:  
39 <https://publications.csiro.au/rpr/download?pid=csiro:EP154327&dsid=DS2>.
- 40 CSIRO, and BOM (2016). State of the Climate 2016. Australia: Commonwealth Scientific and Industrial Research  
41 Organisation (CSIRO) and Bureau of Meteorology (BOM) Available at: <http://www.bom.gov.au/state-of-the-climate/State-of-the-Climite-2016.pdf>.
- 42 Dadaser-Celik, F., and Cengiz, E. (2014). Wind speed trends over Turkey from 1975 to 2006. *Int. J. Climatol.* 34,  
43 1913–1927. doi:10.1002/joc.3810.
- 44 Dai, A. (2013). Increasing drought under global warming in observations and models. *Nat. Clim. Chang.* 3, 52.  
45 doi:10.1038/nclimate1633.
- 46 Dai, A. (2021). Hydroclimatic trends during 1950–2018 over global land. *Clim. Dyn.* doi:10.1007/s00382-021-05684-1.
- 47 Dai, A., and Zhao, T. (2017). Uncertainties in historical changes and future projections of drought. Part I: estimates of  
48 historical drought changes. *Clim. Change* 144, 519–533. doi:10.1007/s10584-016-1705-2.
- 49 Dai, A., Zhao, T., and Chen, J. (2018). Climate Change and Drought: a Precipitation and Evaporation Perspective.  
50 *Curr. Clim. Chang. Reports* 4, 301–312. doi:10.1007/s40641-018-0101-6.
- 51 Daloz, A. S., and Camargo, S. J. (2018). Is the poleward migration of tropical cyclone maximum intensity associated  
52 with a poleward migration of tropical cyclone genesis? *Clim. Dyn.* 50, 705–715. doi:10.1007/s00382-017-3636-7.
- 53 Dankers, R., Arnell, N. W., Clark, D. B., Falloon, P. D., Fekete, B. M., Gosling, S. N., et al. (2014). First look at  
54 changes in flood hazard in the Inter-Sectoral Impact Model Intercomparison Project ensemble. *Proc. Natl. Acad.*  
55 *Sci.* 111, 3257–3261. doi:10.1073/pnas.1302078110.
- 56 Dashkhuu, D., Kim, J. P., Chun, J. A., and Lee, W.-S. (2015). Long-term trends in daily temperature extremes over  
57 Mongolia. *Weather Clim. Extrem.* 8, 26–33. doi:10.1016/j.wace.2014.11.003.
- 58 Davin, E. L., Seneviratne, S. I., Ciais, P., Olliso, A., and Wang, T. (2014). Preferential cooling of hot extremes from  
59 cropland albedo management. *Proc. Natl. Acad. Sci.* 111, 9757–9761. doi:10.1073/pnas.1317323111.
- 60 de Lima, M. I. P., Santo, F. E., Ramos, A. M., and Trigo, R. M. (2015). Trends and correlations in annual extreme

- 1 precipitation indices for mainland Portugal, 1941–2007. *Theor. Appl. Climatol.* 119, 55–75. doi:10.1007/s00704-  
2 013-1079-6.
- 3 De Luca, P., Messori, G., Pons, F. M. E., and Faranda, D. (2020a). Dynamical systems theory sheds new light on  
4 compound climate extremes in Europe and Eastern North America. *Q. J. R. Meteorol. Soc.* 146, 1636–1650.  
5 doi:10.1002/qj.3757.
- 6 De Luca, P., Messori, G., Wilby, R. L., Mazzoleni, M., and Di Baldassarre, G. (2020b). Concurrent wet and dry  
7 hydrological extremes at the global scale. *Earth Syst. Dyn.* 11, 251–266. doi:10.5194/esd-11-251-2020.
- 8 de Vrese, P., Hagemann, S., and Claussen, M. (2016). Asian irrigation, African rain: Remote impacts of irrigation.  
9 *Geophys. Res. Lett.* 43, 3737–3745. doi:10.1002/2016GL068146.
- 10 DeAngelis, A. M., Wang, H., Koster, R. D., Schubert, S. D., Chang, Y., and Marshak, J. (2020). Prediction Skill of the  
11 2012 U.S. Great Plains Flash Drought in Subseasonal Experiment (SubX) Models. *J. Clim.* 33, 6229–6253.  
12 doi:10.1175/JCLI-D-19-0863.1.
- 13 Degeffie, D. T., Fleischer, E., Klemm, O., Soromotin, A. V., Soromotina, O. V., Tolstikov, A. V., et al. (2014). Climate  
14 extremes in South Western Siberia: past and future. *Stoch. Environ. Res. Risk Assess.* 28, 2161–2173.  
15 doi:10.1007/s00477-014-0872-9.
- 16 Delworth, T. L., and Zeng, F. (2014). Regional rainfall decline in Australia attributed to anthropogenic greenhouse  
17 gases and ozone levels. *Nat. Geosci.* 7, 583–587. doi:10.1038/ngeo2201.
- 18 Deng, Y., Jiang, W., He, B., Chen, Z., and Jia, K. (2018). Change in Intensity and Frequency of Extreme Precipitation  
19 and its Possible Teleconnection With Large-Scale Climate Index Over the China From 1960 to 2015. *J. Geophys.*  
20 *Res. Atmos.* 123, 2068–2081. doi:10.1002/2017JD027078.
- 21 Denniston, R. F., and Luetscher, M. (2017). Speleothems as high-resolution paleoflood archives. *Quat. Sci. Rev.* 170,  
22 1–13. doi:10.1016/j.quascirev.2017.05.006.
- 23 Déqué, M., Calmanti, S., Christensen, O. B., Dell Aquila, A., Maule, C. F., Haensler, A., et al. (2017). A multi-model  
24 climate response over tropical Africa at +2 °C. *Clim. Serv.* 7, 87–95. doi:10.1016/j.cliser.2016.06.002.
- 25 Dereczynski, C., Chan Chou, S., Lyra, A., Sondermann, M., Regoto, P., Tavares, P., et al. (2020). Downscaling of  
26 climate extremes over South America – Part I: Model evaluation in the reference climate. *Weather Clim. Extrem.*  
27 29, 100273. doi:10.1016/j.wace.2020.100273.
- 28 Deshpande, N. R., Kothawale, D. R., and Kulkarni, A. (2016). Changes in climate extremes over major river basins of  
29 India. *Int. J. Climatol.* 36, 4548–4559. doi:10.1002/joc.4651.
- 30 Devanand, A., Huang, M., Ashfaq, M., Barik, B., and Ghosh, S. (2019). Choice of Irrigation Water Management  
31 Practice Affects Indian Summer Monsoon Rainfall and Its Extremes. *Geophys. Res. Lett.* 46, 9126–9135.  
32 doi:10.1029/2019GL083875.
- 33 Dey, R., Lewis, S. C., and Abram, N. J. (2019a). Investigating observed northwest Australian rainfall trends in Coupled  
34 Model Intercomparison Project phase 5 detection and attribution experiments. *Int. J. Climatol.* 39, 112–127.  
35 doi:10.1002/joc.5788.
- 36 Dey, R., Lewis, S. C., Arblaster, J. M., and Abram, N. J. (2019b). A review of past and projected changes in Australia’s  
37 rainfall. *Wiley Interdiscip. Rev. Clim. Chang.* 10, e577. doi:10.1002/wcc.577.
- 38 Di Luca, A., de Elía, R., Bador, M., and Argüeso, D. (2020a). Contribution of mean climate to hot temperature  
39 extremes for present and future climates. *Weather Clim. Extrem.* 28, 100255. doi:10.1016/j.wace.2020.100255.
- 40 Di Luca, A., de Elía, R., and Laprise, R. (2015). Challenges in the Quest for Added Value of Regional Climate  
41 Dynamical Downscaling. *Curr. Clim. Chang. Reports* 1, 10–21. doi:10.1007/s40641-015-0003-9.
- 42 Di Luca, A., Evans, J. J. P., Pepler, A. S. A., Alexander, L. V. L., and Argüeso, D. (2016). Evaluating the representation  
43 of Australian East Coast Lows in a regional climate model ensemble. *J. South. Hemisph. Earth Syst. Sci.* 66, 108–  
44 124. doi:10.22499/3.6602.003.
- 45 Di Luca, A., Pitman, A. J., and de Elía, R. (2020b). Decomposing Temperature Extremes Errors in CMIP5 and CMIP6  
46 Models. *Geophys. Res. Lett.* 47, 1–9. doi:10.1029/2020GL088031.
- 47 Diaconescu, E. P., Mailhot, A., Brown, R., and Chaumont, D. (2018). Evaluation of CORDEX-Arctic daily  
48 precipitation and temperature-based climate indices over Canadian Arctic land areas. *Clim. Dyn.* 50, 2061–2085.  
49 doi:10.1007/s00382-017-3736-4.
- 50 Diallo, I., Giorgi, F., Deme, A., Tall, M., Mariotti, L., and Gaye, A. T. (2016). Projected changes of summer monsoon  
51 extremes and hydroclimatic regimes over West Africa for the twenty-first century. *Clim. Dyn.*  
52 doi:10.1007/s00382-016-3052-4.
- 53 Diallo, I., Giorgi, F., Sukumaran, S., Stordal, F., and Giuliani, G. (2015). Evaluation of RegCM4 driven by CAM4 over  
54 Southern Africa: mean climatology, interannual variability and daily extremes of wet season temperature and  
55 precipitation. *Theor. Appl. Climatol.* 121, 749–766. doi:10.1007/s00704-014-1260-6.
- 56 Diedhiou, A., Bichet, A., Wartenburger, R., Seneviratne, S. I., Rowell, D. P., Sylla, M. B., et al. (2018). Changes in  
57 climate extremes over West and Central Africa at 1.5 °C and 2 °C global warming. *Environ. Res. Lett.* 13,  
58 065020. doi:10.1088/1748-9326/aac3e5.
- 59 Dierauer, J. R., Allen, D. M., and Whitfield, P. H. (2019). Snow Drought Risk and Susceptibility in the Western United  
60 States and Southwestern Canada. *Water Resour. Res.* 55, 3076–3091. doi:10.1029/2018WR023229.
- 61 Diffenbaugh, N. S., Scherer, M., and Trapp, R. J. (2013). Robust increases in severe thunderstorm environments in

- 1 response to greenhouse forcing. *Proc. Natl. Acad. Sci.* 110, 16361 LP-16366. doi:10.1073/pnas.1307758110.
- 2 Diffenbaugh, N. S., Singh, D., Mankin, J. S., Horton, D. E., Swain, D. L., Touma, D., et al. (2017). Quantifying the  
3 influence of global warming on unprecedented extreme climate events. *Proc. Natl. Acad. Sci.* 114, 4881–4886.
- 4 Diffenbaugh, N. S., Swain, D. L., and Touma, D. (2015). Anthropogenic warming has increased drought risk in  
5 California. *Proc. Natl. Acad. Sci.* 112, 3931 LP-3936. doi:10.1073/pnas.1422385112.
- 6 Dikšaitytė, A., Viršilė, A., Žaltauskaitė, J., Januškaitienė, I., and Juozapaitienė, G. (2019). Growth and photosynthetic  
7 responses in *Brassica napus* differ during stress and recovery periods when exposed to combined heat, drought  
8 and elevated CO<sub>2</sub>. *Plant Physiol. Biochem.* 142, 59–72. doi:10.1016/j.plaphy.2019.06.026.
- 9 Dimri, A. P. (2019). Comparison of regional and seasonal changes and trends in daily surface temperature extremes  
10 over India and its subregions. *Theor. Appl. Climatol.* 136. doi:10.1007/s00704-018-2486-5.
- 11 Dimri, A. P., Chevuturi, A., Niyogi, D., Thayyen, R. J., Ray, K., Tripathi, S. N., et al. (2017). Cloudbursts in Indian  
12 Himalayas: A review. *Earth-Science Rev.* 168, 1–23. doi:10.1016/J.EARSCIREV.2017.03.006.
- 13 Dimri, A. P., Niyogi, D., Barros, A. P., Ridley, J., Mohanty, U. C., Yasunari, T., et al. (2015). Western Disturbances: A  
14 review. *Rev. Geophys.* 53, 225–246. doi:10.1002/2014RG000460.
- 15 Dirmeyer, P. A., Gao, X., Zhao, M., Guo, Z., Oki, T., and Hanasaki, N. (2006). GSWP-2: Multimodel Analysis and  
16 Implications for Our Perception of the Land Surface. *Bull. Am. Meteorol. Soc.* 87, 1381–1398.  
17 doi:10.1175/BAMS-87-10-1381.
- 18 Dirmeyer, P. A., Jin, Y., Singh, B., and Yan, X. (2013). Trends in land-atmosphere interactions from CMIP5  
19 simulations. *J. Hydrometeorol.* 14, 829–849. doi:10.1175/JHM-D-12-0107.1.
- 20 Diro, G. T., Sushama, L., and Huziy, O. (2018). Snow-atmosphere coupling and its impact on temperature variability  
21 and extremes over North America. *Clim. Dyn.* 50, 2993–3007. doi:10.1007/s00382-017-3788-5.
- 22 Dittus, A. J., Karoly, D. J., Donat, M. G., Lewis, S. C., and Alexander, L. V. (2018). Understanding the role of sea  
23 surface temperature-forcing for variability in global temperature and precipitation extremes. *Weather Clim.*  
24 *Extrem.* 21, 1–9. doi:10.1016/j.wace.2018.06.002.
- 25 Dittus, A. J., Karoly, D. J., Lewis, S. C., and Alexander, L. V. (2014). An investigation of some unexpected frost day  
26 increases in southern Australia. *Aust. Meteorol. Oceanogr. J.* 64, 261–271. doi:10.22499/2.6404.002.
- 27 Dittus, A. J., Karoly, D. J., Lewis, S. C., Alexander, L. V., and Donat, M. G. (2016). A multiregion model evaluation  
28 and attribution study of historical changes in the area affected by temperature and precipitation extremes. *J. Clim.*  
29 29, 8285–8299. doi:10.1175/JCLI-D-16-0164.1.
- 30 Do, H. X., Mei, Y., and Gronewold, A. D. (2020). To What Extent Are Changes in Flood Magnitude Related to  
31 Changes in Precipitation Extremes? *Geophys. Res. Lett.* 47. doi:10.1029/2020GL088684.
- 32 Do, H. X., Westra, S., and Leonard, M. (2017). A global-scale investigation of trends in annual maximum streamflow.  
33 *J. Hydrol.* 552, 28–43. doi:10.1016/j.jhydrol.2017.06.015.
- 34 Dobricic, S., Russo, S., Pozzoli, L., Wilson, J., and Vignati, E. (2020). Increasing occurrence of heat waves in the  
35 terrestrial Arctic. *Environ. Res. Lett.* 15, 024022. doi:10.1088/1748-9326/ab6398.
- 36 Döll, P., Douville, H., Güntner, A., Müller Schmied, H., and Wada, Y. (2016). Modelling Freshwater Resources at the  
37 Global Scale: Challenges and Prospects. *Surv. Geophys.* 37, 195–221. doi:10.1007/s10712-015-9343-1.
- 38 Döll, P., Trautmann, T., Gerten, D., Schmied, H. M., Ostberg, S., Saaed, F., et al. (2018). Risks for the global  
39 freshwater system at 1.5 °C and 2 °C global warming. *Environ. Res. Lett.* 13, 044038. doi:10.1088/1748-  
40 9326/aab792.
- 41 Domínguez-Castro, F., García-Herrera, R., and Vicente-Serrano, S. M. (2018). Wet and dry extremes in Quito  
42 (Ecuador) since the 17th century. *Int. J. Climatol.* 38. doi:10.1002/joc.5312.
- 43 Donat, M. G., Alexander, L. V., Herold, N., and Dittus, A. J. (2016a). Temperature and precipitation extremes in  
44 century-long gridded observations, reanalyses, and atmospheric model simulations. *J. Geophys. Res. Atmos.* 121,  
45 11,174–11,189. doi:10.1002/2016JD025480.
- 46 Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., Dunn, R. J. H. H., et al. (2013a). Updated analyses of  
47 temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset.  
48 *J. Geophys. Res. Atmos.* 118, 2098–2118. doi:10.1002/jgrd.50150.
- 49 Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., and Caesar, J. (2013b). Global Land-Based Datasets for  
50 Monitoring Climatic Extremes. *Bull. Am. Meteorol. Soc.* 94, 997–1006. doi:10.1175/BAMS-D-12-00109.1.
- 51 Donat, M. G., Lowry, A. L., Alexander, L. V., O’Gorman, P. A., and Maher, N. (2016b). More extreme precipitation in  
52 the world’s dry and wet regions. *Nat. Clim. Chang.* 6, 508–513. doi:10.1038/nclimate2941.
- 53 Donat, M. G., Peterson, T. C., Brunet, M., King, A. D., Almazroui, M., Kolli, R. K., et al. (2014a). Changes in extreme  
54 temperature and precipitation in the Arab region: long-term trends and variability related to ENSO and NAO. *Int.*  
55 *J. Climatol.* 34, 581–592. doi:10.1002/joc.3707.
- 56 Donat, M. G., Pitman, A. J., and Angéilil, O. (2018). Understanding and Reducing Future Uncertainty in Midlatitude  
57 Daily Heat Extremes Via Land Surface Feedback Constraints. *Geophys. Res. Lett.* 45, 10,627–10,636.  
58 doi:10.1029/2018GL079128.
- 59 Donat, M. G., Pitman, A. J., and Seneviratne, S. I. (2017). Regional warming of hot extremes accelerated by surface  
60 energy fluxes. *Geophys. Res. Lett.* 44, 7011–7019.
- 61 Donat, M. G., Sillmann, J., Wild, S., Alexander, L. V., Lippmann, T., and Zwiers, F. W. (2014b). Consistency of

- 1 Temperature and Precipitation Extremes across Various Global Gridded In Situ and Reanalysis Datasets. *J. Clim.*  
2 27, 5019–5035. doi:10.1175/JCLI-D-13-00405.1.
- 3 Dong, B., and Dai, A. (2015). The influence of the Interdecadal Pacific Oscillation on Temperature and Precipitation  
4 over the Globe. *Clim. Dyn.* 45, 2667–2681. doi:10.1007/s00382-015-2500-x.
- 5 Dong, B., Sutton, R., and Shaffrey, L. (2014). The 2013 hot, dry, summer in western Europe [in “Explaining Extreme  
6 Events of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S62–S66.
- 7 Dong, B., Sutton, R., Shaffrey, L., and Wilcox, L. (2016a). The 2015 European Heat Wave. *Bull. Am. Meteorol. Soc.*  
8 97, S57–S62. doi:10.1175/BAMS-D-16-0140.1.
- 9 Dong, B., Sutton, R. T., Chen, W., Liu, X., Lu, R., and Sun, Y. (2016b). Abrupt summer warming and changes in  
10 temperature extremes over Northeast Asia since the mid-1990s: Drivers and physical processes. *Adv. Atmos. Sci.*  
11 33, 1005–1023. doi:10.1007/s00376-016-5247-3.
- 12 Dong, B., Sutton, R. T., and Shaffrey, L. (2017). Understanding the rapid summer warming and changes in temperature  
13 extremes since the mid-1990s over Western Europe. *Clim. Dyn.* 48, 1537–1554. doi:10.1007/s00382-016-3158-8.
- 14 Dong, S., Sun, Y., Aguilar, E., Zhang, X., Peterson, T. C., Song, L., et al. (2018). Observed changes in temperature  
15 extremes over Asia and their attribution. *Clim. Dyn.* 51, 339–353. doi:10.1007/s00382-017-3927-z.
- 16 Dong, S., Sun, Y., Li, C., Zhang, X., Min, S.-K., and Kim, Y.-H. (2021). Attribution of Extreme Precipitation with  
17 Updated Observations and CMIP6 Simulations. *J. Clim.* 34, 871–881. doi:10.1175/JCLI-D-19-1017.1.
- 18 Dong, S., Xu, Y., Zhou, B., and Shi, Y. (2015). Assessment of indices of temperature extremes simulated by multiple  
19 CMIP5 models over China. *Adv. Atmos. Sci.* 32, 1077–1091. doi:10.1007/s00376-015-4152-5.
- 20 Donnelly, C., Greuell, W., Andersson, J., Gerten, D., Pisacane, G., Roudier, P., et al. (2017). Impacts of climate change  
21 on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level. *Clim. Change*  
22 143, 13–26. doi:10.1007/s10584-017-1971-7.
- 23 Donnelly, J. P., Smith Bryant, S., Butler, J., Dowling, J., Fan, L., Hausmann, N., et al. (2001). 700 yr sedimentary  
24 record of intense hurricane landfalls in southern New England. *Geol. Soc. Am. Bull.* 113, 714–727.  
25 doi:10.1130/0016-7606(2001)113<0714:YSROIH>2.0.CO;2.
- 26 Dookie, N., Chadee, X. T., and Clarke, R. M. (2019). Trends in extreme temperature and precipitation indices for the  
27 Caribbean small islands: Trinidad and Tobago. *Theor. Appl. Climatol.* 136, 31–44. doi:10.1007/s00704-018-  
28 2463-z.
- 29 Dorigo, W. A., Gruber, A., De Jeu, R. A. M., Wagner, W., Stacke, T., Loew, A., et al. (2015). Evaluation of the ESA  
30 CCI soil moisture product using ground-based observations. *Remote Sens. Environ.* 162, 380–395.  
31 doi:10.1016/j.rse.2014.07.023.
- 32 Dorigo, W. A., Wagner, W., Hohensinn, R., Hahn, S., Paulik, C., Xaver, A., et al. (2011). The International Soil  
33 Moisture Network: a data hosting facility for global in situ soil moisture measurements. *Hydrol. Earth Syst. Sci.*  
34 15, 1675–1698. doi:10.5194/hess-15-1675-2011.
- 35 Dorigo, W., de Jeu, R., Chung, D., Parinussa, R., Liu, Y., Wagner, W., et al. (2012). Evaluating global trends (1988-  
36 2010) in harmonized multi-satellite surface soil moisture. *Geophys. Res. Lett.* 39. doi:10.1029/2012GL052988.
- 37 Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., et al. (2017). ESA CCI Soil Moisture for  
38 improved Earth system understanding: State-of-the art and future directions. *Remote Sens. Environ.* 203, 185–  
39 215. doi:10.1016/j.rse.2017.07.001.
- 40 Dosio, A. (2016). Projections of climate change indices of temperature and precipitation from an ensemble of bias-  
41 adjusted high-resolution EURO-CORDEX regional climate models. *J. Geophys. Res. Atmos.* 121, 5488–5511.  
42 doi:10.1002/2015JD024411.
- 43 Dosio, A. (2017). Projection of temperature and heat waves for Africa with an ensemble of CORDEX Regional Climate  
44 Models. *Clim. Dyn.* 49, 493–519. doi:10.1007/s00382-016-3355-5.
- 45 Dosio, A., and Fischer, E. M. (2018). Will Half a Degree Make a Difference? Robust Projections of Indices of Mean  
46 and Extreme Climate in Europe Under 1.5°C, 2°C, and 3°C Global Warming. *Geophys. Res. Lett.* 45, 935–944.  
47 doi:10.1002/2017GL076222.
- 48 Dosio, A., Jones, R. G., Jack, C., Lennard, C., Nikulin, G., and Hewitson, B. (2019). What can we know about future  
49 precipitation in Africa? Robustness, significance and added value of projections from a large ensemble of  
50 regional climate models. *Clim. Dyn.*, 1–26. doi:10.1007/s00382-019-04900-3.
- 51 Dosio, A., Mentaschi, L., Fischer, E. M., and Wyser, K. (2018). Extreme heat waves under 1.5 °C and 2 °C global  
52 warming. *Environ. Res. Lett.* 13, 054006. doi:10.1088/1748-9326/aab827.
- 53 Dosio, A., Panitz, H.-J., Schubert-Frisius, M., and Lüthi, D. (2015). Dynamical downscaling of CMIP5 global  
54 circulation models over CORDEX-Africa with COSMO-CLM: evaluation over the present climate and analysis  
55 of the added value. *Clim. Dyn.* 44, 2637–2661. doi:10.1007/s00382-014-2262-x.
- 56 Douville, H., Colin, J., Krug, E., Cattiaux, J., and Thao, S. (2016). Midlatitude daily summer temperatures reshaped by  
57 soil moisture under climate change. *Geophys. Res. Lett.* 43, 812–818. doi:10.1002/2015GL066222.
- 58 Douville, H., and Plazzotta, M. (2017). Midlatitude Summer Drying: An Underestimated Threat in CMIP5 Models?  
59 *Geophys. Res. Lett.* 44, 9967–9975. doi:10.1002/2017GL075353.
- 60 Dowdy, A. J. (2018). Climatological Variability of Fire Weather in Australia. *J. Appl. Meteorol. Climatol.* 57, 221–234.  
61 doi:10.1175/JAMC-D-17-0167.1.

- 1 Dowdy, A. J., and Catto, J. L. (2017). Extreme weather caused by concurrent cyclone, front and thunderstorm  
2 occurrences. *Sci. Rep.* 7, 1–8. doi:10.1038/srep40359.
- 3 Dowdy, A. J., Pepler, A., Di Luca, A., Cavicchia, L., Mills, G., Evans, J. P., et al. (2019). Review of Australian east  
4 coast low pressure systems and associated extremes. *Clim. Dyn.* doi:10.1007/s00382-019-04836-8.
- 5 Drijfhout, S., Bathiany, S., Beaulieu, C., Brovkin, V., Claussen, M., Huntingford, C., et al. (2015). Catalogue of abrupt  
6 shifts in Intergovernmental Panel on Climate Change climate models. *Proc. Natl. Acad. Sci.* 112, E5777–E5786.  
7 doi:10.1073/pnas.1511451112.
- 8 Driouech, F., ElRhaz, K., Moufouma-Okia, W., Arjdal, K., and Balhane, S. (2020). Assessing Future Changes of  
9 Climate Extreme Events in the CORDEX-MENA Region Using Regional Climate Model ALADIN-Climate.  
10 *Earth Syst. Environ.* 4, 477–492. doi:10.1007/s41748-020-00169-3.
- 11 Driouech, F., Stafi, H., Khouakhi, A., Moutia, S., Badi, W., ElRhaz, K., et al. (2021). Recent observed country-wide  
12 climate trends in Morocco. *Int. J. Climatol.* 41, 1–20. doi:10.1002/joc.6734.
- 13 Drobinski, P., Silva, N. Da, Panthou, G., Bastin, S., Muller, C., Ahrens, B., et al. (2018). Scaling precipitation extremes  
14 with temperature in the Mediterranean: past climate assessment and projection in anthropogenic scenarios. *Clim.*  
15 *Dyn.* 51, 1237–1257. doi:10.1007/s00382-016-3083-x.
- 16 Drouard, M., Kornhuber, K., and Woollings, T. (2019). Disentangling Dynamic Contributions to Summer 2018  
17 Anomalous Weather Over Europe. *Geophys. Res. Lett.* 46, 12537–12546. doi:10.1029/2019GL084601.
- 18 Drumond, A., Stojanovic, M., Nieto, R., Vicente-Serrano, S. M., and Gimeno, L. (2019). Linking Anomalous Moisture  
19 Transport And Drought Episodes in the IPCC Reference Regions. *Bull. Am. Meteorol. Soc.* 100, 1481–1498.  
20 doi:10.1175/BAMS-D-18-0111.1.
- 21 Du, H., Alexander, L. V., Donat, M. G., Lippmann, T., Srivastava, A., Salinger, J., et al. (2019). Precipitation From  
22 Persistent Extremes is Increasing in Most Regions and Globally. *Geophys. Res. Lett.* 46, 6041–6049.  
23 doi:10.1029/2019GL081898.
- 24 Ducrocq, V., Nuissier, O., Ricard, D., Lebeauupin, C., and Thouvenin, T. (2008). A numerical study of three catastrophic  
25 precipitating events over southern France. II: Mesoscale triggering and stationarity factors. *Q. J. R. Meteorol. Soc.*  
26 134, 131–145. doi:10.1002/qj.199.
- 27 Dudley, R. W., Hirsch, R. M., Archfield, S. A., Blum, A. G., and Renard, B. (2020). Low streamflow trends at human-  
28 impacted and reference basins in the United States. *J. Hydrol.* 580, 124254. doi:10.1016/j.jhydrol.2019.124254.
- 29 Dudley, R. W., Hodgkins, G. A., McHale, M. R., Kolian, M. J., and Renard, B. (2017). Trends in snowmelt-related  
30 streamflow timing in the conterminous United States. *J. Hydrol.* 547, 208–221.  
31 doi:10.1016/j.jhydrol.2017.01.051.
- 32 Duffy, P. B., Brando, P., Asner, G. P., and Field, C. B. (2015). Projections of future meteorological drought and wet  
33 periods in the Amazon. *Proc. Natl. Acad. Sci.* 112, 13172–13177. doi:10.1073/pnas.1421010112.
- 34 Duke, N. C., Kovacs, J. M., Griffiths, A. D., Preece, L., Hill, D. J. E., van Oosterzee, P., et al. (2017). Large-scale  
35 dieback of mangroves in Australia’s Gulf of Carpentaria: a severe ecosystem response, coincidental with an  
36 unusually extreme weather event. *Mar. Freshw. Res.* 68, 1816–1829. doi:10.1071/MF16322.
- 37 Dunn, R. J. H., Alexander, L. V., Donat, M. G., Zhang, X., Bador, M., Herold, N., et al. (2020). Development of an  
38 Updated Global Land In Situ-Based Data Set of Temperature and Precipitation Extremes: HadEX3. *J. Geophys.*  
39 *Res. Atmos.* 125. doi:10.1029/2019JD032263.
- 40 Dunning, C. M., Black, E., and Allan, R. P. (2018). Later Wet Seasons with More Intense Rainfall over Africa under  
41 Future Climate Change. *J. Clim.* 31, 9719–9738. doi:10.1175/JCLI-D-18-0102.1.
- 42 Dunstone, N. J., Smith, D. M., Booth, B. B. B., Hermanson, L., and Eade, R. (2013). Anthropogenic aerosol forcing of  
43 Atlantic tropical storms. *Nat. Geosci.* 6, 534–539. doi:10.1038/ngeo1854.
- 44 Durkee, J. D., and Mote, T. L. (2010). A climatology of warm-season mesoscale convective complexes in subtropical  
45 South America. *Int. J. Climatol.* 30, 418–431. doi:10.1002/joc.1893.
- 46 Easterling, D. R., Arnold, J. R., Knutson, T., Kunkel, K. E., LeGrande, A. N., Leung, L. R., et al. (2017). Precipitation  
47 change in the United States. , eds. D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and  
48 T. K. Maycock Washington, DC, USA: U.S. Global Change Research Program doi:10.7930/J0H993CC.
- 49 Easterling, D. R., Kunkel, K. E., Wehner, M. F., and Sun, L. (2016). Detection and attribution of climate extremes in  
50 the observed record. *Weather Clim. Extrem.* 11, 17–27. doi:10.1016/j.wace.2016.01.001.
- 51 Eden, J. M., Wolter, K., Otto, F. E. L., and Jan van Oldenborgh, G. (2016). Multi-method attribution analysis of  
52 extreme precipitation in Boulder, Colorado. *Environ. Res. Lett.* 11, 124009. doi:10.1088/1748-  
53 9326/11/12/124009.
- 54 Edossa, D. C., Woyessa, Y. E., and Welderufael, W. A. (2016). Spatiotemporal analysis of droughts using self-  
55 calibrating Palmer’s Drought Severity Index in the central region of South Africa. *Theor. Appl. Climatol.* 126,  
56 643–657. doi:10.1007/s00704-015-1604-x.
- 57 El Kenawy, A., López-Moreno, J. I., and Vicente-Serrano, S. M. (2013). Summer temperature extremes in northeastern  
58 Spain: Spatial regionalization and links to atmospheric circulation (1960-2006). *Theor. Appl. Climatol.* 113, 387–  
59 405. doi:10.1007/s00704-012-0797-5.
- 60 Elsner, J. B., Elsner, S. C., and Jagger, T. H. (2015). The increasing efficiency of tornado days in the United States.  
61 *Clim. Dyn.* 45, 651–659. doi:10.1007/s00382-014-2277-3.



- 1 Elsner, J. B., Fricker, T., and Schroder, Z. (2019). Increasingly powerful tornadoes in the United States. *Geophys. Res. Lett.*, 1–10. doi:10.1029/2018GL080819.
- 2
- 3 Elsner, J. B., Kossin, J. P., and Jagger, T. H. (2008). The increasing intensity of the strongest tropical cyclones. *Nature* 455, 92–95. doi:10.1038/nature07234.
- 4
- 5 Emanuel, K. (2017). Assessing the present and future probability of Hurricane Harvey's rainfall. *Proc. Natl. Acad. Sci.* 114, 12681–12684. doi:10.1073/pnas.1716222114.
- 6
- 7 Emanuel, K. (2021). Response of Global Tropical Cyclone Activity to Increasing CO<sub>2</sub>: Results from Downscaling CMIP6 Models. *J. Clim.* 34, 57–70. doi:10.1175/JCLI-D-20-0367.1.
- 8
- 9 Emanuel, K. A. (1987). The dependence of hurricane intensity on climate. *Nature* 326, 483–485. doi:10.1038/326483a0.
- 10
- 11 Emanuel, K. A. (2013). Downscaling CMIP5 climate models shows increased tropical cyclone activity over the 21st century. doi:10.1073/pnas.1301293110.
- 12
- 13 Emanuel, K., Caroff, P., Delgado, S., Guard, C. “Chip,” Guishard, M., Hennon, C., et al. (2018). On the Desirability and Feasibility of a Global Reanalysis of Tropical Cyclones. *Bull. Am. Meteorol. Soc.* 99, 427–429. doi:10.1175/BAMS-D-17-0226.1.
- 14
- 15 Emanuel, K., Ravela, S., Vivant, E., and Risi, C. (2006). A statistical deterministic approach to hurricane risk assessment. *Bull. Am. Meteorol. Soc.* 87, 299–314. doi:10.1175/BAMS-87-3-299.
- 16
- 17 Emanuel, K., Sundararajan, R., and Williams, J. (2008). Hurricanes and global warming: Results from downscaling IPCC AR4 simulations. *Bull. Am. Meteorol. Soc.* 89, 347–367. doi:10.1175/BAMS-89-3-347.
- 18
- 19 Endo, H., Kitoh, A., Mizuta, R., and Ishii, M. (2017). Future Changes in Precipitation Extremes in East Asia and Their Uncertainty Based on Large Ensemble Simulations with a High-Resolution AGCM. 13, 0–5. doi:10.2151/sola.2017-002.
- 20
- 21 Engelbrecht, F. A., McGregor, J. L., and Engelbrecht, C. J. (2009). Dynamics of the Conformal-Cubic Atmospheric Model projected climate-change signal over southern Africa. *Int. J. Climatol.* 29, 1013–1033. doi:10.1002/joc.1742.
- 22
- 23 Engelbrecht, F., Adegoke, J., Bopape, M.-J., Naidoo, M., Garland, R., Thatcher, M., et al. (2015). Projections of rapidly rising surface temperatures over Africa under low mitigation. *Environ. Res. Lett.* 10, 085004. doi:10.1088/1748-9326/10/8/085004.
- 24
- 25 Engström, J., and Keellings, D. (2018). Drought in the Southeastern USA: an assessment of downscaled CMIP5 models. *Clim. Res.* 74, 251–262. doi:10.3354/cr01502.
- 26
- 27 Erdenebat, E., and Sato, T. (2016). Recent increase in heat wave frequency around Mongolia: role of atmospheric forcing and possible influence of soil moisture deficit. *Atmos. Sci. Lett.* 17, 135–140. doi:10.1002/asl.616.
- 28
- 29 Erfanian, A., Wang, G., and Fomenko, L. (2017). Unprecedented drought over tropical South America in 2016: significantly under-predicted by tropical SST. *Sci. Rep.* 7, 5811. doi:10.1038/s41598-017-05373-2.
- 30
- 31 Erlat, E., and Türkeş, M. (2016). Dates of frost onset, frost end and the frost-free season in Turkey: Trends, variability and links to the North Atlantic and Arctic Oscillation indices, 1950–2013. *Clim. Res.* 69, 155–176. doi:10.3354/cr01397.
- 32
- 33 Escalante-Sandoval, C., and Nuñez-García, P. (2017). Meteorological drought features in northern and northwestern parts of Mexico under different climate change scenarios. *J. Arid Land* 9, 65–75. doi:10.1007/s40333-016-0022-y.
- 34
- 35 Evan, A. T., Kossin, J. P., Chung, C. E., and Ramanathan, V. (2011). Arabian Sea tropical cyclones intensified by emissions of black carbon and other aerosols. *Nature* 479, 94–97. doi:10.1038/nature10552.
- 36
- 37 Evans, J. P., Argueso, D., Olson, R., and Di Luca, A. (2017). Bias-corrected regional climate projections of extreme rainfall in south-east Australia. *Theor. Appl. Climatol.* 130, 1085–1098. doi:10.1007/s00704-016-1949-9.
- 38
- 39 Evans, J. P., Di Virgilio, G., Hirsch, A. L., Hoffmann, P., Remedio, A. R., Ji, F., et al. (2020). The CORDEX-Australasia ensemble: evaluation and future projections. *Clim. Dyn.* doi:10.1007/s00382-020-05459-0.
- 40
- 41 Fadnavis, S., Sabin, T. P., Roy, C., Rowlinson, M., Rap, A., Vernier, J.-P., et al. (2019). Elevated aerosol layer over South Asia worsens the Indian droughts. *Sci. Rep.* 9, 10268. doi:10.1038/s41598-019-46704-9.
- 42
- 43 Falconer, R. H., Cobby, D., Smyth, P., Astle, G., Dent, J., and Golding, B. (2009). Pluvial flooding: new approaches in flood warning, mapping and risk management. *J. Flood Risk Manag.* 2, 198–208. doi:10.1111/j.1753-318X.2009.01034.x.
- 44
- 45 Fantini, A., Raffaele, F., Torma, C., Bacer, S., Coppola, E., Giorgi, F., et al. (2018). Assessment of multiple daily precipitation statistics in ERA-Interim driven Med-CORDEX and EURO-CORDEX experiments against high resolution observations. *Clim. Dyn.* 51, 877–900. doi:10.1007/s00382-016-3453-4.
- 46
- 47 Fazel, N., Torabi Haghighi, A., and Kløve, B. (2017). Analysis of land use and climate change impacts by comparing river flow records for headwaters and lowland reaches. *Glob. Planet. Change* 158, 47–56. doi:10.1016/j.gloplacha.2017.09.014.
- 48
- 49 Feng, R., Yu, R., Zheng, H., and Gan, M. (2018). Spatial and temporal variations in extreme temperature in Central Asia. *Int. J. Climatol.* 38, e388–e400. doi:10.1002/joc.5379.
- 50
- 51 Feng, S., Trnka, M., Hayes, M., and Zhang, Y. (2017). Why Do Different Drought Indices Show Distinct Future Drought Risk Outcomes in the U.S. Great Plains? *J. Clim.* 30, 265–278. doi:10.1175/JCLI-D-15-0590.1.
- 52

- 1 Feng, Z., Leung, L. R., Hagos, S., Houze, R. A., Burleyson, C. D., and Balaguru, K. (2016). More frequent intense and  
2 long-lived storms dominate the springtime trend in central US rainfall. *Nat. Commun.* 7, 13429.  
3 doi:10.1038/ncomms13429.
- 4 Fenta, A. A., Yasuda, H., Shimizu, K., and Haregeweyn, N. (2017). Response of streamflow to climate variability and  
5 changes in human activities in the semiarid highlands of northern Ethiopia. *Reg. Environ. Chang.* 17, 1229–1240.  
6 doi:10.1007/s10113-017-1103-y.
- 7 Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., and Xia, L. (2015). Storminess over the North Atlantic  
8 and northwestern Europe – A review. *Q. J. R. Meteorol. Soc.* 141, 350–382. doi:10.1002/qj.2364.
- 9 Ficklin, D. L., Abatzoglou, J. T., Robeson, S. M., Null, S. E., and Knouft, J. H. (2018). Natural and managed  
10 watersheds show similar responses to recent climate change. *Proc. Natl. Acad. Sci.*, 201801026.  
11 doi:10.1073/PNAS.1801026115.
- 12 Field, R. D., van der Werf, G. R., Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., et al. (2016). Indonesian fire activity  
13 and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought. *Proc. Natl. Acad.  
14 Sci.* 113, 9204–9209. doi:10.1073/pnas.1524888113.
- 15 Filahi, S., Tanarhte, M., Mouhir, L., El Morhit, M., and Trambalay, Y. (2016). Trends in indices of daily temperature  
16 and precipitations extremes in Morocco. *Theor. Appl. Climatol.* 124, 959–972. doi:10.1007/s00704-015-1472-4.
- 17 Findell, K. L., Berg, A., Gentine, P., Krasting, J. P., Lintner, B. R., Malyshev, S., et al. (2017). The impact of  
18 anthropogenic land use and land cover change on regional climate extremes. *Nat. Commun.* 8, 989.  
19 doi:10.1038/s41467-017-01038-w.
- 20 Findell, K. L., Keys, P. W., van der Ent, R. J., Lintner, B. R., Berg, A., and Krasting, J. P. (2019). Rising Temperatures  
21 Increase Importance of Oceanic Evaporation as a Source for Continental Precipitation. *J. Clim.* 32, 7713–7726.  
22 doi:10.1175/JCLI-D-19-0145.1.
- 23 Finney, D. L., Marsham, J. H., Jackson, L. S., Kendon, E. J., Rowell, D. P., Boorman, P. M., et al. (2019). Implications  
24 of Improved Representation of Convection for the East Africa Water Budget Using a Convection-Permitting  
25 Model. *J. Clim.* 32, 2109–2129. doi:10.1175/JCLI-D-18-0387.1.
- 26 Finney, D. L., Marsham, J. H., Rowell, D. P., Kendon, E. J., Tucker, S. O., Stratton, R. A., et al. (2020). Effects of  
27 Explicit Convection on Future Projections of Mesoscale Circulations, Rainfall, and Rainfall Extremes over  
28 Eastern Africa. *J. Clim.* 33, 2701–2718. doi:10.1175/JCLI-D-19-0328.1.
- 29 Fioravanti, G., Piervitali, E., and Desiato, F. (2016). Recent changes of temperature extremes over Italy: an index-based  
30 analysis. *Theor. Appl. Climatol.* 123, 473–486. doi:10.1007/s00704-014-1362-1.
- 31 Fischer, A. M., Keller, D. E., Liniger, M. A., Rajczak, J., Schär, C., and Appenzeller, C. (2015). Projected changes in  
32 precipitation intensity and frequency in Switzerland: a multi-model perspective. *Int. J. Climatol.* 35, 3204–3219.  
33 doi:10.1002/joc.4162.
- 34 Fischer, E. M., and Knutti, R. (2014). Detection of spatially aggregated changes in temperature and precipitation  
35 extremes. *Geophys. Res. Lett.* 41, 547–554. doi:10.1002/2013GL058499.
- 36 Fischer, E. M., and Knutti, R. (2015). Anthropogenic contribution to global occurrence of heavy-precipitation and high-  
37 temperature extremes. *Nat. Clim. Chang.* 5, 560–564. doi:10.1038/nclimate2617.
- 38 Fischer, E. M., and Knutti, R. (2016). Observed heavy precipitation increase confirms theory and early models. *Nat.  
39 Clim. Chang.* 6, 986–991. doi:10.1038/nclimate3110.
- 40 Fischer, E. M., Sedláček, J., Hawkins, E., and Knutti, R. (2014). Models agree on forced response pattern of  
41 precipitation and temperature extremes. *Geophys. Res. Lett.* 41, 8554–8562. doi:10.1002/2014GL062018.
- 42 Fitchett, J. M. (2018). Recent emergence of CAT5 tropical cyclones in the South Indian Ocean. *S. Afr. J. Sci.* 114.  
43 doi:10.17159/sajs.2018/4426.
- 44 Fitzpatrick, R. G. J., Parker, D. J., Marsham, J. H., Rowell, D. P., Guichard, F. M., Taylor, C. M., et al. (2020). What  
45 Drives the Intensification of Mesoscale Convective Systems over the West African Sahel under Climate Change?  
46 *J. Clim.* 33, 3151–3172. doi:10.1175/JCLI-D-19-0380.1.
- 47 Flach, M., Gans, F., Brenning, A., Denzler, J., Reichstein, M., Rodner, E., et al. (2017). Multivariate anomaly detection  
48 for Earth observations: a comparison of algorithms and feature extraction techniques. *Earth Syst. Dyn.* 8, 677–  
49 696. doi:10.5194/esd-8-677-2017.
- 50 Flach, M., Sippel, S., Gans, F., Bastos, A., Brenning, A., Reichstein, M., et al. (2018). Contrasting biosphere responses  
51 to hydrometeorological extremes: revisiting the 2010 western Russian heatwave. *Biogeosciences* 15, 6067–6085.  
52 doi:10.5194/bg-15-6067-2018.
- 53 Flannigan, M., Wotton, M., Marshall, G., Groot, W., Johnston, J., Jurko, N., et al. (2016). Fuel moisture sensitivity to  
54 temperature and precipitation: climate change implications. *Clim. Change* 134, 59–71. doi:10.1007/s10584-015-  
55 1521-0.
- 56 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., et al. (2013). “Evaluation of Climate  
57 Models,” in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth  
58 Assessment Report of the Intergovernmental Panel on Climate Change*, eds. T. F. Stocker, D. Qin, G.-K. Plattner,  
59 M. Tignor, S. K. Allen, J. Boschung, et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge  
60 University Press), 741–866. doi:10.1017/CBO9781107415324.020.
- 61 Fontaine, B., Janicot, S., and Monerie, P.-A. (2013). Recent changes in air temperature, heat waves occurrences, and

- 1 atmospheric circulation in Northern Africa. *J. Geophys. Res. Atmos.* 118, 8536–8552. doi:10.1002/jgrd.50667.
- 2 Fontes, C. G., Dawson, T. E., Jardine, K., McDowell, N., Gimenez, B. O., Anderegg, L., et al. (2018). Dry and hot: The  
3 hydraulic consequences of a climate change–type drought for Amazonian trees. *Philos. Trans. R. Soc. B Biol. Sci.*  
4 373. doi:10.1098/rstb.2018.0209.
- 5 Ford, T. W., and Quiring, S. M. (2019). Comparison of Contemporary In Situ, Model, and Satellite Remote Sensing  
6 Soil Moisture With a Focus on Drought Monitoring. *Water Resour. Res.* 55, 1565–1582.  
7 doi:10.1029/2018WR024039.
- 8 Formayer, H., and Fritz, A. (2017). Temperature dependency of hourly precipitation intensities – surface versus cloud  
9 layer temperature. *Int. J. Climatol.* 37, 1–10. doi:10.1002/joc.4678.
- 10 Formetta, G., and Feyen, L. (2019). Empirical evidence of declining global vulnerability to climate-related hazards.  
11 *Glob. Environ. Chang.* 57, 101920. doi:10.1016/j.gloenvcha.2019.05.004.
- 12 Forzieri, G., Feyen, L., Rojas, R., Flörke, M., Wimmer, F., and Bianchi, A. (2014). Ensemble projections of future  
13 streamflow droughts in Europe. *Hydrol. Earth Syst. Sci.* 18, 85–108. doi:10.5194/hess-18-85-2014.
- 14 Forzieri, G., Feyen, L., Russo, S., Voutsoukas, M., Alfieri, L., Outten, S., et al. (2016). Multi-hazard assessment in  
15 Europe under climate change. *Clim. Change* 137, 105–119. doi:10.1007/s10584-016-1661-x.
- 16 Fotso-Nguemo, T. C., Chamani, R., Yepdo, Z. D., Sonkoué, D., Matsaguim, C. N., Vondou, D. A., et al. (2018).  
17 Projected trends of extreme rainfall events from CMIP5 models over Central Africa. *Atmos. Sci. Lett.* 19, e803.  
18 doi:10.1002/asl.803.
- 19 Fotso-Nguemo, T. C., Diallo, I., Diakhaté, M., Vondou, D. A., Mbaye, M. L., Haensler, A., et al. (2019). Projected  
20 changes in the seasonal cycle of extreme rainfall events from CORDEX simulations over Central Africa. *Clim.*  
21 *Change* 155, 339–357.
- 22 Fowler, H. J., Lenderink, G., Prein, A. F., Westra, S., Allan, R. P., Ban, N., et al. (2021). Anthropogenic intensification  
23 of short-duration rainfall extremes. *Nat. Rev. Earth Environ.* 2, 107–122. doi:10.1038/s43017-020-00128-6.
- 24 Francis, J. A., and Vavrus, S. J. (2012). Evidence linking Arctic amplification to extreme weather in mid-latitudes.  
25 *Geophys. Res. Lett.* 39. doi:10.1029/2012GL051000.
- 26 Frank, D. C., Poulter, B., Saurer, M., Esper, J., Huntingford, C., Helle, G., et al. (2015). Water-use efficiency and  
27 transpiration across European forests during the Anthropocene. *Nat. Clim. Chang.* 5, 579–583.  
28 doi:10.1038/nclimate2614.
- 29 Freund, M., Henley, B. J., Karoly, D. J., Allen, K. J., and Baker, P. J. (2017). Multi-century cool- and warm-season  
30 rainfall reconstructions for Australia’s major climatic regions. *Clim. Past* 13, 1751–1770. doi:10.5194/cp-13-  
31 1751-2017.
- 32 Freychet, N., Hsu, H.-H., Chou, C., and Wu, C.-H. (2015). Asian Summer Monsoon in CMIP5 Projections: A Link  
33 between the Change in Extreme Precipitation and Monsoon Dynamics. *J. Clim.* 28, 1477–1493.  
34 doi:10.1175/JCLI-D-14-00449.1.
- 35 Freychet, N., Tett, S. F. B., Hegerl, G. C., and Wang, J. (2018). Central-Eastern China Persistent Heat Waves:  
36 Evaluation of the AMIP Models. *J. Clim.* 31, 3609–3624. doi:10.1175/JCLI-D-17-0480.1.
- 37 Friedrich, K., Grossman, R. L., Huntington, J., Blanken, P. D., Lenters, J., Holman, K. D., et al. (2018). Reservoir  
38 evaporation in the Western United States. *Bull. Am. Meteorol. Soc.* 99, 167–187. doi:10.1175/BAMS-D-15-  
39 00224.1.
- 40 Frieler, K., Meinshausen, M., Mengel, M., Braun, N., and Hare, W. (2012). A Scaling Approach to Probabilistic  
41 Assessment of Regional Climate Change. *J. Clim.* 25, 3117–3144. doi:10.1175/JCLI-D-11-00199.1.
- 42 Frölicher, T. L., Fischer, E. M., and Gruber, N. (2018). Marine heatwaves under global warming. *Nature* 560, 360–364.  
43 doi:10.1038/s41586-018-0383-9.
- 44 Fu, G., Yu, J., Yu, X., Ouyang, R., Zhang, Y., Wang, P., et al. (2013a). Temporal variation of extreme rainfall events in  
45 China, 1961–2009. *J. Hydrol.* 487, 48–59. doi:10.1016/J.JHYDROL.2013.02.021.
- 46 Fu, Q., and Feng, S. (2014). Responses of terrestrial aridity to global warming. *J. Geophys. Res.* 119, 7863–7875.  
47 doi:10.1002/2014JD021608.
- 48 Fu, R., Yin, L., Li, W., Arias, P. A., Dickinson, R. E., Huang, L., et al. (2013b). Increased dry-season length over  
49 southern Amazonia in recent decades and its implication for future climate projection. *Proc. Natl. Acad. Sci. U. S.*  
50 *A.* 110, 18110–18115. doi:10.1073/pnas.1302584110.
- 51 Fuhrmann, C. M., Konrad II, C. E., Kovach, M. M., McLeod, J. T., Schmitz, W. G., and Dixon, P. G. (2014). Ranking  
52 of Tornado Outbreaks across the United States and Their Climatological Characteristics. *Weather Forecast.* 29,  
53 684–701. doi:10.1175/WAF-D-13-00128.1.
- 54 Fundel, F., Jörg-Hess, S., and Zappa, M. (2013). Monthly hydrometeorological ensemble prediction of streamflow  
55 droughts and corresponding drought indices. *Hydrol. Earth Syst. Sci.* 17, 395–407. doi:10.5194/hess-17-395-  
56 2013.
- 57 Funk, C., Davenport, F., Harrison, L., Magadzire, T., Galu, G., Artan, G. A., et al. (2018a). Anthropogenic  
58 Enhancement of Moderate-to-Strong El Niño Events Likely Contributed to Drought and Poor Harvests in  
59 Southern Africa During 2016. *Bull. Am. Meteorol. Soc.* 99, S91–S96. doi:10.1175/BAMS-D-17-0112.1.
- 60 Funk, C., Harrison, L., Shukla, S., Korecha, D., Magadzire, T., Husak, G., et al. (2016). Assessing the Contributions of  
61 Local and East Pacific Warming to the 2015 Droughts in Ethiopia and Southern Africa. *Bull. Am. Meteorol. Soc.*

- 1 97, S75–S80. doi:10.1175/BAMS-D-16-0167.1.
- 2 Funk, C., Harrison, L., Shukla, S., Pomposi, C., Galu, G., Korecha, D., et al. (2018b). Examining the role of unusually  
3 warm Indo-Pacific sea surface temperatures in recent African droughts. *Q. J. R. Meteorol. Soc.*, 1–20.  
4 doi:10.1002/acr.
- 5 Funk, C., Hoell, A., and Stone, D. (2014). Examining the contribution of the observed global warming trend to the  
6 California droughts of 2012/13 and 2013/14 [in “Explaining Extreme Events of 2013 from a Climate  
7 Perspective”]. *Bull. Amer. Meteor. Soc.* 95, S11–S13. doi:10.1175/1520-0477-95.9.S1.1.
- 8 Funk, C., Nicholson, S. E., Landsfeld, M., Klotter, D., Peterson, P., and Harrison, L. (2015a). The Centennial Trends  
9 Greater Horn of Africa precipitation dataset. *Sci. Data* 2, 150050. doi:10.1038/sdata.2015.50.
- 10 Funk, C., Peterson, P., Landsfeld, M., Davenport, F., Becker, A., Schneider, U., et al. (2020). “Algorithm and Data  
11 Improvements for Version 2.1 of the Climate Hazards Center’s InfraRed Precipitation with Stations Data Set,” in  
12 *Satellite Precipitation Measurement: Volume 1*, eds. V. Levizzani, C. Kidd, D. Kirschbaum, C. Kummerow, K.  
13 Nakamura, and F. Turk (Cham, Switzerland: Springer), 409–427. doi:10.1007/978-3-030-24568-9\_23.
- 14 Funk, C., Shukla, S., Hoell, A., and Livneh, B. (2015b). Assessing the contributions of east African and west pacific  
15 warming to the 2014 boreal spring east African drought. *Bull. Am. Meteorol. Soc.* 96, S77–S82.  
16 doi:10.1175/BAMS-D-15-00106.1.
- 17 Furrer, E., Katz, R., Walter, M., and Furrer, R. (2010). Statistical modeling of hot spells and heat waves. *Clim. Res.* 43,  
18 191–205. doi:10.3354/cr00924.
- 19 Gaertner, M. Á., González-Alemán, J. J., Romera, R., Domínguez, M., Gil, V., Sánchez, E., et al. (2018). Simulation of  
20 medicanes over the Mediterranean Sea in a regional climate model ensemble: impact of ocean–atmosphere  
21 coupling and increased resolution. *Clim. Dyn.* 51, 1041–1057. doi:10.1007/s00382-016-3456-1.
- 22 Gaertner, M. A., Jacob, D., Gil, V., Domínguez, M., Padorno, E., Sánchez, E., et al. (2007). Tropical cyclones over the  
23 Mediterranean Sea in climate change simulations. *Geophys. Res. Lett.* 34, L14711. doi:10.1029/2007GL029977.
- 24 Gajić-Čapka, M., Cindrić, K., and Pasarić, Z. (2015). Trends in precipitation indices in Croatia, 1961–2010. *Theor.*  
25 *Appl. Climatol.* 121, 167–177. doi:10.1007/s00704-014-1217-9.
- 26 Galarneau, T. J., Bosart, L. F., and Schumacher, R. S. (2010). Predecessor rain events ahead of tropical cyclones. *Mon.*  
27 *Weather Rev.* 138, 3272–3297. doi:10.1175/2010MWR3243.1.
- 28 Gallant, A. J. E., Reeder, M. J., Risbey, J. S., and Hennessy, K. J. (2013). The characteristics of seasonal-scale droughts  
29 in Australia, 1911–2009. *Int. J. Climatol.* 33, 1658–1672. doi:10.1002/joc.3540.
- 30 Gallo, F., Daron, J., Macadam, I., Cinco, T., Villafuerte II, M., Buonomo, E., et al. (2019). High-resolution regional  
31 climate model projections of future tropical cyclone activity in the Philippines. *Int. J. Climatol.* 39, 1181–1194.  
32 doi:10.1002/joc.5870.
- 33 Gálos, B., Hagemann, S., Hänsler, A., Kindermann, G., Rechid, D., Sieck, K., et al. (2013). Case study for the  
34 assessment of the biogeophysical effects of a potential afforestation in Europe. *Carbon Balance Manag.* 8, 3.  
35 doi:10.1186/1750-0680-8-3.
- 36 Gálos, B., Mátyás, C., and Jacob, D. (2011). Regional characteristics of climate change altering effects of afforestation.  
37 *Environ. Res. Lett.* 6, 044010. doi:10.1088/1748-9326/6/4/044010.
- 38 Gao, J., Shoshiro, M., Roberts, M. J., Haarsma, R., Putrasahan, D., Roberts, C. D., et al. (2020). Influence of model  
39 resolution on bomb cyclones revealed by HighResMIP-PRIMAVERA simulations. *Environ. Res. Lett.* 15,  
40 084001. doi:10.1088/1748-9326/ab88fa.
- 41 Gao, X., Shi, Y., Han, Z., Wang, M., Wu, J., Zhang, D., et al. (2017a). Performance of RegCM4 over major river basins  
42 in China. *Adv. Atmos. Sci.* 34, 441–455. doi:10.1007/s00376-016-6179-7.
- 43 Gao, X., Zhao, Q., Zhao, X., Wu, P., Pan, W., Gao, X., et al. (2017b). Temporal and spatial evolution of the  
44 standardized precipitation evapotranspiration index (SPEI) in the Loess Plateau under climate change from 2001  
45 to 2050. *Sci. Total Environ.* 595, 191–200. doi:10.1016/j.scitotenv.2017.03.226.
- 46 Gao, Y., Xiao, L., Chen, D., Xu, J., and Zhang, H. (2018). Comparison between past and future extreme precipitations  
47 simulated by global and regional climate models over the Tibetan Plateau. *Int. J. Climatol.* 38, 1285–1297.  
48 doi:10.1002/joc.5243.
- 49 García-Cueto, O. R., Santillán-Soto, N., López-Velázquez, E., Reyes-López, J., Cruz-Sotelo, S., and Ojeda-Benítez, S.  
50 (2019). Trends of climate change indices in some Mexican cities from 1980 to 2010. *Theor. Appl. Climatol.* 137,  
51 775–790. doi:10.1007/s00704-018-2620-4.
- 52 García-Garizábal, I., Causapé, J., Abrahao, R., and Merchan, D. (2014). Impact of Climate Change on Mediterranean  
53 Irrigation Demand: Historical Dynamics of Climate and Future Projections. *Water Resour. Manag.* 28, 1449–  
54 1462. doi:10.1007/s11269-014-0565-7.
- 55 García-Herrera, R., Garrido-Pérez, J.M. Barriopedro, D., Ordóñez, C., Vicente-Serrano, S.M. Nieto, R., Gimeno, L.,  
56 Sorí, R., et al. (2019). The European 2016/2017 drought. *J. Clim.* 32, 3169–3187.
- 57 Garner, A. J., Mann, M. E., Emanuel, K. A., Kopp, R. E., Lin, N., Alley, R. B., et al. (2017). Impact of climate change  
58 on New York City’s coastal flood hazard: Increasing flood heights from the preindustrial to 2300 CE. *Proc. Natl.*  
59 *Acad. Sci.* 114, 11861 LP-11866. doi:10.1073/pnas.1703568114.
- 60 Garreaud, R. D., Alvarez-Garretón, C., Barichivich, J., Boisier, J. P., Christie, D., Galleguillos, M., et al. (2017). The  
61 2010–2015 megadrought in central Chile: impacts on regional hydroclimate and vegetation. *Hydrol. Earth Syst.*

- 1 *Sci.* 21, 6307–6327. doi:10.5194/hess-21-6307-2017.
- 2 Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D. (2020). The  
3 Central Chile Mega Drought (2010–2018): A climate dynamics perspective. *Int. J. Climatol.* 40, 421–439.  
4 doi:10.1002/joc.6219.
- 5 Gaupp, F., Hall, J., Mitchell, D., and Dadson, S. (2019). Increasing risks of multiple breadbasket failure under 1.5 and 2  
6 °C global warming. *Agric. Syst.* 175, 34–45. doi:10.1016/j.agry.2019.05.010.
- 7 Ge, G., Shi, Z., Yang, X., Hao, Y., Guo, H., Kossi, F., et al. (2017). Analysis of precipitation extremes in the Qinghai-  
8 Tibetan plateau, China: Spatio-temporal characteristics and topography effects. *Atmosphere (Basel)*. 8, 1–16.  
9 doi:10.3390/atmos8070127.
- 10 Gebrechorkos, S. H., Hülsmann, S., and Bernhofer, C. (2018). Changes in temperature and precipitation extremes in  
11 Ethiopia, Kenya, and Tanzania. *Int. J. Climatol.* 39, 18–30. doi:10.1002/joc.5777.
- 12 Gebremeskel Haile, G., Tang, Q., Leng, G., Jia, G., Wang, J., Cai, D., et al. (2020). Long-term spatiotemporal variation  
13 of drought patterns over the Greater Horn of Africa. *Sci. Total Environ.* 704, 135299.  
14 doi:10.1016/j.scitotenv.2019.135299.
- 15 Geirinhas, J. L., Trigo, R. M., Libonati, R., Coelho, C. A. S., and Palmeira, A. C. (2018). Climatic and synoptic  
16 characterization of heat waves in Brazil. *Int. J. Climatol.* 38, 1760–1776. doi:10.1002/joc.5294.
- 17 Geng, X., Zhang, W., Stuecker, M. F., and Jin, F. F. (2017). Strong sub-seasonal wintertime cooling over East Asia and  
18 Northern Europe associated with super El Niño events. *Sci. Rep.* 7, 1–9. doi:10.1038/s41598-017-03977-2.
- 19 Gensini, V. A., and Brooks, H. E. (2018). Spatial trends in United States tornado frequency. *npj Clim. Atmos. Sci.* 1, 38.  
20 doi:10.1038/s41612-018-0048-2.
- 21 Gensini, V. A., and Mote, T. L. (2015). Downscaled estimates of late 21st century severe weather from CCSM3. *Clim.*  
22 *Change* 129, 307–321. doi:10.1007/s10584-014-1320-z.
- 23 Gergel, D. R., Nijssen, B., Abatzoglou, J. T., Lettenmaier, D. P., and Stumbaugh, M. R. (2017). Effects of climate  
24 change on snowpack and fire potential in the western USA. *Clim. Change* 141, 287–299. doi:10.1007/s10584-  
25 017-1899-y.
- 26 Gervais, M., Tremblay, L. B., Gyakum, J. R., and Atallah, E. (2014). Representing Extremes in a Daily Gridded  
27 Precipitation Analysis over the United States: Impacts of Station Density, Resolution, and Gridding Methods. *J.*  
28 *Clim.* 27, 5201–5218. doi:10.1175/JCLI-D-13-00319.1.
- 29 Ghausi, S. A., and Ghosh, S. (2020). Diametrically Opposite Scaling of Extreme Precipitation and Streamflow to  
30 Temperature in South and Central Asia. *Geophys. Res. Lett.* 47, e2020GL089386. doi:10.1029/2020GL089386.
- 31 Gibba, P., Sylla, M. B., Okogbue, E. C., Gaye, A. T., Nikiema, M., and Kebe, I. (2019). State-of-the-art climate  
32 modeling of extreme precipitation over Africa: analysis of CORDEX added-value over CMIP5. *Theor. Appl.*  
33 *Climatol.* 137, 1041–1057. doi:10.1007/s00704-018-2650-y.
- 34 Gimeno, L., Stohl, A., Trigo, R. M., Dominguez, F., Yoshimura, K., Yu, L., et al. (2012). Oceanic and terrestrial  
35 sources of continental precipitation. *Rev. Geophys.* 50. doi:10.1029/2012RG000389.
- 36 Gimeno, L., Vázquez, M., Eiras-Barca, J., Sorí, R., Stojanovic, M., Algarra, I., et al. (2020). Recent progress on the  
37 sources of continental precipitation as revealed by moisture transport analysis. *Earth-Science Rev.* 201.  
38 doi:10.1016/j.earscirev.2019.103070.
- 39 Giorgi, F., Coppola, E., Raffaele, F., Diro, G. T., Fuentes-Franco, R., Giuliani, G., et al. (2014). Changes in extremes  
40 and hydroclimatic regimes in the CREMA ensemble projections. *Clim. Change* 125, 39–51. doi:10.1007/s10584-  
41 014-1117-0.
- 42 Giorgi, F., Jones, C., and Asrar, G. (2009). Addressing climate information needs at the regional level: the CORDEX  
43 framework. *WMO Bull.* 58, 175.
- 44 Giorgi, F., Raffaele, F., and Coppola, E. (2019). The response of precipitation characteristics to global warming from  
45 climate projections. *Earth Syst. Dyn.* 10, 73–89. doi:10.5194/esd-10-73-2019.
- 46 Giuntoli, I., Renard, B., Vidal, J.-P., and Bard, A. (2013). Low flows in France and their relationship to large-scale  
47 climate indices. *J. Hydrol.* 482, 105–118. doi:10.1016/j.jhydrol.2012.12.038.
- 48 Giuntoli, I., Vidal, J.-P., Prudhomme, C., and Hannah, D. M. (2015). Future hydrological extremes: The uncertainty  
49 from multiple global climate and global hydrological models. *Earth Syst. Dyn.* 6, 267–285. doi:10.5194/esd-6-  
50 267-2015.
- 51 Glas, R., Burns, D., and Lautz, L. (2019). Historical changes in New York State streamflow: Attribution of temporal  
52 shifts and spatial patterns from 1961 to 2016. *J. Hydrol.* 574, 308–323. doi:10.1016/j.jhydrol.2019.04.060.
- 53 Gleixner, S., Demissie, T., and Diro, G. T. (2020). Did ERA5 Improve Temperature and Precipitation Reanalysis over  
54 East Africa? *Atmosphere (Basel)*. 11, 996. doi:10.3390/atmos11090996.
- 55 Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., and Stoffel, M. (2014). 21st century climate change in  
56 the European Alps – A review. *Sci. Total Environ.* 493, 1138–1151. doi:10.1016/j.scitotenv.2013.07.050.
- 57 Gocic, M., and Trajkovic, S. (2014). Analysis of trends in reference evapotranspiration data in a humid climate. *Hydrol.*  
58 *Sci. J.* 59, 165–180. doi:10.1080/02626667.2013.798659.
- 59 González-Alemán, J. J., Pascale, S., Gutierrez-Fernandez, J., Murakami, H., Gaertner, M. A., and Vecchi, G. A. (2019).  
60 Potential Increase in Hazard From Mediterranean Hurricane Activity With Global Warming. *Geophys. Res. Lett.*  
61 46, 1754–1764. doi:10.1029/2018GL081253.

- 1 González-Hidalgo, J. C., Vicente-Serrano, S. M., Peña-Angulo, D., Salinas, C., Tomas-Burguera, M., and Beguería, S.  
2 (2018). High-resolution spatio-temporal analyses of drought episodes in the western Mediterranean basin  
3 (Spanish mainland, Iberian Peninsula). *Acta Geophys.* 66, 381–392. doi:10.1007/s11600-018-0138-x.
- 4 Gosling, S. N., Zaherpour, J., Mount, N. J., Hattermann, F. F., Dankers, R., Arheimer, B., et al. (2017). A comparison  
5 of changes in river runoff from multiple global and catchment-scale hydrological models under global warming  
6 scenarios of 1 °C, 2 °C and 3 °C. *Clim. Change* 141, 577–595. doi:10.1007/s10584-016-1773-3.
- 7 Gou, X., Deng, Y., Gao, L., Chen, F., Cook, E., Yang, M., et al. (2015). Millennium tree-ring reconstruction of drought  
8 variability in the eastern Qilian Mountains, northwest China. *Clim. Dyn.* 45, 1761–1770. doi:10.1007/s00382-  
9 014-2431-y.
- 10 Govekar, P. D., Jakob, C., and Catto, J. (2014). The relationship between clouds and dynamics in Southern Hemisphere  
11 extratropical cyclones in the real world and a climate model. *J. Geophys. Res. Atmos.*  
12 doi:10.1002/2013JD020699.
- 13 Graham, R. M., Cohen, L., Petty, A. A., Boisvert, L. N., Rinke, A., Hudson, S. R., et al. (2017). Increasing frequency  
14 and duration of Arctic winter warming events. *Geophys. Res. Lett.* 44, 6974–6983. doi:10.1002/2017GL073395.
- 15 Green, J. K., Seneviratne, S. I., Berg, A. M., Findell, K. L., Hagemann, S., Lawrence, D. M., et al. (2019). Large  
16 influence of soil moisture on long-term terrestrial carbon uptake. *Nature* 565, 476–479. doi:10.1038/s41586-018-  
17 0848-x.
- 18 Greenbaum, N., Harden, T. M., Baker, V. R., Weisheit, J., Cline, M. L., Porat, N., et al. (2014). A 2000 year natural  
19 record of magnitudes and frequencies for the largest Upper Colorado River floods near Moab, Utah. *Water*  
20 *Resour. Res.* 50, 5249–5269. doi:10.1002/2013WR014835.
- 21 Greve, P., Gudmundsson, L., and Seneviratne, S. I. (2018). Regional scaling of annual mean precipitation and water  
22 availability with global temperature change. *Earth Syst. Dyn.* 9, 227–240. doi:10.3929/ethz-b-000251688.
- 23 Greve, P., Orłowsky, B., Mueller, B., Sheffield, J., Reichstein, M., and Seneviratne, S. I. (2014). Global assessment of  
24 trends in wetting and drying over land. *Nat. Geosci.* 7, 716–721. doi:10.1038/NGEO2247.
- 25 Greve, P., Roderick, M. L., and Seneviratne, S. I. (2017). Simulated changes in aridity from the last glacial maximum to  
26 4xCO<sub>2</sub>. *Environ. Res. Lett.* 12, 114021. doi:10.1088/1748-9326/aa89a3.
- 27 Greve, P., Roderick, M. L., Ukkola, A. M., and Wada, Y. (2019). The aridity Index under global warming. *Environ.*  
28 *Res. Lett.* 14, 124006. doi:10.1088/1748-9326/ab5046.
- 29 Griffin, D., and Anchukaitis, K. J. (2014). How unusual is the 2012–2014 California drought? *Geophys. Res. Lett.* 41,  
30 9017–9023. doi:10.1002/2014GL062433.
- 31 Grillakis, M. G., Koutroulis, A. G., Komma, J., Tsanis, I. K., Wagner, W., and Blöschl, G. (2016). Initial soil moisture  
32 effects on flash flood generation – A comparison between basins of contrasting hydro-climatic conditions. *J.*  
33 *Hydrol.* 541, 206–217. doi:10.1016/J.JHYDROL.2016.03.007.
- 34 Grinsted, A., Ditlevsen, P., and Christensen, J. H. (2019). Normalized US hurricane damage estimates using area of  
35 total destruction, 1900–2018. *Proc. Natl. Acad. Sci.* 116, 23942 LP-23946. doi:10.1073/pnas.1912277116.
- 36 Grose, M. R., Black, M., Risbey, J. S., Uhe, P., Hope, P. K., Haustein, K., et al. (2018). Severe Frosts in Western  
37 Australia in September 2016. *Bull. Am. Meteorol. Soc.* 99, S150–S154. doi:10.1175/BAMS-D-17-0088.1.
- 38 Grose, M. R., Narsey, S., Delage, F. P., Dowdy, A. J., Bador, M., Boschat, G., et al. (2020). Insights From CMIP6 for  
39 Australia’s Future Climate. *Earth’s Futur.* 8. doi:10.1029/2019EF001469.
- 40 Gross, M. H., Donat, M. G., Alexander, L. V., and Sherwood, S. C. (2020). Amplified warming of seasonal cold  
41 extremes relative to the mean in the Northern Hemisphere extratropics. *Earth Syst. Dyn.* 11, 97–111.  
42 doi:10.5194/esd-11-97-2020.
- 43 Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., et al. (2020). Plant  
44 responses to rising vapor pressure deficit. *New Phytol.* 226, 1550–1566. doi:10.1111/nph.16485.
- 45 Grotjahn, R., Black, R., Leung, R., Wehner, M. F., Barlow, M., Bosilovich, M., et al. (2016). North American extreme  
46 temperature events and related large scale meteorological patterns: a review of statistical methods, dynamics,  
47 modeling, and trends. *Clim. Dyn.* 46, 1151–1184. doi:10.1007/s00382-015-2638-6.
- 48 Gu, G., and Adler, R. F. (2018). Precipitation Intensity Changes in the Tropics from Observations and Models. *J. Clim.*  
49 31, 4775–4790. doi:10.1175/JCLI-D-17-0550.1.
- 50 Gu, L., Chen, J., Yin, J., Sullivan, S. C., Wang, H.-M., Guo, S., et al. (2020). Projected increases in magnitude and  
51 socioeconomic exposure of global droughts in 1.5 and 2 °C warmer climates. *Hydrol. Earth Syst. Sci.* 24, 451–  
52 472. doi:10.5194/hess-24-451-2020.
- 53 Gu, X., Li, J., Chen, Y. D., Kong, D., and Liu, J. (2019a). Consistency and Discrepancy of Global Surface Soil  
54 Moisture Changes From Multiple Model-Based Data Sets Against Satellite Observations. *J. Geophys. Res. Atmos.*  
55 124, 1474–1495. doi:10.1029/2018JD029304.
- 56 Gu, X., Zhang, Q., Li, J., Singh, V. P., Liu, J., Sun, P., et al. (2019b). Attribution of Global Soil Moisture Drying to  
57 Human Activities: A Quantitative Viewpoint. *Geophys. Res. Lett.* 46, 2573–2582. doi:10.1029/2018GL080768.
- 58 Gudmundsson, L., Boulange, J., Do, H. X., Gosling, S. N., Grillakis, M. G., Koutroulis, A. G., et al. (2021). Globally  
59 observed trends in mean and extreme river flow attribution to climate change. *Science (80- )*.  
60 doi:10.1126/science.aba3996.
- 61 Gudmundsson, L., Leonard, M., Do, H. X., Westra, S., and Seneviratne, S. I. (2019). Observed Trends in Global

- 1 Indicators of Mean and Extreme Streamflow. *Geophys. Res. Lett.* 46, 756–766. doi:10.1029/2018GL079725.
- 2 Gudmundsson, L., and Seneviratne, S. I. (2016). Anthropogenic climate change affects meteorological drought risk in  
3 Europe. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/4/044005.
- 4 Gudmundsson, L., Seneviratne, S. I., and Zhang, X. (2017). Anthropogenic climate change detected in European  
5 renewable freshwater resources. *Nat. Clim. Chang.* 7, 813–816. doi:10.1038/nclimate3416.
- 6 Guerreiro, S. B., Dawson, R. J., Kilsby, C., Lewis, E., and Ford, A. (2018a). Future heat-waves, droughts and floods in  
7 571 European cities. *Environ. Res. Lett.* 13, 034009. doi:10.1088/1748-9326/aaaad3.
- 8 Guerreiro, S. B., Fowler, H. J., Barbero, R., Westra, S., Lenderink, G., Blenkinsop, S., et al. (2018b). Detection of  
9 continental-scale intensification of hourly rainfall extremes. *Nat. Clim. Chang.* 8, 803–807. doi:10.1038/s41558-  
10 018-0245-3.
- 11 Guhathakurta, P., Menon, P., Inkane, P. M., Krishnan, U., and Sable, S. T. (2017). Trends and variability of  
12 meteorological drought over the districts of India using standardized precipitation index. *J. Earth Syst. Sci.* 126,  
13 120. doi:10.1007/s12040-017-0896-x.
- 14 Guichard, F., and Couvreur, F. (2017). A short review of numerical cloud-resolving models. *Tellus, Ser. A Dyn.*  
15 *Meteorol. Oceanogr.* 69, 1–36. doi:10.1080/16000870.2017.1373578.
- 16 Guillod, B. P., Orłowsky, B., Miralles, D. G., Teuling, A. J., and Seneviratne, S. I. (2015). Reconciling spatial and  
17 temporal soil moisture effects on afternoon rainfall. *Nat. Commun.* 6, 6443. doi:10.1038/ncomms7443.
- 18 Guimberteau, M., Ronchail, J., Espinoza, J. C. C., Lengaigne, M., Sultan, B., Polcher, J., et al. (2013). Future changes  
19 in precipitation and impacts on extreme streamflow over Amazonian sub-basins. *Environ. Res. Lett.* 8, 014035.  
20 doi:10.1088/1748-9326/8/1/014035.
- 21 Guo, J., Huang, G., Wang, X., Li, Y., and Lin, Q. (2018). Dynamically-downscaled projections of changes in  
22 temperature extremes over China. *Clim. Dyn.* 50, 1045–1066. doi:10.1007/s00382-017-3660-7.
- 23 Guo, X., Huang, J., Luo, Y., Zhao, Z., and Xu, Y. (2016). Projection of precipitation extremes for eight global warming  
24 targets by 17 CMIP5 models. *Nat. Hazards* 84, 2299–2319. doi:10.1007/s11069-016-2553-0.
- 25 Guo, X., Huang, J., Luo, Y., Zhao, Z., and Xu, Y. (2017). Projection of heat waves over China for eight different global  
26 warming targets using 12 CMIP5 models. *Theor. Appl. Climatol.* 128, 507–522. doi:10.1007/s00704-015-1718-1.
- 27 Gusain, A., Ghosh, S., and Karmakar, S. (2020). Added value of CMIP6 over CMIP5 models in simulating Indian  
28 summer monsoon rainfall. *Atmos. Res.* 232, 104680. doi:10.1016/j.atmosres.2019.104680.
- 29 Gutmann, E. D., Rasmussen, R. M., Liu, C., Ikeda, K., Bruyere, C. L., Done, J. M., et al. (2018). Changes in hurricanes  
30 from a 13-Yr convection-permitting pseudo- global warming simulation. *J. Clim.* 31, 3643–3657.  
31 doi:10.1175/JCLI-D-17-0391.1.
- 32 Haarsma, R. J., Roberts, M. J., Vidale, P. L., Senior, C. A., Bellucci, A., Bao, Q., et al. (2016). High Resolution Model  
33 Intercomparison Project (HighResMIP v1.0) for CMIP6. *Geosci. Model Dev.* 9, 4185–4208. doi:10.5194/gmd-9-  
34 4185-2016.
- 35 Habib, S. M. A., Sato, T., and Hatsuzuka, D. (2019). Decreasing number of propagating mesoscale convective systems  
36 in Bangladesh and surrounding area during 1998–2015. *Atmos. Sci. Lett.* 20, e879. doi:10.1002/asl.879.
- 37 Haig, J., Nott, J., and Reichert, G.-J. (2014). Australian tropical cyclone activity lower than at any time over the past  
38 550–1,500 years. *Nature* 505, 667–671. doi:10.1038/nature12882.
- 39 Hall, A., and Qu, X. (2006). Using the current seasonal cycle to constrain snow albedo feedback in future climate  
40 change. *Geophys. Res. Lett.* 33, L03502. doi:10.1029/2005GL025127.
- 41 Hall, J., Arheimer, B., Borga, M., Brázdil, R., Claps, P., Kiss, A., et al. (2014). Understanding flood regime changes in  
42 Europe: A state-of-the-art assessment. *Hydrol. Earth Syst. Sci.* 18, 2735–2772. doi:10.5194/hess-18-2735-2014.
- 43 Hall, T. C., Sealy, A. M., Stephenson, T. S., Kusunoki, S., Taylor, M. A., Chen, A. A., et al. (2013). Future climate of  
44 the Caribbean from a super-high-resolution atmospheric general circulation model. *Theor. Appl. Climatol.* 113,  
45 271–287. doi:10.1007/s00704-012-0779-7.
- 46 Hall, T. M., and Kossin, J. P. (2019). Hurricane stalling along the North American coast and implications for rainfall.  
47 *npj Clim. Atmos. Sci.* 2, 1–9. doi:10.1038/s41612-019-0074-8.
- 48 Hamada, A., and Takayabu, Y. N. (2018). Large-scale environmental conditions related to midsummer extreme rainfall  
49 events around Japan in the TRMM region. *J. Clim.* 0, null. doi:10.1175/JCLI-D-17-0632.1.
- 50 Hamada, A., Takayabu, Y. N., Liu, C., and Zipser, E. J. (2015). Weak linkage between the heaviest rainfall and tallest  
51 storms. *Nat. Commun.* 6, 1–6. doi:10.1038/ncomms7213.
- 52 Han, F., Cook, K. H., and Vizy, E. K. (2019). Changes in intense rainfall events and dry periods across Africa in the  
53 twenty-first century. *Clim. Dyn.*, 1–21. doi:10.1007/s00382-019-04653-z.
- 54 Han, J.-Y., Baik, J.-J., and Khain, A. P. (2011). A Numerical Study of Urban Aerosol Impacts on Clouds and  
55 Precipitation. *J. Atmos. Sci.* 69, 504–520. doi:10.1175/JAS-D-11-071.1.
- 56 Han, T., Chen, H., Hao, X., and Wang, H. (2018). Projected changes in temperature and precipitation extremes over the  
57 Silk Road Economic Belt regions by the Coupled Model Intercomparison Project Phase 5 multi-model ensembles.  
58 *Int. J. Climatol.* 38, 4077–4091. doi:10.1002/joc.5553.
- 59 Hande, L. B., Siems, S. T., and Manton, M. J. (2012). Observed Trends in Wind Speed over the Southern Ocean.  
60 *Geophys. Res. Lett.* 39, 1–5. doi:10.1029/2012GL051734.
- 61 Hanel, M., Rakovec, O., Markonis, Y., Máca, P., Samaniego, L., Kyselý, J., et al. (2018). Revisiting the recent

- 1 European droughts from a long-term perspective. *Sci. Rep.* 8. doi:10.1038/s41598-018-27464-4.
- 2 Hannaford, J. (2015). Climate-driven changes in UK river flows. *Prog. Phys. Geogr. Earth Environ.* 39, 29–48.
- 3 doi:10.1177/0309133314536755.
- 4 Hao, Y., Liu, Q., Li, C., Kharel, G., An, L., Stebler, E., et al. (2019). Interactive effect of meteorological drought and
- 5 vegetation types on root zone soil moisture and runoff in Rangeland Watersheds. *Water (Switzerland)* 11.
- 6 doi:10.3390/w11112357.
- 7 Hao, Z., Hao, F., Singh, V. P., and Zhang, X. (2018). Changes in the severity of compound drought and hot extremes
- 8 over global land areas. *Environ. Res. Lett.* 13, 124022. doi:10.1088/1748-9326/aace96.
- 9 Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R. (2020). Increased future occurrences of the exceptional
- 10 2018–2019 Central European drought under global warming. *Sci. Rep.* 10, 12207. doi:10.1038/s41598-020-
- 11 68872-9.
- 12 Harrigan, S., Hannaford, J., Muchan, K., and Marsh, T. J. (2018). Designation and trend analysis of the updated UK
- 13 Benchmark Network of river flow stations: The UKBN2 dataset. *Hydrol. Res.* 49, 552–567.
- 14 doi:10.2166/nh.2017.058.
- 15 Harrington, L. J. (2017). Investigating differences between event-as-class and probability density-based attribution
- 16 statements with emerging climate change. *Clim. Change* 141, 641–654. doi:10.1007/s10584-017-1906-3.
- 17 Harrington, L. J. (2020). Rethinking extreme heat in a cool climate: a New Zealand case study. *Environ. Res. Lett.*
- 18 doi:10.1088/1748-9326/abbd61.
- 19 Harrington, L. J., Gibson, P. B., Dean, S. M., Mitchell, D., Rosier, S. M., and Frame, D. J. (2016). Investigating event-
- 20 specific drought attribution using self-organizing maps. *J. Geophys. Res. Atmos.* 121, 12,712–766,780.
- 21 doi:10.1002/2016JD025602.
- 22 Harrington, L. J., and Otto, F. E. L. (2018a). Adapting attribution science to the climate extremes of tomorrow. *Environ.*
- 23 *Res. Lett.* 13, 123006. doi:10.1088/1748-9326/aaf4cc.
- 24 Harrington, L. J., and Otto, F. E. L. (2018b). Changing population dynamics and uneven temperature emergence
- 25 combine to exacerbate regional exposure to heat extremes under 1.5 °C and 2 °C of warming. *Environ. Res. Lett.*
- 26 13, 034011. doi:10.1088/1748-9326/aaaa99.
- 27 Harrington, L., Rosier, S., Dean, S. M., Stuart, S., and Scahill, A. (2014). The Role of Anthropogenic Climate Change
- 28 in the 2013 Drought Over North Island, New Zealand [in “Explaining Extreme Events of 2013 from a Climate
- 29 Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S45–S48. doi:10.1175/1520-0477-95.9.S1.1.
- 30 Harrington, and Renwick (2014). Secular changes in New Zealand rainfall characteristics 1950–2009. *Weather Clim.*
- 31 34, 50. doi:10.2307/26169744.
- 32 Harris, L. M., Lin, S. J., and Tu, C. Y. (2016). High-resolution climate simulations using GFDL HiRAM with a
- 33 stretched global grid. *J. Clim.* 29, 4293–4314. doi:10.1175/JCLI-D-15-0389.1.
- 34 Harrison, L., Funk, C., and Peterson, P. (2019). Identifying changing precipitation extremes in Sub-Saharan Africa with
- 35 gauge and satellite products. *Environ. Res. Lett.* 14, 085007. doi:10.1088/1748-9326/ab2cae.
- 36 Hartmann, D. J., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Brönnimann, S., Charabi, Y. A.-R., et al.
- 37 (2013). “Observations: Atmosphere and Surface,” in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
- 38 *Change*, eds. T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, et al. (Cambridge, United
- 39 Kingdom and New York, NY, USA: Cambridge University Press), 159–254.
- 40 doi:10.1017/CBO9781107415324.008.
- 41
- 42 Hartmann, F., Merten, J., Fink, M., and Faust, H. (2018). Indonesia’s Fire Crisis 2015 A Twofold Perturbation on the
- 43 Ground. *Pacific Geogr.* 49, 4–11. doi:10.23791/490411.
- 44 Hartmann, H. (2015). Carbon starvation during drought-induced tree mortality – are we chasing a myth? *J. Plant*
- 45 *Hydraul.* 2, e005. doi:10.20870/jph.2015.e005.
- 46 Hartzell, S., Bartlett, M. S., and Porporato, A. (2017). The role of plant water storage and hydraulic strategies in relation
- 47 to soil moisture availability. *Plant Soil* 419, 503–521. doi:10.1007/s11104-017-3341-7.
- 48 Hasan, H. H., Mohd Razali, S. F., Muhammad, N. S., and Ahmad, A. (2019). Research Trends of Hydrological
- 49 Drought: A Systematic Review. *Water* 11. doi:10.3390/w11112252.
- 50 Hashimoto, A., Done, J. M., Fowler, L. D., and Bruyère, C. L. (2016). Tropical cyclone activity in nested regional and
- 51 global grid-refined simulations. *Clim. Dyn.* 47, 497–508. doi:10.1007/s00382-015-2852-2.
- 52 Haslinger, K., Hofstätter, M., Kroisleitner, C., Schöner, W., Laaha, G., Holawe, F., et al. (2019). Disentangling Drivers
- 53 of Meteorological Droughts in the European Greater Alpine Region During the Last Two Centuries. *J. Geophys.*
- 54 *Res. Atmos.* doi:10.1029/2018JD029527.
- 55 Hassanzadeh, P., Lee, C.-Y., Nabizadeh, E., Camargo, S. J., Ma, D., and Yeung, L. Y. (2020). Effects of climate change
- 56 on the movement of future landfalling Texas tropical cyclones. *Nat. Commun.* 11, 3319. doi:10.1038/s41467-020-
- 57 17130-7.
- 58 Hatsuzuka, D., and Sato, T. (2019). Future Changes in Monthly Extreme Precipitation in Japan Using Large-Ensemble
- 59 Regional Climate Simulations. *J. Hydrometeorol.* 20, 563–574. doi:10.1175/JHM-D-18-0095.1.
- 60 Hatsuzuka, D., Sato, T., Yoshida, K., Ishii, M., and Mizuta, R. (2020). Regional projection of tropical-cyclone-induced
- 61 extreme precipitation around Japan based on large ensemble simulations. *SOLA advpub*, 23–29.



- 1 doi:10.2151/sola.2020-005.
- 2 Hauser, M. (2021). Mean temperature anomalies for CMIP5 and CMIP6 (Version v0.1.0).
- 3 doi:10.5281/ZENODO.4600696.
- 4 Hauser, M., Engelbrecht, F., and Fischer, E. M. (2021). Transient global warming levels for CMIP5 and CMIP6
- 5 (Version v0.2.0). doi:10.5281/ZENODO.4600706.
- 6 Hauser, M., Gudmundsson, L., Orth, R., Jézéquel, A., Haustein, K., Vautard, R., et al. (2017). Methods and Model
- 7 Dependency of Extreme Event Attribution: The 2015 European Drought. *Earth's Futur.* 5, 1034–1043.
- 8 doi:10.1002/2017EF000612.
- 9 Hauser, M., Orth, R., and Seneviratne, S. I. (2016). Role of soil moisture versus recent climate change for the 2010 heat
- 10 wave in western Russia. *Geophys. Res. Lett.* 43, 2819–2826. doi:10.1002/2016GL068036.
- 11 Hawcroft, M., Walsh, E., Hodges, K., and Zappa, G. (2018). Significantly increased extreme precipitation expected in
- 12 Europe and North America from extratropical cyclones. *Environ. Res. Lett.* 13. doi:10.1088/1748-9326/aaed59.
- 13 Hayat, F., Ahmed, M. A., Zarebanadkouki, M., Cai, G., and Carminati, A. (2019). Measurements and simulation of leaf
- 14 xylem water potential and root water uptake in heterogeneous soil water contents. *Adv. Water Resour.* 124, 96–
- 15 105. doi:10.1016/j.advwatres.2018.12.009.
- 16 Hazeleger, W., van den Hurk, B. J. J. M., Min, E., van Oldenborgh, G. J., Petersen, A. C., Stainforth, D. A., et al.
- 17 (2015). Tales of future weather. *Nat. Clim. Chang.* 5, 107–113. doi:10.1038/nclimate2450.
- 18 He, B.-R., and Zhai, P.-M. (2018). Changes in persistent and non-persistent extreme precipitation in China from 1961 to
- 19 2016. *Adv. Clim. Chang. Res.* 9, 177–184. doi:10.1016/j.accre.2018.08.002.
- 20 He, F., and Posselt, D. J. (2015). Impact of Parameterized Physical Processes on Simulated Tropical Cyclone
- 21 Characteristics in the Community Atmosphere Model. *J. Clim.* 28, 9857–9872. doi:10.1175/JCLI-D-15-0255.1.
- 22 He, X., Wada, Y., Wanders, N., and Sheffield, J. (2017). Intensification of hydrological drought in California by human
- 23 water management. *Geophys. Res. Lett.* 44, 1777–1785. doi:10.1002/2016GL071665.
- 24 Helama, S., Sohar, K., Läänelaid, A., Bijak, S., and Jaagus, J. (2018). Reconstruction of precipitation variability in
- 25 Estonia since the eighteenth century, inferred from oak and spruce tree rings. *Clim. Dyn.* 50, 4083–4101.
- 26 doi:10.1007/s00382-017-3862-z.
- 27 Held, I. M., and Soden, B. J. (2006). Robust responses of the hydrological cycle to global warming. *J. Clim.* 19, 5686–
- 28 5699. doi:10.1175/JCLI3990.1.
- 29 Held, I. M., and Zhao, M. (2011). The Response of Tropical Cyclone Statistics to an Increase in CO<sub>2</sub> with Fixed Sea
- 30 Surface Temperatures. *J. Clim.* 24, 5353–5364. doi:10.1175/JCLI-D-11-00050.1.
- 31 Helsen, S., van Lipzig, N. P. M., Demuzere, M., Vanden Broucke, S., Caluwaerts, S., De Cruz, L., et al. (2020).
- 32 Consistent scale-dependency of future increases in hourly extreme precipitation in two convection-permitting
- 33 climate models. *Clim. Dyn.* 54, 1267–1280. doi:10.1007/s00382-019-05056-w.
- 34 Herger, N., Angéllil, O., Abramowitz, G., Donat, M., Stone, D., and Lehmann, K. (2018). Calibrating Climate Model
- 35 Ensembles for Assessing Extremes in a Changing Climate. *J. Geophys. Res. Atmos.* 123, 5988–6004.
- 36 doi:10.1029/2018JD028549.
- 37 Herger, N., Sanderson, B. M., and Knutti, R. (2015). Improved pattern scaling approaches for the use in climate impact
- 38 studies. *Geophys. Res. Lett.* 42, 3486–3494. doi:10.1002/2015GL063569.
- 39 Herold, N., Ekström, M., Kala, J., Goldie, J., and Evans, J. P. (2018). Australian climate extremes in the 21st century
- 40 according to a regional climate model ensemble: Implications for health and agriculture. *Weather Clim. Extrem.*
- 41 20, 54–68. doi:10.1016/j.wace.2018.01.001.
- 42 Herrera-Estrada, J. E., Martinez, J. A., Dominguez, F., Findell, K. L., Wood, E. F., and Sheffield, J. (2019). Reduced
- 43 Moisture Transport Linked to Drought Propagation Across North America. *Geophys. Res. Lett.* 46, 5243–5253.
- 44 doi:10.1029/2019GL082475.
- 45 Herrera-Estrada, J. E., and Sheffield, J. (2017). Uncertainties in future projections of summer droughts and heat waves
- 46 over the contiguous United States. *J. Clim.* 30, 6225–6246. doi:10.1175/JCLI-D-16-0491.1.
- 47 Herrera-Estrada, J. E., Satoh, Y., and Sheffield, J. (2017). Spatiotemporal dynamics of global drought. *Geophys. Res.*
- 48 *Lett.* 44, 2254–2263. doi:10.1002/2016GL071768.
- 49 Herring, S. C., Christidis, N., Hoell, A., Hoerling, M. P., and Stott, P. A. (2019). Explaining Extreme Events of 2017
- 50 from a Climate Perspective. *Bull. Am. Meteorol. Soc.* 100, S1–S117. doi:10.1175/bams-
- 51 explainingextremeevents2017.1.
- 52 Herring, S. C., Christidis, N., Hoell, A., Hoerling, M. P., and Stott, P. A. (2020). Explaining Extreme Events of 2018
- 53 from a Climate Perspective. *Bull. Am. Meteorol. Soc.* 101, S1–S140. doi:10.1175/BAMS-
- 54 ExplainingExtremeEvents2018.1.
- 55 Herring, S. C., Christidis, N., Hoell, A., Kossin, J. P., Schreck, C. J., and Stott, P. A. (2018). Explaining Extreme
- 56 Events of 2016 from a Climate Perspective. *Bull. Am. Meteorol. Soc.* 99, S1–S157. doi:10.1175/BAMS-
- 57 ExplainingExtremeEvents2016.1.
- 58 Herring, S. C., Hoell, A., Hoerling, M. P., Kossin, J. P., Schreck, C. J., and Stott, P. A. (2016). Explaining Extreme
- 59 Events of 2015 from a Climate Perspective. *Bull. Am. Meteorol. Soc.* 97, S1–S145. doi:10.1175/BAMS-
- 60 ExplainingExtremeEvents2015.1.
- 61 Herring, S. C., Hoerling, M. P., Kossin, J. P., Peterson, T. C., and Stott, P. A. (2015). Explaining Extreme Events of

- 1 2014 from a Climate Perspective. *Bull. Am. Meteorol. Soc.* 96, S1–S172. doi:10.1175/BAMS-  
2 ExplainingExtremeEvents2014.1.
- 3 Herring, S. C., Hoerling, M. P., Peterson, T. C., and Stott, P. a. (2014). Explaining Extreme Events of 2013 from a  
4 Climate Perspective. *Bull. Am. Meteorol. Soc.* 95, S1–S104. doi:10.1175/1520-0477-95.9.S1.1.
- 5 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. (2020). The ERA5 global  
6 reanalysis. *Q. J. R. Meteorol. Soc.* 146, 1999–2049. doi:10.1002/qj.3803.
- 7 Hertig, E., and Trambly, Y. (2017). Regional downscaling of Mediterranean droughts under past and future climatic  
8 conditions. *Glob. Planet. Change* 151, 36–48. doi:10.1016/j.gloplacha.2016.10.015.
- 9 Hettiarachchi, S., Wasko, C., and Sharma, A. (2018). Increase in flood risk resulting from climate change in a  
10 developed urban watershed - The role of storm temporal patterns. *Hydrol. Earth Syst. Sci.* 22, 2041–2056.  
11 doi:10.5194/hess-22-2041-2018.
- 12 Hidalgo, H. G., Alfaro, E. J., and Quesada-Montano, B. (2017). Observed (1970–1999) climate variability in Central  
13 America using a high-resolution meteorological dataset with implication to climate change studies. *Clim. Change*  
14 141, 13–28. doi:10.1007/s10584-016-1786-y.
- 15 Hillier, J. K., Matthews, T., Wilby, R. L., and Murphy, C. (2020). Multi-hazard dependencies can increase or decrease  
16 risk. *Nat. Clim. Chang.* 10, 595–598. doi:10.1038/s41558-020-0832-y.
- 17 Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., et al. (2013). Global flood  
18 risk under climate change. *Nat. Clim. Chang.* 3, 816–821. doi:10.1038/nclimate1911.
- 19 Hirsch, A. L., Evans, J. P., Di Virgilio, G., Perkins-Kirkpatrick, S. E., Argüeso, D., Pitman, A. J., et al. (2019).  
20 Amplification of Australian Heatwaves via Local Land-Atmosphere Coupling. *J. Geophys. Res. Atmos.* 124,  
21 13625–13647. doi:10.1029/2019JD030665.
- 22 Hirsch, A. L., Guillod, B. P., Seneviratne, S. I., Beyerle, U., Boysen, L. R., Brovkin, V., et al. (2018). Biogeophysical  
23 Impacts of Land-Use Change on Climate Extremes in Low-Emission Scenarios: Results From HAPPI-Land.  
24 *Earth's Futur.* 6, 396–409. doi:10.1002/2017EF000744.
- 25 Hirsch, A. L., Wilhelm, M., Davin, E. L., Thiery, W., and Seneviratne, S. I. (2017). Can climate-effective land  
26 management reduce regional warming? *J. Geophys. Res. Atmos.* 122, 2269–2288. doi:10.1002/2016JD026125.
- 27 Hirschi, M., Seneviratne, S. I., Alexandrov, V., Boberg, F., Boroneant, C., Christensen, O. B., et al. (2011).  
28 Observational evidence for soil-moisture impact on hot extremes in southeastern Europe. *Nat. Geosci.* 4, 17–21.  
29 doi:10.1038/ngeo1032.
- 30 Hobbins, M., McEvoy, D., and Hain, C. (2017). “Evapotranspiration, Evaporative Demand, and Drought,” in *Drought*  
31 *and Water Crises: Integrating Science, Management, and Policy*, 259–287.
- 32 Hobbins, M. T., Wood, A., McEvoy, D. J., Huntington, J. L., Morton, C., Anderson, M., et al. (2016). The evaporative  
33 demand drought index. Part I: Linking drought evolution to variations in evaporative demand. *J. Hydrometeorol.*  
34 17, 1745–1761. doi:10.1175/JHM-D-15-0121.1.
- 35 Hobbins, M., Wood, A., Streubel, D., and Werner, K. (2012). What drives the variability of evaporative demand across  
36 the conterminous United States? *J. Hydrometeorol.* 13, 1195–1214. doi:10.1175/JHM-D-11-0101.1.
- 37 Hodgkins, G. A., Whitfield, P. H., Burn, D. H., Hannaford, J., Renard, B., Stahl, K., et al. (2017). Climate-driven  
38 variability in the occurrence of major floods across North America and Europe. *J. Hydrol.* 552, 704–717.  
39 doi:10.1016/j.jhydrol.2017.07.027.
- 40 Hodnebrog, Ø., Marelle, L., Alterskjær, K., Wood, R. R., Ludwig, R., Fischer, E. M., et al. (2019). Intensification of  
41 summer precipitation with shorter time-scales in Europe. *Environ. Res. Lett.* 14, 124050. doi:10.1088/1748-  
42 9326/ab549c.
- 43 Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., et al. (2018). “Impacts of 1.5°C Global  
44 Warming on Natural and Human Systems,” in *Global Warming of 1.5°C. An IPCC Special Report on the impacts*  
45 *of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in*  
46 *the context of strengthening the global response to the threat of climate change*, eds. V. Masson-Delmotte, P.  
47 Zhai, H. O. Pörtner, D. Roberts, J. Skea, P. R. Shukla, et al. (In Press), 175–312. Available at:  
48 <https://www.ipcc.ch/sr15/chapter/chapter-3>.
- 49 Hoerling, M., Eischeid, J., Kumar, A., Leung, R., Mariotti, A., Mo, K., et al. (2014). Causes and Predictability of the  
50 2012 Great Plains Drought. *Bull. Am. Meteorol. Soc.* 95, 269–282. doi:10.1175/BAMS-D-13-00055.1.
- 51 Hoerling, M., Eischeid, J., Perlwitz, J., Quan, X.-W., Wolter, K., and Cheng, L. (2016). Characterizing Recent Trends  
52 in U.S. Heavy Precipitation. *J. Clim.* 29, 2313–2332. doi:10.1175/JCLI-D-15-0441.1.
- 53 Hoerling, M., Eischeid, J., Perlwitz, J., Quan, X., Zhang, T., Pegion, P., et al. (2012). On the Increased Frequency of  
54 Mediterranean Drought. *J. Clim.* 25, 2146–2161. doi:10.1175/JCLI-D-11-00296.1.
- 55 Hoerling, M., Kumar, A., Dole, R., Nielsen-Gammon, J. W., Eischeid, J., Perlwitz, J., et al. (2013). Anatomy of an  
56 extreme event. *J. Clim.* 26, 2811–2832. doi:10.1175/JCLI-D-12-00270.1.
- 57 Hogeboom, R. J., Knook, L., and Hoekstra, A. Y. (2018). The blue water footprint of the world’s artificial reservoirs for  
58 hydroelectricity, irrigation, residential and industrial water supply, flood protection, fishing and recreation. *Adv.*  
59 *Water Resour.* 113, 285–294. doi:10.1016/j.advwatres.2018.01.028.
- 60 Holland, G., and Bruyère, C. L. (2014). Recent intense hurricane response to global climate change. *Clim. Dyn.* 42,  
61 617–627. doi:10.1007/s00382-013-1713-0.

- 1 Holloway, C. E., Wing, A. A., Bony, S., Muller, C., Masunaga, H., L'Ecuyer, T. S., et al. (2017). Observing Convective  
2 Aggregation. *Surv. Geophys.* 38, 1199–1236. doi:10.1007/s10712-017-9419-1.
- 3 Holmes, C. R., Woollings, T., Hawkins, E., and de Vries, H. (2015). Robust Future Changes in Temperature Variability  
4 under Greenhouse Gas Forcing and the Relationship with Thermal Advection. *J. Clim.* 29, 2221–2236.  
5 doi:10.1175/JCLI-D-14-00735.1.
- 6 Hong, C.-C., Lee, M.-Y., Hsu, H.-H., and Tseng, W.-L. (2018). Distinct Influences of the ENSO-Like and PMM-Like  
7 SST Anomalies on the Mean TC Genesis Location in the Western North Pacific: The 2015 Summer as an  
8 Extreme Example. *J. Clim.* 31, 3049–3059. doi:10.1175/JCLI-D-17-0504.1.
- 9 Hoogewind, K. A., Baldwin, M. E., and Trapp, R. J. (2017). The Impact of Climate Change on Hazardous Convective  
10 Weather in the United States: Insight from High-Resolution Dynamical Downscaling. *J. Clim.* 30, 10081–10100.  
11 doi:10.1175/JCLI-D-16-0885.1.
- 12 Hoogewind, K. A., Chavas, D. R., Schenkel, B. A., and O'Neill, M. E. (2020). Exploring Controls on Tropical Cyclone  
13 Count through the Geography of Environmental Favorability. *J. Clim.* 33, 1725–1745. doi:10.1175/JCLI-D-18-  
14 0862.1.
- 15 Hope, P., Black, M. T., Lim, E.-P., Dowdy, A., Wang, G., Fawcett, R. J. B., et al. (2019). On Determining the Impact of  
16 Increasing Atmospheric CO<sub>2</sub> on the Record Fire Weather in Eastern Australia in February 2017. *Bull. Am.*  
17 *Meteorol. Soc.* 100, S111–S117. doi:10.1175/BAMS-D-18-0135.1.
- 18 Hope, P., Lim, E.-P., Wang, G., Hendon, H. H., and Arblaster, J. M. (2015). Contributors to the Record High  
19 Temperatures Across Australia in Late Spring 2014. *Bull. Am. Meteorol. Soc.* 96, S149–S153.  
20 doi:10.1175/BAMS-D-15-00096.1.
- 21 Hope, P., Wang, G., Lim, E.-P., Hendon, H. H., and Arblaster, J. M. (2016). What caused the record-breaking heat  
22 across Australia in October 2015? [in “Explaining Extreme Events of 2015 from a Climate Perspective”]. *Bull.*  
23 *Am. Meteorol. Soc.* 97, S122–S126. doi:10.1175/BAMS-ExplainingExtremeEvents2015.1.
- 24 Horton, D. E., Johnson, N. C., Singh, D., Swain, D. L., Rajaratnam, B., and Diffenbaugh, N. S. (2015). Contribution of  
25 changes in atmospheric circulation patterns to extreme temperature trends. *Nature* 522, 465–469.  
26 doi:10.1038/nature14550.
- 27 Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., et al. (2014). The Global  
28 Precipitation Measurement Mission. *Bull. Am. Meteorol. Soc.* 95, 701–722. doi:10.1175/BAMS-D-13-00164.1.
- 29 Howarth, M. E., Thorncroft, C. D., and Bosart, L. F. (2019). Changes in Extreme Precipitation in the Northeast United  
30 States: 1979–2014. *J. Hydrometeorol.* 20, 673–689. doi:10.1175/JHM-D-18-0155.1.
- 31 Hsu, W.-C., Patricola, C. M., and Chang, P. (2019). The impact of climate model sea surface temperature biases on  
32 tropical cyclone simulations. *Clim. Dyn.* 53, 173–192. doi:10.1007/s00382-018-4577-5.
- 33 Hu, T., Sun, Y., Zhang, X., Min, S.-K., and Kim, Y.-H. (2020). Human influence on frequency of temperature  
34 extremes. *Environ. Res. Lett.* 15, 064014. doi:10.1088/1748-9326/ab8497.
- 35 Hu, Z., Li, Q., Chen, X., Teng, Z., Chen, C., Yin, G., et al. (2016). Climate changes in temperature and precipitation  
36 extremes in an alpine grassland of Central Asia. *Theor. Appl. Climatol.* 126, 519–531. doi:10.1007/s00704-015-  
37 1568-x.
- 38 Hua, W., Zhou, L., Chen, H., Nicholson, S. E., Raghavendra, A., and Jiang, Y. (2016). Possible causes of the Central  
39 Equatorial African long-term drought. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/12/124002.
- 40 Huang, H., Winter, J. M., Osterberg, E. C., Horton, R. M., and Beckage, B. (2017a). Total and Extreme Precipitation  
41 Changes over the Northeastern United States. *J. Hydrometeorol.* 18, 1783–1798. doi:10.1175/JHM-D-16-0195.1.
- 42 Huang, J., Zhai, J., Jiang, T., Wang, Y., Li, X., Wang, R., et al. (2018a). Analysis of future drought characteristics in  
43 China using the regional climate model CCLM. *Clim. Dyn.* 50, 507–525. doi:10.1007/s00382-017-3623-z.
- 44 Huang, S., Kumar, R., Flörke, M., Yang, T., Hundecha, Y., Kraft, P., et al. (2017b). Evaluation of an ensemble of  
45 regional hydrological models in 12 large-scale river basins worldwide. *Clim. Change* 141, 381–397.  
46 doi:10.1007/s10584-016-1841-8.
- 47 Huang, S., Kumar, R., Rakovec, O., Aich, V., Wang, X., Samaniego, L., et al. (2018b). Multimodel assessment of flood  
48 characteristics in four large river basins at global warming of 1.5, 2.0 and 3.0 K above the pre-industrial level.  
49 *Environ. Res. Lett.* 13, 124005. doi:10.1088/1748-9326/aae94b.
- 50 Huang, S., Li, P., Huang, Q., Leng, G., Hou, B., and Ma, L. (2017c). The propagation from meteorological to  
51 hydrological drought and its potential influence factors. *J. Hydrol.* 547, 184–195.  
52 doi:10.1016/j.jhydrol.2017.01.041.
- 53 Huerta, A., and Lavado-Casimiro, W. (2020). Trends and variability of precipitation extremes in the Peruvian Altiplano  
54 (1971–2013). *Int. J. Climatol.* n/a. doi:10.1002/joc.6635.
- 55 Hui, P., Tang, J., Wang, S., Niu, X., Zong, P., and Dong, X. (2018). Climate change projections over China using  
56 regional climate models forced by two CMIP5 global models. Part I: evaluation of historical simulations. *Int. J.*  
57 *Climatol.* 38, e57–e77. doi:10.1002/joc.5351.
- 58 Huijnen, V., Wooster, M. J., Kaiser, J. W., Gaveau, D. L. A., Flemming, J., Parrington, M., et al. (2016). Fire carbon  
59 emissions over maritime southeast Asia in 2015 largest since 1997. *Sci. Rep.* 6, 1–8. doi:10.1038/srep26886.
- 60 Humphrey, V., Zscheischler, J., Ciais, P., Gudmundsson, L., Sitch, S., and Seneviratne, S. I. (2018). Sensitivity of  
61 atmospheric CO<sub>2</sub> growth rate to observed changes in terrestrial water storage. *Nature* 560, 628–631.

- 1 doi:10.1038/s41586-018-0424-4.
- 2 Hundecha, Y., Sunyer, M. A., Lawrence, D., Madsen, H., Willems, P., Bürger, G., et al. (2016). Inter-comparison of  
3 statistical downscaling methods for projection of extreme flow indices across Europe. *J. Hydrol.* 541, 1273–1286.  
4 doi:10.1016/j.jhydrol.2016.08.033.
- 5 Huning, L. S., and AghaKouchak, A. (2020). Global snow drought hot spots and characteristics. *Proc. Natl. Acad. Sci.*  
6 *U. S. A.* 117, 19753–19759. doi:10.1073/pnas.1915921117.
- 7 Hunt, E. D., Svoboda, M., Wardlow, B., Hubbard, K., Hayes, M., and Arkebauer, T. (2014). Monitoring the effects of  
8 rapid onset of drought on non-irrigated maize with agronomic data and climate-based drought indices. *Agric. For.*  
9 *Meteorol.* 191, 1–11. doi:10.1016/j.agrformet.2014.02.001.
- 10 Hunt, K. M. R., Turner, A. G., and Shaffrey, L. C. (2018). Extreme Daily Rainfall in Pakistan and North India: Scale  
11 Interactions, Mechanisms, and Precursors. *Mon. Weather Rev.* 146, 1005–1022. doi:10.1175/MWR-D-17-0258.1.
- 12 Hussain, M. S., and Lee, S. (2013). The regional and the seasonal variability of extreme precipitation trends in Pakistan.  
13 *Asia-Pacific J. Atmos. Sci.* 49, 421–441. doi:10.1007/s13143-013-0039-5.
- 14 Imada, Y., Maeda, S., Watanabe, M., Shiogama, H., Mizuta, R., Ishii, M., et al. (2017). Recent enhanced seasonal  
15 temperature contrast in Japan from large ensemble high-resolution climate simulations. *Atmosphere (Basel)*. 8.  
16 doi:10.3390/atmos8030057.
- 17 Imada, Y., Shiogama, H., Takahashi, C., Watanabe, M., Mori, M., Kamae, Y., et al. (2018). Climate Change Increased  
18 the Likelihood of the 2016 Heat Extremes in Asia [in “Explaining Extreme Events of 2016 from a Climate  
19 Perspective”]. *Bull. Am. Meteorol. Soc.* 99, S97–S101. doi:10.1175/BAMS-D-17-0109.1.
- 20 Imada, Y., Shiogama, H., Watanabe, M., Mori, M., Ishii, M., and Kimoto, M. (2014). The Contribution of  
21 anthropogenic forcing to the Japanese heat waves of 2013 [in “Explaining Extreme Events of 2013 from a  
22 Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S52–S54. doi:10.1175/1520-0477-95.9.S1.1.
- 23 Imada, Y., Watanabe, M., Kawase, H., Shiogama, H., and Arai, M. (2019). The July 2018 High Temperature Event in  
24 Japan Could Not Have Happened without Human-Induced Global Warming. *SOLA* 15A, 8–12.  
25 doi:10.2151/sola.15A-002.
- 26 Imada, Y., Watanabe, M., Mori, M., Kimoto, M., Shiogama, H., and Ishii, M. (2013). Contribution of Atmospheric  
27 Circulation Change to the 2012 Heavy Rainfall in Southwestern Japan. *Bull. Am. Meteorol. Soc.* 96, 52–54.
- 28 Imbach, P., Chou, S. C., Lyra, A., Rodrigues, D., Rodriguez, D., Latinovic, D., et al. (2018). Future climate change  
29 scenarios in Central America at high spatial resolution. *PLoS One* 13, e0193570.  
30 doi:10.1371/journal.pone.0193570.
- 31 INMET (2017). Situação da Seca Observada nas Regiões Norte e Nordeste do Brasil em 2016. Brasília, Brazil: Instituto  
32 Nacional de Meteorologia (INMET) Available at:  
33 [https://portal.inmet.gov.br/uploads/notastecnicas/trabalho\\_tecnico\\_02-2017.pdf](https://portal.inmet.gov.br/uploads/notastecnicas/trabalho_tecnico_02-2017.pdf).
- 34 Innocenti, S., Maillhot, A., Leduc, M., Cannon, A. J., and Frigon, A. (2019). Projected Changes in the Probability  
35 Distributions, Seasonality, and Spatiotemporal Scaling of Daily and Subdaily Extreme Precipitation Simulated by  
36 a 50-Member Ensemble Over Northeastern North America. *J. Geophys. Res. Atmos.* 124, 10427–10449.  
37 doi:10.1029/2019JD031210.
- 38 IPCC (2012). Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special  
39 Report of Working Groups I and II of the Intergovernmental Panel on Climate Change. , eds. C. B. Field, V.  
40 Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, et al. Cambridge, United Kingdom and New York, NY ,  
41 USA: Cambridge University Press Available at: [https://www.ipcc.ch/report/managing-the-risks-of-extreme-](https://www.ipcc.ch/report/managing-the-risks-of-extreme-events-and-disasters-to-advance-climate-change-adaptation)  
42 [events-and-disasters-to-advance-climate-change-adaptation](https://www.ipcc.ch/report/managing-the-risks-of-extreme-events-and-disasters-to-advance-climate-change-adaptation).
- 43 IPCC (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth  
44 Assessment Report of the Intergovernmental Panel on Climate Change. , eds. T. F. Stocker, D. Qin, G.-K.  
45 Plattner, M. Tignor, S. K. Allen, J. Boschung, et al. Cambridge, United Kingdom and New York, NY, USA:  
46 Cambridge University Press doi:10.1017/CBO9781107415324.
- 47 IPCC (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects.*  
48 *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
49 *Change.* , eds. C. B. Field, V. R. Barros, D. J. Dokken, K. J. Mach, M. D. Mastrandrea, T. E. Bilir, et al. United  
50 Kingdom and New York, NY, USA,: ambridge University Press, Cambridge.
- 51 IPCC (2018). “Summary for Policymakers,” in *Global warming of 1.5°C. An IPCC Special Report on the impacts of*  
52 *global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the*  
53 *context of strengthening the global response to,* eds. V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J.  
54 Skea, P. R. Shukla, et al. (In Press), 32. Available at: <https://www.ipcc.ch/sr15/chapter/spm>.
- 55 IPCC (2019). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. , eds. H.-O. Pörtner, D. C.  
56 Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, et al. In Press Available at:  
57 <https://www.ipcc.ch/srocc>.
- 58 Irannezhad, M., Chen, D., Kløve, B., and Moradkhani, H. (2017). Analysing the variability and trends of precipitation  
59 extremes in Finland and their connection to atmospheric circulation patterns. *Int. J. Climatol.* 37, 1053–1066.  
60 doi:10.1002/joc.5059.
- 61 Ishak, E. H., Rahman, A., Westra, S., Sharma, A., and Kuczera, G. (2013). Evaluating the non-stationarity of australian

- 1 annual maximum flood. *J. Hydrol.* 494, 134–145. doi:10.1016/j.jhydrol.2013.04.021.
- 2 Ishiyama, T., Satoh, M., and Yamada, Y. (2018). Potential differences in North Pacific tropical cyclone activity  
3 between the super El Niño years of 1997 and 2015. *J. Meteorol. Soc. Japan.*
- 4 Ivancic, T. J., and Shaw, S. B. (2015). Examining why trends in very heavy precipitation should not be mistaken for  
5 trends in very high river discharge. *Clim. Change* 133, 681–693. doi:10.1007/s10584-015-1476-1.
- 6 Jacob, D., Kotova, L., Teichmann, C., Sobolowski, S. P., Vautard, R., Donnelly, C., et al. (2018). Climate Impacts in  
7 Europe Under +1.5°C Global Warming. *Earth's Futur.* 6, 264–285. doi:10.1002/2017EF000710.
- 8 Jacob, D., Teichmann, C., Sobolowski, S., Katragkou, E., Anders, I., Belda, M., et al. (2020). Regional climate  
9 downscaling over Europe: perspectives from the EURO-CORDEX community. *Reg. Environ. Chang.* 20, 51.  
10 doi:10.1007/s10113-020-01606-9.
- 11 Jakob, D., and Walland, D. (2016). Variability and long-term change in Australian temperature and precipitation  
12 extremes. *Weather Clim. Extrem.* 14, 36–55. doi:10.1016/j.wace.2016.11.001.
- 13 James, R., Washington, R., Schleussner, C.-F., Rogelj, J., and Conway, D. (2017). Characterizing half-a-degree  
14 difference: a review of methods for identifying regional climate responses to global warming targets. *Wiley*  
15 *Interdiscip. Rev. Clim. Chang.* 8, e457. doi:10.1002/wcc.457.
- 16 Jehanzaib, M., Shah, S. A., Yoo, J., and Kim, T.-W. (2020). Investigating the impacts of climate change and human  
17 activities on hydrological drought using non-stationary approaches. *J. Hydrol.* 588.  
18 doi:10.1016/j.jhydrol.2020.125052.
- 19 Jeon, S., Paciorek, C. J., and Wehner, M. F. (2016). Quantile-based bias correction and uncertainty quantification of  
20 extreme event attribution statements. *Weather Clim. Extrem.* 12, 24–32. doi:10.1016/j.wace.2016.02.001.
- 21 Jeong, J.-H., Fan, J., Homeyer, C. R., and Hou, Z. (2020). Understanding Hailstone Temporal Variability and  
22 Contributing Factors over the U.S. Southern Great Plains. *J. Clim.* 33, 3947–3966. doi:10.1175/JCLI-D-19-  
23 0606.1.
- 24 Jeong, S.-J., Ho, C.-H., Piao, S., Kim, J., Ciais, P., Lee, Y.-B., et al. (2014). Effects of double cropping on summer  
25 climate of the North China Plain and neighbouring regions. *Nat. Clim. Chang.* 4, 615–619.  
26 doi:10.1038/nclimate2266.
- 27 Jézéquel, A., Dépoues, V., Guillemot, H., Trolliet, M., Vanderlinden, J.-P., and Yiou, P. (2018). Behind the veil of  
28 extreme event attribution. *Clim. Change.* doi:10.1007/s10584-018-2252-9.
- 29 Jhahharia, D., Kumar, R., Dabral, P. P., Singh, V. P., Choudhary, R. R., and Dinpashoh, Y. (2015). Reference  
30 evapotranspiration under changing climate over the Thar Desert in India. *Meteorol. Appl.* 22, 425–435.  
31 doi:10.1002/met.1471.
- 32 Ji, P., Yuan, X., Jiao, Y., Wang, C., Han, S., and Shi, C. (2020). Anthropogenic contributions to the 2018 extreme  
33 flooding over upper Yellow River basin in China. *Bull. Am. Meteorol. Soc.* doi:10.1175/BAMS-D-19-0105.1.
- 34 Ji, Z., and Kang, S. (2015). Evaluation of extreme climate events using a regional climate model for China. *Int. J.*  
35 *Climatol.* 35, 888–902. doi:10.1002/joc.4024.
- 36 Jia, B., Liu, J., Xie, Z., and Shi, C. (2018). Interannual Variations and Trends in Remotely Sensed and Modeled Soil  
37 Moisture in China. *J. Hydrometeorol.* 19, 831–847. doi:10.1175/JHM-D-18-0003.1.
- 38 Jia, G., E. Shevliakova, Artaxo, P., Noblet-Ducoudré, N. De, Houghton, R., House, J., et al. (2019a). Land–climate  
39 interactions. In: *Climate Change and Land: an IPCC special report on climate change, desertification, land*  
40 *degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems.* ,  
41 eds. P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, et al.
- 42 Jia, G., Shevliakova, E., Artaxo, P., Noblet-Ducoudré, N. De, Houghton, R., House, J., et al. (2019b). “Land–climate  
43 interactions,” in *Climate Change and Land: an IPCC special report on climate change, desertification, land*  
44 *degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems,*  
45 eds. P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, et al. (In Press),  
46 131–248. Available at: <https://www.ipcc.ch/srcl/chapter/chapter-2>.
- 47 Jiang, F., Hu, R.-J., Wang, S.-P., Zhang, Y.-W., and Tong, L. (2013). Trends of precipitation extremes during 1960–  
48 2008 in Xinjiang, the Northwest China. *Theor. Appl. Climatol.* 111, 133–148. doi:10.1007/s00704-012-0657-3.
- 49 Jiménez-Muñoz, J. C., Mattar, C., Barichivich, J., Santamaría-Artigas, A., Takahashi, K., Malhi, Y., et al. (2016).  
50 Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–  
51 2016. *Sci. Rep.* 6, 33130. doi:10.1038/srep33130.
- 52 Jimenez Cisneros, B. E., Oki, T., Arnell, N. W., Benito, G., Cogley, J. G., Petra, D., et al. (2014). “IPCC WGIIAR5  
53 Chapter 3 Freshwater resources,” in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A:*  
54 *Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*  
55 *Intergovernmental Panel on Climate Change*, eds. T. E. B. Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach,  
56 M.D. Mastrandrea, S. M. M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy,  
57 and L. L. W. P.R. Mastrandrea (Cambridge University Press, Cambridge, United Kingdom and New York,  
58 NY, USA), 229–269.
- 59 Johnson, N. C., Xie, S. P., Kosaka, Y., and Li, X. (2018). Increasing occurrence of cold and warm extremes during the  
60 recent global warming slowdown. *Nat. Commun.* 9, 4–6. doi:10.1038/s41467-018-04040-y.
- 61 Jolly, W. M., Cochrane, M. A., Freeborn, P. H., Holden, Z. A., Brown, T. J., Williamson, G. J., et al. (2015). Climate-

- 1 induced variations in global wildfire danger from 1979 to 2013. *Nat. Commun.* 6, 7537.  
2 doi:10.1038/ncomms8537.
- 3 Junk, J., Goergen, K., and Krein, A. (2019). Future Heat Waves in Different European Capitals Based on Climate  
4 Change Indicators. *Int. J. Environ. Res. Public Health* 16, 3959. doi:10.3390/ijerph16203959.
- 5 Kamae, Y., Mei, W., Xie, S. P., Naoi, M., and Ueda, H. (2017a). Atmospheric rivers over the Northwestern Pacific:  
6 Climatology and interannual variability. *J. Clim.* 30, 5605–5619. doi:10.1175/JCLI-D-16-0875.1.
- 7 Kamae, Y., Shiogama, H., Imada, Y., Mori, M., Arakawa, O., Mizuta, R., et al. (2017b). Forced response and internal  
8 variability of summer climate over western North America. *Clim. Dyn.* 49, 403–417. doi:10.1007/s00382-016-  
9 3350-x.
- 10 Kamae, Y., Shiogama, H., Watanabe, M., and Kimoto, M. (2014). Attributing the increase in Northern Hemisphere hot  
11 summers since the late 20th century. *Geophys. Res. Lett.* 41, 5192–5199. doi:10.1002/2014GL061062.
- 12 Kambezidis, H. D., Kaskaoutis, D. G., Kharol, S. K., Moorthy, K. K., Satheesh, S. K., Kalapureddy, M. C. R., et al.  
13 (2012). Multi-decadal variation of the net downward shortwave radiation over south Asia: The solar dimming  
14 effect. *Atmos. Environ.* 50, 360–372. doi:10.1016/j.atmosenv.2011.11.008.
- 15 Kanada, S., Takemi, T., Kato, M., Yamasaki, S., Fudeyasu, H., Tsuboki, K., et al. (2017a). A multimodel  
16 intercomparison of an intense typhoon in future, warmer climates by Four 5-km-Mesh models. *J. Clim.* 30, 6017–  
17 6036. doi:10.1175/JCLI-D-16-0715.1.
- 18 Kanada, S., Tsujino, S., Aiki, H., Yoshioka, M. K., Miyazawa, Y., Tsuboki, K., et al. (2017b). Impacts of SST Patterns  
19 on Rapid Intensification of Typhoon Megi (2010). *J. Geophys. Res. Atmos.* 122, 13,213-245,262.  
20 doi:10.1002/2017JD027252.
- 21 Kanada, S., and Wada, A. (2016). Sensitivity to Horizontal Resolution of the Simulated Intensifying Rate and Inner-  
22 Core Structure of Typhoon Ida, an Extremely Intense Typhoon. *J. Meteorol. Soc. Japan. Ser. II* 94A, 181–190.  
23 doi:10.2151/jmsj.2015-037.
- 24 Kanada, S., Wada, A., and Sugi, M. (2013). Future changes in structures of extremely intense tropical cyclones using a  
25 2-km mesh nonhydrostatic model. *J. Clim.* 26, 9986–10005. doi:10.1175/JCLI-D-12-00477.1.
- 26 Kang, N. Y., and Elsner, J. B. (2012). Consensus on climate trends in Western North Pacific tropical cyclones. *J. Clim.*  
27 25, 7564–7573. doi:10.1175/JCLI-D-11-00735.1.
- 28 Kaniewski, D., Van Campo, E., and Weiss, H. (2012). Drought is a recurring challenge in the Middle East. *Proc. Natl.*  
29 *Acad. Sci. U. S. A.* 109, 3862–3867. doi:10.1073/pnas.1116304109.
- 30 Kar-Man Chang, E. (2018). CMIP5 Projected Change in Northern Hemisphere Winter Cyclones with Associated  
31 Extreme Winds. *J. Clim.* 31, 6527–6542. doi:10.1175/JCLI-D-17-0899.1.
- 32 Karoly, D. J., Black, M. T., Grose, M. R., and King, A. D. (2016). 25. The roles of climate change and El Niño in the  
33 record low rainfall in October 2015 in Tasmania, Australia. *Bull. Am. Meteorol. Soc.* 97, S127–S130.  
34 doi:10.1175/BAMS-D-16-0139.1.
- 35 Kato, T. (2020). Quasi-stationary band-shaped precipitation systems, named as “senjo-kousuitai”, causing localized  
36 heavy rainfall in Japan. *J. Meteorol. Soc. Japan* 98, 485–509. doi:10.2151/jmsj.2020-029.
- 37 Kattsov, V. M., Shkolnik, I. M., and Efimov, S. V. (2017). Climate change projections in Russian regions: The detailing  
38 in physical and probability spaces. *Russ. Meteorol. Hydrol.* 42, 452–460. doi:10.3103/S1068373917070044.
- 39 Kawase, H., Imada, Y., Tsuguti, H., Nakaegawa, T., Seinino, N., Murata, A., et al. (2019). The heavy rain event of July  
40 2019 in Japan enhanced by historical warming. *Bull Am Meteorol Soc.* doi:10.1175/BAMS-D-19-0173.1.
- 41 Kay, A. L., Bell, V. A., Guillod, B. P., Jones, R. G., and Rudd, A. C. (2018). National-scale analysis of low flow  
42 frequency: historical trends and potential future changes. *Clim. Change* 147, 585–599. doi:10.1007/s10584-018-  
43 2145-y.
- 44 Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., et al. (2015). The Community Earth System Model  
45 (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the Presence of  
46 Internal Climate Variability. *Bull. Am. Meteorol. Soc.* 96, 1333–1349. doi:10.1175/BAMS-D-13-00255.1.
- 47 Keellings, D., and Waylen, P. (2014). Increased risk of heat waves in Florida: Characterizing changes in bivariate heat  
48 wave risk using extreme value analysis. *Appl. Geogr.* 46, 90–97. doi:10.1016/j.apgeog.2013.11.008.
- 49 Kelley, C. P., Mohtadi, S., Cane, M. A., Seager, R., and Kushnir, Y. (2015). Climate change in the Fertile Crescent and  
50 implications of the recent Syrian drought. *Proc. Natl. Acad. Sci. U. S. A.* 112, 3241–3246.  
51 doi:10.1073/pnas.1421533112.
- 52 Kelley, C., Ting, M., Seager, R., and Kushnir, Y. (2012). The relative contributions of radiative forcing and internal  
53 climate variability to the late 20th Century winter drying of the Mediterranean region. *Clim. Dyn.* 38, 2001–2015.  
54 doi:10.1007/s00382-011-1221-z.
- 55 Kendon, E. J., Ban, N., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C., et al. (2017). Do Convection-  
56 Permitting Regional Climate Models Improve Projections of Future Precipitation Change? *Bull. Am. Meteorol.*  
57 *Soc.* 98, 79–93. doi:10.1175/BAMS-D-15-0004.1.
- 58 Kendon, E. J., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C., and Senior, C. A. (2014). Heavier summer  
59 downpours with climate change revealed by weather forecast resolution model. *Nat. Clim. Chang.* 4, 570–576.  
60 doi:10.1038/nclimate2258.
- 61 Kendon, E. J., Stratton, R. A., Tucker, S., Marsham, J. H., Berthou, S., Rowell, D. P., et al. (2019). Enhanced future

- 1 changes in wet and dry extremes over Africa at convection-permitting scale. *Nat. Commun.* 10, 1794.  
2 doi:10.1038/s41467-019-09776-9.
- 3 Kendon, M. (2014). Has there been a recent increase in UK weather records? *Weather* 69, 327–332.  
4 doi:10.1002/wea.2439.
- 5 Kew, S. F., Philip, S. Y., Hauser, M., Hobbins, M., Wanders, N., van Oldenborgh, G. J., et al. (2021). Impact of  
6 precipitation and increasing temperatures on drought trends in eastern Africa. *Earth Syst. Dyn.* 12, 17–35.  
7 doi:10.5194/esd-12-17-2021.
- 8 Kew, S. F., Philip, S. Y., Jan van Oldenborgh, G., van der Schrier, G., Otto, F. E. L., and Vautard, R. (2019). The  
9 Exceptional Summer Heat Wave in Southern Europe 2017 [in “Explaining Extreme Events of 2017 from a  
10 Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 100, S49–S53. doi:10.1175/BAMS-D-18-0109.1.
- 11 Kew, S. F., Selten, F. M., Lenderink, G., and Hazeleger, W. (2013). The simultaneous occurrence of surge and  
12 discharge extremes for the Rhine delta. *Nat. Hazards Earth Syst. Sci.* 13, 2017–2029. doi:10.5194/nhess-13-2017-  
13 2013.
- 14 Khan, N., Shahid, S., Ismail, T. bin, and Wang, X.-J. (2019). Spatial distribution of unidirectional trends in temperature  
15 and temperature extremes in Pakistan. *Theor. Appl. Climatol.* 136, 899–913. doi:10.1007/s00704-018-2520-7.
- 16 Kharin, V. V., Flato, G. M., Zhang, X., Gillett, N. P., Zwiers, F., and Anderson, K. J. (2018). Risks from Climate  
17 Extremes Change Differently from 1.5°C to 2.0°C Depending on Rarity. *Earth’s Futur.* 6, 704–715.  
18 doi:10.1002/2018EF000813.
- 19 Kharin, V. V., Zwiers, F. W., Zhang, X., and Wehner, M. (2013). Changes in temperature and precipitation extremes in  
20 the CMIP5 ensemble. *Clim. Change* 119, 345–357. doi:10.1007/s10584-013-0705-8.
- 21 Khlebnikova, E. I., Rudakova, Y. L., Sall’ I. A., Efimov, S. V., and Shkolnik, I. M. (2019a). Changes in indicators of  
22 temperature extremes in the 21st century: ensemble projections for the territory of Russia. *Russ. Meteorol.*  
23 *Hydrol.* 44, 159–168. doi:10.3103/S1068373919030014.
- 24 Khlebnikova, E. I., Rudakova, Y. L., and Shkolnik, I. M. (2019b). Changes in precipitation regime over the territory of  
25 Russia: Data of regional climate modeling and observations. *Russ. Meteorol. Hydrol.* 44, 431–439.  
26 doi:10.3103/S106837391907001X.
- 27 Khong, A., Wang, J. K., Quiring, S. M., and Ford, T. W. (2015). Soil moisture variability in Iowa. *Int. J. Climatol.* 35,  
28 2837–2848. doi:10.1002/joc.4176.
- 29 Khouakhi, A., Villarini, G., and Vecchi, G. A. (2016). Contribution of Tropical Cyclones to Rainfall at the Global  
30 Scale. *J. Clim.* 30, 359–372. doi:10.1175/JCLI-D-16-0298.1.
- 31 Kim, B., and Sanders, B. F. (2016). Dam-Break Flood Model Uncertainty Assessment: Case Study of Extreme Flooding  
32 with Multiple Dam Failures in Gangneung, South Korea. *J. Hydraul. Eng.* 142, 05016002.  
33 doi:10.1061/(ASCE)HY.1943-7900.0001097.
- 34 Kim, D., Moon, Y., Camargo, S. J., Wing, A. A., Sobel, A. H., Murakami, H., et al. (2018a). Process-Oriented  
35 Diagnosis of Tropical Cyclones in High-Resolution GCMs. *J. Clim.* 31, 1685–1702. doi:10.1175/JCLI-D-17-  
36 0269.1.
- 37 Kim, G., Cha, D.-H., Park, C., Lee, G., Jin, C.-S., Lee, D.-K., et al. (2018b). Future changes in extreme precipitation  
38 indices over Korea. *Int. J. Climatol.* 38, e862–e874. doi:10.1002/joc.5414.
- 39 Kim, H. S., Vecchi, G. A., Knutson, T. R., Anderson, W. G., Delworth, T. L., Rosati, A., et al. (2014a). Tropical  
40 cyclone simulation and response to CO<sub>2</sub> doubling in the GFDL CM2.5 high-resolution coupled climate model. *J.*  
41 *Clim.* 27, 8034–8054. doi:10.1175/JCLI-D-13-00475.1.
- 42 Kim, I.-W. W., Oh, J., Woo, S., and Kripalani, R. H. (2019). Evaluation of precipitation extremes over the Asian  
43 domain: observation and modelling studies. *Clim. Dyn.* 52, 1317–1342. doi:10.1007/s00382-018-4193-4.
- 44 Kim, J., Waliser, D. E., Matmann, C. A., Goodale, C. E., Hart, A. F., Zimdars, P. A., et al. (2014b). Evaluation of the  
45 CORDEX-Africa multi-RCM hindcast: systematic model errors. *Clim. Dyn.* 42, 1189–1202. doi:10.1007/s00382-  
46 013-1751-7.
- 47 Kim, Y.-H., Min, S.-K., Stone, D. A. D. A., Shiogama, H., and Wolski, P. (2018c). Multi-model event attribution of the  
48 summer 2013 heat wave in Korea. *Weather Clim. Extrem.* 20, 33–44. doi:10.1016/j.wace.2018.03.004.
- 49 Kim, Y.-H., Min, S.-K., Zhang, X., Sillmann, J., and Sandstad, M. (2020). Evaluation of the CMIP6 multi-model  
50 ensemble for climate extreme indices. *Weather Clim. Extrem.* 29, 100269. doi:10.1016/j.wace.2020.100269.
- 51 Kim, Y. H., Min, S. K., Zhang, X., Zwiers, F., Alexander, L. V., Donat, M. G., et al. (2016). Attribution of extreme  
52 temperature changes during 1951–2010. *Clim. Dyn.* 46, 1769–1782. doi:10.1007/s00382-015-2674-2.
- 53 King, A. D. (2017). Attributing Changing Rates of Temperature Record Breaking to Anthropogenic Influences. *Earth’s*  
54 *Futur.* 5, 1156–1168. doi:10.1002/2017EF000611.
- 55 King, A. D., Black, M. T., Karoly, D. J., and Donat, M. G. (2015a). Increased Likelihood of Brisbane, Australia, G20  
56 Heat Event Due to Anthropogenic Climate Change. *Bull. Am. Meteorol. Soc.* 96, S141–S144.  
57 doi:10.1175/BAMS-D-15-00098.1.
- 58 King, A. D., Black, M. T., Min, S.-K., Fischer, E. M., Mitchell, D. M., Harrington, L. J., et al. (2016a). Emergence of  
59 heat extremes attributable to anthropogenic influences. *Geophys. Res. Lett.* 43, 3438–3443.  
60 doi:10.1002/2015GL067448.
- 61 King, A. D., Jan van Oldenborgh, G., Karoly, D. J., Lewis, S. C., and Cullen, H. (2015b). Attribution of the record high

- 1 Central England temperature of 2014 to anthropogenic influences. *Environ. Res. Lett.* 10, 054002.  
2 doi:10.1088/1748-9326/10/5/054002.
- 3 King, A. D., Karoly, D. J., Donat, M. G., and Alexander, L. V. (2014). Climate change turns Australia's 2013 big dry  
4 into a year of record-breaking heat [in "Explaining Extreme Events of 2013 from a Climate Perspective"]. *Bull.*  
5 *Am. Meteorol. Soc.* 95, S41–S45. doi:10.1175/1520-0477-95.9.S41.
- 6 King, A. D., Karoly, D. J., and van Oldenborgh, G. J. (2016b). Climate Change and El Niño Increase Likelihood of  
7 Indonesian Heat and Drought. *Bull. Am. Meteorol. Soc.* 97, S113–S117. doi:10.1175/BAMS-D-16-0164.1.
- 8 King, A. D., Knutti, R., Uhe, P., Mitchell, D. M., Lewis, S. C., Arblaster, J. M., et al. (2018). On the linearity of local  
9 and regional temperature changes from 1.5°C to 2°C of global warming. *J. Clim.*, JCLI-D-17-0649.1.  
10 doi:10.1175/JCLI-D-17-0649.1.
- 11 Kingston, D. G., Stagge, J. H., Tallaksen, L. M., and Hannah, D. M. (2015). European-scale drought: Understanding  
12 connections between atmospheric circulation and meteorological drought indices. *J. Clim.* 28, 505–516.  
13 doi:10.1175/JCLI-D-14-00001.1.
- 14 Kinter, J. L., Cash, B., Achuthavari, D., Adams, J., Altshuler, E., Dirmeyer, P., et al. (2013). Revolutionizing climate  
15 modeling with project athena: A multi-institutional, international collaboration. *Bull. Am. Meteorol. Soc.* 94, 231–  
16 245. doi:10.1175/BAMS-D-11-00043.1.
- 17 Kirchmeier-Young, M. C., and Zhang, X. (2020). Human influence has intensified extreme precipitation in North  
18 America. *Proc. Natl. Acad. Sci.* 117, 13308–13313. doi:10.1073/pnas.1921628117.
- 19 Kirchmeier-Young, M. C., Wan, H., Zhang, X., and Seneviratne, S. I. (2019). Importance of Framing for Extreme  
20 Event Attribution: The Role of Spatial and Temporal Scales. *Earth's Futur.* 7, 1192–1204.  
21 doi:10.1029/2019EF001253.
- 22 Kirono, D. G. C., Hennessy, K. J., and Grose, M. R. (2017). Increasing risk of months with low rainfall and high  
23 temperature in southeast Australia for the past 150 years. *Clim. Risk Manag.* 16, 10–21.  
24 doi:10.1016/j.crm.2017.04.001.
- 25 Kirono, D. G. C., Round, V., Heady, C., Chiew, F. H. S., and Osbrough, S. (2020). Drought projections for Australia:  
26 Updated results and analysis of model simulations. *Weather Clim. Extrem.* 30, 100280.  
27 doi:10.1016/j.wace.2020.100280.
- 28 Kirshbaum, D. J., Merlis, T. M., Gyakum, J. R., and McTaggart-Cowan, R. (2017). Sensitivity of Idealized Moist  
29 Baroclinic Waves to Environmental Temperature and Moisture Content. *J. Atmos. Sci.* 75, 337–360.  
30 doi:10.1175/jas-d-17-0188.1.
- 31 Kishitawala, C. M., Jaiswal, N., Singh, R., and Niyogi, D. (2012). Tropical cyclone intensification trends during satellite  
32 era (1986–2010). *Geophys. Res. Lett.* 39, 1–6. doi:10.1029/2012GL051700.
- 33 Kitoh, A., and Endo, H. (2019). Future Changes in Precipitation Extremes Associated with Tropical Cyclones Projected  
34 by Large-Ensemble Simulations. *J. Meteorol. Soc. Japan. Ser. II* 97, 141–152. doi:10.2151/jmsj.2019-007.
- 35 Kjeldsen, T. R., Macdonald, N., Lang, M., Mediero, L., Albuquerque, T., Bogdanowicz, E., et al. (2014). Documentary  
36 evidence of past floods in Europe and their utility in flood frequency estimation. *J. Hydrol.* 517, 963–973.  
37 doi:10.1016/j.jhydrol.2014.06.038.
- 38 Klerk, W. J., Winsemius, H. C., van Verseveld, W. J., Bakker, A. M. R., and Diermanse, F. L. M. (2015). The co-  
39 incidence of storm surges and extreme discharges within the Rhine–Meuse Delta. *Environ. Res. Lett.* 10, 035005.  
40 doi:10.1088/1748-9326/10/3/035005.
- 41 Klutse, N. A. B., Ajayi, V. O., Gbobaniyi, E. O., Egbebiyi, T. S., Kouadio, K., Nkrumah, F., et al. (2018). Potential  
42 impact of 1.5 °C and 2 °C global warming on consecutive dry and wet days over West Africa. *Environ. Res. Lett.*  
43 13, 055013. doi:10.1088/1748-9326/aab37b.
- 44 Klutse, N. A. B., Sylla, M. B., Diallo, I., Sarr, A., Dosio, A., Diedhiou, A., et al. (2016). Daily characteristics of West  
45 African summer monsoon precipitation in CORDEX simulations. *Theor. Appl. Climatol.* 123, 369–386.  
46 doi:10.1007/s00704-014-1352-3.
- 47 Knapp, K. R., Velden, C. S., Wimmers, A. J., Knapp, K. R., Velden, C. S., and Wimmers, A. J. (2018). A Global  
48 Climatology of Tropical Cyclone Eyes. *Mon. Weather Rev.* 146, 2089–2101. doi:10.1175/MWR-D-17-0343.1.
- 49 Knighton, J., Conneely, J., and Walter, M. T. (2019). Possible Increases in Flood Frequency Due to the Loss of Eastern  
50 Hemlock in the Northeastern United States: Observational Insights and Predicted Impacts. *Water Resour. Res.* 55,  
51 5342–5359. doi:10.1029/2018WR024395.
- 52 Knutson, T., Camargo, S. J., Chan, J. C. L., Emanuel, K., Ho, C.-H., Kossin, J., et al. (2019). Tropical Cyclones and  
53 Climate Change Assessment: Part I: Detection and Attribution. *Bull. Am. Meteorol. Soc.* 100, 1987–2007.  
54 doi:10.1175/BAMS-D-18-0189.1.
- 55 Knutson, T., Camargo, S. J., Chan, J. C. L., Emanuel, K., Ho, C.-H., Kossin, J., et al. (2020). Tropical Cyclones and  
56 Climate Change Assessment: Part II. Projected Response to Anthropogenic Warming. *Bull. Am. Meteorol. Soc.*  
57 101, E303–E322. doi:10.1175/BAMS-D-18-0194.1.
- 58 Knutson, T. R., Sirutis, J. J., Vecchi, G. A., Garner, S., Zhao, M., Kim, H.-S., et al. (2013a). Dynamical Downscaling  
59 Projections of Twenty-First-Century Atlantic Hurricane Activity: CMIP3 and CMIP5 Model-Based Scenarios. *J.*  
60 *Clim.* 26, 6591–6617. doi:10.1175/JCLI-D-12-00539.1.
- 61 Knutson, T. R., Sirutis, J. J., Zhao, M., Tuleya, R. E., Bender, M., Vecchi, G. A., et al. (2015). Global Projections of



- 1 Intense Tropical Cyclone Activity for the Late Twenty-First Century from Dynamical Downscaling of  
2 CMIP5/RCP4.5 Scenarios. *J. Clim.* 28, 7203–7224. doi:10.1175/JCLI-D-15-0129.1.
- 3 Knutson, T. R., and Zeng, F. (2018). Model Assessment of Observed Precipitation Trends over Land Regions:  
4 Detectable Human Influences and Possible Low Bias in Model Trends. *J. Clim.* 31, 4617–4637.  
5 doi:10.1175/JCLI-D-17-0672.1.
- 6 Knutson, T. R., Zeng, F., and Wittenberg, A. T. (2013b). Multimodel assessment of regional surface temperature trends:  
7 CMIP3 and CMIP5 twentieth-century simulations. *J. Clim.* 26, 8709–8743. doi:10.1175/JCLI-D-12-00567.1.
- 8 Knutson, T. R., Zeng, F., and Wittenberg, A. T. (2014a). Multimodel assessment of extreme annual-mean warm  
9 anomalies during 2013 over regions of Australia and the western tropical Pacific [in “Explaining Extreme Events  
10 of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S26–S30. doi:10.1175/1520-0477-95.9.S1.1.
- 11 Knutson, T., Zeng, F., and Wittenberg, A. (2014b). Seasonal and annual mean precipitation extremes during during  
12 2013: A U.S. focused analysis [in “Explaining Extreme Events of 2013 from a Climate Perspective”]. *Bull. Am.*  
13 *Meteorol. Soc.* 95, S19–S23. doi:10.1175/1520-0477-95.9.S1.1.
- 14 Kodama, C., Stevens, B., Mauritsen, T., Seiki, T., and Satoh, M. (2019). A New Perspective for Future Precipitation  
15 Change from Intense Extratropical Cyclones. *Geophys. Res. Lett.* 46. doi:10.1029/2019GL084001.
- 16 Kodama, C., Yamada, Y., Noda, A. T., Kikuchi, K., Kajikawa, Y., Nasuno, T., et al. (2015). A 20-year climatology of a  
17 NICAM AMIP-type simulation. *J. Meteorol. Soc. Japan* 93, 393–424. doi:10.2151/jmsj.2015-024.
- 18 Kogan, F., and Guo, W. (2017). Strong 2015–2016 El Niño and implication to global ecosystems from space data. *Int.*  
19 *J. Remote Sens.* 38, 161–178. doi:10.1080/01431161.2016.1259679.
- 20 Konapala, G., and Mishra, A. (2017). Review of complex networks application in hydroclimatic extremes with an  
21 implementation to characterize spatio-temporal drought propagation in continental USA. *J. Hydrol.* 555, 600–  
22 620. doi:10.1016/j.jhydrol.2017.10.033.
- 23 Konapala, G., and Mishra, A. (2020). Quantifying Climate and Catchment Control on Hydrological Drought in the  
24 Continental United States. *Water Resour. Res.* 56. doi:10.1029/2018WR024620.
- 25 Kong, Q., Guerreiro, S. B., Blenkinsop, S., Li, X.-F., and Fowler, H. J. (2020). Increases in summertime concurrent  
26 drought and heatwave in Eastern China. *Weather Clim. Extrem.* 28, 100242. doi:10.1016/j.wace.2019.100242.
- 27 Konings, A. G., Williams, A. P., and Gentine, P. (2017). Sensitivity of grassland productivity to aridity controlled by  
28 stomatal and xylem regulation. *Nat. Geosci.* 10, 284. doi:10.1038/ngeo2903.
- 29 Kooperman, G. J., Fowler, M. D., Hoffman, F. M., Koven, C. D., Lindsay, K., Pritchard, M. S., et al. (2018). Plant  
30 Physiological Responses to Rising CO2 Modify Simulated Daily Runoff Intensity With Implications for Global-  
31 Scale Flood Risk Assessment. *Geophys. Res. Lett.* 45, 12,412–457,466. doi:10.1029/2018GL079901.
- 32 Kornhuber, K., Coumou, D., Vogel, E., Lesk, C., Donges, J. F., Lehmann, J., et al. (2020). Amplified Rossby waves  
33 enhance risk of concurrent heatwaves in major breadbasket regions. *Nat. Clim. Chang.* 10, 48–53.  
34 doi:10.1038/s41558-019-0637-z.
- 35 Kornhuber, K., Osprey, S., Coumou, D., Petri, S., Petoukhov, V., Rahmstorf, S., et al. (2019). Extreme weather events  
36 in early summer 2018 connected by a recurrent hemispheric wave-7 pattern. *Environ. Res. Lett.* 14, 54002.  
37 doi:10.1088/1748-9326/ab13bf.
- 38 Kossin, J. P. (2017). Hurricane intensification along United States coast suppressed during active hurricane periods.  
39 *Nature* 541, 390–393. doi:10.1038/nature20783.
- 40 Kossin, J. P. (2018). A global slowdown of tropical-cyclone translation speed. *Nature* 558, 104–107.  
41 doi:10.1038/s41586-018-0158-3.
- 42 Kossin, J. P. (2019). Reply to: Moon, I.-J. et al.; Lanzante, J. R. *Nature* 570, E16–E22. doi:10.1038/s41586-019-1224-  
43 1.
- 44 Kossin, J. P., Emanuel, K. A., and Camargo, S. J. (2016a). Past and Projected Changes in Western North Pacific  
45 Tropical Cyclone Exposure. *J. Clim.* 29, 5725–5739. doi:10.1175/JCLI-D-16-0076.1.
- 46 Kossin, J. P., Emanuel, K. A., and Vecchi, G. A. (2014). The poleward migration of the location of tropical cyclone  
47 maximum intensity. *Nature*, 4–7. doi:10.1038/nature13278.
- 48 Kossin, J. P., Emanuel, K. A., and Vecchi, G. A. (2016b). Comment on ‘Roles of interbasin frequency changes in the  
49 poleward shifts of the maximum intensity location of tropical cyclones.’ *Environ. Res. Lett.* 11, 068001.  
50 doi:10.1088/1748-9326/11/6/068001.
- 51 Kossin, J. P., Hall, T., Knutson, T., Kunkel, K. E., Trapp, R. J., Waliser, D. E., et al. (2017). “Extreme Storms. Climate  
52 Science Special Report: Fourth National Climate Assessment, Volume I,” in *Climate Science Special Report:*  
53 *Fourth National Climate Assessment, Volume I*, eds. D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken,  
54 B. C. Stewart, and T. K. Maycock (Washington, DC, USA: U.S. Global Change Research Program), 257–276.  
55 doi:10.7930/J07S7KXX.
- 56 Kossin, J. P., Knapp, K. R., Olander, T. L., and Velden, C. S. (2020). Global increase in major tropical cyclone  
57 exceedance probability over the past four decades. *Proc. Natl. Acad. Sci.* 117, 11975–11980.  
58 doi:10.1073/pnas.1920849117.
- 59 Kossin, J. P., Olander, T. L., and Knapp, K. R. (2013). Trend analysis with a new global record of tropical cyclone  
60 intensity. *J. Clim.* 26, 9960–9976. doi:10.1175/JCLI-D-13-00262.1.
- 61 Koster, R. D., Chang, Y., Wang, H., and Schubert, S. D. (2016). Impacts of Local Soil Moisture Anomalies on the  
**Do Not Cite, Quote or Distribute** 11-268 Total pages: 345

- 1 Atmospheric Circulation and on Remote Surface Meteorological Fields during Boreal Summer: A  
2 Comprehensive Analysis over North America. *J. Clim.* 29, 7345–7364. doi:10.1175/JCLI-D-16-0192.1.
- 3 Koster, R. D., Guo, Z., Yang, R., Dirmeyer, P. A., Mitchell, K., and Puma, M. J. (2009). On the Nature of Soil Moisture  
4 in Land Surface Models. *J. Clim.* 22, 4322–4335. doi:10.1175/2009JCLI2832.1.
- 5 Koster, R. D., Mahanama, S. P. P., Yamada, T. J., Balsamo, G., Berg, A. A., Boisserie, M., et al. (2011). The second  
6 phase of the global land-atmosphere coupling experiment: Soil moisture contributions to subseasonal forecast  
7 skill. *J. Hydrometeorol.* 12, 805–822. doi:10.1175/2011JHM1365.1.
- 8 Krishnan, R., Sabin, T. P., Vellore, R., Mujumdar, M., Sanjay, J., Goswami, B. N., et al. (2016). Deciphering the  
9 desiccation trend of the South Asian monsoon hydroclimate in a warming world. *Clim. Dyn.* 47, 1007–1027.  
10 doi:10.1007/s00382-015-2886-5.
- 11 Krueger, O., Schenk, F., Feser, F., and Weisse, R. (2013). Inconsistencies between long-term trends in storminess  
12 derived from the 20CR reanalysis and observations. *J. Clim.* 26, 868–874. doi:10.1175/JCLI-D-12-00309.1.
- 13 Krug, A., Primo, C., Fischer, S., Schumann, A., and Ahrens, B. (2020). On the temporal variability of widespread rain-  
14 on-snow floods. *Meteorol. Zeitschrift* 29, 147–163. doi:10.1127/METZ/2020/0989.
- 15 Kruger, A. C., and Nxumalo, M. (2017). Surface temperature trends from homogenized time series in South Africa:  
16 1931–2015. *Int. J. Climatol.* 37, 2364–2377.
- 17 Kruger, A. C., Rautenbach, H., Mbatha, S., Ngwenya, S., and Makgoale, T. E. (2019). Historical and projected trends in  
18 near-surface temperature indices for 22 locations in South Africa. *S. Afr. J. Sci.* 115.  
19 doi:10.17159/sajs.2019/4846.
- 20 Kruger, A. C., and Sekele, S. S. (2013). Trends in extreme temperature indices in South Africa: 1962–2009. *Int. J.*  
21 *Climatol.* 33, 661–676. doi:10.1002/joc.3455.
- 22 Krysanova, V., Vetter, T., Eisner, S., Huang, S., Pechlivanidis, I., Strauch, M., et al. (2017). Intercomparison of  
23 regional-scale hydrological models and climate change impacts projected for 12 large river basins worldwide - A  
24 synthesis. *Environ. Res. Lett.* 12. doi:10.1088/1748-9326/aa8359.
- 25 Kubota, H., Matsumoto, J., Zaiki, M., Tsukahara, T., Mikami, T., Allan, R., et al. (2021). Tropical cyclones over the  
26 western north Pacific since the mid-nineteenth century. *Clim. Change* 164, 29. doi:10.1007/s10584-021-02984-7.
- 27 Kuleshov, Y., Fawcett, R., Qi, L., Trewin, B., Jones, D., McBride, J., et al. (2010). Trends in tropical cyclones in the  
28 South Indian Ocean and the South Pacific Ocean. *J. Geophys. Res. Atmos.* 115. doi:10.1029/2009JD012372.
- 29 Kumar, S., Allan, R. P., Zwiers, F., Lawrence, D. M., and Dirmeyer, P. A. (2015). Revisiting trends in wetness and  
30 dryness in the presence of internal climate variability and water limitations over land. *Geophys. Res. Lett.* 42,  
31 10,867–10,875. doi:10.1002/2015GL066858.
- 32 Kumar, S. P. and R. S. N. and D. N. (2017). Linkage between global sea surface temperature and hydroclimatology of a  
33 major river basin of India before and after 1980. *Environ. Res. Lett.* 12, 124002. doi:10.1088/1748-9326/aa9664.
- 34 Kundzewicz, Z. W., Pin'skwar, I., and Brakenridge, G. R. (2018). Changes in river flood hazard in Europe: A review.  
35 *Hydrol. Res.* 49, 294–302. doi:10.2166/nh.2017.016.
- 36 Kunii, M., Otsuka, M., Shimoji, K., and Seko, H. (2016). Ensemble Data Assimilation and Forecast Experiments for the  
37 September 2015 Heavy Rainfall Event in Kanto and Tohoku Regions with Atmospheric Motion Vectors from  
38 Himawari-8. *SOLA* 12, 209–214. doi:10.2151/sola.2016-042.
- 39 Kunkel, K. E., Karl, T. R., Brooks, H., Kossin, J., Lawrimore, J. H., Arndt, D., et al. (2013). Monitoring and  
40 understanding trends in extreme storms: State of knowledge. *Bull. Am. Meteorol. Soc.* 94, 499–514.  
41 doi:10.1175/BAMS-D-11-00262.1.
- 42 Kusunoki, S. (2017). Future Changes in Global Precipitation Projected by the Atmospheric Model MRI-AGCM3.2H  
43 with a 60-km Size. *Atmos.* 8. doi:10.3390/atmos8050093.
- 44 Kusunoki, S. (2018a). Future changes in precipitation over East Asia projected by the global atmospheric model MRI-  
45 AGCM3.2. *Clim. Dyn.* 51, 4601–4617. doi:10.1007/s00382-016-3499-3.
- 46 Kusunoki, S. (2018b). Is the global atmospheric model MRI-AGCM3.2 better than the CMIP5 atmospheric models in  
47 simulating precipitation over East Asia? *Clim. Dyn.* 51, 4489–4510. doi:10.1007/s00382-016-3335-9.
- 48 Kusunoki, S., and Mizuta, R. (2013). Changes in precipitation intensity over East Asia during the 20th and 21st  
49 centuries simulated by a global atmospheric model with a 60 km grid size. *J. Geophys. Res. Atmos.* 118, 11,007-  
50 11,016. doi:10.1002/jgrd.50877.
- 51 Kusunoki, S., Nakaegawa, T., Pinzón, R., Sanchez-Galan, J. E., and Fábrega, J. R. (2019). Future precipitation changes  
52 over Panama projected with the atmospheric global model MRI-AGCM3.2. *Clim. Dyn.* 53, 5019–5034.  
53 doi:10.1007/s00382-019-04842-w.
- 54 Lackmann, G. M. (2013). The south-central U.S. flood of may 2010: Present and future. *J. Clim.* 26, 4688–4709.  
55 doi:10.1175/JCLI-D-12-00392.1.
- 56 Lackmann, G. M. (2015). Hurricane Sandy before 1900 and after 2100. *Bull. Am. Meteorol. Soc.* 96, 547–560.  
57 doi:10.1175/BAMS-D-14-00123.1.
- 58 Laliberté, F., Howell, S. E. L., and Kushner, P. J. (2015). Regional variability of a projected sea ice-free Arctic during  
59 the summer months. *Geophys. Res. Lett.* 43, 256–263. doi:10.1002/2015GL066855.
- 60 Landsea, C. W. (2015). Comments on “Monitoring and understanding trends in extreme storms: state of knowledge.”.  
61 *Bull. Am. Meteorol. Soc.* 96, 1175–1182. doi:10.1175/1520-0477-96.7.1175.

- 1 Langerwisch, F., Rost, S., Gerten, D., Poulter, B., Rammig, A., and Cramer, W. (2013). Potential effects of climate  
2 change on inundation patterns in the Amazon Basin. *Hydrol. Earth Syst. Sci.* 17, 2247–2262. doi:10.5194/hess-  
3 17-2247-2013.
- 4 Lanzante, J. R. (2019). Uncertainties in tropical-cyclone translation speed. *Nature* 570, E6–E15. doi:10.1038/s41586-  
5 019-1223-2.
- 6 Lau, N.-C., and Nath, M. J. (2014). Model simulation and projection of European heat waves in present-day and future  
7 climates. *J. Clim.* 27, 3713–3730. doi:10.1175/JCLI-D-13-00284.1.
- 8 Lawal, K. A., Abatan, A. A., Angélil, O., Olaniyan, E., Olusoji, V. H., Oguntunde, P. G., et al. (2016). The Late Onset  
9 of the 2015 Wet Season in Nigeria. *Bull. Am. Meteorol. Soc.* 97, S63–S69. doi:10.1175/BAMS-D-16-0131.1.
- 10 Leach, N. J., Li, S., Sparrow, S., van Oldenborgh, G. J., Lott, F. C., Weisheimer, A., et al. (2020). Anthropogenic  
11 Influence on the 2018 Summer Warm Spell in Europe: The Impact of Different Spatio-Temporal Scales. *Bull.*  
12 *Am. Meteorol. Soc.* 101, S41–S46. doi:10.1175/BAMS-D-19-0201.1.
- 13 Lee, C.-Y., Camargo, S. J., Sobel, A. H., and Tippet, M. K. (2020a). Statistical–Dynamical Downscaling Projections of  
14 Tropical Cyclone Activity in a Warming Climate: Two Diverging Genesis Scenarios. *J. Clim.* 33, 4815–4834.  
15 doi:10.1175/JCLI-D-19-0452.1.
- 16 Lee, C. Y., Tippet, M. K., Sobel, A. H., and Camargo, S. J. (2018a). An environmentally forced tropical cyclone  
17 hazard model. *J. Adv. Model. Earth Syst.* 10, 223–241. doi:10.1002/2017MS001186.
- 18 Lee, D., Min, S., Fischer, E., Shioyama, H., and Bethke, I. (2018b). Impacts of half a degree additional warming on the  
19 Asian summer monsoon rainfall characteristics. *Environ. Res. Lett.* 13. doi:10.1088/1748-9326/aab55d.
- 20 Lee, J., Li, S., Lund, R., Lee, J., Li, S., and Lund, R. (2014). Trends in Extreme U.S. Temperatures. *J. Clim.* 27, 4209–  
21 4225. doi:10.1175/JCLI-D-13-00283.1.
- 22 Lee, T.-C., Knutson, T. R., Nakaegawa, T., Ying, M., and Cha, E. J. (2020b). Third assessment on impacts of climate  
23 change on tropical cyclones in the Typhoon Committee Region – Part I: Observed changes, detection and  
24 attribution. *Trop. Cyclone Res. Rev.* 9, 1–22. doi:10.1016/j.tccr.2020.03.001.
- 25 Lee, W., Choi, H. M., Kim, D., Honda, Y., Guo, Y.-L. L., and Kim, H. (2018c). Temporal changes in mortality  
26 attributed to heat extremes for 57 cities in Northeast Asia. *Sci. Total Environ.* 616–617, 703–709.  
27 doi:10.1016/j.scitotenv.2017.10.258.
- 28 Lehner, F., Coats, S., Stocker, T. F., Pendergrass, A. G., Sanderson, B. M., Raible, C. C., et al. (2017). Projected  
29 drought risk in 1.5°C and 2°C warmer climates. *Geophys. Res. Lett.* 44, 7419–7428. doi:10.1002/2017GL074117.
- 30 Lehner, F., Deser, C., Simpson, I. R., and Terray, L. (2018). Attributing the U.S. Southwest’s Recent Shift Into Drier  
31 Conditions. *Geophys. Res. Lett.* 45, 6251–6261. doi:10.1029/2018GL078312.
- 32 Lehner, F., and Stocker, T. F. (2015). From local perception to global perspective. *Nat. Clim. Chang.* 5, 731–734.  
33 doi:10.1038/nclimate2660.
- 34 Lejeune, Q., Davin, E. L., Gudmundsson, L., Winckler, J., and Seneviratne, S. I. (2018). Historical deforestation locally  
35 increased the intensity of hot days in northern mid-latitudes. *Nat. Clim. Chang.* 8, 386–390. doi:10.1038/s41558-  
36 018-0131-z.
- 37 Lejeune, Q., Seneviratne, S. I., and Davin, E. L. (2017). Historical Land-Cover Change Impacts on Climate:  
38 Comparative Assessment of LUCID and CMIP5 Multimodel Experiments. *J. Clim.* 30, 1439–1459.  
39 doi:10.1175/JCLI-D-16-0213.1.
- 40 Lelieveld, J., Proestos, Y., Hadjinicolaou, P., Tanarhte, M., Tyrllis, E., and Zittis, G. (2016). Strongly increasing heat  
41 extremes in the Middle East and North Africa (MENA) in the 21st century. *Clim. Change* 137, 245–260.  
42 doi:10.1007/s10584-016-1665-6.
- 43 Lemordant, L., Gentine, P., Stéfanon, M., Drobinski, P., and Fatichi, S. (2016). Modification of land-atmosphere  
44 interactions by CO<sub>2</sub> effects: Implications for summer dryness and heat wave amplitude. *Geophys. Res. Lett.* 43,  
45 10,240-10,248. doi:10.1002/2016GL069896.
- 46 Lemordant, L., Gentine, P., Swann, A. S., Cook, B. I., and Scheff, J. (2018). Critical impact of vegetation physiology  
47 on the continental hydrologic cycle in response to increasing CO<sub>2</sub>. *Proc. Natl. Acad. Sci.* 115, 4093–4098.  
48 doi:10.1073/pnas.1720712115.
- 49 Lenderink, G., Barbero, R., Loriaux, J. M., and Fowler, H. J. (2017). Super-Clausius–Clapeyron Scaling of Extreme  
50 Hourly Convective Precipitation and Its Relation to Large-Scale Atmospheric Conditions. *J. Clim.* 30, 6037–  
51 6052. doi:10.1175/JCLI-D-16-0808.1.
- 52 Lewis, E., Fowler, H., Alexander, L., Dunn, R., McClean, F., Barbero, R., et al. (2019a). GSDR: A Global Sub-Daily  
53 Rainfall Dataset. *J. Clim.* 32, 4715–4729. doi:10.1175/JCLI-D-18-0143.1.
- 54 Lewis, S. C., Blake, S. A. P., Trewin, B., Black, M. T., Dowdy, A. J., Perkins-Kirkpatrick, S. E., et al. (2019b).  
55 Deconstructing factors contributing to the 2018 fire weather in Queensland, Australia. *Bull. Am. Meteorol. Soc.*,  
56 BAMS-D-19-0144.1. doi:10.1175/BAMS-D-19-0144.1.
- 57 Lewis, S. C., and Karoly, D. J. (2013). Anthropogenic contributions to Australia’s record summer temperatures of 2013.  
58 *Geophys. Res. Lett.* 40, 3705–3709. doi:10.1002/grl.50673.
- 59 Lewis, S. C., and Karoly, D. J. (2014). The Role of Anthropogenic Forcing in the Record 2013 Australia-Wide Annual  
60 and Spring Temperatures [in “Explaining Extreme Events of 2013 from a Climate Perspective”]. *Bull. Am.*  
61 *Meteorol. Soc.* 95, S31–S34. doi:10.1175/1520-0477-95.9.S1.1.

- 1 Lewis, S. C., and King, A. D. (2015). Dramatically increased rate of observed hot record breaking in recent Australian  
2 temperatures. *Geophys. Res. Lett.* 42, 7776–7784. doi:10.1002/2015GL065793.
- 3 Lewis, S. C., King, A. D., and Mitchell, D. M. (2017a). Australia’s Unprecedented Future Temperature Extremes Under  
4 Paris Limits to Warming. *Geophys. Res. Lett.* 44, 9947–9956.
- 5 Lewis, S. C., King, A. D., and Perkins-Kirkpatrick, S. E. (2017b). Defining a New Normal for Extremes in a Warming  
6 World. *Bull. Am. Meteorol. Soc.* 98, 1139–1151. doi:10.1175/BAMS-D-16-0183.1.
- 7 Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. F., and Nepstad, D. (2011). The 2010 Amazon  
8 Drought. *Science (80-. )*. 331, 554–554. doi:10.1126/science.1200807.
- 9 Lhotka, O., Kysely, J., and Farda, A. (2018). Climate change scenarios of heat waves in Central Europe and their  
10 uncertainties. *Theor. Appl. Climatol.* 131, 1043–1054. doi:10.1007/s00704-016-2031-3.
- 11 Li, C., Fang, Y., Caldeira, K., Zhang, X., Diffenbaugh, N. S., and Michalak, A. M. (2018a). Widespread persistent  
12 changes to temperature extremes occurred earlier than predicted. *Sci. Rep.* 8, 1007. doi:10.1038/s41598-018-  
13 19288-z.
- 14 Li, C., Zhang, X., Zwiers, F., Fang, Y., and Michalak, A. M. (2017a). Recent Very Hot Summers in Northern  
15 Hemispheric Land Areas Measured by Wet Bulb Globe Temperature Will Be the Norm Within 20 Years. *Earth’s*  
16 *Futur.* 5, 1203–1216. doi:10.1002/2017EF000639.
- 17 Li, C., Zwiers, F., Zhang, X., Chen, G., Lu, J., Li, G., et al. (2019a). Larger Increases in More Extreme Local  
18 Precipitation Events as Climate Warms. *Geophys. Res. Lett.* 46, 6885–6891. doi:10.1029/2019GL082908.
- 19 Li, C., Zwiers, F., Zhang, X., and Li, G. (2018b). How much information is required to well-constrain local estimates of  
20 future precipitation extremes? *Earth’s Futur.*, 2018EF001001. doi:10.1029/2018EF001001.
- 21 Li, C., Zwiers, F., Zhang, X., Li, G., Sun, Y., and Wehner, M. (2020a). Changes in annual extremes of daily  
22 temperature and precipitation in CMIP6 models. *J. Clim.* (submitted, 1–61. doi:10.1175/JCLI-D-19-1013.1.
- 23 Li, D., Lettenmaier, D. P., Margulis, S. A., and Andreadis, K. (2019b). The Role of Rain-on-Snow in Flooding Over the  
24 Conterminous United States. *Water Resour. Res.* 55, 8492–8513. doi:10.1029/2019WR024950.
- 25 Li, D., Zhou, T., Zou, L., Zhang, W., and Zhang, L. (2018c). Extreme High-Temperature Events Over East Asia in  
26 1.5°C and 2°C Warmer Futures: Analysis of NCAR CESM Low-Warming Experiments. *Geophys. Res. Lett.* 45,  
27 1541–1550. doi:10.1002/2017GL076753.
- 28 Li, G., Zhang, X., Cannon, A. J., Murdock, T., Sobie, S., Zwiers, F., et al. (2018d). Indices of Canada’s future climate  
29 for general and agricultural adaptation applications. *Clim. Change* 148, 249–263. doi:10.1007/s10584-018-2199-  
30 x.
- 31 Li, H., Chen, H., and Wang, H. (2017b). Effects of anthropogenic activity emerging as intensified extreme precipitation  
32 over China. *J. Geophys. Res. Atmos.* 122, 6899–6914. doi:10.1002/2016JD026251.
- 33 Li, H., and Sriver, R. L. (2018). Tropical Cyclone Activity in the High-Resolution Community Earth System Model and  
34 the Impact of Ocean Coupling. *J. Adv. Model. Earth Syst.* 10, 165–186. doi:10.1002/2017MS001199.
- 35 Li, L., She, D., Zheng, H., Lin, P., and Yang, Z.-L. (2020b). Elucidating diverse drought characteristics from two  
36 meteorological drought indices (SPI and SPEI) in China. *J. Hydrometeorol.* 21, 1513–1530. doi:10.1175/JHM-D-  
37 19-0290.s1.
- 38 Li, L., Yao, N., Li, Y., Liu, D. L., Wang, B., and Ayantobo, O. O. (2019c). Future projections of extreme temperature  
39 events in different sub-regions of China. *Atmos. Res.* 217, 150–164. doi:10.1016/j.atmosres.2018.10.019.
- 40 Li, M. (2017). Temperature reconstruction and volcanic eruption signal from tree-ring width and maximum latewood  
41 density over the past 304 years in the southeastern Tibetan Plateau. 2021–2032. doi:10.1007/s00484-017-1395-0.
- 42 Li, M., Woollings, T., Hodges, K., and Masato, G. (2014). Extratropical cyclones in a warmer, moister climate: A  
43 recent Atlantic analogue. *Geophys. Res. Lett.* 41, 8594–8601. doi:10.1002/2014GL062186.
- 44 Li, S., Otto, F. E. L., Harrington, L. J., Sparrow, S. N., Wallom, D., Wallom, S. L. and F. E. L. O. and L. J. H. and S. N.  
45 S. and D., et al. (2019d). A pan-South-America assessment of avoided exposure to dangerous extreme  
46 precipitation by limiting to 1.5 °C warming. *Environ. Res. Lett.* doi:10.1088/1748-9326/ab50a2.
- 47 Li, W., Jiang, Z., Zhang, X., and Li, L. (2018e). On the Emergence of Anthropogenic Signal in Extreme Precipitation  
48 Change Over China. *Geophys. Res. Lett.* doi:10.1029/2018GL079133.
- 49 Li, W., Jiang, Z., Zhang, X., Li, L., and Sun, Y. (2018f). Additional risk in extreme precipitation in China from 1.5 °C  
50 to 2.0 °C global warming levels. *Sci. Bull.* 63, 228–234. doi:10.1016/j.scib.2017.12.021.
- 51 Li, X., Liu, L., Li, H., Wang, S., and Heng, J. (2020c). Spatiotemporal soil moisture variations associated with hydro-  
52 meteorological factors over the Yarlung Zangbo River basin in Southeast Tibetan Plateau. *Int. J. Climatol.* 40,  
53 188–206. doi:10.1002/joc.6202.
- 54 Li, X., Wang, X., and Babovic, V. (2018g). Analysis of variability and trends of precipitation extremes in Singapore  
55 during 1980–2013. *Int. J. Climatol.* 38, 125–141. doi:10.1002/joc.5165.
- 56 Li, X., You, Q., Ren, G., Wang, S., Zhang, Y., Yang, J., et al. (2019e). Concurrent droughts and hot extremes in  
57 northwest China from 1961 to 2017. *Int. J. Climatol.* 39, 2186–2196. doi:10.1002/joc.5944.
- 58 Li, Y., Zhao, M., Motesharrei, S., Mu, Q., Kalnay, E., and Li, S. (2015). Local cooling and warming effects of forests  
59 based on satellite observations. *Nat. Commun.* 6, 6603. doi:10.1038/ncomms7603.
- 60 Li, Z., Chen, Y., Fang, G., and Li, Y. (2017c). Multivariate assessment and attribution of droughts in Central Asia. *Sci.*  
61 *Rep.* 7, 1316. doi:10.1038/s41598-017-01473-1.

- 1 Li, Z., Chen, Y., Shen, Y., Liu, Y., and Zhang, S. (2013). Analysis of changing pan evaporation in the arid region of  
2 Northwest China. *Water Resour. Res.* 49, 2205–2212. doi:10.1002/wrcr.20202.
- 3 Lian, X., Piao, S., Li, L. Z. X., Li, Y., Huntingford, C., Ciais, P., et al. (2020). Summer soil drying exacerbated by  
4 earlier spring greening of northern vegetation. *Sci. Adv.* 6, eaax0255. doi:10.1126/sciadv.aax0255.
- 5 Liang, C., Chen, T., Dolman, H., Shi, T., Wei, X., Xu, J., et al. (2020). Drying and wetting trends and vegetation  
6 covariations in the drylands of China. *Water (Switzerland)* 12. doi:10.3390/W12040933.
- 7 Liang, J., Wang, C., and Hodges, K. I. (2017). Evaluation of tropical cyclones over the South China Sea simulated by  
8 the 12 km MetUM regional climate model. *Q. J. R. Meteorol. Soc.* 143, 1641–1656. doi:10.1002/qj.3035.
- 9 Libertino, A., Ganora, D., and Claps, P. (2019). Evidence for Increasing Rainfall Extremes Remains Elusive at Large  
10 Spatial Scales: The Case of Italy. *Geophys. Res. Lett.* 46, 7437–7446. doi:10.1029/2019GL083371.
- 11 Lin, C., Yang, K., Qin, J., and Fu, R. (2013). Observed Coherent Trends of Surface and Upper-Air Wind Speed over  
12 China since 1960. *J. Clim.* 26, 2891–2903. doi:10.1175/JCLI-D-12-00093.1.
- 13 Lin, L., Wang, Z., Xu, Y., and Fu, Q. (2016). Sensitivity of precipitation extremes to radiative forcing of greenhouse  
14 gases and aerosols. *Geophys. Res. Lett.* 43, 9860–9868. doi:10.1002/2016GL070869.
- 15 Lin, L., Wang, Z., Xu, Y., Fu, Q., and Dong, W. (2018). Larger Sensitivity of Precipitation Extremes to Aerosol Than  
16 Greenhouse Gas Forcing in CMIP5 Models. *J. Geophys. Res. Atmos.* doi:10.1029/2018JD028821.
- 17 Lin, N., Lane, P., Emanuel, K. A., Sullivan, R. M., and Donnelly, J. P. (2014). Heightened hurricane surge risk in  
18 northwest Florida revealed from climatological-hydrodynamic modeling and paleorecord reconstruction. *J.*  
19 *Geophys. Res. Atmos.* 119, 8606–8623. doi:10.1002/2014JD021584.
- 20 Lin, P., He, Z., Du, J., Chen, L., Zhu, X., and Li, J. (2017). Recent changes in daily climate extremes in an arid  
21 mountain region, a case study in northwestern China’s Qilian Mountains. *Sci. Rep.* 7, 1–15. doi:10.1038/s41598-  
22 017-02345-4.
- 23 Lionello, P., and Scarascia, L. (2020). The relation of climate extremes with global warming in the Mediterranean  
24 region and its north versus south contrast. *Reg. Environ. Chang.* 20, 31. doi:10.1007/s10113-020-01610-z.
- 25 Littell, J. S., Peterson, D. L., Riley, K. L., Liu, Y., and Luce, C. H. (2016). A review of the relationships between  
26 drought and forest fire in the United States. *Glob. Chang. Biol.* 22, 2353–2369. doi:10.1111/gcb.13275.
- 27 Little, C. M., Horton, R. M., Kopp, R. E., Oppenheimer, M., Vecchi, G. A., and Villarini, G. (2015). Joint projections  
28 of US East Coast sea level and storm surge. *Nat. Clim. Chang.* 5, 1114. doi:10.1038/nclimate2801.
- 29 Liu, B., Zhu, C., Su, J., Ma, S., and Xu, K. (2019a). Record-Breaking Northward Shift of the Western North Pacific  
30 Subtropical High in July 2018. *J. Meteorol. Soc. Japan. Ser. II* 97, 913–925. doi:10.2151/jmsj.2019-047.
- 31 Liu, C., and Zipser, E. J. (2015). The global distribution of largest, deepest, and most intense precipitation systems.  
32 *Geophys. Res. Lett.* 42, 3591–3595. doi:10.1002/2015GL063776.
- 33 Liu, L., Gudmundsson, L., Hauser, M., Qin, D., Li, S., and Seneviratne, S. I. (2020). Soil moisture dominates dryness  
34 stress on ecosystem production globally. *Nat. Commun.* 11, 4892. doi:10.1038/s41467-020-18631-1.
- 35 Liu, M., Shen, Y., Qi, Y., Wang, Y., and Geng, X. (2019b). Changes in precipitation and drought extremes over the past  
36 half century in China. *Atmosphere (Basel)*. 10. doi:10.3390/atmos10040203.
- 37 Liu, M., Vecchi, G. A., Smith, J. A., and Knutson, T. R. (2019c). Causes of large projected increases in hurricane  
38 precipitation rates with global warming. *npj Clim. Atmos. Sci.* 2, 1–5. doi:10.1038/s41612-019-0095-3.
- 39 Liu, Q., Babanin, A. V., Zieger, S., Young, I. R., and Guan, C. (2016a). Wind and wave climate in the Arctic Ocean as  
40 observed by altimeters. *J. Clim.* 29, 7957–7975. doi:10.1175/JCLI-D-16-0219.1.
- 41 Liu, T., Huang, C. C., Pang, J., Zhou, Y., Zhang, Y., Ji, L., et al. (2014). Extraordinary hydro-climatic events during  
42 1800 – 1600 yr BP in the Jin – Shaan Gorges along the middle Yellow River, China. *Palaeogeogr.*  
43 *Palaeoclimatol. Palaeoecol.* 410, 143–152. doi:10.1016/j.palaeo.2014.05.039.
- 44 Liu, W., and Sun, F. (2016). Assessing estimates of evaporative demand in climate models using observed pan  
45 evaporation over China. *J. Geophys. Res.* 121, 8329–8349. doi:10.1002/2016JD025166.
- 46 Liu, W., and Sun, F. (2017). Projecting and attributing future changes of evaporative demand over china in CMIP5  
47 climate models. *J. Hydrometeorol.* 18, 977–991. doi:10.1175/JHM-D-16-0204.1.
- 48 Liu, Y., Pan, Z., Zhuang, Q., Miralles, D. G., Teuling, A. J., Zhang, T., et al. (2015). Agriculture intensifies soil  
49 moisture decline in Northern China. *Sci. Rep.* 5, 11261. doi:10.1038/srep11261.
- 50 Liu, Y., Wu, G., Guo, R., and Wan, R. (2016b). Changing landscapes by damming: the Three Gorges Dam causes  
51 downstream lake shrinkage and severe droughts. *Landsc. Ecol.* 31, 1883–1890. doi:10.1007/s10980-016-0391-9.
- 52 Liuzzo, L., Viola, F., and Noto, L. (2014). Wind speed and temperature trends impacts on reference evapotranspiration  
53 in Southern Italy. *Theor. Appl. Climatol.* 123. doi:10.1007/s00704-014-1342-5.
- 54 Livneh, B., and Badger, A. M. (2020). Drought less predictable under declining future snowpack. *Nat. Clim. Chang.* 10,  
55 452–458. doi:10.1038/s41558-020-0754-8.
- 56 Llasat, M. C., Marcos, R., Turco, M., Gilabert, J., and Llasat-Botija, M. (2016). Trends in flash flood events versus  
57 convective precipitation in the Mediterranean region: The case of Catalonia. *J. Hydrol.* 541, 24–37.  
58 doi:10.1016/j.jhydrol.2016.05.040.
- 59 Lloyd-Hughes, B. (2014). The impracticality of a universal drought definition. *Theor. Appl. Climatol.* 117, 607–611.  
60 doi:10.1007/s00704-013-1025-7.
- 61 Lok, C. C. F., and Chan, J. C. L. (2018). Changes of tropical cyclone landfalls in South China throughout the twenty-

- 1 first century. *Clim. Dyn.* 51, 2467–2483. doi:10.1007/s00382-017-4023-0.
- 2 López-Franca, N., Zaninelli, P., Carril, A., Menéndez, C., and Sánchez, E. (2016). Changes in temperature extremes for  
3 21st century scenarios over South America derived from a multi-model ensemble of regional climate models.  
4 *Clim. Res.* 68, 151–167. doi:10.3354/cr01393.
- 5 Lorenz, R., Argüeso, D., Donat, M. G., Pitman, A. J., Hurk, B. Van Den, Berg, A., et al. (2016). Influence of land-  
6 atmosphere feedbacks on temperature and precipitation extremes in the GLACE-CMIP5 ensemble. *J. Geophys.*  
7 *Res.* 121, 607–623. doi:10.1002/2015JD024053.
- 8 Lorenz, R., Stalhandske, Z., and Fischer, E. M. (2019). Detection of a Climate Change Signal in Extreme Heat, Heat  
9 Stress, and Cold in Europe From Observations. *Geophys. Res. Lett.* 46, 8363–8374. doi:10.1029/2019GL082062.
- 10 Lorenzo-Lacruz, J., Garcia, C., and Morán-Tejeda, E. (2017). Groundwater level responses to precipitation variability  
11 in Mediterranean insular aquifers. *J. Hydrol.* 552, 516–531. doi:10.1016/j.jhydrol.2017.07.011.
- 12 Lorenzo-Lacruz, J., Morán-Tejeda, E., Vicente-Serrano, S. M., and López-Moreno, J. I. (2013). Streamflow droughts in  
13 the Iberian Peninsula between 1945 and 2005: Spatial and temporal patterns. *Hydrol. Earth Syst. Sci.* 17,  
14 doi:10.5194/hess-17-119-2013.
- 15 Lott, F. C., Christidis, N., and Stott, P. A. (2013). Can the 2011 East African drought be attributed to human-induced  
16 climate change? *Geophys. Res. Lett.* 40, 1177–1181. doi:10.1002/grl.50235.
- 17 Lott, F. C., and Stott, P. A. (2016). Evaluating Simulated Fraction of Attributable Risk Using Climate Observations. *J.*  
18 *Clim.* 29, 4565–4575. doi:10.1175/JCLI-D-15-0566.1.
- 19 Louise, B., Cayan, D., Franco, G., Fisher, L., and Ziaja, S. (2018). Statewide Summary Report [in “California’s Fourth  
20 Climate Change Assessment”]. Available at: [https://www.energy.ca.gov/sites/default/files/2019-](https://www.energy.ca.gov/sites/default/files/2019-11/Statewide_Reports-SUM-CCCA4-2018-013_Statewide_Summary_Report_ADA.pdf)  
21 [11/Statewide\\_Reports-SUM-CCCA4-2018-013\\_Statewide\\_Summary\\_Report\\_ADA.pdf](https://www.energy.ca.gov/sites/default/files/2019-11/Statewide_Reports-SUM-CCCA4-2018-013_Statewide_Summary_Report_ADA.pdf).
- 22 Lovino, M. A., Müller, O. V., Berbery, E. H., and Müller, G. V. (2018). How have daily climate extremes changed in  
23 the recent past over northeastern Argentina? *Glob. Planet. Change* 168, 78–97.  
24 doi:10.1016/J.GLOPLACHA.2018.06.008.
- 25 Lu, C., Sun, Y., Wan, H., Zhang, X., and Yin, H. (2016). Anthropogenic influence on the frequency of extreme  
26 temperatures in China. *Geophys. Res. Lett.* 43, 6511–6518. doi:10.1002/2016GL069296.
- 27 Lu, C., Sun, Y., and Zhang, X. (2018). Multimodel detection and attribution of changes in warm and cold spell  
28 durations. *Environ. Res. Lett.* 13, 074013. doi:10.1088/1748-9326/aacb3e.
- 29 Lu, J., Carbone, G. J., and Grego, J. M. (2019). Uncertainty and hotspots in 21st century projections of agricultural  
30 drought from CMIP5 models. *Sci. Rep.* 9, 4922. doi:10.1038/s41598-019-41196-z.
- 31 Lucas, C., Timbal, B., and Nguyen, H. (2014). The expanding tropics: a critical assessment of the observational and  
32 modeling studies. *Wiley Interdiscip. Rev. Clim. Chang.* 5, 89–112. doi:10.1002/wcc.251.
- 33 Luiz Silva, W., Xavier, L. N. R., Maceira, M. E. P., and Rotunno, O. C. (2018). Climatological and hydrological  
34 patterns and verified trends in precipitation and streamflow in the basins of Brazilian hydroelectric plants. *Theor.*  
35 *Appl. Climatol.* doi:10.1007/s00704-018-2600-8.
- 36 Luo, M., and Lau, N.-C. (2016). Heat Waves in Southern China: Synoptic Behavior, Long-Term Change, and  
37 Urbanization Effects. *J. Clim.* 30, 703–720. doi:10.1175/JCLI-D-16-0269.1.
- 38 Luterbacher, J., Werner, J. P., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F. J., Barriopedro, D., et al.  
39 (2016). European summer temperatures since Roman times. *Environ. Res. Lett.* 11, 024001. doi:10.1088/1748-  
40 9326/11/2/024001.
- 41 Ma, S., Zhou, T., Dai, A., and Han, Z. (2015). Observed Changes in the Distributions of Daily Precipitation Frequency  
42 and Amount over China from 1960 to 2013. *J. Clim.* 28, 6960–6978. doi:10.1175/JCLI-D-15-0011.1.
- 43 Ma, S., Zhou, T., Stone, D. A., Polson, D., Dai, A., Stott, P. A., et al. (2017). Detectable Anthropogenic Shift toward  
44 Heavy Precipitation over Eastern China. *J. Clim.* 30, 1381–1396. doi:10.1175/JCLI-D-16-0311.1.
- 45 Macdonald, N., and Sangster, H. (2017). High-magnitude flooding across Britain since AD 1750. *Hydrol. Earth Syst.*  
46 *Sci.* 21, 1631–1650. doi:10.5194/hess-21-1631-2017.
- 47 Maček, U., Bezak, N., and Šraj, M. (2018). Reference evapotranspiration changes in Slovenia, Europe. *Agric. For.*  
48 *Meteorol.* 260–261, 183–192. doi:10.1016/j.agrformet.2018.06.014.
- 49 Madhura, R. K., Krishnan, R., Revadekar, J. V., Mujumdar, M., and Goswami, B. N. (2015). Changes in western  
50 disturbances over the Western Himalayas in a warming environment. *Clim. Dyn.* 44, 1157–1168.  
51 doi:10.1007/s00382-014-2166-9.
- 52 Madsen, H., Lawrence, D., Lang, M., Martinkova, M., and Kjeldsen, T. R. (2014). Review of trend analysis and climate  
53 change projections of extreme precipitation and floods in Europe. *J. Hydrol.* 519, 3634–3650.  
54 doi:10.1016/j.jhydrol.2014.11.003.
- 55 Maggioni, V., Meyers, P. C., and Robinson, M. D. (2016). A Review of Merged High-Resolution Satellite Precipitation  
56 Product Accuracy during the Tropical Rainfall Measuring Mission (TRMM) Era. *J. Hydrometeorol.* 17, 1101–  
57 1117. doi:10.1175/JHM-D-15-0190.1.
- 58 Magnusson, L., Bidlot, J.-R., Lang, S. T. K., Thorpe, A., Wedi, N., and Yamaguchi, M. (2014). Evaluation of Medium-  
59 Range Forecasts for Hurricane Sandy. *Mon. Weather Rev.* 142, 1962–1981. doi:10.1175/MWR-D-13-00228.1.
- 60 Mahoney, K. (2020). Extreme Hail Storms and Climate Change: Foretelling the Future In Tiny, Turbulent Crystal  
61 Balls? *Bull. Am. Meteorol. Soc.* 95, S17–S22. doi:10.1175/BAMS-D-19-0233.1.

- 1 Mahoney, K., Alexander, M. A., Thompson, G., Barsugli, J. J., and Scott, J. D. (2012). Changes in hail and flood risk in  
2 high-resolution simulations over Colorado's mountains. *Nat. Clim. Chang.* 2, 125–131.  
3 doi:10.1038/nclimate1344.
- 4 Mahony, C. R., and Cannon, A. J. (2018). Wetter summers can intensify departures from natural variability in a  
5 warming climate. *Nat. Commun.* 9, 783. doi:10.1038/s41467-018-03132-z.
- 6 Mahto, S. S., and Mishra, V. (2019). Does ERA-5 Outperform Other Reanalysis Products for Hydrologic Applications  
7 in India? *J. Geophys. Res. Atmos.* 124, 9423–9441. doi:10.1029/2019JD031155.
- 8 Maier, N., Breuer, L., Chamorro, A., Kraft, P., and Houska, T. (2018). Multi-Source Uncertainty Analysis in Simulating  
9 Floodplain Inundation under Climate Change. *Water* 10, 809. doi:10.3390/w10060809.
- 10 Maksimović, Č., Prodanović, D., Boonya-Aroonnet, S., Leitão, J. P., Djordjević, S., and Allitt, R. (2009). Overland  
11 flow and pathway analysis for modelling of urban pluvial flooding. *J. Hydraul. Res.* 47, 512–523.  
12 doi:10.1080/00221686.2009.9522027.
- 13 Malik, N., Bookhagen, B., and Mucha, P. J. (2016). Spatiotemporal patterns and trends of Indian monsoonal rainfall  
14 extremes. *Geophys. Res. Lett.* 43, 1710–1717. doi:10.1002/2016GL067841.
- 15 Mallakpour, I., and Villarini, G. (2015). The changing nature of flooding across the central United States. *Nat. Clim.*  
16 *Chang.* 5, 250–254. doi:10.1038/nclimate2516.
- 17 Manda, A., Nakamura, H., Asano, N., Iizuka, S., Miyama, T., Moteki, Q., et al. (2014). Impacts of a warming marginal  
18 sea on torrential rainfall organized under the Asian summer monsoon. *Sci. Rep.* 4, 1–6. doi:10.1038/srep05741.
- 19 Mandapaka, P. V., and Lo, E. Y. M. (2018). Assessment of future changes in Southeast Asian precipitation using the  
20 NASA Earth Exchange Global Daily Downscaled Projections data set. *Int. J. Climatol.* 38, 5231–5244.  
21 doi:10.1002/joc.5724.
- 22 Mangini, W., Viglione, A., Hall, J., Hundecha, Y., Ceola, S., Montanari, A., et al. (2018). Detection of trends in  
23 magnitude and frequency of flood peaks across Europe. *Hydrol. Sci. J.* 63, 493–512.  
24 doi:10.1080/02626667.2018.1444766.
- 25 Mankin, J. S., Seager, R., Smerdon, J. E., Cook, B. I., and Williams, A. P. (2019). Mid-latitude freshwater availability  
26 reduced by projected vegetation responses to climate change. *Nat. Geosci.* 12, 983–988. doi:10.1038/s41561-019-  
27 0480-x.
- 28 Mann, M. E., and Gleick, P. H. (2015). Climate change and California drought in the 21st century. *Proc. Natl. Acad.*  
29 *Sci. U. S. A.* 112, 3858–3859. doi:10.1073/pnas.1503667112.
- 30 Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K., and Coumou, D. (2017). Influence of  
31 Anthropogenic Climate Change on Planetary Wave Resonance and Extreme Weather Events. *Sci. Rep.* 7, 45242.  
32 doi:10.1038/srep45242.
- 33 Marciano, C. G., Lackmann, G. M., and Robinson, W. A. (2015). Changes in U.S. East Coast cyclone dynamics with  
34 climate change. *J. Clim.* 28, 468–484. doi:10.1175/JCLI-D-14-00418.1.
- 35 Marengo, J. A., and Espinoza, J. C. (2016). Extreme seasonal droughts and floods in Amazonia: causes, trends and  
36 impacts. *Int. J. Climatol.* 36, 1033–1050. doi:10.1002/joc.4420.
- 37 Marengo, J. A., Torres, R. R., and Alves, L. M. (2017). Drought in Northeast Brazil – past, present, and future. *Theor.*  
38 *Appl. Climatol.* 129, 1189–1200. doi:10.1007/s00704-016-1840-8.
- 39 Marin, L., Birsan, M. V., Bojariu, R., Dumitrescu, A., Micu, D. M., and Manea, A. (2014). An overview of annual  
40 climatic changes in Romania: Trends in air temperature, precipitation, sunshine hours, cloud cover, relative  
41 humidity and wind speed during the 1961–2013 period. *Carpathian J. Earth Environ. Sci.* 9, 253–258. Available  
42 at: <http://www.cjees.ro/viewTopic.php?topicId=489>.
- 43 Mariotti, A., Pan, Y., Zeng, N., and Alessandri, A. (2015). Long-term climate change in the Mediterranean region in the  
44 midst of decadal variability. *Clim. Dyn.* 44, 1437–1456. doi:10.1007/s00382-015-2487-3.
- 45 Markonis, Y., Kumar, R., Hanel, M., Rakovec, O., Máca, P., and AghaKouchak, A. (2021). The rise of compound  
46 warm-season droughts in Europe. *Sci. Adv.* 7, eabb9668. doi:10.1126/sciadv.abb9668.
- 47 Martens, B., Waegeman, W., Dorigo, W. A., Verhoest, N. E. C., and Miralles, D. G. (2018). Terrestrial evaporation  
48 response to modes of climate variability. *npj Clim. Atmos. Sci.* 1, 43. doi:10.1038/s41612-018-0053-5.
- 49 Marthews, T. R., Otto, F. E. L., Mitchell, D., Dadson, S. J., and Jones, R. G. (2015). The 2014 Drought in the Horn of  
50 Africa: Attribution of Meteorological Drivers [in “Explaining Extreme Events of 2014 from a Climate  
51 Perspective”]. *Bull. Am. Meteorol. Soc.* 96, S83–S88. doi:10.1175/BAMS-D-15-00115.1.
- 52 Martin, E. R. (2018). Future Projections of Global Pluvial and Drought Event Characteristics. *Geophys. Res. Lett.*  
53 doi:10.1029/2018GL079807.
- 54 Martin, J. T., Pederson, G. T., Woodhouse, C. A., Cook, E. R., McCabe, G. J., Anchukaitis, K. J., et al. (2020).  
55 Increased drought severity tracks warming in the United States' largest river basin. *Proc. Natl. Acad. Sci. U. S. A.*  
56 117. doi:10.1073/pnas.1916208117.
- 57 Martinez-Austria, P., and Bandala, E. (2017). Temperature and Heat-Related Mortality Trends in the Sonoran and  
58 Mojave Desert Region. *Atmosphere (Basel)*. 8, 53. doi:10.3390/atmos8030053.
- 59 Martins, E. S. P. R., Coelho, C. A. S., Haarsma, R., Otto, F. E. L., King, A. D., Jan van Oldenborgh, G., et al. (2018). A  
60 Multimethod Attribution Analysis of the Prolonged Northeast Brazil Hydrometeorological Drought (2012–16).  
61 *Bull. Am. Meteorol. Soc.* 99, S65–S69. doi:10.1175/BAMS-D-17-0102.1.

- 1 Martius, O., Pfahl, S., and Chevalier, C. (2016). A global quantification of compound precipitation and wind extremes.  
2 *Geophys. Res. Lett.* 43, 7709–7717. doi:10.1002/2016GL070017.
- 3 Marvel, K., Cook, B. I., Bonfils, C. J. W., Durack, P. J., Smerdon, J. E., and Williams, A. P. (2019). Twentieth-century  
4 hydroclimate changes consistent with human influence. *Nature* 569, 59–65. doi:10.1038/s41586-019-1149-8.
- 5 Marx, A., Kumar, R., Thober, S., Rakovec, O., Wanders, N., Zink, M., et al. (2018). Climate change alters low flows in  
6 Europe under global warming of 1.5, 2, and 3°C. *Hydrol. Earth Syst. Sci.* 22, 1017–1032. doi:10.5194/hess-22-  
7 1017-2018.
- 8 Marzin, C., Rahmat, R., Bernie, D., Bricheno, L., Buonomo, E., Calvert, D., et al. (2015). Singapore 2nd National  
9 Climate Change Study – Phase 1. Singapore: Meteorological Service Singapore Available at:  
10 <http://ccrs.weather.gov.sg/Publications-Second-National-Climate-Change-Science-Reports/>.
- 11 Mascioli, N. R., Fiore, A. M., Previdi, M., and Correa, G. (2016). Temperature and Precipitation Extremes in the United  
12 States: Quantifying the Responses to Anthropogenic Aerosols and Greenhouse Gases. *J. Clim.* 29, 2689–2701.  
13 doi:10.1175/JCLI-D-15-0478.1.
- 14 Masseroni, D., Camici, S., Cislighi, A., Vacchiano, G., Massari, C., and Brocca, L. (2020). 65-year changes of annual  
15 streamflow volumes across Europe with a focus on the Mediterranean basin. *Hydrol. Earth Syst. Sci. Discuss.*  
16 2020, 1–16. doi:10.5194/hess-2020-21.
- 17 Massey, N., Aina, T., Rye, C. J., Otto, F. E. L., Wilson, S., Jones, R. G., et al. (2012). Have the odds of warm  
18 November temperature and of cold December temperatures in central England changed? [in “Explaining Extreme  
19 Events of 2011 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 93, 1057–1059. doi:10.1175/BAMS-D-12-  
20 00021.1.
- 21 Masson-Delmotte, V., M. S., Abe-Ouchi, A., Beer, J., Ganopolski, A., Rouco, J. F. G., et al. (2013). “Information from  
22 Paleoclimate Archives,” in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I  
23 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. T. Stocker, D. Qin, G.-  
24 K. Plattner, M. Tignor, S. K. Allen, J. Boschung, et al. (Cambridge, United Kingdom and New York, NY, USA:  
25 Cambridge University Press), 383–464. doi:10.1017/CBO9781107415324.013.
- 26 Mateo, C. M. R., Yamazaki, D., Kim, H., Champathong, A., Vaze, J., and Oki, T. (2017). Impacts of spatial resolution  
27 and representation of flow connectivity on large-scale simulation of floods. *Hydrol. Earth Syst. Sci.* 21, 5143–  
28 5163. doi:10.5194/hess-21-5143-2017.
- 29 Mathbout, S., Lopez-Bustins, J. A., Martin-Vide, J., Bech, J., and Rodrigo, F. S. (2018a). Spatial and temporal analysis  
30 of drought variability at several time scales in Syria during 1961–2012. *Atmos. Res.* 200, 153–168.  
31 doi:10.1016/j.atmosres.2017.09.016.
- 32 Mathbout, S., Lopez-Bustins, J. A., Royé, D., Martin-Vide, J., Bech, J., and Rodrigo, F. S. (2018b). Observed Changes  
33 in Daily Precipitation Extremes at Annual Timescale Over the Eastern Mediterranean During 1961–2012. *Pure  
34 Appl. Geophys.* 175, 3875–3890. doi:10.1007/s00024-017-1695-7.
- 35 Matthes, H., Rinke, A., and Dethloff, K. (2015). Recent changes in Arctic temperature extremes: Warm and cold spells  
36 during winter and summer. *Environ. Res. Lett.* 10. doi:10.1088/1748-9326/10/11/114020.
- 37 Matthews, T. K. R., Wilby, R. L., and Murphy, C. (2017). Communicating the deadly consequences of global warming  
38 for human heat stress. *Proc. Natl. Acad. Sci. U. S. A.* 114, 3861–3866. doi:10.1073/pnas.1617526114.
- 39 Mátyás, C., and Sun, G. (2014). Forests in a water limited world under climate change. *Environ. Res. Lett.* 9.  
40 doi:10.1088/1748-9326/9/8/085001.
- 41 Maúre, G., Pinto, I., Ndebele-Murisa, M., Muthige, M., Lennard, C., Nikulin, G., et al. (2018). The southern African  
42 climate under 1.5 °C and 2 °C of global warming as simulated by CORDEX regional climate models. *Environ.  
43 Res. Lett.* doi:10.1088/1748-9326/aab190.
- 44 Maxwell, R. M., and Condon, L. E. (2016). Connections between groundwater flow and transpiration partitioning.  
45 *Science (80- )*. 353, 377–380. doi:10.1126/science.aaf7891.
- 46 Mazdiyasi, O., AghaKouchak, A., Davis, S. J., Madadgar, S., Mehran, A., Ragno, E., et al. (2017). Increasing  
47 probability of mortality during Indian heat waves. *Sci. Adv.* 3, e1700066. doi:10.1126/sciadv.1700066.
- 48 Mba, W. P., Longandjo, G. N. T., Moufouma-Okia, W., Bell, J. P., James, R., Vondou, D. A., et al. (2018).  
49 Consequences of 1.5 °C and 2 °C global warming levels for temperature and precipitation changes over Central  
50 Africa. *Environ. Res. Lett.* 13. doi:10.1088/1748-9326/aab048.
- 51 Mbokodo, I., Bopape, M.-J., Chikoore, H., Engelbrecht, F., and Nethengwe, N. (2020). Heatwaves in the Future  
52 Warmer Climate of South Africa. *Atmosphere (Basel)*. 11, 712.
- 53 McBride, J. I., Sahany, S., Hassim, M. E. E., Nguyen, C. M., Lim, S.-Y., Rahmat, R., et al. (2015). The 2014 Record  
54 Dry Spell at Singapore: An Intertropical Convergence Zone (ITCZ) Drought. *Bull. Am. Meteorol. Soc.* 96, S126–  
55 S130. doi:10.1175/BAMS-D-15-00117.1.
- 56 McCollum, D. L., Gambhir, A., Rogelj, J., and Wilson, C. (2020). Energy modellers should explore extremes more  
57 systematically in scenarios. *Nat. Energy* 5, 104–107. doi:10.1038/s41560-020-0555-3.
- 58 McDowell, N. G., and Allen, C. D. (2015). Darcy’s law predicts widespread forest mortality under climate warming.  
59 *Nat. Clim. Chang.* 5, 669–672. doi:10.1038/nclimate2641.
- 60 McDowell, N. G., Allen, C. D., Anderson-Teixeira, K., Aukema, B. H., Bond-Lamberty, B., Chini, L., et al. (2020).  
61 Pervasive shifts in forest dynamics in a changing world. *Science (80- )*. 368. doi:10.1126/science.aaz9463.



- 1 McDowell, N. G., Williams, A. P., Xu, C., Pockman, W. T., Dickman, L. T., Sevanto, S., et al. (2016). Multi-scale  
2 predictions of massive conifer mortality due to chronic temperature rise. *Nat. Clim. Chang.* 6, 295–300.  
3 doi:10.1038/nclimate2873.
- 4 McEvoy, D. J., Huntington, J. L., Hobbins, M. T., Wood, A., Morton, C., Anderson, M., et al. (2016). The evaporative  
5 demand drought index. Part II: CONUS-wide assessment against common drought indicators. *J. Hydrometeorol.*  
6 17, 1763–1779. doi:10.1175/JHM-D-15-0122.1.
- 7 McGree, S., Herold, N., Alexander, L., Schreider, S., Kuleshov, Y., Ene, E., et al. (2019). Recent changes in mean and  
8 extreme temperature and precipitation in the western Pacific Islands. *J. Clim.*, JCLI-D-18-0748.1.  
9 doi:10.1175/JCLI-D-18-0748.1.
- 10 McInnes, K. L., Walsh, K. J. E., Hoeke, R. K., O’Grady, J. G., Colberg, F., and Hubbert, G. D. (2014). Quantifying  
11 storm tide risk in Fiji due to climate variability and change. *Glob. Planet. Change* 116, 115–129.  
12 doi:10.1016/j.gloplacha.2014.02.004.
- 13 McInnes, K. L., White, C. J., Haigh, I. D., Hemer, M. A., Hoeke, R. K., Holbrook, N. J., et al. (2016). Natural hazards  
14 in Australia: sea level and coastal extremes. *Clim. Change* 139, 69–83. doi:10.1007/s10584-016-1647-8.
- 15 McKenzie, D., and Littell, J. S. (2017). Climate change and the eco-hydrology of fire: Will area burned increase in a  
16 warming western USA. *Ecol. Appl.* 27, 26–36. doi:10.1002/eap.1420.
- 17 McLean, N. M., Stephenson, T. S., Taylor, M. A., and Campbell, J. D. (2015). Characterization of Future Caribbean  
18 Rainfall and Temperature Extremes across Rainfall Zones. *Adv. Meteorol.* 2015, 1–18. doi:10.1155/2015/425987.
- 19 McMahan, T. A., Peel, M. C., Lowe, L., Srikanthan, R., and McVicar, T. R. (2013). Estimating actual, potential,  
20 reference crop and pan evaporation using standard meteorological data: A pragmatic synthesis. *Hydrol. Earth  
21 Syst. Sci.* 17, 1331–1363. doi:10.5194/hess-17-1331-2013.
- 22 McVicar, T. R., Roderick, M. L., Donohue, R. J., Li, L. T., Van Niel, T. G., Thomas, A., et al. (2012a). Global review  
23 and synthesis of trends in observed terrestrial near-surface wind speeds: Implications for evaporation. *J. Hydrol.*  
24 416–417, 182–205. doi:10.1016/j.jhydrol.2011.10.024.
- 25 McVicar, T. R., Roderick, M. L., Donohue, R. J., and Van Niel, T. G. (2012b). Less bluster ahead? ecohydrological  
26 implications of global trends of terrestrial near-surface wind speeds. *Ecohydrology* 5, 381–388.  
27 doi:10.1002/eco.1298.
- 28 MedECC (2020). “MedECC 2020 Summary for Policymakers,” in *Climate and Environmental Change in the  
29 Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report*, eds.  
30 W. Cramer, J. Guiot, K. Marini, M. V. Balzan, S. Cherif, E. Doblas-Miranda, et al. (Marseille, France: Union for  
31 the Mediterranean, Plan Bleu, UNEP/MAP), 34. Available at: [https://www.medecc.org/first-mediterranean-  
32 assessment-report-mar1/](https://www.medecc.org/first-mediterranean-assessment-report-mar1/).
- 33 Mediero, L., Kjeldsen, T. R., Macdonald, N., Kohnova, S., Merz, B., Vorogushyn, S., et al. (2015). Identification of  
34 coherent flood regions across Europe by using the longest streamflow records. *J. Hydrol.* 528, 341–360.  
35 doi:10.1016/j.jhydrol.2015.06.016.
- 36 Meehl, G. A., Tebaldi, C., and Adams-Smith, D. (2016). US daily temperature records past, present, and future. *Proc.  
37 Natl. Acad. Sci.* 113, 13977–13982. doi:10.1073/pnas.1606117113.
- 38 Mei, W., and Xie, S. P. (2016). Intensification of landfalling typhoons over the northwest Pacific since the late 1970s.  
39 *Nat. Geosci.* 9, 753–757. doi:10.1038/ngeo2792.
- 40 Mekis, É., Vincent, L. A., Shephard, M. W., and Zhang, X. (2015). Observed Trends in Severe Weather Conditions  
41 Based on Humidex, Wind Chill, and Heavy Rainfall Events in Canada for 1953–2012. *Atmosphere-Ocean* 53,  
42 383–397. doi:10.1080/07055900.2015.1086970.
- 43 Menezes-Silva, P. E., Loram-Lourenço, L., Alves, R. D. F. B., Sousa, L. F., Almeida, S. E. da S., and Farnese, F. S.  
44 (2019). Different ways to die in a changing world: Consequences of climate change for tree species performance  
45 and survival through an ecophysiological perspective. *Ecol. Evol.* 0. doi:10.1002/ece3.5663.
- 46 Menkes, C. E., Lengaigne, M., Marchesiello, P., Jourdain, N. C., Vincent, E. M., Lefèvre, J., et al. (2012). Comparison  
47 of tropical cyclogenesis indices on seasonal to interannual timescales. *Clim. Dyn.* 38, 301–321.  
48 doi:10.1007/s00382-011-1126-x.
- 49 Meredith, E. P., Semenov, V. A., Maraun, D., Park, W., and Chernokulsky, A. V. (2015). Crucial role of Black Sea  
50 warming in amplifying the 2012 Krymsk precipitation extreme. *Nat. Geosci.* 8, 615–619. doi:10.1038/ngeo2483.
- 51 Mernild, S. H., Hanna, E., Yde, J. C., Cappelen, J., and Malmros, J. K. (2014). Coastal Greenland air temperature  
52 extremes and trends 1890–2010: annual and monthly analysis. *Int. J. Climatol.* 34. doi:10.1002/joc.3777.
- 53 Meseguer-Ruiz, O., Ponce-Philimon, P. I., Quispe-Jofré, A. S., Guijarro, J. A., and Sarricolea, P. (2018). Spatial  
54 behaviour of daily observed extreme temperatures in Northern Chile (1966–2015): data quality, warming trends,  
55 and its orographic and latitudinal effects. *Stoch. Environ. Res. Risk Assess.* 32, 3503–3523. doi:10.1007/s00477-  
56 018-1557-6.
- 57 MfE (2018). Climate Change Projections for New Zealand: Atmosphere Projections Based on Simulations from the  
58 IPCC Fifth Assessment, 2nd Edition. Wellington, NZ: New Zealand Ministry for the Environment (MfE)  
59 Available at: [http://www.mfe.govt.nz/sites/default/files/media/Climate Change/climate-projections-snapshot.pdf](http://www.mfe.govt.nz/sites/default/files/media/Climate%20Change/climate-projections-snapshot.pdf).
- 60 MfE, and Stats NZ (2020). New Zealand’s Environmental Reporting Series: Our atmosphere and climate 2020. New  
61 Zealand Ministry for the Environment (MfE) and Stats NZ Available at:

- 1 <https://www.mfe.govt.nz/publications/environmental-reporting/our-atmosphere-and-climate-2020>.
- 2 Michaelis, A. C., Willison, J., Lackmann, G. M., and Robinson, W. A. (2017). Changes in winter North Atlantic  
3 extratropical cyclones in high-resolution regional pseudo-global warming simulations. *J. Clim.* 30, 6905–6925.  
4 doi:10.1175/JCLI-D-16-0697.1.
- 5 Miglietta, M. M., and Rotunno, R. (2019). Development mechanisms for Mediterranean tropical-like cyclones  
6 (medicanes). *Q. J. R. Meteorol. Soc.* 145, 1444–1460. doi:10.1002/qj.3503.
- 7 Milly, P. C. D., and Dunne, K. A. (2016). Potential evapotranspiration and continental drying. *Nat. Clim. Chang.* 6,  
8 946–949. doi:10.1038/nclimate3046.
- 9 Milly, P. C. D., and Dunne, K. A. (2020). Colorado River flow dwindles as warming-driven loss of reflective snow  
10 energizes evaporation. *Science (80-. )* 367, 1252–1255. doi:10.1126/science.aax0194.
- 11 Min, S., Kim, Y.-H., Lee, S.-M., Sparrow, S., Li, S., Lott, F. C., et al. (2020). Quantifying human impact on the 2018  
12 summer longest heat wave in south korea. *Bull Am Meteorol Soc.* doi:10.1175/BAMS-D-19-0151.1.
- 13 Miralles, D. G., Gentine, P., Seneviratne, S. I., and Teuling, A. J. (2019). Land-atmospheric feedbacks during droughts  
14 and heatwaves: state of the science and current challenges. *Ann. N. Y. Acad. Sci.* 1436, 19–35.  
15 doi:10.1111/nyas.13912.
- 16 Miralles, D. G., Teuling, A. J., Van Heerwaarden, C. C., and De Arellano, J. V. G. (2014a). Mega-heatwave  
17 temperatures due to combined soil desiccation and atmospheric heat accumulation. *Nat. Geosci.* 7, 345–349.  
18 doi:10.1038/ngeo2141.
- 19 Miralles, D. G., van den Berg, M. J., Gash, J. H., Parinussa, R. M., de Jeu, R. A. M., Beck, H. E., et al. (2014b). El  
20 Niño–La Niña cycle and recent trends in continental evaporation. *Nat. Clim. Chang.* 4, 122–126.  
21 doi:10.1038/nclimate2068.
- 22 Mishra, V. (2020). Long-term (1870–2018) drought reconstruction in context of surface water security in India. *J.*  
23 *Hydrol.* 580. doi:10.1016/j.jhydrol.2019.124228.
- 24 Mishra, V., Kumar, D., Ganguly, A. R., Sanjay, J., Mujumdar, M., Krishnan, R., et al. (2014a). Reliability of regional  
25 and global climate models to simulate precipitation extremes over India. *J. Geophys. Res. Atmos.* 119, 9301–  
26 9323. doi:10.1002/2014JD021636.
- 27 Mishra, V., Mukherjee, S., Kumar, R., and Stone, D. A. (2017). Heat wave exposure in India in current, 1.5 °C, and  
28 2.0 °C worlds. *Environ. Res. Lett.* 12, 124012. doi:10.1088/1748-9326/aa9388.
- 29 Mishra, V., Shah, R., and Thrasher, B. (2014b). Soil Moisture Droughts under the Retrospective and Projected Climate  
30 in India. *J. Hydrometeorol.* 15, 2267–2292. doi:10.1175/JHM-D-13-0177.1.
- 31 Mitchell, P. J., O’Grady, A. P., Hayes, K. R., and Pinkard, E. A. (2014). Exposure of trees to drought-induced die-off is  
32 defined by a common climatic threshold across different vegetation types. *Ecol. Evol.* 4, 1088–1101.  
33 doi:10.1002/ece3.1008.
- 34 Mitchell, T. D. (2003). Pattern Scaling. An Examination of the Accuracy of the Technique for Describing Future  
35 Climates. *Clim. Change* 60, 217–242. doi:10.1023/A:1026035305597.
- 36 Mizuta, R., and Endo, H. (2020). Projected Changes in Extreme Precipitation in a 60-km AGCM Large Ensemble and  
37 Their Dependence on Return Periods. *Geophys. Res. Lett.* 47, 1–8. doi:10.1029/2019GL086855.
- 38 Mo, K. C., and Lettenmaier, D. P. (2018). Drought variability and trends over the central United States in the  
39 instrumental record. *J. Hydrometeorol.* 19, 1149–1166. doi:10.1175/JHM-D-17-0225.1.
- 40 Moftakhari, H. R., Salvadori, G., AghaKouchak, A., Sanders, B. F., and Matthew, R. A. (2017). Compounding effects  
41 of sea level rise and fluvial flooding. *Proc. Natl. Acad. Sci.* 114, 9785–9790. doi:10.1073/pnas.1620325114.
- 42 Mohammed, R., and Scholz, M. (2016). Impact of climate variability and streamflow alteration on groundwater  
43 contribution to the base flow of the Lower Zab River (Iran and Iraq). *Environ. Earth Sci.* 75. doi:10.1007/s12665-  
44 016-6205-1.
- 45 Molina, M. O., Sánchez, E., and Gutiérrez, C. (2020). Future heat waves over the Mediterranean from an Euro-  
46 CORDEX regional climate model ensemble. *Sci. Rep.* 10, 8801. doi:10.1038/s41598-020-65663-0.
- 47 Molnar, P., Fatichi, S., Gaál, L., Szolgay, J., and Burlando, P. (2015). Storm type effects on super Clausius-Clapeyron  
48 scaling of intense rainstorm properties with air temperature. *Hydrol. Earth Syst. Sci.* 19, 1753–1766.  
49 doi:10.5194/hess-19-1753-2015.
- 50 Monjo, R., Gaitán, E., Pórtoles, J., Ribalaygua, J., and Torres, L. (2016). Changes in extreme precipitation over Spain  
51 using statistical downscaling of CMIP5 projections. *Int. J. Climatol.* 36, 757–769. doi:10.1002/joc.4380.
- 52 Montero-Martínez, M. J., Santana-Sepúlveda, J. S., Pérez-Ortiz, N. I., Pita-Díaz, Ó., and Castillo-Liñan, S. (2018).  
53 Comparing climate change indices between a northern (arid) and a southern (humid) basin in Mexico during the  
54 last decades. *Adv. Sci. Res.* 15, 231–237. doi:10.5194/asr-15-231-2018.
- 55 Moon, H., Gudmundsson, L., and Seneviratne, S. I. (2018). Drought Persistence Errors in Global Climate Models. *J.*  
56 *Geophys. Res. Atmos.* 123, 3483–3496. doi:10.1002/2017JD027577.
- 57 Moon, H., Guillod, B. P., Gudmundsson, L., and Seneviratne, S. I. (2019). Soil Moisture Effects on Afternoon  
58 Precipitation Occurrence in Current Climate Models. *Geophys. Res. Lett.* 46, 1861–1869.  
59 doi:10.1029/2018GL080879.
- 60 Moon, I.-J., Kim, S.-H., Klotzbach, P., and Chan, J. C. L. (2015). Roles of interbasin frequency changes in the poleward  
61 shifts of the maximum intensity location of tropical cyclones. *Environ. Res. Lett.* 10, 104004. doi:10.1088/1748-

- 1 9326/10/10/104004.
- 2 Moon, I.-J., Kim, S.-H., Klotzbach, P., and Chan, J. C. L. (2016). Reply to Comment on ‘Roles of interbasin frequency
- 3 changes in the poleward shifts of maximum intensity location of tropical cyclones.’ *Environ. Res. Lett.* 11,
- 4 068002. doi:10.1088/1748-9326/11/6/068002.
- 5 Moon, Y., Kim, D., Camargo, S. J., Wing, A. A., Sobel, A. H., Murakami, H., et al. (2020). Azimuthally Averaged
- 6 Wind and Thermodynamic Structures of Tropical Cyclones in Global Climate Models and Their Sensitivity to
- 7 Horizontal Resolution. *J. Clim.* 33, 1575–1595. doi:10.1175/JCLI-D-19-0172.1.
- 8 Moore, G. W. K. (2016). The December 2015 North Pole Warming Event and the Increasing Occurrence of Such
- 9 Events. *Sci. Rep.* 6. doi:10.1038/srep39084.
- 10 Mora, C., Spirandelli, D., Franklin, E. C., Lynham, J., Kantar, M. B., Miles, W., et al. (2018). Broad threat to humanity
- 11 from cumulative climate hazards intensified by greenhouse gas emissions. *Nat. Clim. Chang.* 8, 1062–1071.
- 12 doi:10.1038/s41558-018-0315-6.
- 13 Moravec, V., Markonis, Y., Rakovec, O., Kumar, R., and Hanel, M. (2019). A 250-Year European Drought Inventory
- 14 Derived From Ensemble Hydrologic Modeling. *Geophys. Res. Lett.* 46, 5909–5917. doi:10.1029/2019GL082783.
- 15 Morgan, J. A., LeCain, D. R., Pendall, E., Blumenthal, D. M., Kimball, B. A., Carrillo, Y., et al. (2011). C4 grasses
- 16 prosper as carbon dioxide eliminates desiccation in warmed semi-arid grassland. *Nature* 476, 202–205.
- 17 doi:10.1038/nature10274.
- 18 Mori, N., Ariyoshi, N., Shimura, T., Miyashita, T., and Ninomiya, J. (2021). Future projection of maximum potential
- 19 storm surge height at three major bays in Japan using the maximum potential intensity of a tropical cyclone. *Clim.*
- 20 *Change* 164, 25. doi:10.1007/s10584-021-02980-x.
- 21 Mori, N., Shimura, T., Yoshida, K., Mizuta, R., Okada, Y., Fujita, M., et al. (2019). Future changes in extreme storm
- 22 surges based on mega-ensemble projection using 60-km resolution atmospheric global circulation model. *Coast.*
- 23 *Eng. J.* 61, 295–307. doi:10.1080/21664250.2019.1586290.
- 24 Moron, V., Oueslati, B., Pohl, B., Rome, S., and Janicot, S. (2016). Trends of mean temperatures and warm extremes in
- 25 northern tropical Africa (1961–2014) from observed and PPCA-reconstructed time series. *J. Geophys. Res. Atmos.*
- 26 121, 5298–5319. doi:10.1002/2015JD024303.
- 27 Morrison, A., Villarini, G., Zhang, W., and Scoccimarro, E. (2019). Projected changes in extreme precipitation at sub-
- 28 daily and daily time scales. *Glob. Planet. Change* 182, 103004. doi:10.1016/j.gloplacha.2019.103004.
- 29 Mostafa, A. N., Wheida, A., El Nazer, M., Adel, M., El Leithy, L., Siour, G., et al. (2019). Past (1950–2017) and future
- 30 (–2100) temperature and precipitation trends in Egypt. *Weather Clim. Extrem.* 26, 100225.
- 31 doi:10.1016/j.wace.2019.100225.
- 32 Mote, P. W., Rupp, D. E., Li, S., Sharp, D. J., Otto, F., Uhe, P. F., et al. (2016). Perspectives on the causes of
- 33 exceptionally low 2015 snowpack in the western United States. *Geophys. Res. Lett.* 43, 10,910-980,988.
- 34 doi:10.1002/2016GL069965.
- 35 Mouhamed, L., Traore, S. B., Alhassane, A., and Sarr, B. (2013). Evolution of some observed climate extremes in the
- 36 West African Sahel. *Weather Clim. Extrem.* doi:10.1016/j.wace.2013.07.005.
- 37 Mozny, M., Trnka, M., Vlach, V., Vizina, A., Potopova, V., Zahradnicek, P., et al. (2020). Past (1971–2018) and future
- 38 (2021–2100) pan evaporation rates in the Czech Republic. *J. Hydrol.* 590. doi:10.1016/j.jhydrol.2020.125390.
- 39 Mtongori, H. I., Stordal, F., and Benestad, R. E. (2016). Evaluation of Empirical Statistical Downscaling Models’ Skill
- 40 in Predicting Tanzanian Rainfall and Their Application in Providing Future Downscaled Scenarios. *J. Clim.* 29,
- 41 3231–3252. doi:10.1175/JCLI-D-15-0061.1.
- 42 Mueller, B., and Seneviratne, S. (2014). Systematic land climate and evapotranspiration biases in CMIP5 simulations.
- 43 *Geophys. Res. Lett.* 41, 128–134. doi:10.1002/2013GL058055.
- 44 Mueller, B., and Seneviratne, S. I. (2012). Hot days induced by precipitation deficits at the global scale. *Proc. Natl.*
- 45 *Acad. Sci.* 109, 12398–12403. doi:10.1073/pnas.1204330109.
- 46 Mueller, B., and Zhang, X. (2016). Causes of drying trends in northern hemispheric land areas in reconstructed soil
- 47 moisture data. *Clim. Change* 134, 255–267. doi:10.1007/s10584-015-1499-7.
- 48 Mueller, B., Zhang, X., and Zwiers, F. W. (2016a). Historically hottest summers projected to be the norm for more than
- 49 half of the world’s population within 20 years. *Environ. Res. Lett.* 11, 1–15. doi:10.1088/1748-9326/11/4/044011.
- 50 Mueller, N. D., Butler, E. E., McKinnon, K. A., Rhines, A., Tingley, M., Holbrook, N. M., et al. (2016b). Cooling of
- 51 US Midwest summer temperature extremes from cropland intensification. *Nat. Clim. Chang.* 6, 317–322.
- 52 doi:10.1038/nclimate2825.
- 53 Muerth, M. J., Gauvin St-Denis, B., Ricard, S., Velázquez, J. A., Schmid, J., Minville, M., et al. (2013). On the need for
- 54 bias correction in regional climate scenarios to assess climate change impacts on river runoff. *Hydrol. Earth Syst.*
- 55 *Sci.* 17, 1189–1204. doi:10.5194/hess-17-1189-2013.
- 56 Mukherjee, S., Aadhar, S., Stone, D., and Mishra, V. (2018a). Increase in extreme precipitation events under
- 57 anthropogenic warming in India. *Weather Clim. Extrem.* 20, 45–53. doi:10.1016/J.WACE.2018.03.005.
- 58 Mukherjee, S., Mishra, A., and Trenberth, K. E. (2018b). Climate Change and Drought: a Perspective on Drought
- 59 Indices. *Curr. Clim. Chang. Reports* 4, 145–163. doi:10.1007/s40641-018-0098-x.
- 60 Muller, J., Collins, J. M., Gibson, S., and Paxton, L. (2017). “Recent Advances in the Emerging Field of
- 61 Paleotempestology,” in *Hurricanes and Climate Change: Volume 3*, eds. J. M. Collins and K. Walsh (Cham,

- 1 Switzerland: Springer), 1–33. doi:10.1007/978-3-319-47594-3\_1.
- 2 Müller, W. A., Borchert, L., and Ghosh, R. (2020). Observed Subdecadal Variations of European Summer  
3 Temperatures. *Geophys. Res. Lett.* 47, e2019GL086043. doi:10.1029/2019GL086043.
- 4 Murakami, H., Delworth, T. L., Cooke, W. F., Zhao, M., Xiang, B., and Hsu, P.-C. (2020). Detected climatic change in  
5 global distribution of tropical cyclones. *Proc. Natl. Acad. Sci.* 117, 10706–10714. doi:10.1073/pnas.1922500117.
- 6 Murakami, H., Levin, E., Delworth, T. L., Gudgel, R., and Hsu, P.-C. (2018). Dominant effect of relative tropical  
7 Atlantic warming on major hurricane occurrence. *Science (80-. )*. 362, 794–799. doi:10.1126/science.aat6711.
- 8 Murakami, H., Li, T., and Hsu, P. C. (2014). Contributing factors to the recent high level of accumulated cyclone  
9 energy (ACE) and power dissipation index (PDI) in the North Atlantic. *J. Clim.* 27, 3023–3034.  
10 doi:10.1175/JCLI-D-13-00394.1.
- 11 Murakami, H., Vecchi, G. A., Delworth, T. L., Wittenberg, A. T., Underwood, S., Gudgel, R., et al. (2017a). Dominant  
12 Role of Subtropical Pacific Warming in Extreme Eastern Pacific Hurricane Seasons: 2015 and the Future. *J. Clim.*  
13 30, 243–264. doi:10.1175/JCLI-D-16-0424.1.
- 14 Murakami, H., Vecchi, G. A., and Underwood, S. (2017b). Increasing frequency of extremely severe cyclonic storms  
15 over the Arabian Sea. *Nat. Clim. Chang.* 7, 885–889. doi:10.1038/s41558-017-0008-6.
- 16 Murakami, H., Vecchi, G. A., Underwood, S., Delworth, T. L., Wittenberg, A. T., Anderson, W. G., et al. (2015).  
17 Simulation and prediction of category 4 and 5 hurricanes in the high-resolution GFDL HiFLOR coupled climate  
18 model. *J. Clim.* 28, 9058–9079. doi:10.1175/JCLI-D-15-0216.1.
- 19 Murakami, H., Wang, Y., Yoshimura, H., Mizuta, R., Sugi, M., Shindo, E., et al. (2012). Future changes in tropical  
20 cyclone activity projected by the new high-resolution MRI-AGCM. *J. Clim.* 25, 3237–3260. doi:10.1175/JCLI-D-  
21 11-00415.1.
- 22 Muramatsu, T., Kato, T., Nakazato, M., Endo, H., and Kitoh, A. (2016). Future Change of Tornadogenesis-Favorable  
23 Environmental Conditions in Japan Estimated by a 20-km-Mesh Atmospheric General Circulation Model. *J.*  
24 *Meteorol. Soc. Japan. Ser. II* 94A, 105–120. doi:10.2151/jmsj.2015-053.
- 25 Murari, K. K., Ghosh, S., Patwardhan, A., Daly, E., and Salvi, K. (2015). Intensification of future severe heat waves in  
26 India and their effect on heat stress and mortality. *Reg. Environ. Chang.* 15. doi:10.1007/s10113-014-0660-6.
- 27 Murata, A., Sasaki, H., Kawase, H., and Nosaka, M. (2017). Evaluation of precipitation over an oceanic region of Japan  
28 in convection-permitting regional climate model simulations. *Clim. Dyn.* 48, 1779–1792. doi:10.1007/s00382-  
29 016-3172-x.
- 30 Murata, A., Sasaki, H., Kawase, H., Nosaka, M., Oh'izumi, M., Kato, T., et al. (2015). Projection of Future Climate  
31 Change over Japan in Ensemble Simulations with a High-Resolution Regional Climate Model. *SOLA* 11, 90–94.  
32 doi:10.2151/sola.2015-022.
- 33 Myhre, G., Alterskjær, K., Stjern, C. W., Hodnebrog, Ø., Marelle, L., Samset, B. H., et al. (2019). Frequency of  
34 extreme precipitation increases extensively with event rareness under global warming. *Sci. Rep.* 9, 16063.  
35 doi:10.1038/s41598-019-52277-4.
- 36 Nabat, P., Somot, S., Mallet, M., Sanchez-Lorenzo, A., and Wild, M. (2014). Contribution of anthropogenic sulfate  
37 aerosols to the changing Euro-Mediterranean climate since 1980. *Geophys. Res. Lett.* 41, 5605–5611.  
38 doi:10.1002/2014GL060798.
- 39 Nackley, L. L., Betzelberger, A., Skowno, A., G. West, A., Ripley, B. S., Bond, W. J., et al. (2018). CO2 enrichment  
40 does not entirely ameliorate Vachellia karroo drought inhibition: A missing mechanism explaining savanna bush  
41 encroachment. *Environ. Exp. Bot.* 155, 98–106. doi:10.1016/j.envexpbot.2018.06.018.
- 42 Naik, M., and Abiodun, B. J. (2020). Projected changes in drought characteristics over the Western Cape, South Africa.  
43 *Meteorol. Appl.* 27. doi:10.1002/met.1802.
- 44 Nakaegawa, T., Kitoh, A., Murakami, H., and Kusunoki, S. (2013). Annual maximum 5-day rainfall total and  
45 maximum number of consecutive dry days over Central America and the Caribbean in the late twenty-first  
46 century projected by an atmospheric general circulation model with three different horizontal resolutions. *Theor.*  
47 *Appl. Climatol.* 116. doi:10.1007/s00704-013-0934-9.
- 48 Nakamura, J., Camargo, S. J., Sobel, A. H., Henderson, N., Emanuel, K. A., Kumar, A., et al. (2017). Western North  
49 Pacific Tropical Cyclone Model Tracks in Present and Future Climates. *J. Geophys. Res. Atmos.* 122, 9721–9744.  
50 doi:10.1002/2017JD027007.
- 51 Nakano, M., Wada, A., Sawada, M., Yoshimura, H., Onishi, R., Kawahara, S., et al. (2017). Global 7km mesh  
52 nonhydrostatic Model Intercomparison Project for improving Typhoon forecast (TYMIP-G7): Experimental  
53 design and preliminary results. *Geosci. Model Dev.* 10, 1368–1381. doi:10.5194/gmd-10-1363-2017.
- 54 Nakayama, T., and Shankman, D. (2013). Impact of the Three-Gorges Dam and water transfer project on Changjiang  
55 floods. *Glob. Planet. Change* 100, 38–50. doi:10.1016/j.gloplacha.2012.10.004.
- 56 Nangombe, S., Zho, T., Zhang, L., and Zhang, W. (2020). Attribution of the 2018 october-december drought over south  
57 southern africa. *Bull. Am. Meteorol. Soc.* 101, S135–S140. doi:10.1175/BAMS-D-19-0179.1.
- 58 Nangombe, S., Zhou, T., Zhang, W., Wu, B., Hu, S., Zou, L., et al. (2018). Record-breaking climate extremes in Africa  
59 under stabilized 1.5 °C and 2 °C global warming scenarios. *Nat. Clim. Chang.* 8, 375–380. doi:10.1038/s41558-  
60 018-0145-6.
- 61 NASEM (2016). Attribution of Extreme Weather Events in the Context of Climate Change. Washington, DC, USA:

- 1 National Academies of Sciences Engineering and Medicine (NASEM). The National Academies Press  
2 doi:10.17226/21852.
- 3 Nasim, W., Amin, A., Fahad, S., Awais, M., Khan, N., Mubeen, M., et al. (2018). Future risk assessment by estimating  
4 historical heat wave trends with projected heat accumulation using SimCLIM climate model in Pakistan. *Atmos.*  
5 *Res.* 205, 118–133. doi:10.1016/j.atmosres.2018.01.009.
- 6 Nasrollahi, N., Aghakouchak, A., Cheng, L., Damberg, L., Phillips, T. J., Miao, C., et al. (2015). How well do CMIP5  
7 climate simulations replicate historical trends and patterns of meteorological droughts? *Water Resour. Res.* 51,  
8 2847–2864. doi:10.1002/2014WR016318.
- 9 Nastos, P. T., and Kapsomenakis, J. (2015). Regional climate model simulations of extreme air temperature in Greece.  
10 Abnormal or common records in the future climate? *Atmos. Res.* 152, 43–60. doi:10.1016/j.atmosres.2014.02.005.
- 11 Naumann, G., Alfieri, L., Wyser, K., Mentaschi, L., Betts, R. A., Carrao, H., et al. (2018). Global Changes in Drought  
12 Conditions Under Different Levels of Warming. *Geophys. Res. Lett.* 45, 3285–3296. doi:10.1002/2017GL076521.
- 13 Naveendrakumar, G., Vithanage, M., Kwon, H.-H., Chandrasekara, S. S. K., Iqbal, M. C. M., Pathmarajah, S., et al.  
14 (2019). South Asian perspective on temperature and rainfall extremes: A review. *Atmos. Res.* 225, 110–120.  
15 doi:10.1016/j.atmosres.2019.03.021.
- 16 Nayak, S., and Dairaku, K. (2016). Future changes in extreme precipitation intensities associated with temperature  
17 under SRES A1B scenario. *Hydrol. Res. Lett.* 10, 139–144. doi:10.3178/hr.l.10.139.
- 18 Nayak, S., Dairaku, K., Takayabu, I., Suzuki-Parker, A., and Ishizaki, N. N. (2017). Extreme precipitation linked to  
19 temperature over Japan: current evaluation and projected changes with multi-model ensemble downscaling. *Clim.*  
20 *Dyn.* 51, 1–17. doi:10.1007/s00382-017-3866-8.
- 21 Nazrul Islam, M., Almazroui, M., Dambul, R., Jones, P. D., and Alamoudi, A. O. (2015). Long-term changes in  
22 seasonal temperature extremes over Saudi Arabia during 1981–2010. *Int. J. Climatol.* 35, 1579–1592.  
23 doi:10.1002/joc.4078.
- 24 Neri, A., Villarini, G., Slater, L. J., and Napolitano, F. (2019). On the statistical attribution of the frequency of flood  
25 events across the U.S. Midwest. *Adv. Water Resour.* 127, 225–236. doi:10.1016/j.advwatres.2019.03.019.
- 26 Neu, U., Akperov, M. G., Bellenbaum, N., Benestad, R., Blender, R., Caballero, R., et al. (2013). Imilast: A community  
27 effort to intercompare extratropical cyclone detection and tracking algorithms. *Bull. Am. Meteorol. Soc.*  
28 doi:10.1175/BAMS-D-11-00154.1.
- 29 Neukom, R., Gergis, J., Karoly, D. J., Wanner, H., Curran, M., Elbert, J., et al. (2014). Inter-hemispheric temperature  
30 variability over the past millennium. *Nat. Clim. Chang.* 4, 362–367. doi:10.1038/nclimate2174.
- 31 Neumann, B., Vafeidis, A. T., Zimmermann, J., and Nicholls, R. J. (2015). Future Coastal Population Growth and  
32 Exposure to Sea-Level Rise and Coastal Flooding – A Global Assessment. *PLoS One* 10, e0118571.  
33 doi:10.1371/journal.pone.0118571.
- 34 Newman, M., Wittenberg, A. T., Cheng, L., Compo, G. P., and Smith, C. A. (2018). The Extreme 2015/16 El Niño, in  
35 the Context of Historical Climate Variability and Change. *Bull. Am. Meteorol. Soc.* 99, S16–S20.  
36 doi:10.1175/BAMS-D-17-0116.1.
- 37 Nguvava, M., Abiodun, B. J., and Otieno, F. (2019). Projecting drought characteristics over East African basins at  
38 specific global warming levels. *Atmos. Res.* 228, 41–54. doi:10.1016/j.atmosres.2019.05.008.
- 39 Nicholls, R. J., Brown, S., Goodwin, P., Wahl, T., Lowe, J., Solan, M., et al. (2018). Stabilization of global temperature  
40 at 1.5°C and 2.0°C: implications for coastal areas. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 376, 20160448.  
41 doi:10.1098/rsta.2016.0448.
- 42 Nicholson, S. E. (2017). Climate and climatic variability of rainfall over eastern Africa. *Rev. Geophys.* 55, 590–635.
- 43 Nicolai-Shaw, N., Hirschi, M., Mittelbach, H., and Seneviratne, S. I. (2015). Spatial representativeness of soil moisture  
44 using in situ, remote sensing, and land reanalysis data. *J. Geophys. Res. Atmos.* 120, 9955–9964.  
45 doi:10.1002/2015JD023305.
- 46 Nie, J., Sobel, A. H., Shaevitz, D. A., and Wang, S. (2018). Dynamic amplification of extreme precipitation sensitivity.  
47 *Proc. Natl. Acad. Sci.* 115, 9467 LP-9472. doi:10.1073/pnas.1800357115.
- 48 Nied, M., Pardowitz, T., Nissen, K., Ulbrich, U., Hundecha, Y., and Merz, B. (2014). On the relationship between  
49 hydro-meteorological patterns and flood types. *J. Hydrol.* 519, 3249–3262. doi:10.1016/j.jhydrol.2014.09.089.
- 50 Nikulin, G., Lennard, C., Dosio, A., Kjellström, E., Chen, Y., Hänsler, A., et al. (2018). The effects of 1.5 and 2 degrees  
51 of global warming on Africa in the CORDEX ensemble. *Environ. Res. Lett.* 13, 065003. doi:10.1088/1748-  
52 9326/aab1b1.
- 53 Niranjana Kumar, K., Rajeevan, M., Pai, D. S., Srivastava, A. K., and Preethi, B. (2013). On the observed variability of  
54 monsoon droughts over India. *Weather Clim. Extrem.* 1, 42–50. doi:10.1016/J.WACE.2013.07.006.
- 55 Nissen, K. M., and Ulbrich, U. (2017). Increasing frequencies and changing characteristics of heavy precipitation events  
56 threatening infrastructure in Europe under climate change. *Nat. Hazards Earth Syst. Sci.* 17, 1177–1190.  
57 doi:10.5194/nhess-17-1177-2017.
- 58 Niu, X., Wang, S., Tang, J., Lee, D.-K., Gutowski, W., Dairaku, K., et al. (2018). Ensemble evaluation and projection  
59 of climate extremes in China using RMIP models. *Int. J. Climatol.* 38, 2039–2055. doi:10.1002/joc.5315.
- 60 Nolan, R. H., Boer, M. M., Collins, L., Resco de Dios, V., Clarke, H., Jenkins, M., et al. (2020). Causes and  
61 consequences of eastern Australia’s 2019–20 season of mega-fires. *Glob. Chang. Biol.* 26, 1039–1041.

- 1 doi:10.1111/gcb.14987.
- 2 Norrant, C., and Douguédroit, A. (2006). Monthly and daily precipitation trends in the Mediterranean (1950-2000).  
3 *Theor. Appl. Climatol.* 83, 89–106. doi:10.1007/s00704-005-0163-y.
- 4 Nott, J., Smithers, S., Walsh, K., and Rhodes, E. (2009). Sand beach ridges record 6000 year history of extreme tropical  
5 cyclone activity in northeastern Australia. *Quat. Sci. Rev.* 28, 1511–1520. doi:10.1016/j.quascirev.2009.02.014.
- 6 O’Gorman, P. (2014). Contrasting responses of mean and extreme snowfall to climate change. *Nature* 512, 416–418.
- 7 OB, C., Yang, S., Boberg, F., C, F. M., Thejll, P., Olesen, M., et al. (2015). Scalability of regional climate change in  
8 Europe for high-end scenarios. *Clim. Res.* 64, 25–38. doi:10.3354/cr01286.
- 9 Odoulami, R. C., Abiodun, B. J., and Ajayi, A. E. (2019). Modelling the potential impacts of afforestation on extreme  
10 precipitation over West Africa. *Clim. Dyn.* 52, 2185–2198. doi:10.1007/s00382-018-4248-6.
- 11 Oey, L.-Y., and Chou, S. C. (2016). Evidence of rising and poleward shift of storm surge in western North Pacific in  
12 recent decades. *J. Geophys. Res. Ocean.* 121, 5181–5192. doi:10.1002/2015JC011516.
- 13 Ogata, T., Mizuta, R., Adachi, Y., Murakami, H., and Ose, T. (2015). Effect of air-sea coupling on the frequency  
14 distribution of intense tropical cyclones over the northwestern Pacific. *Geophys. Res. Lett.* 42, 410–415, 421.  
15 doi:10.1002/2015GL066774.
- 16 Ogata, T., Mizuta, R., Adachi, Y., Murakami, H., and Ose, T. (2016). Atmosphere-Ocean Coupling Effect on Intense  
17 Tropical Cyclone Distribution and its Future Change with 60 km-AOGCM. *Sci. Rep.* 6, 29800.  
18 doi:10.1038/srep29800.
- 19 Oguntunde, P. G., Abiodun, B. J., Lischeid, G., and Abatan, A. A. (2020). Droughts projection over the Niger and Volta  
20 River basins of West Africa at specific global warming levels. *Int. J. Climatol.* 40, 5688–5699.  
21 doi:10.1002/joc.6544.
- 22 Oh, H., Ha, K.-J., and Timmermann, A. (2018). Disentangling Impacts of Dynamic and Thermodynamic Components  
23 on Late Summer Rainfall Anomalies in East Asia. *J. Geophys. Res. Atmos.* 123, 8623–8633.  
24 doi:10.1029/2018JD028652.
- 25 Ohba, M., and Sugimoto, S. (2019). Differences in climate change impacts between weather patterns: possible effects  
26 on spatial heterogeneous changes in future extreme rainfall. *Clim. Dyn.* 52, 4177–4191. doi:10.1007/s00382-018-  
27 4374-1.
- 28 Ohba, M., and Sugimoto, S. (2020). Impacts of climate change on heavy wet snowfall in Japan. *Clim. Dyn.* 54, 3151–  
29 3164. doi:10.1007/s00382-020-05163-z.
- 30 Oizumi, T., Saito, K., Ito, J., Kuroda, T., Doc, L., DUC, L., et al. (2018). Ultra-High-Resolution Numerical Weather  
31 Prediction with a Large Domain Using the K Computer: A Case Study of the Izu Oshima Heavy Rainfall Event  
32 on October 15-16, 2013. *J. Meteorol. Soc. Japan. Ser. II* 96, 25–54. doi:10.2151/jmsj.2018-006.
- 33 Olmo, M., Bettolli, M. L., and Rusticucci, M. (2020). Atmospheric circulation influence on temperature and  
34 precipitation individual and compound daily extreme events: Spatial variability and trends over southern South  
35 America. *Weather Clim. Extrem.* 29, 100267. doi:10.1016/j.wace.2020.100267.
- 36 Olsson, J., and Foster, K. (2013). Short-term precipitation extremes in regional climate simulations for Sweden. *Hydrol.*  
37 *Res.* 45, 479–489. doi:10.2166/nh.2013.206.
- 38 Omondi, P. A., Awange, J. L., Forootan, E., Ogallo, L. A., Barakiza, R., Girmaw, G. B., et al. (2014). Changes in  
39 temperature and precipitation extremes over the Greater Horn of Africa region from 1961 to 2010. *Int. J.*  
40 *Climatol.* 34, 1262–1277. doi:10.1002/joc.3763.
- 41 Ongoma, V., Chen, H., and Gao, C. (2018a). Projected changes in mean rainfall and temperature over East Africa based  
42 on CMIP5 models. *Int. J. Climatol.* 38, 1375–1392. doi:10.1002/joc.5252.
- 43 Ongoma, V., Chen, H., Gao, C., Nyongesa, A. M., and Polong, F. (2018b). Future changes in climate extremes over  
44 Equatorial East Africa based on CMIP5 multimodel ensemble. *Nat. Hazards* 90, 901–920. doi:10.1007/s11069-  
45 017-3079-9.
- 46 Orlowsky, B., and Seneviratne, S. I. (2013). Elusive drought: Uncertainty in observed trends and short-and long-term  
47 CMIP5 projections. *Hydrol. Earth Syst. Sci.* 17, 1765–1781. doi:10.5194/hess-17-1765-2013.
- 48 Ortega, J. A., Razola, L., and Garzón, G. (2014). Recent human impacts and change in dynamics and morphology of  
49 ephemeral rivers. *Nat. Hazards Earth Syst. Sci.* 14, 713–730. doi:10.5194/nhess-14-713-2014.
- 50 Orth, R., Vogel, M. M., Luterbacher, J., Pfister, C., and Seneviratne, S. I. (2016a). Did European temperatures in 1540  
51 exceed present-day records? *Environ. Res. Lett.* 11, 114021. doi:10.1088/1748-9326/11/11/114021.
- 52 Orth, R., Zscheischler, J., and Seneviratne, S. I. (2016b). Record dry summer in 2015 challenges precipitation  
53 projections in Central Europe. *Sci. Rep.* 6, 28334. doi:10.1038/srep28334.
- 54 Osima, S., Indasi, V. S., Zaroug, M., Endris, H. S., Gudoshava, M., Misiani, H. O., et al. (2018). Projected climate over  
55 the Greater Horn of Africa under 1.5 °C and 2 °C global warming. *Environ. Res. Lett.* 13, 065004.  
56 doi:10.1088/1748-9326/aaba1b.
- 57 Otkin, J. A., Anderson, M. C., Hain, C., Svoboda, M., Johnson, D., Mueller, R., et al. (2016). Assessing the evolution of  
58 soil moisture and vegetation conditions during the 2012 United States flash drought. *Agric. For. Meteorol.* 218–  
59 219, 230–242. doi:10.1016/j.agrformet.2015.12.065.
- 60 Otkin, J. A., Svoboda, M., Hunt, E. D., Ford, T. W., Anderson, M. C., Hain, C., et al. (2018). Flash droughts: A review  
61 and assessment of the challenges imposed by rapid-onset droughts in the United States. *Bull. Am. Meteorol. Soc.*

- 1 99, 911–919. doi:10.1175/BAMS-D-17-0149.1.
- 2 Otto, F. E. L. (2017). Attribution of Weather and Climate Events. *Annu. Rev. Environ. Resour.* 42, null.
- 3 doi:10.1146/annurev-environ-102016-060847.
- 4 Otto, F. E. L., Boyd, E., Jones, R. G., Cornforth, R. J., James, R., Parker, H. R., et al. (2015a). Attribution of extreme
- 5 weather events in Africa: a preliminary exploration of the science and policy implications. *Clim. Change* 132,
- 6 531–543. doi:10.1007/s10584-015-1432-0.
- 7 Otto, F. E. L., Harrington, L., Schmitt, K., Philip, S., Kew, S., van Oldenborgh, G. J., et al. (2020). Challenges to
- 8 Understanding Extreme Weather Changes in Lower Income Countries. *Bull. Am. Meteorol. Soc.* 101, E1851–
- 9 E1860. doi:10.1175/BAMS-D-19-0317.1.
- 10 Otto, F. E. L., Haustein, K., Uhe, P., Coelho, C. A. S., Aravequia, J. A., Almeida, W., et al. (2015b). Factors Other
- 11 Than Climate Change, Main Drivers of 2014/15 Water Shortage in Southeast Brazil. *Bull. Am. Meteorol. Soc.* 96,
- 12 S35–S40. doi:10.1175/BAMS-D-15-00120.1.
- 13 Otto, F. E. L., Jones, R. G., Halladay, K., and Allen, M. R. (2013). Attribution of changes in precipitation patterns in
- 14 African rainforests. *Philos. Trans. R. Soc. B Biol. Sci.* 368, 20120299. doi:10.1098/rstb.2012.0299.
- 15 Otto, F. E. L., Massey, N., van Oldenborgh, G. J., Jones, R. G., and Allen, M. R. (2012). Reconciling two approaches to
- 16 attribution of the 2010 Russian heat wave. *Geophys. Res. Lett.* 39, n/a-n/a. doi:10.1029/2011GL050422.
- 17 Otto, F. E. L., Philip, S., Kew, S., Li, S., King, A., and Cullen, H. (2018a). Attributing high-impact extreme events
- 18 across timescales – a case study of four different types of events. *Clim. Change* 149, 399–412.
- 19 doi:10.1007/s10584-018-2258-3.
- 20 Otto, F. E. L., Rosier, S. M., Allen, M. R., Massey, N. R., Rye, C. J., and Quintana, J. I. (2015c). Attribution analysis of
- 21 high precipitation events in summer in England and Wales over the last decade. *Clim. Change* 132, 77–91.
- 22 doi:10.1007/s10584-014-1095-2.
- 23 Otto, F. E. L., van der Wiel, K., van Oldenborgh, G. J., Philip, S., Kew, S. F., Uhe, P., et al. (2018b). Climate change
- 24 increases the probability of heavy rains in Northern England/Southern Scotland like those of storm Desmond—a
- 25 real-time event attribution revisited. *Environ. Res. Lett.* 13, 024006. doi:10.1088/1748-9326/aa9663.
- 26 Otto, F. E. L., van Oldenborgh, G. J., Eden, J., Stott, P. A., Karoly, D. J., and Allen, M. R. (2016). The attribution
- 27 question. *Nat. Clim. Chang.* 6, 813–816. doi:10.1038/nclimate3089.
- 28 Otto, F. E. L., Wolski, P., Lehner, F., Tebaldi, C., van Oldenborgh, G. J., Hogesteeger, S., et al. (2018c). Anthropogenic
- 29 influence on the drivers of the Western Cape drought 2015–2017. *Environ. Res. Lett.* 13, 124010.
- 30 doi:10.1088/1748-9326/aae9f9.
- 31 Ozturk, T., Ceber, Z. P., Türkeş, M., and Kurnaz, M. L. (2015). Projections of climate change in the Mediterranean
- 32 Basin by using downscaled global climate model outputs. *Int. J. Climatol.* 35, 4276–4292. doi:10.1002/joc.4285.
- 33 Paciorek, C. J., Stone, D. A., and Wehner, M. F. (2018). Quantifying statistical uncertainty in the attribution of human
- 34 influence on severe weather. *Weather Clim. Extrem.* 20, 69–80. doi:10.1016/j.wace.2018.01.002.
- 35 Padrón, R. S., Gudmundsson, L., Decharme, B., Ducharne, A., Lawrence, D. M., Mao, J., et al. (2020). Observed
- 36 changes in dry-season water availability attributed to human-induced climate change. *Nat. Geosci.* 13, 477–481.
- 37 doi:10.1038/s41561-020-0594-1.
- 38 Padrón, R. S., Gudmundsson, L., and Seneviratne, S. I. (2019). Observational Constraints Reduce Likelihood of
- 39 Extreme Changes in Multidecadal Land Water Availability. *Geophys. Res. Lett.* 46, 736–744.
- 40 doi:10.1029/2018GL080521.
- 41 Pai, D. S., Sridhar, L., Badwaik, M. R., and Rajeevan, M. (2015). Analysis of the daily rainfall events over India using a
- 42 new long period (1901–2010) high resolution (0.25° × 0.25°) gridded rainfall data set. *Clim. Dyn.* 45, 755–776.
- 43 doi:10.1007/s00382-014-2307-1.
- 44 Paik, S., and Min, S.-K. (2018). Assessing the Impact of Volcanic Eruptions on Climate Extremes Using CMIP5
- 45 Models. *J. Clim.* 31, 5333–5349. doi:10.1175/JCLI-D-17-0651.1.
- 46 Paik, S., Min, S., Zhang, X., Donat, M. G., King, A. D., and Sun, Q. (2020). Determining the Anthropogenic
- 47 Greenhouse Gas Contribution to the Observed Intensification of Extreme Precipitation. *Geophys. Res. Lett.* 47,
- 48 e2019GL086875. doi:10.1029/2019GL086875.
- 49 Pal, J. S., and Eltahir, E. A. B. (2016). Future temperature in southwest Asia projected to exceed a threshold for human
- 50 adaptability. *Nat. Clim. Chang.* 6, 197–200. doi:10.1038/nclimate2833.
- 51 Palazzi, E., Hardenberg, J., and Provenzale, A. (2013). Precipitation in the Hindu-Kush Karakoram Himalaya:
- 52 Observations and future scenarios. *J. Geophys. Res. Atmos.* 118, 85–100. doi:10.1029/2012JD018697.
- 53 Palipane, E., and Grotjahn, R. (2018). Future Projections of the Large-Scale Meteorology Associated with California
- 54 Heat Waves in CMIP5 Models. *J. Geophys. Res. Atmos.* 123, 8500–8517. doi:10.1029/2018JD029000.
- 55 Pall, P., Patricola, C. M., Wehner, M. F., Stone, D. A., Paciorek, C. J., and Collins, W. D. (2017). Diagnosing
- 56 conditional anthropogenic contributions to heavy Colorado rainfall in September 2013. *Weather Clim. Extrem.*
- 57 17, 1–6. doi:10.1016/j.wace.2017.03.004.
- 58 Pall, P., Tallaksen, L. M., and Stordal, F. (2019). A climatology of rain-on-snow events for Norway. *J. Clim.* 32, 6995–
- 59 7016. doi:10.1175/JCLI-D-18-0529.1.
- 60 Paltan, H., Allen, M., Haustein, K., Fuldaer, L., and Dadson, S. (2018). Global implications of 1.5 °C and 2 °C warmer
- 61 worlds on extreme river flows. *Environ. Res. Lett.* 13, 94003. doi:10.1088/1748-9326/aad985.

- 1 Panisset, J. S., Libonati, R., Gouveia, C. M. P., Machado-Silva, F., França, D. A., França, J. R. A., et al. (2018).  
2 Contrasting patterns of the extreme drought episodes of 2005, 2010 and 2015 in the Amazon Basin. *Int. J.*  
3 *Climatol.* 38, 1096–1104. doi:10.1002/joc.5224.
- 4 Park, C., and Min, S.-K. (2018). Multi-RCM near-term projections of summer climate extremes over East Asia. *Clim.*  
5 *Dyn.* 0, 1–16. doi:10.1007/s00382-018-4425-7.
- 6 Park, C., Min, S.-K., Lee, D., Cha, D.-H., Suh, M.-S., Kang, H.-S., et al. (2016). Evaluation of multiple regional climate  
7 models for summer climate extremes over East Asia. *Clim. Dyn.* 46, 2469–2486. doi:10.1007/s00382-015-2713-z.
- 8 Park Williams, A., Cook, B. I., Smerdon, J. E., Bishop, D. A., Seager, R., and Mankin, J. S. (2017). The 2016  
9 Southeastern U.S. Drought: An Extreme Departure From Centennial Wetting and Cooling. *J. Geophys. Res.*  
10 *Atmos.* 122, 10,810–888,905. doi:10.1002/2017JD027523.
- 11 Parker, H. R., Lott, F. C., Cornforth, R. J., Mitchell, D. M., Sparrow, S., and Wallom, D. (2017). A comparison of  
12 model ensembles for attributing 2012 West African rainfall. *Environ. Res. Lett.* 12, 14019. doi:10.1088/1748-  
13 9326/aa5386.
- 14 Parker, T. J., Berry, G. J., Reeder, M. J., and Nicholls, N. (2014). Modes of climate variability and heat waves in  
15 Victoria, southeastern Australia. *Geophys. Res. Lett.* 41, 6926–6934. doi:10.1002/2014GL061736.
- 16 Pascale, S., Kapnick, S. B., Delworth, T. L., and Cooke, W. F. (2020). Increasing risk of another Cape Town “Day  
17 Zero” drought in the 21st century. *Proc. Natl. Acad. Sci.* 117, 29495 LP-29503. doi:10.1073/pnas.2009144117.
- 18 Pascale, S., Lucarini, V., Feng, X., Porporato, A., and ul Hasson, S. (2016). Projected changes of rainfall seasonality  
19 and dry spells in a high greenhouse gas emissions scenario. *Clim. Dyn.* 46, 1331–1350. doi:10.1007/s00382-015-  
20 2648-4.
- 21 Paschalis, A., Faticchi, S., Molnar, P., Rimkus, S., and Burlando, P. (2014). On the effects of small scale space-time  
22 variability of rainfall on basin flood response. *J. Hydrol.* 514, 313–327. doi:10.1016/j.jhydrol.2014.04.014.
- 23 Patarčić, M., Gajić-Čapka, M., Cindrić, K., and Branković, C. (2014). Recent and near-future changes in  
24 precipitation extreme indices over the Croatian Adriatic coast. *Clim. Res.* 61, 157–176. doi:10.3354/cr01250.
- 25 Patra, P. K., Crisp, D., Kaiser, J. W., Wunch, D., Saeki, T., Ichii, K., et al. (2017). The Orbiting Carbon Observatory  
26 (OCO-2) tracks 2-3 peta-gram increase in carbon release to the atmosphere during the 2014-2016 El Niño. *Sci.*  
27 *Rep.* 7, 1–12. doi:10.1038/s41598-017-13459-0.
- 28 Patricola, C. M., Saravanan, R., and Chang, P. (2018). The Response of Atlantic Tropical Cyclones to Suppression of  
29 African Easterly Waves. *Geophys. Res. Lett.* 45, 471–479. doi:10.1002/2017GL076081.
- 30 Patricola, C. M., and Wehner, M. F. (2018). Anthropogenic influences on major tropical cyclone events. *Nature* 563,  
31 339–346. doi:10.1038/s41586-018-0673-2.
- 32 Paul, S., Ghosh, S., Mathew, M., Devanand, A., Karmakar, S., and Niyogi, D. (2018). Increased Spatial Variability and  
33 Intensification of Extreme Monsoon Rainfall due to Urbanization. *Sci. Rep.* 8, 3918. doi:10.1038/s41598-018-  
34 22322-9.
- 35 Paulo, A., Martins, D., and Pereira, L. S. (2016). Influence of Precipitation Changes on the SPI and Related Drought  
36 Severity. An Analysis Using Long-Term Data Series. *Water Resour. Manag.* 30, 5737–5757.  
37 doi:10.1007/s11269-016-1388-5.
- 38 Paxian, A., Hertig, E., Seubert, S., Vogt, G., Jacobeit, J., and Paeth, H. (2014). Present-day and future mediterranean  
39 precipitation extremes assessed by different statistical approaches. *Clim. Dyn.* 44, 845–860. doi:10.1007/s00382-  
40 014-2428-6.
- 41 Pedron, I. T., Silva Dias, M. A. F., de Paula Dias, S., Carvalho, L. M. V., and Freitas, E. D. (2017). Trends and  
42 variability in extremes of precipitation in Curitiba – Southern Brazil. *Int. J. Climatol.* 37, 1250–1264.  
43 doi:10.1002/joc.4773.
- 44 Peña-Angulo, D., Reig-Gracia, F., Domínguez-Castro, F., Revuelto, J., Aguilar, E., van der Schrier, G., et al. (2020a).  
45 ECTACI: European Climatology and Trend Atlas of Climate Indices (1979–2017). *J. Geophys. Res. Atmos.* 125.  
46 doi:10.1029/2020JD032798.
- 47 Peña-Angulo, D., Vicente-Serrano, S. M., Domínguez-Castro, F., Murphy, C., Reig, F., Trambly, Y., et al. (2020b).  
48 Long-term precipitation in Southwestern Europe reveals no clear trend attributable to anthropogenic forcing.  
49 *Environ. Res. Lett.* 15, 094070. doi:10.1088/1748-9326/ab9c4f.
- 50 Pendergrass, A. G. (2018). What precipitation is extreme? *Science (80-. )*. 360, 1072–1073.  
51 doi:10.1126/science.aat1871.
- 52 Pendergrass, A. G., Coleman, D. B., Deser, C., Lehner, F., Rosenbloom, N., and Simpson, I. R. (2019). Nonlinear  
53 Response of Extreme Precipitation to Warming in CESM1. *Geophys. Res. Lett.* 46, 10551–10560.  
54 doi:10.1029/2019GL084826.
- 55 Pendergrass, A. G., Lehner, F., Sanderson, B. M., and Xu, Y. (2015). Does extreme precipitation intensity depend on  
56 the emissions scenario? *Geophys. Res. Lett.* 42, 8767–8774. doi:10.1002/2015GL065854.
- 57 Pendergrass, A. G., Meehl, G. A., Pulwarty, R., Hobbins, M., Hoell, A., AghaKouchak, A., et al. (2020). Flash droughts  
58 present a new challenge for subseasonal-to-seasonal prediction. *Nat. Clim. Chang.* 10, 191–199.  
59 doi:10.1038/s41558-020-0709-0.
- 60 Pepler, A., Ashcroft, L., and Trewin, B. (2018). The relationship between the subtropical ridge and Australian  
61 temperatures. *J. South. Hemisph. Earth Syst. Sci.* 68, 201–214. doi:10.22499/3.6801.011.



- 1 Pepler, A. S., Di Luca, A., Ji, F., Alexander, L. V., Evans, J. P., and Sherwood, S. C. (2015). Impact of Identification  
2 Method on the Inferred Characteristics and Variability of Australian East Coast Lows. *Mon. Weather Rev.* 143,  
3 864–877. doi:10.1175/MWR-D-14-00188.1.
- 4 Pepler, A. S., Luca, A. Di, Ji, F., Alexander, L. V., Evans, J. P., and Sherwood, S. C. (2016). Projected changes in east  
5 Australian midlatitude cyclones during the 21st century. *Geophys. Res. Lett.* 43, 334–340.  
6 doi:10.1002/2015GL067267.Received.
- 7 Pereira, L. S., Allen, R. G., Smith, M., and Raes, D. (2015). Crop evapotranspiration estimation with FAO56: Past and  
8 future. *Agric. Water Manag.* 147, 4–20. doi:10.1016/j.agwat.2014.07.031.
- 9 Perkins-Kirkpatrick, S. E., and Gibson, P. B. (2018). Changes in regional heatwave characteristics as a function of  
10 increasing global temperature. *Sci. Rep.* 7, 1–12. doi:10.1038/s41598-017-12520-2.
- 11 Perkins-Kirkpatrick, S. E., and Lewis, S. C. (2020). Increasing trends in regional heatwaves. *Nat. Commun.* 11, 1–8.  
12 doi:10.1038/s41467-020-16970-7.
- 13 Perkins-Kirkpatrick, S. E., White, C. J., Alexander, L. V., Argüeso, D., Boschat, G., Cowan, T., et al. (2016). Natural  
14 hazards in Australia: heatwaves. *Clim. Change* 139, 101–114. doi:10.1007/s10584-016-1650-0.
- 15 Perkins, S. E. (2015). A review on the scientific understanding of heatwaves – Their measurement, driving mechanisms,  
16 and changes at the global scale. *Atmos. Res.* 164–165, 242–267. doi:10.1016/j.atmosres.2015.05.014.
- 17 Perkins, S. E., and Alexander, L. V. (2013). On the Measurement of Heat Waves. *J. Clim.* 26, 4500–4517.  
18 doi:10.1175/JCLI-D-12-00383.1.
- 19 Perkins, S. E., and Gibson, P. B. (2015). Increased Risk of the 2014 Australian May Heatwave Due to Anthropogenic  
20 Activity. *Bull. Am. Meteorol. Soc.* 96, S154–S157. doi:10.1175/BAMS-D-15-00074.1.
- 21 Perkins, S. E., Lewis, S. C., King, A. D., and Alexander, L. V. (2014). Increased simulated risk of the hot Australian  
22 summer of 2012–2013 due to anthropogenic activity as measured by heatwave frequency and intensity [in  
23 “Explaining Extreme Events of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S34–S37.  
24 doi:10.1175/1520-0477-95.9.S1.1.
- 25 Persad, G. G., and Caldeira, K. (2018). Divergent global-scale temperature effects from identical aerosols emitted in  
26 different regions. *Nat. Commun.* 9, 3289. doi:10.1038/s41467-018-05838-6.
- 27 Peterson, T. C., Heim, R. R., Hirsch, R., Kaiser, D. P., Brooks, H., Diffenbaugh, N. S., et al. (2013a). Monitoring and  
28 Understanding Changes in Heat Waves, Cold Waves, Floods, and Droughts in the United States: State of  
29 Knowledge. *Bull. Am. Meteorol. Soc.* 94, 821–834. doi:10.1175/BAMS-D-12-00066.1.
- 30 Peterson, T. C., Hoerling, M. P., Stott, P. A., and Herring, S. C. (2013b). Explaining Extreme Events of 2012 from a  
31 Climate Perspective. *Bull. Am. Meteorol. Soc.* 94, S1–S74. doi:10.1175/BAMS-D-13-00085.1.
- 32 Peterson, T. C., Stott, P. A., and Herring, S. (2012). Explaining Extreme Events of 2011 from a Climate Perspective.  
33 *Bull. Am. Meteorol. Soc.* 93, 1041–1067. doi:10.1175/BAMS-D-12-00021.1.
- 34 Pfahl, S., O’Gorman, P. A., and Fischer, E. M. (2017). Understanding the regional pattern of projected future changes  
35 in extreme precipitation. *Nat. Clim. Chang.* 7, 423. doi:10.1038/nclimate3287.
- 36 Pfahl, S., and Wernli, H. (2012). Quantifying the relevance of cyclones for precipitation extremes. *J. Clim.* 25, 6770–  
37 6780. doi:10.1175/JCLI-D-11-00705.1.
- 38 Phelan, P. E., Kaloush, K., Miner, M., Golden, J., Phelan, B., Silva, H., et al. (2015). Urban Heat Island: Mechanisms,  
39 Implications, and Possible Remedies. *Annu. Rev. Environ. Resour.* 40, 285–307. doi:10.1146/annurev-environ-  
40 102014-021155.
- 41 Phibbs, S., and Toumi, R. (2016). The dependence of precipitation and its footprint on atmospheric temperature in  
42 idealized extratropical cyclones. *J. Geophys. Res.* 121, 8743–8754. doi:10.1002/2015JD024286.
- 43 Philip, S., Kew, S. F., Jan van Oldenborgh, G., Aalbers, E., Vautard, R., Otto, F., et al. (2018a). Validation of a Rapid  
44 Attribution of the May/June 2016 Flood-Inducing Precipitation in France to Climate Change. *J. Hydrometeorol.*  
45 19, 1881–1898. doi:10.1175/JHM-D-18-0074.1.
- 46 Philip, S., Kew, S. F., Jan van Oldenborgh, G., Otto, F., O’Keefe, S., Haustein, K., et al. (2018b). Attribution Analysis  
47 of the Ethiopian Drought of 2015. *J. Clim.* 31, 2465–2486. doi:10.1175/JCLI-D-17-0274.1.
- 48 Philip, S., Kew, S., van Oldenborgh, G. J., Otto, F., Vautard, R., van der Wiel, K., et al. (2020). A protocol for  
49 probabilistic extreme event attribution analyses. *Adv. Stat. Clim. Meteorol. Ocean.* 6, 177–203.  
50 doi:10.5194/ascmo-6-177-2020.
- 51 Philip, S., Sparrow, S., Kew, S. F., van der Wiel, K., Wanders, N., Singh, R., et al. (2019). Attributing the 2017  
52 Bangladesh floods from meteorological and hydrological perspectives. *Hydrol. Earth Syst. Sci.* 23, 1409–1429.  
53 doi:10.5194/hess-23-1409-2019.
- 54 Piaget, N., Froidevaux, P., Giannakaki, P., Gierth, F., Martius, O., Riemer, M., et al. (2015). Dynamics of a local Alpine  
55 flooding event in October 2011: moisture source and large-scale circulation. *Q. J. R. Meteorol. Soc.* 141, 1922–  
56 1937. doi:10.1002/qj.2496.
- 57 Pinto, I., Jack, C., and Hewitson, B. (2018). Process-based model evaluation and projections over southern Africa from  
58 Coordinated Regional Climate Downscaling Experiment and Coupled Model Intercomparison Project Phase 5  
59 models. *Int. J. Climatol.*, 1–11. doi:10.1002/joc.5666.
- 60 Pinto, I., Lennard, C., Tadross, M., Hewitson, B., Dosio, A., Nikulin, G., et al. (2016). Evaluation and projections of  
61 extreme precipitation over southern Africa from two CORDEX models. *Clim. Change* 135. doi:10.1007/s10584-

- 1 015-1573-1.
- 2 Pisaniello, J. D., Tingey-Holyoak, J., and Burritt, R. L. (2012). Appropriate small dam management for minimizing  
3 catchment-wide safety threats: International benchmarked guidelines and demonstrative cases studies. *Water*  
4 *Resour. Res.* 48. doi:10.1029/2011WR011155.
- 5 Pithan, F., and Mauritsen, T. (2014). Arctic amplification dominated by temperature feedbacks in contemporary climate  
6 models. *Nat. Geosci.* 7, 181–184. doi:10.1038/ngeo2071.
- 7 Piticar, A. (2018). Changes in heat waves in Chile. *Glob. Planet. Change* 169, 234–246.  
8 doi:10.1016/j.gloplacha.2018.08.007.
- 9 Piticar, A., Mihăilă, D., Lazurca, L. G., Bistricean, P.-I., Puțuntică, A., and Briciu, A.-E. (2016). Spatiotemporal  
10 distribution of reference evapotranspiration in the Republic of Moldova. *Theor. Appl. Climatol.* 124, 1133–1144.  
11 doi:10.1007/s00704-015-1490-2.
- 12 Podschwit, H. R., Larkin, N. K., Steel, E. A., Cullen, A., and Alvarado, E. (2018). Multi-model forecasts of very-large  
13 fire occurrences during the end of the 21st century. *Climate* 6, 1–21. doi:10.3390/cli6040100.
- 14 Pokhrel, Y., Felfelani, F., Satoh, Y., Boulange, J., Burek, P., Gädeke, A., et al. (2021). Global terrestrial water storage  
15 and drought severity under climate change. *Nat. Clim. Chang.* doi:10.1038/s41558-020-00972-w.
- 16 Poshtiri, M. P., and Pal, I. (2016). Patterns of hydrological drought indicators in major U.S. River basins. *Clim. Change*  
17 134, 549–563. doi:10.1007/s10584-015-1542-8.
- 18 Potopová, V., Štěpánek, P., Zahradníček, P., Farda, A., Türkott, L., and Soukup, J. (2018). Projected changes in the  
19 evolution of drought on various timescales over the Czech Republic according to Euro-CORDEX models. *Int. J.*  
20 *Climatol.* 38, e939–e954. doi:10.1002/joc.5421.
- 21 Preimesberger, W., Scanlon, T., Su, C.-H., Gruber, A., and Dorigo, W. (2020). Homogenization of Structural Breaks in  
22 the Global ESA CCI Soil Moisture Multisatellite Climate Data Record. *IEEE Trans. Geosci. Remote Sens.*, 1–18.  
23 doi:10.1109/tgrs.2020.3012896.
- 24 Prein, A. F., Gobiet, A., Truhetz, H., Keuler, K., Goergen, K., Teichmann, C., et al. (2016a). Precipitation in the  
25 EURO-CORDEX 0.11° and 0.44° simulations: high resolution, high benefits? *Clim. Dyn.* 46, 383–412.  
26 doi:10.1007/s00382-015-2589-y.
- 27 Prein, A. F., and Holland, G. J. (2018). Global estimates of damaging hail hazard. *Weather Clim. Extrem.* 22, 10–23.  
28 doi:10.1016/j.wace.2018.10.004.
- 29 Prein, A. F., Langhans, W., Fossier, G., Ferrone, A., Ban, N., Goergen, K., et al. (2015). A review on regional  
30 convection-permitting climate modeling: Demonstrations, prospects, and challenges. *Rev. Geophys.* 53, 323–361.  
31 doi:10.1002/2014RG000475.
- 32 Prein, A. F., Liu, C., Ikeda, K., Bullock, R., Rasmussen, R. M., Holland, G. J., et al. (2017a). Simulating North  
33 American mesoscale convective systems with a convection-permitting climate model. *Clim. Dyn.* 0, 1–16.  
34 doi:10.1007/s00382-017-3993-2.
- 35 Prein, A. F., Liu, C., Ikeda, K., Trier, S. B., Rasmussen, R. M., Holland, G. J., et al. (2017b). Increased rainfall volume  
36 from future convective storms in the US. *Nat. Clim. Chang.* 7, 880–884. doi:10.1038/s41558-017-0007-7.
- 37 Prein, A. F., Rasmussen, R. M., Ikeda, K., Liu, C., Clark, M. P., and Holland, G. J. (2016b). The future intensification  
38 of hourly precipitation extremes. *Nat. Clim. Chang.* 7, 48. doi:10.1038/nclimate3168.
- 39 Priestley, M. D. K. K., Ackerley, D., Catto, J. L., Hodges, K. I., McDonald, R. E., and Lee, R. W. (2020). An Overview  
40 of the Extratropical Storm Tracks in CMIP6 Historical Simulations. *J. Clim.* 33, 6315–6343. doi:10.1175/JCLI-D-  
41 19-0928.1.
- 42 Priya, P., Krishnan, R., Mujumdar, M., and Houze, R. A. (2017). Changing monsoon and midlatitude circulation  
43 interactions over the Western Himalayas and possible links to occurrences of extreme precipitation. *Clim. Dyn.*  
44 49, 2351–2364. doi:10.1007/s00382-016-3458-z.
- 45 Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., et al. (2014). Hydrological  
46 droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment. *Proc.*  
47 *Natl. Acad. Sci. U. S. A.* 111, 3262–3267. doi:10.1073/pnas.1222473110.
- 48 Púčik, T., Groenemeijer, P., Rädler, A. T., Tijssen, L., Nikulin, G., Prein, A. F., et al. (2017). Future Changes in  
49 European Severe Convection Environments in a Regional Climate Model Ensemble. *J. Clim.* 30, 6771–6794.  
50 doi:10.1175/JCLI-D-16-0777.1.
- 51 Qian, C., Wang, J., Dong, S., Yin, H., Burke, C., Ciavarella, A., et al. (2018). Human Influence on the Record-breaking  
52 Cold Event in January of 2016 in Eastern China [in “Explaining Extreme Events of 2016 from a Climate  
53 Perspective”]. *Bull. Am. Meteorol. Soc.* 99, S118–S122. doi:10.1175/BAMS-D-17-0095.1.
- 54 Qian, C., Zhang, X., and Li, Z. (2019). Linear trends in temperature extremes in China, with an emphasis on non-  
55 Gaussian and serially dependent characteristics. *Clim. Dyn.* 53, 533–550. doi:10.1007/s00382-018-4600-x.
- 56 Qin, N., Wang, J., Yang, G., Chen, X., Liang, H., and Zhang, J. (2015a). Spatial and temporal variations of extreme  
57 precipitation and temperature events for the Southwest China in 1960–2009. *Geoenvironmental Disasters* 2.  
58 doi:10.1186/s40677-015-0014-9.
- 59 Qin, Y., Yang, D., Lei, H., Xu, K., and Xu, X. (2015b). Comparative analysis of drought based on precipitation and soil  
60 moisture indices in Haihe basin of North China during the period of 1960–2010. *J. Hydrol.* 526, 55–67.  
61 doi:10.1016/j.jhydrol.2014.09.068.

- 1 Qiu, J., Gao, Q., Wang, S., and Su, Z. (2016). Comparison of temporal trends from multiple soil moisture data sets and  
2 precipitation: The implication of irrigation on regional soil moisture trend. *Int. J. Appl. Earth Obs. Geoinf.* 48,  
3 17–27. doi:10.1016/j.jag.2015.11.012.
- 4 Quesada, B., Vautard, R., Yiou, P., Hirschi, M., and Seneviratne, S. I. (2012). Asymmetric European summer heat  
5 predictability from wet and dry southern winters and springs. *Nat. Clim. Chang.* 2, 736–741.  
6 doi:10.1038/nclimate1536.
- 7 Quintana-Seguí, P., Barella-Ortiz, A., Regueiro-Sanfiz, S., and Miguez-Macho, G. (2020). The Utility of Land-Surface  
8 Model Simulations to Provide Drought Information in a Water Management Context Using Global and Local  
9 Forcing Datasets. *Water Resour. Manag.* 34, 2135–2156. doi:10.1007/s11269-018-2160-9.
- 10 Quiring, S. M., Ford, T. W., Wang, J. K., Khong, A., Harris, E., Lindgren, T., et al. (2016). The North American Soil  
11 Moisture Database: Development and Applications. *Bull. Am. Meteorol. Soc.* 97, 1441–1459.  
12 doi:10.1175/BAMS-D-13-00263.1.
- 13 Raghavan, S. V., Hur, J., and Liang, S.-Y. (2018). Evaluations of NASA NEX-GDDP data over Southeast Asia: present  
14 and future climates. *Clim. Change* 148, 503–518. doi:10.1007/s10584-018-2213-3.
- 15 Ragone, F., Mariotti, M., Parodi, A., von Hardenberg, J., and Pasquero, C. (2018). A Climatological Study of Western  
16 Mediterranean Medicanes in Numerical Simulations with Explicit and Parameterized Convection. *Atmosphere*  
17 (*Basel*). 9, 397. doi:10.3390/atmos9100397.
- 18 Rahimi, M., and Fatemi, S. S. (2019). Mean versus Extreme Precipitation Trends in Iran over the Period 1960–2017.  
19 *Pure Appl. Geophys.* 176, 3717–3735. doi:10.1007/s00024-019-02165-9.
- 20 Rahimi, M., and Hejabi, S. (2018). Spatial and temporal analysis of trends in extreme temperature indices in Iran over  
21 the period 1960–2014. *Int. J. Climatol.* 38, 272–282. doi:10.1002/joc.5175.
- 22 Rahimi, M., Mohammadian, N., Vanashi, A. R., and Whan, K. (2018). Trends in Indices of Extreme Temperature and  
23 Precipitation in Iran over the Period 1960–2014. *Open J. Ecol.* 08, 396–415. doi:10.4236/oje.2018.87024.
- 24 Rai, P., Choudhary, A., and Dimri, A. P. (2019). Future precipitation extremes over India from the CORDEX-South  
25 Asia experiments. *Theor. Appl. Climatol.* doi:10.1007/s00704-019-02784-1.
- 26 Rajbhandari, R., Shrestha, A. B., Kulkarni, A., Patwardhan, S. K., and Bajracharya, S. R. (2015). Projected changes in  
27 climate over the Indus river basin using a high resolution regional climate model (PRECIS). *Clim. Dyn.* 44, 339–  
28 357. doi:10.1007/s00382-014-2183-8.
- 29 Rajczak, J., Pall, P., and Schär, C. (2013). Projections of extreme precipitation events in regional climate simulations  
30 for Europe and the Alpine Region. *J. Geophys. Res. Atmos.* 118, 3610–3626. doi:10.1002/jgrd.502972013.
- 31 Rajczak, J., and Schär, C. (2017). Projections of Future Precipitation Extremes Over Europe: A Multimodel Assessment  
32 of Climate Simulations. *J. Geophys. Res. Atmos.* 122, 10,773–10,800. doi:10.1002/2017JD027176.
- 33 Rajsekhar, D., and Gorelick, S. M. (2017). Increasing drought in Jordan: Climate change and cascading Syrian land-use  
34 impacts on reducing transboundary flow. *Sci. Adv.* 3, 1–15. doi:10.1126/sciadv.1700581.
- 35 Ralph, F. M., Dettinger, M. D., Cairns, M. M., Galarneau, T. J., and Eylander, J. (2018). Defining “Atmospheric  
36 River”: How the Glossary of Meteorology Helped Resolve a Debate. *Bull. Am. Meteorol. Soc.* 99, 837–839.  
37 doi:10.1175/BAMS-D-17-0157.1.
- 38 Ramos, A. M., Tomé, R., Trigo, R. M., Liberato, M. L. R., and Pinto, J. G. (2016). Projected changes in atmospheric  
39 rivers affecting Europe in CMIP5 models. *Geophys. Res. Lett.* 43, 9315–9323. doi:10.1002/2016GL070634.
- 40 Rasmijn, L. M., Van Der Schrier, G., Bintanja, R., Barkmeijer, J., Sterl, A., and Hazeleger, W. (2018). Future  
41 equivalent of 2010 Russian heatwave intensified by weakening soil moisture constraints. *Nat. Clim. Chang.* 8,  
42 381–385. doi:10.1038/s41558-018-0114-0.
- 43 Rasmussen, K. L., and Houze, R. A. (2011). Orographic Convection in Subtropical South America as Seen by the TRMM  
44 Satellite. *Mon. Weather Rev.* 139, 2399–2420. doi:10.1175/MWR-D-10-05006.1.
- 45 Rasmussen, K. L., Prein, A. F., Rasmussen, R. M., Ikeda, K., and Liu, C. (2017). Changes in the convective population  
46 and thermodynamic environments in convection-permitting regional climate simulations over the United States.  
47 *Clim. Dyn.* doi:10.1007/s00382-017-4000-7.
- 48 Ratnam, J. V., Behera, S. K., Ratna, S. B., Rajeevan, M., and Yamagata, T. (2016). Anatomy of Indian heatwaves. *Sci.*  
49 *Rep.* 6, 1–11. doi:10.1038/srep24395.
- 50 Rauniyar, S. P., and Power, S. B. (2020). The impact of anthropogenic forcing and natural processes on past, present,  
51 and future rainfall over Victoria, Australia. *J. Clim.* 33, 8087–8106. doi:10.1175/JCLI-D-19-0759.1.
- 52 Raymond, C., Horton, R. M., Zscheischler, J., Martius, O., AghaKouchak, A., Balch, J., et al. (2020). Understanding  
53 and managing connected extreme events. *Nat. Clim. Chang.* 10, 611–621. doi:10.1038/s41558-020-0790-4.
- 54 Raymond, F., Ullmann, A., Camberlin, P., Oueslati, B., and Drobinski, P. (2018). Atmospheric conditions and weather  
55 regimes associated with extreme winter dry spells over the Mediterranean basin. *Clim. Dyn.* 50, 4437–4453.  
56 doi:10.1007/s00382-017-3884-6.
- 57 Raymond, F., Ullmann, A., Trambly, Y., Drobinski, P., and Camberlin, P. (2019). Evolution of Mediterranean extreme  
58 dry spells during the wet season under climate change. *Reg. Environ. Chang.* 19, 2339–2351.  
59 doi:10.1007/s10113-019-01526-3.
- 60 Reale, M., and Lionello, P. (2013). Synoptic climatology of winter intense precipitation events along the Mediterranean  
61 coasts. *Nat. Hazards Earth Syst. Sci.* 13, 1707–1722. doi:10.5194/nhess-13-1707-2013.

- 1 Reboita, M. S., da Rocha, R. P., Ambrizzi, T., and Gouveia, C. D. (2015). Trend and teleconnection patterns in the  
2 climatology of extratropical cyclones over the Southern Hemisphere. *Clim. Dyn.* 45, 1929–1944.  
3 doi:10.1007/s00382-014-2447-3.
- 4 Reboita, M. S., Reale, M., da Rocha, R. P., Giorgi, F., Giuliani, G., Coppola, E., et al. (2020). Future changes in the  
5 wintertime cyclonic activity over the CORDEX-CORE southern hemisphere domains in a multi-model approach.  
6 *Clim. Dyn.* doi:10.1007/s00382-020-05317-z.
- 7 Redmond, G., Hodges, K. I., Mcsweeney, C., Jones, R., and Hein, D. (2015). Projected changes in tropical cyclones  
8 over Vietnam and the South China Sea using a 25 km regional climate model perturbed physics ensemble. *Clim.*  
9 *Dyn.* 45, 1983–2000. doi:10.1007/s00382-014-2450-8.
- 10 Reed, K. A., Bacmeister, J. T., Huff, J. J. A., Wu, X., Bates, S. C., and Rosenbloom, N. A. (2019). Exploring the Impact  
11 of Dust on North Atlantic Hurricanes in a High-Resolution Climate Model. *Geophys. Res. Lett.* 46, 1105–1112.  
12 doi:10.1029/2018GL080642.
- 13 Reed, K. A., Bacmeister, J. T., Rosenbloom, N. A., Wehner, M. F., Bates, S. C., Lauritzen, P. H., et al. (2015). Impact  
14 of the dynamical core on the direct simulation of tropical cyclones in a high-resolution global model. *Geophys.*  
15 *Res. Lett.* 42, 3603–3608. doi:10.1002/2015GL063974.
- 16 Reed, K. A., and Jablonowski, C. (2011). Impact of physical parameterizations on idealized tropical cyclones in the  
17 Community Atmosphere Model. *Geophys. Res. Lett.* 38. doi:10.1029/2010GL046297.
- 18 Reed, K. A., Stansfield, A. M., Wehner, M. F., and Zarzycki, C. M. (2020). Forecasted attribution of the human  
19 influence on Hurricane Florence. *Sci. Adv.* 6, eaaw9253. doi:10.1126/sciadv.aaw9253.
- 20 Rehbein, A., Ambrizzi, T., and Mechoso, C. R. (2018). Mesoscale convective systems over the Amazon basin. Part I:  
21 climatological aspects. *Int. J. Climatol.* 38, 215–229. doi:10.1002/joc.5171.
- 22 Rehman, S. (2013). Long-term wind speed analysis and detection of its trends using Mann-Kendall test and linear  
23 regression method. *Arab. J. Sci. Eng.* 38, 421–437. doi:10.1007/s13369-012-0445-5.
- 24 Reichle, R. H., Draper, C. S., Liu, Q., Girotto, M., Mahanama, S. P. P., Koster, R. D., et al. (2017). Assessment of  
25 MERRA-2 Land Surface Hydrology Estimates. *J. Clim.* 30, 2937–2960. doi:10.1175/JCLI-D-16-0720.1.
- 26 Ren, L., Ren, L., Zhou, T., Zhou, T., Zhou, T., and Zhang, W. (2020). Attribution of the record-breaking heat event  
27 over Northeast Asia in summer 2018: The role of circulation. *Environ. Res. Lett.* 15. doi:10.1088/1748-  
28 9326/ab8032.
- 29 Reyer, C. P. ., Otto, I. M., Adams, S., Albrecht, T., Baarsch, F., Carlsburg, M., et al. (2017). Climate change impacts in  
30 Central Asia and their implications for development. *Reg. Environ. Chang.* 17, 1639–1650. doi:10.1007/s10113-  
31 015-0893-z.
- 32 Rhoades, A. M., Ullrich, P. A., and Zarzycki, C. M. (2018). Projecting 21st century snowpack trends in western USA  
33 mountains using variable-resolution CESM. *Clim. Dyn.* 50, 261–288. doi:10.1007/s00382-017-3606-0.
- 34 Ribal, A., and Young, I. R. (2019). 33 Years of Globally Calibrated Wave Height and Wind Speed Data Based on  
35 Altimeter Observations. *Sci. data* 6, 77. doi:10.1038/s41597-019-0083-9.
- 36 Ribeiro, I. O., Andreoli, R. V., Kayano, M. T., Sousa, T. R., Medeiros, A. S., Godoi, R. H. M., et al. (2018). Biomass  
37 burning and carbon monoxide patterns in Brazil during the extreme drought years of 2005, 2010, and 2015.  
38 *Environ. Pollut.* 243, 1008–1014. doi:10.1016/j.envpol.2018.09.022.
- 39 Ribes, A., Thao, S., Vautard, R., Dubuisson, B., Somot, S., Colin, J., et al. (2019). Observed increase in extreme daily  
40 rainfall in the French Mediterranean. *Clim. Dyn.* 52, 1095–1114. doi:10.1007/s00382-018-4179-2.
- 41 Ridder, N., de Vries, H., and Drijfhout, S. (2018). The role of atmospheric rivers in compound events consisting of  
42 heavy precipitation and high storm surges along the Dutch coast. *Nat. Hazards Earth Syst. Sci.* 18, 3311–3326.  
43 doi:10.5194/nhess-18-3311-2018.
- 44 Ridder, N. N., Pitman, A. J., Westra, S., Ukkola, A., Do, H. X., Bador, M., et al. (2020). Global hotspots for the  
45 occurrence of compound events. *Nat. Commun.* 11, 5956. doi:10.1038/s41467-020-19639-3.
- 46 Ridley, J., Wiltshire, A., and Mathison, C. (2013). More frequent occurrence of westerly disturbances in Karakoram up  
47 to 2100. *Sci. Total Environ.* 468–469, S31–S35. doi:10.1016/J.SCITOTENV.2013.03.074.
- 48 Ringard, J., Dieppois, B., Rome, S., Diedhiou, A., Pellarin, T., Konaré, A., et al. (2016). The intensification of thermal  
49 extremes in west Africa. *Glob. Planet. Change* 139, 66–77. doi:10.1016/j.gloplacha.2015.12.009.
- 50 Risser, M. D., and Wehner, M. F. (2017). Attributable Human-Induced Changes in the Likelihood and Magnitude of the  
51 Observed Extreme Precipitation during Hurricane Harvey. *Geophys. Res. Lett.* 44, 12,412–457,464.  
52 doi:10.1002/2017GL075888.
- 53 Rivera, J. A., and Arnould, G. (2020). Evaluation of the ability of CMIP6 models to simulate precipitation over  
54 Southwestern South America: Climatic features and long-term trends (1901–2014). *Atmos. Res.* 241, 104953.  
55 doi:https://doi.org/10.1016/j.atmosres.2020.104953.
- 56 Rivera, J. A., and Penalba, O. C. (2018). Spatio-temporal assessment of streamflow droughts over Southern South  
57 America: 1961–2006. *Theor. Appl. Climatol.* 133, 1021–1033. doi:10.1007/s00704-017-2243-1.
- 58 Roberts, M. J., Camp, J., Seddon, J., Vidale, P. L., Hodges, K., Vanniere, B., et al. (2020a). Impact of model resolution  
59 on tropical cyclone simulation using the HighResMIP PRIMAVERA multi-model ensemble. *J. Clim.* 33, 2557–  
60 2583. doi:10.1175/JCLI-D-19-0639.1.
- 61 Roberts, M. J., Camp, J., Seddon, J., Vidale, P. L., Hodges, K., Vanniere, B., et al. (2020b). Projected Future changes of

- 1 tropical cyclones using the CMIP6 HighResMIP multi-model ensemble. *Geophys. Res. Lett.* 47, e2020GL088662.  
2 doi:10.1029/2020GL088662.
- 3 Roberts, M. J., Vidale, P. L., Senior, C., Hewitt, H. T., Bates, C., Berthou, S., et al. (2018). The Benefits of Global High  
4 Resolution for Climate Simulation: Process Understanding and the Enabling of Stakeholder Decisions at the  
5 Regional Scale. *Bull. Am. Meteorol. Soc.* 99, 2341–2359. doi:10.1175/BAMS-D-15-00320.1.
- 6 Rodell, M., Famiglietti, J. S., Wiese, D. N., Reager, J. T., Beaudoin, H. K., Landerer, F. W., et al. (2018). Emerging  
7 trends in global freshwater availability. *Nature* 557, 651–659. doi:10.1038/s41586-018-0123-1.
- 8 Roderick, M. L., Greve, P., and Farquhar, G. D. (2015). On the assessment of aridity with changes in atmospheric CO<sub>2</sub>.  
9 *Water Resour. Res.* 51, 5450–5463. doi:10.1002/2015WR017031.
- 10 Rogelj, J. (2013). Uncertainties of low greenhouse gas emission scenarios. doi:doi.org/10.3929/ethz-a-009915210.
- 11 Rogger, M., Agnoletti, M., Alaoui, A., Bathurst, J. C., Bodner, G., Borga, M., et al. (2017). Land use change impacts on  
12 floods at the catchment scale: Challenges and opportunities for future research. *Water Resour. Res.* 53, 5209–  
13 5219. doi:10.1002/2017WR020723.
- 14 Rohat, G., Flacke, J., Dosio, A., Pedde, S., Dao, H., and van Maarseveen, M. (2019). Influence of changes in  
15 socioeconomic and climatic conditions on future heat-related health challenges in Europe. *Glob. Planet. Change*  
16 172, 45–59. doi:10.1016/j.gloplacha.2018.09.013.
- 17 Rohini, P., Rajeevan, M., and Srivastava, A. K. (2016). On the Variability and Increasing Trends of Heat Waves over  
18 India. *Sci. Rep.*, 1–9.
- 19 Romera, R., Gaertner, M. Á., Sánchez, E., Domínguez, M., González-Alemán, J. J., and Miglietta, M. M. (2017).  
20 Climate change projections of medicanes with a large multi-model ensemble of regional climate models. *Glob.*  
21 *Planet. Change* 151, 134–143. doi:10.1016/j.gloplacha.2016.10.008.
- 22 Romero, R., and Emanuel, K. (2013). Medicanes risk in a changing climate. *J. Geophys. Res. Atmos.* 118, 5992–6001.  
23 doi:10.1002/jgrd.50475.
- 24 Romero, R., and Emanuel, K. (2017). Climate change and hurricane-like extratropical cyclones: Projections for North  
25 Atlantic polar lows and medicanes based on CMIP5 models. *J. Clim.* 30, 279–299. doi:10.1175/JCLI-D-16-  
26 0255.1.
- 27 Romps, D. M. (2016). Clausius-clapeyron scaling of CAPE from analytical solutions to RCE. *J. Atmos. Sci.* 73, 3719–  
28 3737. doi:10.1175/JAS-D-15-0327.1.
- 29 Rosenzweig, B. R., McPhillips, L., Chang, H., Cheng, C., Welty, C., Matsler, M., et al. (2018). Pluvial flood risk and  
30 opportunities for resilience. *WIREs Water* 5, e1302. doi:10.1002/wat2.1302.
- 31 Rosier, S., Dean, S., Stuart, S., Carey-Smith, T., Black, M. T., and Massey, N. (2016). Extreme rainfall in early July  
32 2014 in northland, New Zealand—was there an anthropogenic influence? *Bull. Am. Meteorol. Soc.* 96, S136–S140.  
33 doi:10.1175/BAMS-D-15-00105.1.
- 34 Roth, M., Buishand, T. A., Jongbloed, G., Klein Tank, A. M. G., and van Zanten, J. H. (2014). Projections of  
35 precipitation extremes based on a regional, non-stationary peaks-over-threshold approach: A case study for the  
36 Netherlands and north-western Germany. *Weather Clim. Extrem.* 4, 1–10. doi:10.1016/j.wace.2014.01.001.
- 37 Roth, M., Jongbloed, G., and Buishand, A. (2018). Monotone trends in the distribution of climate extremes. *Theor.*  
38 *Appl. Climatol.* doi:10.1007/s00704-018-2546-x.
- 39 Roudier, P., Andersson, J. C. M., Donnelly, C., Feyen, L., Greuell, W., and Ludwig, F. (2016). Projections of future  
40 floods and hydrological droughts in Europe under a +2°C global warming. *Clim. Change* 135, 341–355.  
41 doi:10.1007/s10584-015-1570-4.
- 42 Rowell, D. P., Booth, B. B. B., Nicholson, S. E., and Good, P. (2015). Reconciling Past and Future Rainfall Trends over  
43 East Africa. *J. Clim.* 28, 9768–9788. doi:10.1175/JCLI-D-15-0140.1.
- 44 Rowland, L., Da Costa, A. C. L., Galbraith, D. R., Oliveira, R. S., Binks, O. J., Oliveira, A. A. R., et al. (2015). Death  
45 from drought in tropical forests is triggered by hydraulics not carbon starvation. *Nature* 528, 119–122.  
46 doi:10.1038/nature15539.
- 47 Roxy, M. K. K., Ghosh, S., Pathak, A., Athulya, R., Mujumdar, M., Murtugudde, R., et al. (2017). A threefold rise in  
48 widespread extreme rain events over central India. *Nat. Commun.* 8, 708. doi:10.1038/s41467-017-00744-9.
- 49 Roy, S. Sen (2019). Spatial patterns of trends in seasonal extreme temperatures in India during 1980–2010. *Weather*  
50 *Clim. Extrem.* 24, 100203. doi:10.1016/j.wace.2019.100203.
- 51 Ruffault, J., Curt, T., Martin-Stpaul, N. K., Moron, V., and Trigo, R. M. (2018). Extreme wildfire events are linked to  
52 global-change-type droughts in the northern Mediterranean. *Nat. Hazards Earth Syst. Sci.* 18, 847–856.  
53 doi:10.5194/nhess-18-847-2018.
- 54 Ruml, M., Gregorić, E., Vujadinović, M., Radovanović, S., Matović, G., Vuković, A., et al. (2017). Observed changes  
55 of temperature extremes in Serbia over the period 1961 – 2010. *Atmos. Res.* 183, 26–41.  
56 doi:10.1016/j.atmosres.2016.08.013.
- 57 Ruosteenoja, K., Markkanen, T., Venäläinen, A., Räisänen, P., and Peltola, H. (2018). Seasonal soil moisture and  
58 drought occurrence in Europe in CMIP5 projections for the 21st century. *Clim. Dyn.* 50, 1177–1192.  
59 doi:10.1007/s00382-017-3671-4.
- 60 Rupp, D. E., Mote, P. W., Massey, N., Otto, F. E. L., and Allen, M. R. (2013). Human influence on the probability of  
61 low precipitation in the central United States in 2012 [in “Explaining Extreme Events of 2012 from a Climate

- 1 Perspective”]. *Bull. Am. Meteorol. Soc.* 94, S2–S6. doi:10.1175/BAMS-D-13-00085.1.
- 2 Ruprich-Robert, Y., Delworth, T., Msadek, R., Castruccio, F., Yeager, S., and Danabasoglu, G. (2018). Impacts of the  
3 Atlantic Multidecadal Variability on North American Summer Climate and Heat Waves. *J. Clim.* 31, 3679–3700.  
4 doi:10.1175/JCLI-D-17-0270.1.
- 5 Russo, S., Dosio, A., Sterl, A., Barbosa, P., and Vogt, J. (2013). Projection of occurrence of extreme dry-wet years and  
6 seasons in Europe with stationary and nonstationary Standardized Precipitation Indices. *J. Geophys. Res. Atmos.*  
7 118, 7628–7639. doi:10.1002/jgrd.50571.
- 8 Russo, S., Marchese, A. F., Sillmann, J., and Immé, G. (2016). When will unusual heat waves become normal in a  
9 warming Africa? *Environ. Res. Lett.* 11, 054016. doi:10.1088/1748-9326/11/5/054016.
- 10 Russo, S., Sillmann, J., and Fischer, E. M. (2015). Top ten European heatwaves since 1950 and their occurrence in the  
11 coming decades. *Environ. Res. Lett.* 10, 124003. doi:10.1088/1748-9326/10/12/124003.
- 12 Rusticucci, M., Barrucand, M., and Collazo, S. (2017). Temperature extremes in the Argentina central region and their  
13 monthly relationship with the mean circulation and ENSO phases. *Int. J. Climatol.* 37, 3003–3017.  
14 doi:10.1002/joc.4895.
- 15 Sabeerali, C. T., Rao, S. A., Dhakate, A. R., Salunke, K., and Goswami, B. N. (2015). Why ensemble mean projection  
16 of south Asian monsoon rainfall by CMIP5 models is not reliable? *Clim. Dyn.* 45, 161–174. doi:10.1007/s00382-  
17 014-2269-3.
- 18 Saha, A., Ghosh, S., Sahana, A. S., and Rao, E. P. (2014). Failure of CMIP5 climate models in simulating post-1950  
19 decreasing trend of Indian monsoon. *Geophys. Res. Lett.* 41, 7323–7330. doi:10.1002/2014GL061573.
- 20 Sahoo, B., and Bhaskaran, P. K. (2016). Assessment on historical cyclone tracks in the Bay of Bengal, east coast of  
21 India. *Int. J. Climatol.* 36, 95–109. doi:10.1002/joc.4331.
- 22 Salinger, M. J., and Porteous, A. S. (2014). New Zealand Climate: Patterns of Drought 1941/42 – 2012/13. *Weather  
23 Clim.* 34, 2–19. doi:10.2307/26169741.
- 24 Salnikov, V., Tutulina, G., Polyakova, S., Petrova, Y., and Skakova, A. (2015). Climate change in Kazakhstan during  
25 the past 70 years. *Quat. Int.* 358, 77–82. doi:10.1016/j.quaint.2014.09.008.
- 26 Salvador, M. de A., and de Brito, J. I. B. (2018). Trend of annual temperature and frequency of extreme events in the  
27 MATOPIBA region of Brazil. *Theor. Appl. Climatol.* 133, 253–261. doi:10.1007/s00704-017-2179-5.
- 28 Salvi, K., and Ghosh, S. (2016). Projections of Extreme Dry and Wet Spells in the 21st Century India Using Stationary  
29 and Non-stationary Standardized Precipitation Indices. *Clim. Change* 139, 667–681. doi:10.1007/s10584-016-  
30 1824-9.
- 31 Samaniego, L., Kumar, R., Breuer, L., Chamorro, A., Flörke, M., Pechlivanidis, I. G., et al. (2017). Propagation of  
32 forcing and model uncertainties on to hydrological drought characteristics in a multi-model century-long  
33 experiment in large river basins. *Clim. Change* 141, 435–449. doi:10.1007/s10584-016-1778-y.
- 34 Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., et al. (2018). Anthropogenic warming  
35 exacerbates European soil moisture droughts. *Nat. Clim. Chang.* 8, 421–426. doi:10.1038/s41558-018-0138-5.
- 36 Samouly, A. A., Luong, C. N., Li, Z., Smith, S., Baetz, B., and Ghaith, M. (2018). Performance of multi-model  
37 ensembles for the simulation of temperature variability over Ontario, Canada. *Environ. Earth Sci.* 77, 524.  
38 doi:10.1007/s12665-018-7701-2.
- 39 Samset, B. H., Sand, M., Smith, C. J., Bauer, S. E., Forster, P. M., Fuglestedt, J. S., et al. (2018). Climate Impacts  
40 From a Removal of Anthropogenic Aerosol Emissions. *Geophys. Res. Lett.* 45, 1020–1029.  
41 doi:10.1002/2017GL076079.
- 42 Samset, B. H., Stjern, C. W., Lund, M. T., Mohr, C. W., Sand, M., and Daloz, A. S. (2019). How Daily Temperature  
43 and Precipitation Distributions Evolve With Global Surface Temperature. *Earth’s Futur.* 7, 1323–1336.  
44 doi:10.1029/2019EF001160.
- 45 Samuels, R., Hochman, A., Baharad, A., Givati, A., Levi, Y., Yosef, Y., et al. (2018). Evaluation and projection of  
46 extreme precipitation indices in the Eastern Mediterranean based on CMIP5 multi-model ensemble. *Int. J.  
47 Climatol.* 38, 2280–2297. doi:10.1002/joc.5334.
- 48 Sánchez-Benítez, A., Barriopedro, D., and García-Herrera, R. (2020). Tracking Iberian heatwaves from a new  
49 perspective. *Weather Clim. Extrem.* 28, 100238. doi:10.1016/j.wace.2019.100238.
- 50 Sanchez-Lorenzo, A., Wild, M., Brunetti, M., Guijarro, J. A., Hakuba, M. Z., Calbó, J., et al. (2015). Reassessment and  
51 update of long-term trends in downward surface shortwave radiation over Europe (1939–2012). *J. Geophys. Res.*  
52 120, 9555–9569. doi:10.1002/2015JD023321.
- 53 Sanderson, B. M., Xu, Y., Tebaldi, C., Wehner, M., O’Neill, B., Jahn, A., et al. (2017a). Community climate  
54 simulations to assess avoided impacts in 1.5 and 2C futures. *Earth Syst. Dyn.* 8, 827–847. doi:10.5194/esd-8-827-  
55 2017.
- 56 Sanderson, M., Economou, T., Salmon, K., and Jones, S. (2017b). Historical Trends and Variability in Heat Waves in  
57 the United Kingdom. *Atmosphere (Basel)*. 8, 117–191. doi:10.3390/atmos8100191.
- 58 Sandvik, M. I., Sorteberg, A., and Rasmussen, R. (2018). Sensitivity of historical orographically enhanced extreme  
59 precipitation events to idealized temperature perturbations. *Clim. Dyn.* 50, 143–157. doi:10.1007/s00382-017-  
60 3593-1.
- 61 Sanginés de Cárcer, P., Vítasse, Y., Peñuelas, J., Jassey, V. E. J., Buttler, A., and Signarbieux, C. (2018). Vapor–

- 1 pressure deficit and extreme climatic variables limit tree growth. *Glob. Chang. Biol.* 24, 1108–1122.  
2 doi:10.1111/gcb.13973.
- 3 Sanogo, S., Fink, A. H., Omotosho, J. A., Ba, A., Redl, R., and Ermert, V. (2015). Spatio-temporal characteristics of the  
4 recent rainfall recovery in West Africa. *Int. J. Climatol.* 35, 4589–4605. doi:10.1002/joc.4309.
- 5 Santanello Jr., J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., et al. (2018). Land–  
6 Atmosphere Interactions: The LoCo Perspective. *Bull. Am. Meteorol. Soc.* 99, 1253–1272. doi:10.1175/BAMS-D-  
7 17-0001.1.
- 8 Sarangi, C., Tripathi, S. N., Kanawade, V. P., Koren, I., and Pai, D. S. (2017). Investigation of the aerosol–cloud–  
9 rainfall association over the Indian summer monsoon region. *Atmos. Chem. Phys.* 17, 5185–5204.  
10 doi:10.5194/acp-17-5185-2017.
- 11 Sarhadi, A., Ausin, M. C., Wiper, M. P., Touma, D., and Diffenbaugh, N. S. (2018). Multidimensional risk in a  
12 nonstationary climate: Joint probability of increasingly severe warm and dry conditions. *Sci. Adv.* 4, eaau3487.  
13 doi:10.1126/sciadv.aau3487.
- 14 Sato, T., and Nakamura, T. (2019). Intensification of hot Eurasian summers by climate change and land–atmosphere  
15 interactions. *Sci. Rep.* 9, 10866. doi:10.1038/s41598-019-47291-5.
- 16 Satoh, M., Stevens, B., Judt, F., Khairoutdinov, M., Lin, S., and Putman, W. M. (2019). Global Cloud-Resolving  
17 Models. *Curr. Clim. Chang. Reports*. doi:10.1007/s40641-019-00131-0.
- 18 Satoh, M., Tomita, H., Yashiro, H., Kajikawa, Y., Miyamoto, Y., Yamaura, T., et al. (2017). Outcomes and challenges  
19 of global high-resolution non-hydrostatic atmospheric simulations using the K computer. *Prog. Earth Planet. Sci.*  
20 4, 13. doi:10.1186/s40645-017-0127-8.
- 21 Satoh, M., Yamada, Y., Sugi, M., Komada, C., and Noda, A. T. (2015). Constraint on Future Change in Global  
22 Frequency of Tropical Cyclones due to Global Warming. *J. Meteorol. Soc. Japan. Ser. II* 93, 489–500.  
23 doi:10.2151/jmsj.2015-025.
- 24 Saurral, R. I., Camilloni, I. A., and Barros, V. R. (2017). Low-frequency variability and trends in centennial  
25 precipitation stations in southern South America. *Int. J. Climatol.* 37, 1774–1793. doi:10.1002/joc.4810.
- 26 Scaife, A. A., Comer, R., Dunstone, N., Fereday, D., Folland, C., Good, E., et al. (2017). Predictability of European  
27 winter 2015/2016. *Atmos. Sci. Lett.* 18, 38–44. doi:10.1002/asl.721.
- 28 Schaller, N., Kay, A. L., Lamb, R., Massey, N. R., van Oldenborgh, G. J., Otto, F. E. L., et al. (2016). Human influence  
29 on climate in the 2014 southern England winter floods and their impacts. *Nat. Clim. Chang.* 6, 627–634.  
30 doi:10.1038/nclimate2927.
- 31 Schaller, N., Otto, F. E. L., van Oldenborgh, G. J., Massey, N. R., Sparrow, S., and Allen, M. R. (2014). The heavy  
32 precipitation event of May–June 2013 in the upper Danube and Elbe basins. *Bull. Amer. Meteor. Soc.* 95, S69–  
33 S72.
- 34 Schaller, N., Sillmann, J., Anstey, J., Fischer, E. M., Grams, C. M., and Russo, S. (2018). Influence of blocking on  
35 Northern European and Western Russian heatwaves in large climate model ensembles. *Environ. Res. Lett.* 13.  
36 doi:10.1088/1748-9326/aaba55.
- 37 Schaller, N., Sillmann, J., Müller, M., Haarsma, R., Hazeleger, W., Hegdahl, T. J., et al. (2020). The role of spatial and  
38 temporal model resolution in a flood event storyline approach in western Norway. *Weather Clim. Extrem.* 29,  
39 100259. doi:10.1016/j.wace.2020.100259.
- 40 Scheff, J. (2018). Drought Indices, Drought Impacts, CO<sub>2</sub>, and Warming: a Historical and Geologic Perspective. *Curr.*  
41 *Clim. Chang. Reports* 4, 202–209. doi:10.1007/s40641-018-0094-1.
- 42 Scheff, J., and Frierson, D. M. W. (2015). Terrestrial aridity and its response to greenhouse warming across CMIP5  
43 climate models. *J. Clim.* 28, 5583–5600. doi:10.1175/JCLI-D-14-00480.1.
- 44 Scheff, J., Mankin, J. S., Coats, S., and Liu, H. (2021). CO<sub>2</sub>-plant effects do not account for the gap between dryness  
45 indices and projected dryness impacts in CMIP6 or CMIP5. *Environ. Res. Lett.* 16, 34018. doi:10.1088/1748-  
46 9326/abd8fd.
- 47 Scheff, J., Seager, R., Liu, H., and Coats, S. (2017). Are glacials dry? Consequences for paleoclimatology and for  
48 greenhouse warming. *J. Clim.* 30, 6593–6609. doi:10.1175/JCLI-D-16-0854.1.
- 49 Scher, S., Haarsma, R. J., de Vries, H., Drijfhout, S. S., and van Delden, A. J. (2017). Resolution dependence of  
50 extreme precipitation and deep convection over the Gulf Stream. *J. Adv. Model. Earth Syst.* 9, 1186–1194.  
51 doi:10.1002/2016MS000903.
- 52 Scherrer, S. C., Fischer, E. M., Posselt, R., Liniger, M. A., Croci-Maspoli, M., and Knutti, R. (2016). Emerging trends  
53 in heavy precipitation and hot temperature extremes in Switzerland. *J. Geophys. Res. Atmos.* 121, 2626–2637.  
54 doi:10.1002/2015JD024634.
- 55 Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., et al. (2014). Multimodel assessment of  
56 water scarcity under climate change. *Proc. Natl. Acad. Sci.* 111, 3245–3250. doi:10.1073/pnas.1222460110.
- 57 Schleussner, C.-F., Lissner, T. K., Fischer, E. M., Wohland, J., Perrette, M., Golly, A., et al. (2016). Differential climate  
58 impacts for policy-relevant limits to global warming: the case of 1.5 °C and 2 °C. *Earth Syst. Dyn.* 7, 327–351.  
59 doi:10.5194/esd-7-327-2016.
- 60 Schmid, P. E., and Niyogi, D. (2017). Modeling Urban Precipitation Modification by Spatially Heterogeneous  
61 Aerosols. *J. Appl. Meteorol. Climatol.* 56, 2141–2153. doi:10.1175/JAMC-D-16-0320.1.

- 1 Schneider, D., Huggel, C., Cochachin, A., Guillén, S., and García, J. (2014). Mapping hazards from glacier lake  
2 outburst floods based on modelling of process cascades at Lake 513, Carhuaz, Peru. *Adv. Geosci.* 35, 145–155.  
3 doi:10.5194/adgeo-35-145-2014.
- 4 Schneider, T., Bischoff, T., and Plotka, H. (2015). Physics of changes in synoptic midlatitude temperature variability. *J.*  
5 *Clim.* 28, 2312–2331. doi:10.1175/JCLI-D-14-00632.1.
- 6 Schoetter, R., Cattiaux, J., and Douville, H. (2015). Changes of western European heat wave characteristics projected  
7 by the CMIP5 ensemble. *Clim. Dyn.* 45, 1601–1616. doi:10.1007/s00382-014-2434-8.
- 8 Schreck, C. J., Knapp, K. R., and Kossin, J. P. (2014). The impact of best track discrepancies on global tropical cyclone  
9 climatologies using IBTrACS. *Mon. Weather Rev.* 142, 3881–3899. doi:10.1175/MWR-D-14-00021.1.
- 10 Schubert, S. D., Stewart, R. E., Wang, H., Barlow, M., Berbery, E. H., Cai, W., et al. (2016). Global Meteorological  
11 Drought: A Synthesis of Current Understanding with a Focus on SST Drivers of Precipitation Deficits. *J. Clim.*  
12 29, 3989–4019. doi:10.1175/JCLI-D-15-0452.1.
- 13 Schubert, S. D., Wang, H., Koster, R. D., Suarez, M. J., and Groisman, P. Y. (2014). Northern Eurasian heat waves and  
14 droughts. *J. Clim.* 27, 3169–3207. doi:10.1175/JCLI-D-13-00360.1.
- 15 Schumacher, D. L., Keune, J., van Heerwaarden, C. C., Vilà-Guerau de Arellano, J., Teuling, A. J., and Miralles, D. G.  
16 (2019). Amplification of mega-heatwaves through heat torrents fuelled by upwind drought. *Nat. Geosci.* 12, 712–  
17 717. doi:10.1038/s41561-019-0431-6.
- 18 Schumacher, R. S., and Johnson, R. H. (2005). Organization and Environmental Properties of Extreme-Rain-Producing  
19 Mesoscale Convective Systems. *Mon. Weather Rev.* 133, 961–976. doi:10.1175/MWR2899.1.
- 20 Schwanghart, W., Worn, R., Huggel, C., Stoffel, M., and Korup, O. (2016). Uncertainty in the Himalayan energy–  
21 water nexus: estimating regional exposure to glacial lake outburst floods. *Environ. Res. Lett.* 11, 074005.  
22 doi:10.1088/1748-9326/11/7/074005.
- 23 Schwingshackl, C., Davin, E. L., Hirschi, M., Sørland, S. L., Wartenburger, R., and Seneviratne, S. I. (2019). Regional  
24 climate model projections underestimate future warming due to missing plant physiological CO<sub>2</sub> response.  
25 *Environ. Res. Lett.* 14, 114019. doi:10.1088/1748-9326/ab4949.
- 26 Scoccimarro, E., Fogli, P. G., Reed, K. A., Gualdi, S., Masina, S., and Navarra, A. (2017). Tropical cyclone interaction  
27 with the ocean: The role of high-frequency (subdaily) coupled processes. *J. Clim.* 30, 145–162.  
28 doi:10.1175/JCLI-D-16-0292.1.
- 29 Seager, R., and Hoerling, M. (2014). Atmosphere and ocean origins of North American droughts. *J. Clim.* 27, 4581–  
30 4606. doi:10.1175/JCLI-D-13-00329.1.
- 31 Seager, R., Hoerling, M., Schubert, S., Wang, H., Lyon, B., Kumar, A., et al. (2015a). Causes of the 2011–14 California  
32 drought. *J. Clim.* 28, 6997–7024. doi:10.1175/JCLI-D-14-00860.1.
- 33 Seager, R., Hooks, A., Williams, A. P., Cook, B., Nakamura, J., and Henderson, N. (2015b). Climatology, Variability,  
34 and Trends in the U.S. Vapor Pressure Deficit, an Important Fire-Related Meteorological Quantity. *J. Appl.*  
35 *Meteorol. Climatol.* 54, 1121–1141. doi:10.1175/JAMC-D-14-0321.1.
- 36 Seager, R., Nakamura, J., and Ting, M. (2019). Mechanisms of seasonal soil moisture drought onset and termination in  
37 the southern Great Plains. *J. Hydrometeorol.* 20, 751–771. doi:10.1175/JHM-D-18-0191.1.
- 38 Sedlmeier, K., Feldmann, H., and Schädler, G. (2018). Compound summer temperature and precipitation extremes over  
39 central Europe. *Theor. Appl. Climatol.* 131, 1493–1501. doi:10.1007/s00704-017-2061-5.
- 40 Seeley, J. T., and Romps, D. M. (2015). Why does tropical convective available potential energy (CAPE) increase with  
41 warming? *Geophys. Res. Lett.* 42, 10,410–429,437. doi:10.1002/2015GL066199.
- 42 Seiler, C., and Zwiers, F. W. (2016a). How well do CMIP5 climate models reproduce explosive cyclones in the  
43 extratropics of the Northern Hemisphere? *Clim. Dyn.* 46, 1241–1256. doi:10.1007/s00382-015-2642-x.
- 44 Seiler, C., and Zwiers, F. W. (2016b). How will climate change affect explosive cyclones in the extratropics of the  
45 Northern Hemisphere? *Clim. Dyn.* 46, 3633–3644. doi:10.1007/s00382-015-2791-y.
- 46 Seiler, C., Zwiers, F. W., Hodges, K. I., and Scinocca, J. F. (2018). How does dynamical downscaling affect model  
47 biases and future projections of explosive extratropical cyclones along North America’s Atlantic coast? *Clim.*  
48 *Dyn.* 50, 677–692. doi:10.1007/s00382-017-3634-9.
- 49 Sekizawa, S., Miyasaka, T., Nakamura, H., Shimpo, A., Takemura, K., and Maeda, S. (2019). Anomalous Moisture  
50 Transport and Oceanic Evaporation during a Torrential Rainfall Event over Western Japan in Early July 2018.  
51 *SOLA* 15A, 25–30. doi:10.2151/sola.15A-005.
- 52 Selten, F. M., Bintanja, R., Vautard, R., and van den Hurk, B. J. J. M. (2020). Future continental summer warming  
53 constrained by the present-day seasonal cycle of surface hydrology. *Sci. Rep.* 10, 4721. doi:10.1038/s41598-020-  
54 61721-9.
- 55 Sen Roy, S., and Rouault, M. (2013). Spatial patterns of seasonal scale trends in extreme hourly precipitation in South  
56 Africa. *Appl. Geogr.* 39, 151–157. doi:10.1016/j.apgeog.2012.11.022.
- 57 Şen, Z. (2018). *Flood Modeling, Prediction and Mitigation*. Cham, Switzerland: Springer doi:10.1007/978-3-319-  
58 52356-9.
- 59 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., et al. (2010). Investigating soil moisture-  
60 climate interactions in a changing climate: A review. *Earth-Science Rev.* 99, 125–161.  
61 doi:10.1016/j.earscirev.2010.02.004.



- 1 Seneviratne, S. I., Donat, M. G., Mueller, B., and Alexander, L. V. (2014). No pause in the increase of hot temperature  
2 extremes. *Nat. Clim. Chang.* 4, 161–163. doi:10.1038/nclimate2145.
- 3 Seneviratne, S. I., Donat, M. G., Pitman, A. J., Knutti, R., and Wilby, R. L. (2016). Allowable CO2 emissions based on  
4 regional and impact-related climate targets. *Nature* 529, 477–483. doi:10.1038/nature16542.
- 5 Seneviratne, S. I., and Hauser, M. (2020). Regional climate sensitivity of climate extremes in CMIP6 vs CMIP5 multi-  
6 model ensembles. *Earth's Futur.* doi:10.1029/2019EF001474.
- 7 Seneviratne, S. I., Nicholls, N., Easterling, D., Goodess, C. M., Kanae, S., Kossin, J., et al. (2012). “Changes in Climate  
8 Extremes and their Impacts on the Natural Physical Environment,” in *Managing the Risks of Extreme Events and*  
9 *Disasters to Advance Climate Change Adaptation*, eds. C. B. Field, V. Barros, T. F. Stocker, and Q. Dahe  
10 (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 109–230.  
11 doi:10.1017/CBO9781139177245.006.
- 12 Seneviratne, S. I., Rogelj, J., S  ferian, R., Wartenburger, R., Allen, M. R., Cain, M., et al. (2018a). The many possible  
13 climates from the Paris Agreement’s aim of 1.5  C warming. *Nature* 558, 41–49. doi:10.1038/s41586-018-0181-  
14 4.
- 15 Seneviratne, S. I., Wartenburger, R., Guillod, B. P., Hirsch, A. L., Vogel, M. M., Brovkin, V., et al. (2018b). Climate  
16 extremes, land–climate feedbacks and land-use forcing at 1.5 C. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*  
17 376, 20160450. doi:10.1098/rsta.2016.0450.
- 18 Seneviratne, S. I., Wilhelm, M., Stanelle, T., Van Den Hurk, B., Hagemann, S., Berg, A., et al. (2013). Impact of soil  
19 moisture-climate feedbacks on CMIP5 projections: First results from the GLACE-CMIP5 experiment. *Geophys.*  
20 *Res. Lett.* 40, 5212–5217. doi:10.1002/grl.50956.
- 21 Seo, Y.-W., Kim, H., Yun, K.-S., Lee, J.-Y., Ha, K.-J., and Moon, J.-Y. (2014). Future change of extreme temperature  
22 climate indices over East Asia with uncertainties estimation in the CMIP5. *Asia-Pacific J. Atmos. Sci.* 50, 609–  
23 624. doi:10.1007/s13143-014-0050-5.
- 24 Seong, M.-G., Min, S.-K., Kim, Y.-H., Zhang, X., and Sun, Y. (2020). Anthropogenic Greenhouse Gas and Aerosol  
25 Contributions to Extreme Temperature Changes During 1951–2015. *J. Clim.*, 1–41. doi:10.1175/JCLI-D-19-  
26 1023.1.
- 27 Serrano-Notivol, R., Beguer  a, S., Saz, M.   ., and de Luis, M. (2018). Recent trends reveal decreasing intensity of  
28 daily precipitation in Spain. *Int. J. Climatol.* 38, 4211–4224. doi:10.1002/joc.5562.
- 29 Sevanto, S., McDowell, N. G., Dickman, L. T., Pangle, R., and Pockman, W. T. (2014). How do trees die? A test of the  
30 hydraulic failure and carbon starvation hypotheses. *Plant, Cell Environ.* 37, 153–161. doi:10.1111/pce.12141.
- 31 Shaevitz, D. A., Camargo, S. J., Sobel, A. H., Jonas, J. A., Kim, D., Kumar, A., et al. (2014). Characteristics of tropical  
32 cyclones in high-resolution models in the present climate. *J. Adv. Model. Earth Syst.* 6, 1154–1172.  
33 doi:10.1002/2014MS000372.
- 34 Shanas, P. R., and Kumar, V. S. (2015). Trends in surface wind speed and significant wave height as revealed by ERA-  
35 Interim wind wave hindcast in the Central Bay of Bengal. *Int. J. Climatol.* 35, 2654–2663. doi:10.1002/joc.4164.
- 36 Sharafati, A., Nabaei, S., and Shahid, S. (2020). Spatial assessment of meteorological drought features over different  
37 climate regions in Iran. *Int. J. Climatol.* 40, 1864–1884. doi:10.1002/joc.6307.
- 38 Sharma, A., Wasko, C., and Lettenmaier, D. P. (2018). If precipitation extremes are increasing, why aren’t floods?  
39 *Water Resour. Res.* 54, 8545–8551. doi:10.1029/2018WR023749.
- 40 Sharma, S., and Mujumdar, P. (2017). Increasing frequency and spatial extent of concurrent meteorological droughts  
41 and heatwaves in India. *Sci. Rep.* 7, 15582. doi:10.1038/s41598-017-15896-3.
- 42 Sharmila, S., and Walsh, K. J. E. (2018). Recent poleward shift of tropical cyclone formation linked to Hadley cell  
43 expansion. *Nat. Clim. Chang.* doi:10.1038/s41558-018-0227-5.
- 44 Shashikanth, K., Ghosh, S., H, V., and Karmakar, S. (2018). Future projections of Indian summer monsoon rainfall  
45 extremes over India with statistical downscaling and its consistency with observed characteristics. *Clim. Dyn.* 51,  
46 1–15. doi:10.1007/s00382-017-3604-2.
- 47 Shastri, H., Paul, S., Ghosh, S., and Karmakar, S. (2015). Impacts of urbanization on Indian summer monsoon rainfall  
48 extremes. *J. Geophys. Res. Atmos.* 120, 496–516. doi:10.1002/2014JD022061.
- 49 Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y.-T., et al. (2016). Storm track  
50 processes and the opposing influences of climate change. *Nat. Geosci.* 9, 656–664. doi:10.1038/ngeo2783.
- 51 Sheffield, J., Wood, E. F., and Roderick, M. L. (2012). Little change in global drought over the past 60 years. *Nature*  
52 491, 435–438. doi:10.1038/nature11575.
- 53 Sheikh, M. M., Manzoor, N., Ashraf, J., Adnan, M., Collins, D., Hameed, S., et al. (2015). Trends in extreme daily  
54 rainfall and temperature indices over South Asia. *Int. J. Climatol.* 35, 1625–1637. doi:10.1002/joc.4081.
- 55 Shephard, M. W., Mekis, E., Morris, R. J., Feng, Y., Zhang, X., Kilcup, K., et al. (2014). Trends in Canadian Short-  
56 Duration Extreme Rainfall: Including an Intensity–Duration–Frequency Perspective. *Atmosphere-Ocean* 52, 398–  
57 417. doi:10.1080/07055900.2014.969677.
- 58 Shepherd, J. M. (2013). Impacts of Urbanization on Precipitation and Storms: Physical Insights and Vulnerabilities.  
59 *Clim. Vulnerability*, 109–125. doi:10.1016/B978-0-12-384703-4.00503-7.
- 60 Shepherd, T. G. (2014). Atmospheric circulation as a source of uncertainty in climate change projections. *Nat. Geosci.*  
61 7, 703–708. doi:10.1038/NCEO2253.

- 1 Shepherd, T. G. (2016). A Common Framework for Approaches to Extreme Event Attribution. *Curr. Clim. Chang.*  
2 *Reports* 2, 28–38. doi:10.1007/s40641-016-0033-y.
- 3 Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., et al. (2018). Storylines: an  
4 alternative approach to representing uncertainty in physical aspects of climate change. *Clim. Change* 151, 555–  
5 571. doi:10.1007/s10584-018-2317-9.
- 6 Sheridan, S. C., Lee, C. C., and Smith, E. T. (2020). A Comparison Between Station Observations and Reanalysis Data  
7 in the Identification of Extreme Temperature Events. *Geophys. Res. Lett.* 47. doi:10.1029/2020GL088120.
- 8 Sherwood, S. C., Bony, S., and Dufresne, J.-L. (2014). Spread in model climate sensitivity traced to atmospheric  
9 convective mixing. *Nature* 505, 37–42. doi:10.1038/nature12829.
- 10 Sherwood, S., and Fu, Q. (2014). A Drier Future? *Science* (80-. ). 343, 737–739. doi:10.1126/science.1247620.
- 11 Shevchenko, O., Lee, H., Snizhko, S., and Mayer, H. (2014). Long-term analysis of heat waves in Ukraine. *Int. J.*  
12 *Climatol.* 34, 1642–1650. doi:10.1002/joc.3792.
- 13 Shi, C., Jiang, Z.-H., Chen, W.-L., and Li, L. (2018). Changes in temperature extremes over China under 1.5 °C and 2  
14 °C global warming targets. *Adv. Clim. Chang. Res.* 9, 120–129. doi:10.1016/j.accre.2017.11.003.
- 15 Shi, L., Feng, P., Wang, B., Liu, D. L., and Yu, Q. (2020). Quantifying future drought change and associated  
16 uncertainty in southeastern Australia with multiple potential evapotranspiration models. *J. Hydrol.* 590, 125394.  
17 doi:https://doi.org/10.1016/j.jhydrol.2020.125394.
- 18 Shi, Y., Wang, G., and Gao, X. (2017). Role of resolution in regional climate change projections over China. *Clim.*  
19 *Dyn.* 51, 2375–2396. doi:10.1007/s00382-017-4018-x.
- 20 Shimpou, A., Takemura, K., Wakamatsu, S., Togawa, H., Mochizuki, Y., Takekawa, M., et al. (2019a). Primary Factors  
21 behind the Heavy Rain Event of July 2018 and the Subsequent Heat Wave in Japan. *SOLA* 15A, 13–18.  
22 doi:10.2151/sola.15A-003.
- 23 Shimpou, A., Takemura, K., Wakamatsu, S., Togawa, H., Mochizuki, Y., Takekawa, M., et al. (2019b). Primary Factors  
24 behind the Heavy Rain Event of July 2018 and the Subsequent Heat Wave in Japan. *SOLA* 15A, 13–18.  
25 doi:10.2151/sola.15A-003.
- 26 Shin, J., Olson, R., and An, S.-I. (2018). Projected Heat Wave Characteristics over the Korean Peninsula During the  
27 Twenty-First Century. *Asia-Pacific J. Atmos. Sci.* 54, 53–61. doi:10.1007/s13143-017-0059-7.
- 28 Shiogama, H., Hirata, R., Hasegawa, T., Fujimori, S., Ishizaki, N. N., Chatani, S., et al. (2020). Historical and future  
29 anthropogenic warming effects on droughts, fires and fire emissions of CO<sub>2</sub> and PM<sub>2.5</sub> in equatorial Asia when  
30 2015-like El Niño events occur. *Earth Syst. Dyn.* 11, 435–445. doi:10.5194/esd-11-435-2020.
- 31 Shiogama, H., Imada, Y., Mori, M., Mizuta, R., Stone, D., Yoshida, K., et al. (2016). Attributing Historical Changes in  
32 Probabilities of Record-Breaking Daily Temperature and Precipitation Extreme Events. *SOLA* 12, 225–231.  
33 doi:10.2151/sola.2016-045.
- 34 Shiogama, H., Watanabe, M., Imada, Y., Mori, M., Ishii, M., and Kimoto, M. (2013). An event attribution of the 2010  
35 drought in the South Amazon region using the MIROC5 model. *Atmos. Sci. Lett.* 14, 170–175.  
36 doi:10.1002/asl2.435.
- 37 Shukla, S., Safeeq, M., AghaKouchak, A., Guan, K., and Funk, C. (2015). Temperature impacts on the water year 2014  
38 drought in California. *Geophys. Res. Lett.* 42, 4384–4393. doi:10.1002/2015GL063666.
- 39 Sigmond, M., Fyfe, J. C., and Swart, N. C. (2018). Ice-free Arctic projections under the Paris Agreement. *Nat. Clim.*  
40 *Chang.* 8, 404–408. doi:10.1038/s41558-018-0124-y.
- 41 Sikorska, A. E., Viviroli, D., and Seibert, J. (2015). Flood-type classification in mountainous catchments using crisp  
42 and fuzzy decision trees. *Water Resour. Res.* 51, 7959–7976. doi:10.1002/2015WR017326.
- 43 Sillmann, J., Donat, M. G., Fyfe, J. C., and Zwiers, F. W. (2014). Observed and simulated temperature extremes during  
44 the recent warming hiatus. *Environ. Res. Lett.* 9, 64023–64029. doi:10.1088/1748-9326/9/6/064023.
- 45 Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F. W., and Bronaugh, D. (2013a). Climate extremes indices in the  
46 CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate. *J. Geophys. Res. Atmos.* 118,  
47 1716–1733. doi:10.1002/jgrd.50203.
- 48 Sillmann, J., Kharin, V. V., Zwiers, F. W., Zhang, X., and Bronaugh, D. (2013b). Climate extremes indices in the  
49 CMIP5 multimodel ensemble: Part 2. Future climate projections. *J. Geophys. Res. Atmos.* 118, 2473–2493.  
50 doi:10.1002/jgrd.50188.
- 51 Sillmann, J., Stjern, C. W., Myhre, G., and Forster, P. M. (2017a). Slow and fast responses of mean and extreme  
52 precipitation to different forcing in CMIP5 simulations. *Geophys. Res. Lett.* 44, 6383–6390.  
53 doi:10.1002/2017GL073229.
- 54 Sillmann, J., Stjern, C. W., Myhre, G., Samset, B. H., Hodnebrog, Ø., Andrews, T., et al. (2019). Extreme wet and dry  
55 conditions affected differently by greenhouse gases and aerosols. *npj Clim. Atmos. Sci.* 2, 24.  
56 doi:10.1038/s41612-019-0079-3.
- 57 Sillmann, J., Thorarindottir, T., Keenlyside, N., Schaller, N., Alexander, L. V., Hegerl, G., et al. (2017b).  
58 Understanding, modeling and predicting weather and climate extremes: Challenges and opportunities. *Weather*  
59 *Clim. Extrem.* 18, 65–74. doi:10.1016/j.wace.2017.10.003.
- 60 Simmons, A. J., Willett, K. M., Jones, P. D., Thorne, P. W., and Dee, D. P. (2010). Low-frequency variations in surface  
61 atmospheric humidity, temperature, and precipitation: Inferences from reanalyses and monthly gridded

- 1 observational data sets. *J. Geophys. Res. Atmos.* 115. doi:10.1029/2009JD012442.
- 2 Singh, D., Horton, D. E., Tsiang, M., Haugen, M., Ashfaq, M., Mei, R., et al. (2014a). Severe precipitation in northern  
3 India in June 2013: Causes, historical context, and changes in probability [in “Explaining Extreme Events of 2013  
4 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S58–S61. doi:10.1175/1520-0477-95.9.S1.1.
- 5 Singh, D., Seager, R., Cook, B. I., Cane, M., Ting, M., Cook, E., et al. (2018). Climate and the Global Famine of 1876–  
6 78. *J. Clim.* 31, 9445–9467. doi:10.1175/JCLI-D-18-0159.1.
- 7 Singh, D., Tsiang, M., Rajaratnam, B., and Diffenbaugh, N. S. (2014b). Observed changes in extreme wet and dry  
8 spells during the South Asian summer monsoon season. *Nat. Clim. Chang.* 4, 456–461.  
9 doi:10.1038/nclimate2208.
- 10 Singh, K., Panda, J., Sahoo, M., and Mohapatra, M. (2019). Variability in Tropical Cyclone Climatology over North  
11 Indian Ocean during the Period 1891 to 2015. *Asia-Pacific J. Atmos. Sci.* 55, 269–287. doi:10.1007/s13143-018-  
12 0069-0.
- 13 Singh, M. S., Kuang, Z., Maloney, E. D., Hannah, W. M., and Wolding, B. O. (2017a). Increasing potential for intense  
14 tropical and subtropical thunderstorms under global warming. *Proc. Natl. Acad. Sci.* 114, 11657 LP-11662.  
15 doi:10.1073/pnas.1707603114.
- 16 Singh, M. S., and O’Gorman, P. A. (2013). Influence of entrainment on the thermal stratification in simulations of  
17 radiative-convective equilibrium. *Geophys. Res. Lett.* 40, 4398–4403. doi:10.1002/grl.50796.
- 18 Singh, S., Ghosh, S., Sahana, A. S., Vittal, H., and Karmakar, S. (2017b). Do dynamic regional models add value to the  
19 global model projections of Indian monsoon? *Clim. Dyn.* 48, 1375–1397. doi:10.1007/s00382-016-3147-y.
- 20 Singh, V., and Goyal, M. K. (2016). Changes in climate extremes by the use of CMIP5 coupled climate models over  
21 eastern Himalayas. *Environ. Earth Sci.* 75, 839. doi:10.1007/s12665-016-5651-0.
- 22 Sippel, S., El-Madany, T. S., Migliavacca, M., Mahecha, M. D., Carrara, A., Flach, M., et al. (2018). Warm Winter,  
23 Wet Spring, and an Extreme Response in Ecosystem Functioning on the Iberian Peninsula. *Bull. Am. Meteorol.*  
24 *Soc.* 99, S80–S85. doi:10.1175/BAMS-D-17-0135.1.
- 25 Sippel, S., and Otto, F. E. L. (2014). Beyond climatological extremes – assessing how the odds of hydrometeorological  
26 extreme events in South-East Europe change in a warming climate. *Clim. Change* 125, 381–398.  
27 doi:10.1007/s10584-014-1153-9.
- 28 Sippel, S., Otto, F. E. L., Flach, M., and van Oldenborgh, G. J. (2016). The role of anthropogenic warming in 2015  
29 central European heat waves. *Bull. Am. Meteorol. Soc.* 97, S51–S56. doi:10.1175/BAMS-D-16-0150.1.
- 30 Sippel, S., Zscheischler, J., Heimann, M., Lange, H., Mahecha, M. D., van Oldenborgh, G. J., et al. (2017a). Have  
31 precipitation extremes and annual totals been increasing in the world’s dry regions over the last 60 years? *Hydrol.*  
32 *Earth Syst. Sci.* 21, 441–458. doi:10.5194/hess-21-441-2017.
- 33 Sippel, S., Zscheischler, J., Heimann, M., Otto, F. E. L., Peters, J., and Mahecha, M. D. (2015). Quantifying changes in  
34 climate variability and extremes: Pitfalls and their overcoming. *Geophys. Res. Lett.* 42, 9990–9998.  
35 doi:10.1002/2015GL066307.
- 36 Sippel, S., Zscheischler, J., Mahecha, M. D., Orth, R., Reichstein, M., Vogel, M., et al. (2017b). Refining multi-model  
37 projections of temperature extremes by evaluation against land–atmosphere coupling diagnostics. *Earth Syst.*  
38 *Dyn.* 8, 387–403. doi:10.5194/esd-8-387-2017.
- 39 Siswanto, Jan van Oldenborgh, G., van der Schrier, G., Lenderink, G., and van den Hurk, B. (2015). Trends in High-  
40 Daily Precipitation Events in Jakarta and the Flooding of January 2014. *Bull. Am. Meteorol. Soc.* 96, S131–S135.  
41 doi:10.1175/BAMS-D-15-00128.1.
- 42 Skansi, M. de los M., Brunet, M., Sigró, J., Aguilar, E., Arevalo Groening, J. A., Bentancur, O. J., et al. (2013).  
43 Warming and wetting signals emerging from analysis of changes in climate extreme indices over South America.  
44 *Glob. Planet. Change* 100, 295–307. doi:10.1016/j.gloplacha.2012.11.004.
- 45 Skouggaard Kaspersen, P., Høegh Ravn, N., Arnbjerg-Nielsen, K., Madsen, H., and Drews, M. (2017). Comparison of  
46 the impacts of urban development and climate change on exposing European cities to pluvial flooding. *Hydrol.*  
47 *Earth Syst. Sci.* 21, 4131–4147. doi:10.5194/hess-21-4131-2017.
- 48 Slater, L. J., Singer, M. B., and Kirchner, J. W. (2015). Hydrologic versus geomorphic drivers of trends in flood hazard.  
49 *Geophys. Res. Lett.* 42, 370–376. doi:10.1002/2014GL062482.
- 50 Slater, L. J., and Villarini, G. (2016). Recent trends in U.S. flood risk. *Geophys. Res. Lett.* 43, 12,428–12,436.  
51 doi:10.1002/2016GL071199.
- 52 Slater, L., and Villarini, G. (2017). On the impact of gaps on trend detection in extreme streamflow time series. *Int. J.*  
53 *Climatol.* 37, 3976–3983. doi:10.1002/joc.4954.
- 54 Smerdon, J. E., and Pollack, H. N. (2016). Reconstructing Earth’s surface temperature over the past 2000 years: the  
55 science behind the headlines. *Wiley Interdiscip. Rev. Clim. Chang.* 7, 746–771. doi:10.1002/wcc.418.
- 56 Smiatek, G., Kunstmann, H., and Senatore, A. (2016). EURO-CORDEX regional climate model analysis for the  
57 Greater Alpine Region: Performance and expected future change. *J. Geophys. Res. Atmos.* 121, 7710–7728.  
58 doi:10.1002/2015JD024727.
- 59 Smith, B. K., Smith, J. A., Baeck, M. L., Villarini, G., and Wright, D. B. (2013). Spectrum of storm event hydrologic  
60 response in urban watersheds. *Water Resour. Res.* 49, 2649–2663. doi:10.1002/wrcr.20223.
- 61 Sobel, A. H., and Camargo, S. J. (2011). Projected Future Seasonal Changes in Tropical Summer Climate. *J. Clim.* 24,

- 1 473–487. doi:10.1175/2010JCLI3748.1.
- 2 Sobel, A. H., Camargo, S. J., Hall, T. M., Lee, C.-Y., Tippett, M. K., and Wing, A. A. (2016). Human influence on  
3 tropical cyclone intensity. *Science* (80-. ). 353, 242 LP-246. doi:10.1126/science.aaf6574.
- 4 Sobel, A. H., Camargo, S. J., and Previdi, M. (2019). Aerosol versus greenhouse gas effects on tropical cyclone  
5 potential intensity and the hydrologic cycle. *J. Clim.* 32, 5511–5527. doi:10.1175/JCLI-D-18-0357.1.
- 6 Sohn, B. J., Ryu, G.-H., Song, H.-J., and Ou, M.-L. (2013). Characteristic Features of Warm-Type Rain Producing  
7 Heavy Rainfall over the Korean Peninsula Inferred from TRMM Measurements. *Mon. Weather Rev.* 141, 3873–  
8 3888. doi:10.1175/MWR-D-13-00075.1.
- 9 Sohrabi, M. M., Ryu, J. H., Abatzoglou, J., and Tracy, J. (2015). Development of soil moisture drought index to  
10 characterize droughts. *J. Hydrol. Eng.* 20. doi:10.1061/(ASCE)HE.1943-5584.0001213.
- 11 Solander, K. C., Newman, B. D., de Araujo, A., Barnard, H. R., Berry, Z. C., Bonal, D., et al. (2020). The pantropical  
12 response of soil moisture to El Niño. *Hydrol. Earth Syst. Sci.* 24, 2303–2322. doi:10.5194/hess-24-2303-2020.
- 13 Song, J., and Klotzbach, P. J. (2018). What Has Controlled the Poleward Migration of Annual Averaged Location of  
14 Tropical Cyclone Lifetime Maximum Intensity Over the Western North Pacific Since 1961? *Geophys. Res. Lett.*  
15 45, 1148–1156. doi:10.1002/2017GL076883.
- 16 Song, X., Song, Y., and Chen, Y. (2020). Secular trend of global drought since 1950. *Environ. Res. Lett.* 15, 094073.  
17 doi:10.1088/1748-9326/aba20d.
- 18 Song, X., Zhang, Z., Chen, Y., Wang, P., Xiang, M., Shi, P., et al. (2014). Spatiotemporal changes of global extreme  
19 temperature events (ETEs) since 1981 and the meteorological causes. 70, 975–994. doi:10.1007/s11069-013-  
20 0856-y.
- 21 Sonkoué, D., Monkam, D., Fotso-Nguemo, T. C., Yepdo, Z. D., and Vondou, D. A. (2019). Evaluation and projected  
22 changes in daily rainfall characteristics over Central Africa based on a multi-model ensemble mean of CMIP5  
23 simulations. *Theor. Appl. Climatol.* 137, 2167–2186. doi:10.1007/s00704-018-2729-5.
- 24 Sorribas, M. V., Paiva, R. C. D., Melack, J. M., Bravo, J. M., Jones, C., Carvalho, L., et al. (2016). Projections of  
25 climate change effects on discharge and inundation in the Amazon basin. *Clim. Change* 136, 555–570.  
26 doi:10.1007/s10584-016-1640-2.
- 27 Sousa, P. M., Trigo, R. M., Barriopedro, D., Soares, P. M. M., Ramos, A. M., and Liberato, M. L. R. (2017). Responses  
28 of European precipitation distributions and regimes to different blocking locations. *Clim. Dyn.* 48, 1141–1160.  
29 doi:10.1007/s00382-016-3132-5.
- 30 Sparrow, S., Su, Q., Tian, F., Li, S., Chen, Y., Chen, W., et al. (2018). Attributing human influence on the July 2017  
31 Chinese heatwave: the influence of sea-surface temperatures. *Environ. Res. Lett.* 13, 114004. doi:10.1088/1748-  
32 9326/aae356.
- 33 Spennemann, P. C., Fernández-Long, M. E., Gattinoni, N. N., Cammalleri, C., and Naumann, G. (2020). Soil moisture  
34 evaluation over the Argentine Pampas using models, satellite estimations and in-situ measurements. *J. Hydrol.*  
35 *Reg. Stud.* 31. doi:10.1016/j.ejrh.2020.100723.
- 36 Sperry, J. S., Wang, Y., Wolfe, B. T., Mackay, D. S., Anderegg, W. R. L., McDowell, N. G., et al. (2016). Pragmatic  
37 hydraulic theory predicts stomatal responses to climatic water deficits. *New Phytol.* 212, 577–589.  
38 doi:10.1111/nph.14059.
- 39 Spinoni, J., Barbosa, P., Buchignani, E., Cassano, J., Cavazos, T., Christensen, J. H., et al. (2020). Future Global  
40 Meteorological Drought Hot Spots: A Study Based on CORDEX Data. *J. Clim.* 33, 3635–3661.  
41 doi:10.1175/JCLI-D-19-0084.1.
- 42 Spinoni, J., Barbosa, P., De Jager, A., McCormick, N., Naumann, G., Vogt, J. V., et al. (2019). A new global database  
43 of meteorological drought events from 1951 to 2016. *J. Hydrol. Reg. Stud.* 22, 100593.  
44 doi:10.1016/J.EJRH.2019.100593.
- 45 Spinoni, J., Naumann, G., Carrao, H., Barbosa, P., and Vogt, J. (2014). World drought frequency, duration, and severity  
46 for 1951–2010. *Int. J. Climatol.* 34, 2792–2804. doi:10.1002/joc.3875.
- 47 Spinoni, J., Naumann, G., and Vogt, J. V. (2017). Pan-European seasonal trends and recent changes of drought  
48 frequency and severity. *Glob. Planet. Change* 148, 113–130. doi:10.1016/j.gloplacha.2016.11.013.
- 49 Spinoni, J., Naumann, G., Vogt, J. V., and Barbosa, P. (2015). The biggest drought events in Europe from 1950 to  
50 2012. *J. Hydrol. Reg. Stud.* 3, 509–524. doi:10.1016/j.ejrh.2015.01.001.
- 51 Spinoni, J., Vogt, J. V., Barbosa, P., Dosio, A., McCormick, N., Bigano, A., et al. (2018a). Changes of heating and  
52 cooling degree-days in Europe from 1981 to 2100. *Int. J. Climatol.* 38, e191–e208. doi:10.1002/joc.5362.
- 53 Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., and Dosio, A. (2018b). Will drought events become more frequent  
54 and severe in Europe? *Int. J. Climatol.* 38, 1718–1736. doi:10.1002/joc.5291.
- 55 Srivastava, A., Grotjahn, R., and Ullrich, P. A. (2020). Evaluation of historical CMIP6 model simulations of extreme  
56 precipitation over contiguous US regions. *Weather Clim. Extrem.* 29, 100268. doi:10.1016/j.wace.2020.100268.
- 57 Stagge, J. H., Kingston, D. G., Tallaksen, L. M., and Hannah, D. M. (2017). Observed drought indices show increasing  
58 divergence across Europe. *Sci. Rep.* 7. doi:10.1038/s41598-017-14283-2.
- 59 Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F., and Stahl, K. (2015). Candidate Distributions for  
60 Climatological Drought Indices (SPI and SPEI). *Int. J. Climatol.* 35, 4027–4040. doi:10.1002/joc.4267.
- 61 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., Van Lanen, H. A. J., Sauquet, E., et al. (2010). Streamflow trends

- 1 in Europe: Evidence from a dataset of near-natural catchments. *Hydrol. Earth Syst. Sci.* 14, 2367–2382.  
2 doi:10.5194/hess-14-2367-2010.
- 3 Stansfield, A. M., Reed, K. A., and Zarzycki, C. M. (2020). Changes in Precipitation From North Atlantic Tropical  
4 Cyclones Under RCP Scenarios in the Variable-Resolution Community Atmosphere Model. *Geophys. Res. Lett.*  
5 47, e2019GL086930. doi:10.1029/2019GL086930.
- 6 Staudinger, M., Weiler, M., and Seibert, J. (2015). Quantifying sensitivity to droughts—an experimental modeling  
7 approach. *Hydrol. Earth Syst. Sci.* 19, 1371–1384. doi:10.5194/hess-19-1371-2015.
- 8 Stéfanon, M., Drobinski, P., D’Andrea, F., Lebeaupin-Brossier, C., and Bastin, S. (2014). Soil moisture-temperature  
9 feedbacks at meso-scale during summer heat waves over Western Europe. *Clim. Dyn.* 42, 1309–1324.  
10 doi:10.1007/s00382-013-1794-9.
- 11 Stegehuis, A. I., Vautard, R., Ciais, P., Teuling, A. J., Jung, M., and Yiou, P. (2013). Summer temperatures in Europe  
12 and land heat fluxes in observation-based data and regional climate model simulations. *Clim. Dyn.* 41, 455–477.  
13 doi:10.1007/s00382-012-1559-x.
- 14 Stennett-Brown, R. K., Jones, J. J. P., Stephenson, T. S., and Taylor, M. A. (2017). Future Caribbean temperature and  
15 rainfall extremes from statistical downscaling. *Int. J. Climatol.* 37, 4828–4845. doi:10.1002/joc.5126.
- 16 Stephens, C. M., McVicar, T. R., Johnson, F. M., and Marshall, L. A. (2018). Revisiting Pan Evaporation Trends in  
17 Australia a Decade on. *Geophys. Res. Lett.* 45, 11,111–164,172. doi:10.1029/2018GL079332.
- 18 Stephenson, T. S., Vincent, L. A., Allen, T., Van Meerbeeck, C. J., McLean, N., Peterson, T. C., et al. (2014). Changes  
19 in extreme temperature and precipitation in the Caribbean region, 1961–2010. *Int. J. Climatol.* 34, 2957–2971.  
20 doi:10.1002/joc.3889.
- 21 Sterling, S. M., Ducharne, A., and Polcher, J. (2013). The impact of global land-cover change on the terrestrial water  
22 cycle. *Nat. Clim. Chang.* 3, 385–390. doi:10.1038/nclimate1690.
- 23 Stillman, S., Zeng, X., and Bosilovich, M. G. (2016). Evaluation of 22 Precipitation and 23 Soil Moisture Products over  
24 a Semiarid Area in Southeastern Arizona. *J. Hydrometeorol.* 17, 211–230. doi:10.1175/JHM-D-15-0007.1.
- 25 Stocker, B. D., Zscheischler, J., Keenan, T. F., Prentice, I. C., Peñuelas, J., and Seneviratne, S. I. (2018). Quantifying  
26 soil moisture impacts on light use efficiency across biomes. *New Phytol.* 218, 1430–1449.  
27 doi:10.1111/nph.15123.
- 28 Stoelzle, M., Stahl, K., and Weiler, M. (2013). Are streamflow recession characteristics really characteristic? *Hydrol.*  
29 *Earth Syst. Sci.* 17, 817–828. doi:10.5194/hess-17-817-2013.
- 30 Stott, P. A., Christidis, N., Otto, F. E. L., Sun, Y., Vanderlinden, J.-P., van Oldenborgh, G. J., et al. (2016). Attribution  
31 of extreme weather and climate-related events. *Wiley Interdiscip. Rev. Clim. Chang.* 7, 23–41.  
32 doi:10.1002/wcc.380.
- 33 Strandberg, G., and Kjellström, E. (2019). Climate Impacts from Afforestation and Deforestation in Europe. *Earth*  
34 *Interact.* 23, 1–27. doi:10.1175/EI-D-17-0033.1.
- 35 Stratton, R. A., Senior, C. A., Vosper, S. B., Folwell, S. S., Boutle, I. A., Earnshaw, P. D., et al. (2018). A Pan-African  
36 Convection-Permitting Regional Climate Simulation with the Met Office Unified Model: CP4-Africa. *J. Clim.* 31,  
37 3485–3508. doi:10.1175/JCLI-D-17-0503.1.
- 38 Strong, J. D. O., Vecchi, G. A., and Ginoux, P. (2018). The Climatological Effect of Saharan Dust on Global Tropical  
39 Cyclones in a Fully Coupled GCM. *J. Geophys. Res. Atmos.* 123, 5538–5559. doi:10.1029/2017JD027808.
- 40 Studholme, J., and Gulev, S. (2018). Concurrent Changes to Hadley Circulation and the Meridional Distribution of  
41 Tropical Cyclones. *J. Clim.* 31, 4367–4389. doi:10.1175/JCLI-D-17-0852.1.
- 42 Suarez-Gutierrez, L., Li, C., Müller, W. A., and Marotzke, J. (2018). Internal variability in European summer  
43 temperatures at 1.5 °C and 2 °C of global warming. *Environ. Res. Lett.* 13, 064026. doi:10.1088/1748-  
44 9326/aaba58.
- 45 Suarez-Gutierrez, L., Müller, W. A., Li, C., and Marotzke, J. (2020a). Dynamical and thermodynamical drivers of  
46 variability in European summer heat extremes. *Clim. Dyn.* 54, 4351–4366. doi:10.1007/s00382-020-05233-2.
- 47 Suarez-Gutierrez, L., Müller, W. A., Li, C., and Marotzke, J. (2020b). Hotspots of extreme heat under global warming.  
48 *Clim. Dyn.* doi:10.1007/s00382-020-05263-w.
- 49 Sugi, M., Murakami, H., and Yoshida, K. (2017). Projection of future changes in the frequency of intense tropical  
50 cyclones. *Clim. Dyn.* 49, 619–632. doi:10.1007/s00382-016-3361-7.
- 51 Sugi, M., Murakami, H., and Yoshimura, J. (2012). On the Mechanism of Tropical Cyclone Frequency Changes Due to  
52 Global Warming. *J. Meteorol. Soc. Japan* 90A, 397–408. doi:10.2151/jmsj.2012-A24.
- 53 Sugi, M., Yamada, Y., Yoshida, K., Mizuta, R., Nakano, M., Kodama, C., et al. (2020). Future changes in frequency of  
54 tropical cyclone seeds. *SOLA* 16, 70–74. doi:10.2151/sola.2020-012.
- 55 Sui, C. H., Satoh, M., and Suzuki, K. (2020). Precipitation efficiency and its role in cloud-radiative feedbacks to climate  
56 variability. *J. Meteorol. Soc. Japan* 98, 261–282. doi:10.2151/jmsj.2020-024.
- 57 Sui, C., Zhang, Z., Yu, L., Li, Y., and Song, M. (2017). Investigation of Arctic air temperature extremes at north of 60N  
58 in winter. *Acta Oceanol. Sin.* 36, 51–60. doi:10.1007/s13131-017-1137-5.
- 59 Sui, Y., Lang, X., and Jiang, D. (2018). Projected signals in climate extremes over China associated with a 2 °C global  
60 warming under two RCP scenarios. *Int. J. Climatol.* 38, e678–e697. doi:10.1002/joc.5399.
- 61 Sun, C., Jiang, Z., Li, W., Hou, Q., and Li, L. (2019a). Changes in extreme temperature over China when global

- 1 warming stabilized at 1.5 °C and 2.0 °C. *Sci. Rep.* 9, 14982. doi:10.1038/s41598-019-50036-z.
- 2 Sun, J., Wang, D., Hu, X., Ling, Z., and Wang, L. (2019b). Ongoing Poleward Migration of Tropical Cyclone  
3 Occurrence Over the Western North Pacific Ocean. *Geophys. Res. Lett.* 46, 9110–9117.  
4 doi:10.1029/2019GL084260.
- 5 Sun, Q., and Miao, C. (2018). Extreme Rainfall (R20mm, RX5day) in Yangtze–Huai, China, in June–July 2016: The  
6 Role of ENSO and Anthropogenic Climate Change. *Bull. Am. Meteorol. Soc.* 99, S102–S106.  
7 doi:10.1175/BAMS-D-17-0091.1.
- 8 Sun, Q., Miao, C., Hanel, M., Borthwick, A. G. L., Duan, Q., Ji, D., et al. (2019c). Global heat stress on health,  
9 wildfires, and agricultural crops under different levels of climate warming. *Environ. Int.* 128, 125–136.  
10 doi:10.1016/j.envint.2019.04.025.
- 11 Sun, Q., Zhang, X., Zwiers, F., Westra, S., and Alexander, L. V (2020). A global, continental and regional analysis of  
12 changes in extreme precipitation. *J. Clim.*, 1–52. doi:10.1175/JCLI-D-19-0892.1.
- 13 Sun, Q., Zwiers, F., Zhang, X., and Li, G. (2019d). A comparison of intra-annual and long-term trend scaling of  
14 extreme precipitation with temperature in a large-ensemble regional climate simulation. *J. Clim.* Under Revi.
- 15 Sun, X. B., Ren, G. Y., Shrestha, A. B., Ren, Y. Y., You, Q. L., Zhan, Y. J., et al. (2017). Changes in extreme  
16 temperature events over the Hindu Kush Himalaya during 1961–2015. *Adv. Clim. Chang. Res.* 8, 157–165.  
17 doi:10.1016/j.accr.2017.07.001.
- 18 Sun, Y., Hu, T., and Zhang, X. (2018a). Substantial Increase in Heat Wave Risks in China in a Future Warmer World.  
19 *Earth's Futur.* 6, 1528–1538. doi:10.1029/2018EF000963.
- 20 Sun, Y., Hu, T., Zhang, X., Li, C., Lu, C., Ren, G., et al. (2019e). Contribution of Global warming and Urbanization to  
21 Changes in Temperature Extremes in Eastern China. *Geophys. Res. Lett.* 46, 11426–11434.  
22 doi:10.1029/2019GL084281.
- 23 Sun, Y., Hu, T., Zhang, X., Wan, H., Stott, P., and Lu, C. (2018b). Anthropogenic Influence on the Eastern China 2016  
24 Super Cold Surge. *Bull. Am. Meteorol. Soc.* 99, S123–S127. doi:10.1175/BAMS-D-17-0092.1.
- 25 Sun, Y., Zhang, X., Zwiers, F. W., Song, L., Wan, H., Hu, T., et al. (2014). Rapid increase in the risk of extreme  
26 summer heat in Eastern China. *Nat. Clim. Chang.* 4, 1082–1085. doi:10.1038/nclimate2410.
- 27 Sun, Z., Ouyang, Z., Zhao, J., Li, S., Zhang, X., and Ren, W. (2018c). Recent rebound in observational large-pan  
28 evaporation driven by heat wave and droughts by the Lower Yellow River. *J. Hydrol.* 565, 237–247.  
29 doi:10.1016/J.JHYDROL.2018.08.014.
- 30 Sunyer, M. A., Hundecha, Y., Lawrence, D., Madsen, H., Willems, P., Martinkova, M., et al. (2015). Inter-comparison  
31 of statistical downscaling methods for projection of extreme precipitation in Europe. *Hydrol. Earth Syst. Sci.* 19,  
32 1827–1847. doi:10.5194/hess-19-1827-2015.
- 33 Supari, Tangang, F., Juneng, L., and Aldrian, E. (2017). Observed changes in extreme temperature and precipitation  
34 over Indonesia. *Int. J. Climatol.* 37, 1979–1997. doi:10.1002/joc.4829.
- 35 Supari, Tangang, F., Juneng, L., Cruz, F., Chung, J. X., Ngai, S. T., et al. (2020). Multi-model projections of  
36 precipitation extremes in Southeast Asia based on CORDEX-Southeast Asia simulations. *Environ. Res.* 184,  
37 109350. doi:10.1016/j.envres.2020.109350.
- 38 Sutton, R. T. (2018). ESD Ideas: a simple proposal to improve the contribution of IPCC WGI to the assessment and  
39 communication of climate change risks. *Earth Syst. Dynam.* 9, 1155–1158. doi:10.5194/esd-9-1155-2018.
- 40 Sutton, R. T. (2019). Climate Science Needs to Take Risk Assessment Much More Seriously. *Bull. Am. Meteorol. Soc.*  
41 100, 1637–1642. doi:10.1175/BAMS-D-18-0280.1.
- 42 Swain, D. L., Langenbrunner, B., Neelin, J. D., and Hall, A. (2018). Increasing precipitation volatility in twenty-first-  
43 century California. *Nat. Clim. Chang.* 8, 427–433. doi:10.1038/s41558-018-0140-y.
- 44 Swain, D. L., Tsiang, M., Haugen, M., Singh, D., Charland, A., Rajaratnam, B., et al. (2014). The Extraordinary  
45 California Drought of 2013/2014: Character, Context, and the Role of Climate Change [in “Explaining Extreme  
46 Events of 2013 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 95, S3–S7. doi:10.1175/1520-0477-  
47 95.9.S1.1.
- 48 Swain, S., and Hayhoe, K. (2015). CMIP5 projected changes in spring and summer drought and wet conditions over  
49 North America. *Clim. Dyn.* 44, 2737–2750. doi:10.1007/s00382-014-2255-9.
- 50 Swann, A. L. S. (2018). Plants and Drought in a Changing Climate. *Curr. Clim. Chang. Reports* 4, 192–201.  
51 doi:10.1007/s40641-018-0097-y.
- 52 Swann, A. L. S., Hoffman, F. M., Koven, C. D., and Randerson, J. T. (2016). Plant responses to increasing CO2 reduce  
53 estimates of climate impacts on drought severity. *Proc. Natl. Acad. Sci.* 113, 10019–10024.  
54 doi:10.1073/pnas.1604581113.
- 55 Swierczynski, T., Lauterbach, S., Dulski, P., Delgado, J., Merz, B., and Brauer, A. (2013). Mid- to late Holocene flood  
56 frequency changes in the northeastern Alps as recorded in varved sediments of Lake Mondsee (Upper Austria).  
57 *Quat. Sci. Rev.* 80, 78–90. doi:10.1016/j.quascirev.2013.08.018.
- 58 Syafrina, A. H., Zalina, M. D., and Juneng, L. (2015). Historical trend of hourly extreme rainfall in Peninsular  
59 Malaysia. *Theor. Appl. Climatol.* 120, 259–285. doi:10.1007/s00704-014-1145-8.
- 60 Sylla, M. B., Elguindi, N., Giorgi, F., and Wisser, D. (2016). Projected robust shift of climate zones over West Africa in  
61 response to anthropogenic climate change for the late 21st century. *Clim. Change* 134, 241–253.

- 1 doi:10.1007/s10584-015-1522-z.
- 2 Szeto, K., Gysbers, P., Brimelow, J., and Stewart, R. (2015). The 2014 Extreme Flood on the Southeastern Canadian  
3 Prairies. *Bull. Am. Meteorol. Soc.* 96, S20–S24. doi:10.1175/BAMS-D-15-00110.1.
- 4 Tabari, H., and Aghajanloo, M.-B. (2013). Temporal pattern of aridity index in Iran with considering precipitation and  
5 evapotranspiration trends. *Int. J. Climatol.* 33, 396–409. doi:10.1002/joc.3432.
- 6 Tabari, H., Madani, K., and Willems, P. (2020). The contribution of anthropogenic influence to more anomalous  
7 extreme precipitation in Europe. *Environ. Res. Lett.* 15, 104077. doi:10.1088/1748-9326/abb268.
- 8 Tabari, H., and Willems, P. (2018). More prolonged droughts by the end of the century in the Middle East. *Environ.*  
9 *Res. Lett.* 13. doi:10.1088/1748-9326/aae09c.
- 10 Takahashi, C., Watanabe, M., and Mori, M. (2017). Significant Aerosol Influence on the Recent Decadal Decrease in  
11 Tropical Cyclone Activity Over the Western North Pacific. *Geophys. Res. Lett.* 44, 9496–9504.  
12 doi:10.1002/2017GL075369.
- 13 Takahashi, C., Watanabe, M., Shiogama, H., Imada, Y., and Mori, M. (2016). A Persistent Japanese Heat Wave in  
14 Early August 2015: Roles of Natural Variability and Human-Induced Warming. *Bull. Am. Meteorol. Soc.* 97,  
15 S107–S112. doi:10.1175/BAMS-D-16-0157.1.
- 16 Takayabu, I., Hibino, K., Sasaki, H., Shiogama, H., Mori, N., Shibutani, Y., et al. (2015). Climate change effects on the  
17 worst-case storm surge: a case study of Typhoon Haiyan. *Environ. Res. Lett.* 10, 064011. doi:10.1088/1748-  
18 9326/10/6/064011.
- 19 Takemi, T., and Unuma, T. (2019). Diagnosing Environmental Properties of the July 2018 Heavy Rainfall Event in  
20 Japan. *SOLA 15A*, 60–65. doi:10.2151/sola.15A-011.
- 21 Takemura, K., Wakamatsu, S., Togawa, H., Shimpo, A., Kobayashi, C., Maeda, S., et al. (2019). Extreme Moisture  
22 Flux Convergence over Western Japan during the Heavy Rain Event of July 2018. *SOLA 15A*, 49–54.  
23 doi:10.2151/sola.15A-009.
- 24 Talchabhadel, R., Karki, R., Thapa, B. R., Maharjan, M., and Parajuli, B. (2018). Spatio-temporal variability of extreme  
25 precipitation in Nepal. *Int. J. Climatol.* 38, 4296–4313. doi:10.1002/joc.5669.
- 26 Tallaksen, L. M., and Stahl, K. (2014). Spatial and temporal patterns of large-scale droughts in Europe: Model  
27 dispersion and performance. *Geophys. Res. Lett.* 41, 429–434. doi:10.1002/2013GL058573.
- 28 Tamura, T., Nicholas, W. A., Oliver, T. S. N., and Brooke, B. P. (2018). Coarse-sand beach ridges at Cowley Beach,  
29 north-eastern Australia: Their formative processes and potential as records of tropical cyclone history.  
30 *Sedimentology* 65, 721–744. doi:10.1111/sed.12402.
- 31 Tandon, N. F., Zhang, X., and Sobel, A. H. (2018). Understanding the Dynamics of Future Changes in Extreme  
32 Precipitation Intensity. *Geophys. Res. Lett.* 45, 2870–2878. doi:10.1002/2017GL076361.
- 33 Tangang, F., Supari, S., Chung, J. X., Cruz, F., Salimun, E., Ngai, S. T., et al. (2018). Future changes in annual  
34 precipitation extremes over Southeast Asia under global warming of 2°C. *APN Sci. Bull.* 8, 6–11.  
35 doi:10.30852/sb.2018.436.
- 36 Tao, Y., Wang, W., Song, S., and Ma, J. (2018). Spatial and Temporal Variations of Precipitation Extremes and  
37 Seasonality over China from 1961–2013. *Water* 10, 719. doi:10.3390/w10060719.
- 38 Taszarek, M., Allen, J., Púčik, T., Groenemeijer, P., Czernecki, B., Kolendowicz, L., et al. (2019). A Climatology of  
39 Thunderstorms across Europe from a Synthesis of Multiple Data Sources. *J. Clim.* 32, 1813–1837.  
40 doi:10.1175/JCLI-D-18-0372.1.
- 41 Taszarek, M., Brooks, H. E., Czernecki, B., Szuster, P., and Fortuniak, K. (2018). Climatological Aspects of  
42 Convective Parameters over Europe: A Comparison of ERA-Interim and Sounding Data. *J. Clim.* 31, 4281–4308.  
43 doi:10.1175/JCLI-D-17-0596.1.
- 44 Tauvale, L., and Tsuboki, K. (2019). Characteristics of Tropical Cyclones in the Southwest Pacific. *J. Meteorol. Soc.*  
45 *Japan. Ser. II* 97, 711–731. doi:10.2151/jmsj.2019-042.
- 46 Taylor, C. M., Belusic, D., Guichard, F., Parker, D. J., Vischel, T., Bock, O., et al. (2017). Frequency of extreme  
47 Sahelian storms tripled since 1982 in satellite observations. *Nature* 544, 475–478. doi:10.1038/nature22069.
- 48 Taylor, C. M., de Jeu, R. A. M., Guichard, F., Harris, P. P., and Dorigo, W. A. (2012). Afternoon rain more likely over  
49 drier soils. *Nature* 489, 423–426. doi:10.1038/nature11377.
- 50 Taylor, M. A., Clarke, L. A., Centella, A., Bezanilla, A., Stephenson, T. S., Jones, J. J., et al. (2018). Future Caribbean  
51 Climates in a World of Rising Temperatures: The 1.5 vs 2.0 Dilemma. *J. Clim.* 31, 2907–2926.  
52 doi:10.1175/JCLI-D-17-0074.1.
- 53 Taylor, R. G., Scanlon, B., Döll, P., Rodell, M., Van Beek, R., Wada, Y., et al. (2013). Ground water and climate  
54 change. *Nat. Clim. Chang.* 3, 322–329. doi:10.1038/nclimate1744.
- 55 Tebaldi, C., Armbruster, A., Engler, H. P., and Link, R. (2020). Emulating climate extreme indices. *Environ. Res. Lett.*  
56 15. doi:10.1088/1748-9326/ab8332.
- 57 Tebaldi, C., and Knutti, R. (2018). Evaluating the accuracy of climate change pattern emulation for low warming  
58 targets. *Environ. Res. Lett.* 13, 055006. doi:10.1088/1748-9326/aabef2.
- 59 Tebaldi, C., and Wehner, M. F. (2018). Benefits of mitigation for future heat extremes under RCP4.5 compared to  
60 RCP8.5. *Clim. Change* 146, 349–361. doi:10.1007/s10584-016-1605-5.
- 61 Tencer, B.; Rusticucci, M. (2012). Analysis of interdecadal variability of temperature extreme events in Argentina

- 1 applying EVT. *Atmosfera* 25, 327–337.
- 2 Tencer, B., Bettolli, M., and Rusticucci, M. (2016). Compound temperature and precipitation extreme events in  
3 southern South America: associated atmospheric circulation, and simulations by a multi-RCM ensemble. *Clim.*  
4 *Res.* 68, 183–199. doi:10.3354/cr01396.
- 5 Tennille, S. A., and Ellis, K. N. (2017). Spatial and temporal trends in the location of the lifetime maximum intensity of  
6 tropical cyclones. *Atmosphere (Basel)*. 8, 1–9. doi:10.3390/atmos8100198.
- 7 Teufel, B., Diro, G. T., Whan, K., Milrad, S. M., Jeong, D. I., Ganji, A., et al. (2017). Investigation of the 2013 Alberta  
8 flood from weather and climate perspectives. *Clim. Dyn.* 48, 2881–2899. doi:10.1007/s00382-016-3239-8.
- 9 Teufel, B., Sushama, L., Huziy, O., Diro, G. T., Jeong, D. I., Winger, K., et al. (2019). Investigation of the mechanisms  
10 leading to the 2017 Montreal flood. *Clim. Dyn.* 52, 4193–4206. doi:10.1007/s00382-018-4375-0.
- 11 Teuling, A. J. (2018). A hot future for European droughts. *Nat. Clim. Chang.* 8, 364–365. doi:10.1038/s41558-018-  
12 0154-5.
- 13 Teuling, A. J., de Badts, E., Jansen, F. A., Fuchs, R., Buitink, J., van Dijke, A. J., et al. (2019). Climate change,  
14 reforestation/afforestation, and urbanisation impacts on evapotranspiration and streamflow in Europe. *Hydrol.*  
15 *Earth Syst. Sci.* 23, 3631–3652. doi:10.5194/hess-2018-634.
- 16 Teuling, A. J., Van Loon, A. F., Seneviratne, S. I., Lehner, I., Aubinet, M., Heinesch, B., et al. (2013).  
17 Evapotranspiration amplifies European summer drought. *Geophys. Res. Lett.* 40, 2071–2075.  
18 doi:10.1002/grl.50495.
- 19 Thackeray, C. W., DeAngelis, A. M., Hall, A., Swain, D. L., and Qu, X. (2018). On the Connection Between Global  
20 Hydrologic Sensitivity and Regional Wet Extremes. *Geophys. Res. Lett.* 45. doi:10.1029/2018GL079698.
- 21 Thiery, W., Davin, E. L., Lawrence, D. M., Hirsch, A. L., Hauser, M., and Seneviratne, S. I. (2017). Present-day  
22 irrigation mitigates heat extremes. *J. Geophys. Res. Atmos.* 122, 1403–1422. doi:10.1002/2016JD025740.
- 23 Thiery, W., Davin, E. L., Seneviratne, S. I., Bedka, K., Lhermitte, S., and van Lipzig, N. P. M. (2016). Hazardous  
24 thunderstorm intensification over Lake Victoria. *Nat. Commun.* 7, 12786. doi:10.1038/ncomms12786.
- 25 Thiery, W., Visser, A. J., Fischer, E. M., Hauser, M., Hirsch, A. L., Lawrence, D. M., et al. (2020). Warming of hot  
26 extremes alleviated by expanding irrigation. *Nat. Commun.* 11, 290. doi:10.1038/s41467-019-14075-4.
- 27 Thober, S., Kumar, R., Wanders, N., Marx, A., Pan, M., Rakovec, O., et al. (2018). Multi-model ensemble projections  
28 of European river floods and high flows at 1.5, 2, and 3 degrees global warming. *Environ. Res. Lett.* 13, 014003.  
29 doi:10.1088/1748-9326/aa9e35.
- 30 Thorarindottir, T. L., Sillmann, J., Haugen, M., Gissibl, N., and Sandstad, M. (2020). Evaluation of CMIP5 and  
31 CMIP6 simulations of historical surface air temperature extremes using proper evaluation methods. *Environ. Res.*  
32 *Lett.* 15, 124041. doi:10.1088/1748-9326/abc778.
- 33 Tian, X., Shu, L., Wang, M., and Zhao, F. (2017). The impact of climate change on fire risk in Daxing'anling, China. *J.*  
34 *For. Res.* 28, 997–1006. doi:10.1007/s11676-017-0383-x.
- 35 Tisdeman, E., Hannaford, J., and Stahl, K. (2018). Human influences on streamflow drought characteristics in England  
36 and Wales. *Hydrol. Earth Syst. Sci.* 22, 1051–1064. doi:10.5194/hess-22-1051-2018.
- 37 Tilinina, N., Gulev, S. K., Rudeva, I., and Koltermann, P. (2013). Comparing cyclone life cycle characteristics and their  
38 interannual variability in different reanalyses. *J. Clim.* doi:10.1175/JCLI-D-12-00777.1.
- 39 Timmermans, B., Patricola, C., and Wehner, M. (2018). Simulation and analysis of extreme hurricane-driven wave  
40 climate under two ocean warming scenarios. *Oceanography* 31, 88–99. doi:10.5670/oceanog.2018.218.
- 41 Timmermans, B., Stone, D., Wehner, M., and Krishnan, H. (2017). Impact of tropical cyclones on modeled extreme  
42 wind-wave climate. *Geophys. Res. Lett.* 44, 1393–1401. doi:10.1002/2016GL071681.
- 43 Timmermans, B., Wehner, M., Cooley, D., O'Brien, T., and Krishnan, H. (2019). An evaluation of the consistency of  
44 extremes in gridded precipitation data sets. *Clim. Dyn.* 52, 6651–6670. doi:10.1007/s00382-018-4537-0.
- 45 Ting, M., Camargo, S. J., Li, C., and Kushnir, Y. (2015). Natural and Forced North Atlantic Hurricane Potential  
46 Intensity Change in CMIP5 Models. *J. Clim.* 28, 3926–3942. doi:10.1175/JCLI-D-14-00520.1.
- 47 Ting, M., Kossin, J. P., Camargo, S. J., and Li, C. (2019). Past and Future Hurricane Intensity Change along the U.S.  
48 East Coast. *Sci. Rep.* 9, 7795. doi:10.1038/s41598-019-44252-w.
- 49 Tippett, M. K., Camargo, S. J., and Sobel, A. H. (2011). A Poisson Regression Index for Tropical Cyclone Genesis and  
50 the Role of Large-Scale Vorticity in Genesis. *J. Clim.* 24, 2335–2357. doi:10.1175/2010JCLI3811.1.
- 51 Tochimoto, E., and Niino, H. (2016). Structural and Environmental Characteristics of Extratropical Cyclones that Cause  
52 Tornado Outbreaks in the Warm Sector: A Composite Study. *Mon. Weather Rev.* 144, 945–969.  
53 doi:10.1175/MWR-D-15-0015.1.
- 54 Tochimoto, E., and Niino, H. (2018). Structure and Environment of Tornado-Spawning Extratropical Cyclones around  
55 Japan. *J. Meteorol. Soc. Japan* 96, 355–380. doi:10.2151/jmsj.2018-043.
- 56 Todzo, S., Bichet, A., and Diedhiou, A. (2020). Intensification of the hydrological cycle expected in West Africa over  
57 the 21st century. *Earth Syst. Dyn.* 11, 319–328. doi:10.5194/esd-11-319-2020.
- 58 Tokinaga, H., and Xie, S.-P. (2011). Wave- and Anemometer-Based Sea Surface Wind (WASWind) for Climate  
59 Change Analysis. *J. Clim.* 24, 267–285. doi:10.1175/2010JCLI3789.1.
- 60 Tölle, M. H., Schefczyk, L., and Gutjahr, O. (2018). Scale dependency of regional climate modeling of current and  
61 future climate extremes in Germany. *Theor. Appl. Climatol.* 134, 829–848. doi:10.1007/s00704-017-2303-6.



- 1 Tomas-Burguera, M., Vicente-Serrano, S. M., Peña-Angulo, D., Domínguez-Castro, F., Noguera, I., and El Kenawy, A.  
2 (2020). Global characterization of the varying responses of the Standardized Evapotranspiration Index (SPEI) to  
3 atmospheric evaporative demand (AED). *J. Geophys. Res. Atmos.* 125, e2020JD0330178.  
4 doi:10.1029/2020JD033017.
- 5 Tomozeiu, R., Agrillo, G., Cacciamani, C., and Pavan, V. (2014). Statistically downscaled climate change projections  
6 of surface temperature over Northern Italy for the periods 2021–2050 and 2070–2099. *Nat. Hazards* 72, 143–168.  
7 doi:10.1007/s11069-013-0552-y.
- 8 Toreti, A., Belward, A., Perez-Dominguez, I., Naumann, G., Luterbacher, J., Cronie, O., et al. (2019). The Exceptional  
9 2018 European Water Seesaw Calls for Action on Adaptation. *Earth's Futur.* 7, 652–663.  
10 doi:10.1029/2019EF001170.
- 11 Touma, D., Ashfaq, M., Nayak, M. A., Kao, S.-C., and Diffenbaugh, N. S. (2015). A multi-model and multi-index  
12 evaluation of drought characteristics in the 21st century. *J. Hydrol.* 526, 196–207.  
13 doi:10.1016/j.jhydrol.2014.12.011.
- 14 Tous, M., Zappa, G., Romero, R., Shaffrey, L., and Vidale, P. L. (2016). Projected changes in medicanes in the  
15 HadGEM3 N512 high-resolution global climate model. *Clim. Dyn.* 47, 1913–1924. doi:10.1007/s00382-015-  
16 2941-2.
- 17 Tozer, C. R., Risbey, J. S., Grose, M., Monselesan, D. P., Squire, D. T., Black, A. S., et al. (2020). A 1-Day Extreme  
18 Rainfall Event in Tasmania: Process Evaluation and Long Tail Attribution [in “Explaining Extreme Events of  
19 2018 from a Climate Perspective”]. *Bull. Am. Meteorol. Soc.* 101, S123–S128. doi:10.1175/BAMS-D-19-0219.1.
- 20 Trambly, Y., and Somot, S. (2018). Future evolution of extreme precipitation in the Mediterranean. *Clim. Change* 151,  
21 289–302. doi:10.1007/s10584-018-2300-5.
- 22 Trambly, Y., Villarini, G., and Zhang, W. (2020). Observed Changes in Flood Hazard in Africa. *Environ. Res. Lett.*  
23 doi:10.1088/1748-9326/abb90b.
- 24 Trancoso, R., Larsen, J. R., McVicar, T. R., Phinn, S. R., and McAlpine, C. A. (2017). CO<sub>2</sub>-vegetation feedbacks and  
25 other climate changes implicated in reducing base flow. *Geophys. Res. Lett.* 44, 2310–2318.  
26 doi:10.1002/2017GL072759.
- 27 Trapp, R. J., Hoogewind, K. A., and Lasher-Trapp, S. (2019). Future Changes in Hail Occurrence in the United States  
28 Determined through Convection-Permitting Dynamical Downscaling. *J. Clim.* 32, 5493–5509. doi:10.1175/JCLI-  
29 D-18-0740.1.
- 30 Trapp, R. J., Tessendorf, S. A., Godfrey, E. S., and Brooks, H. E. (2005). Tornadoes from Squall Lines and Bow  
31 Echoes. Part I: Climatological Distribution. *Weather Forecast.* 20, 23–34. doi:10.1175/WAF-835.1.
- 32 Trenary, L., DelSole, T., Doty, B., and Tippet, M. K. (2015). Was the Cold Eastern US Winter of 2014 Due to  
33 Increased Variability? *Bull. Am. Meteorol. Soc.* 96, S15–S19. doi:10.1175/BAMS-D-15-00138.1.
- 34 Trenary, L., DelSole, T., Tippet, M. K., and Doty, B. (2016). Extreme Eastern U.S. Winter of 2015 Not Symptomatic  
35 of Climate Change. *Bull. Am. Meteorol. Soc.* 97, S31–S35. doi:10.1175/BAMS-D-16-0156.1.
- 36 Trenberth, K. E., Fasullo, J. T., and Shepherd, T. G. (2015). Attribution of climate extreme events. *Nat. Clim. Chang.* 5,  
37 725–730. doi:10.1038/nclimate2657.
- 38 Trenberth, K., Lijing, C., Peter, J., Yongxin, Z., and John, F. (2018). Hurricane Harvey Links to Ocean Heat Content  
39 and Climate Change Adaptation. *Earth's Futur.* 6, 730–744. doi:10.1029/2018EF000825.
- 40 Trigg, M. A., Birch, C. E., Neal, J. C., Bates, P. D., Smith, A., Sampson, C. C., et al. (2016). The credibility challenge  
41 for global fluvial flood risk analysis. *Environ. Res. Lett.* 11, 094014. doi:10.1088/1748-9326/11/9/094014.
- 42 Trinh-Tuan, L., Matsumoto, J., Tangang, F. T., Juneng, L., Cruz, F., Narisma, G., et al. (2019). Application of Quantile  
43 Mapping bias correction for mid-future precipitation projections over Vietnam. *Sci. Online Lett. Atmos.* 15, 1–6.  
44 doi:10.2151/SOLA.2019-001.
- 45 Trismidianto, and Satyawardhana, H. (2018). Mesoscale Convective Complexes (MCCs) over the Indonesian Maritime  
46 Continent during the ENSO events. *IOP Conf. Ser. Earth Environ. Sci.* 149, 012025. doi:10.1088/1755-  
47 1315/149/1/012025.
- 48 Trnka, M., Brázdil, R., Balek, J., Semerádová, D., Hlavinka, P., Možný, M., et al. (2015a). Drivers of soil drying in the  
49 Czech Republic between 1961 and 2012. *Int. J. Climatol.* 35, 2664–2675. doi:10.1002/joc.4167.
- 50 Trnka, M., Brázdil, R., Možný, M., Štěpánek, P., Dobrovolný, P., Zahradníček, P., et al. (2015b). Soil moisture trends  
51 in the Czech Republic between 1961 and 2012. *Int. J. Climatol.* 35, 3733–3747. doi:10.1002/joc.4242.
- 52 Trzeciak, T. M., Knippertz, P., Pirret, J. S. R., and Williams, K. D. (2016). Can we trust climate models to realistically  
53 represent severe European windstorms? *Clim. Dyn.* 46, 3431–3451. doi:10.1007/s00382-015-2777-9.
- 54 Tsuboki, K., Yoshioka, M. K., Shinoda, T., Kato, M., Kanada, S., and Kitoh, A. (2015). Future increase of  
55 supertyphoon intensity associated with climate change. *Geophys. Res. Lett.* 42, 646–652.  
56 doi:10.1002/2014GL061793.
- 57 Tsuguti, H., Seino, N., Kawase, H., Imada, Y., Nakaegawa, T., and Takayabu, I. (2018). Meteorological overview and  
58 mesoscale characteristics of the Heavy Rain Event of July 2018 in Japan. *Landslides* 16, 363–371.  
59 doi:10.1007/s10346-018-1098-6.
- 60 Tsuji, H., Yokoyama, C., and Takayabu, Y. N. (2019). Contrasting features of the July 2018 heavy rainfall event and  
61 the 2017 Northern Kyushu rainfall event in Japan. *J. Meteorol. Soc. Japan* in review.

- 1 Turco, M., Jerez, S., Augusto, S., Tarín-Carrasco, P., Ratola, N., Jiménez-Guerrero, P., et al. (2019). Climate drivers of the  
2 the 2017 devastating fires in Portugal. *Sci. Rep.* 9. doi:10.1038/s41598-019-50281-2.
- 3 Türkeş, M., and Erlat, E. (2018). Variability and trends in record air temperature events of Turkey and their associations  
4 with atmospheric oscillations and anomalous circulation patterns. *Int. J. Climatol.* 38, 5182–5204.  
5 doi:10.1002/joc.5720.
- 6 Tuttle, S., and Salvucci, G. (2016). Atmospheric science: Empirical evidence of contrasting soil moisture-precipitation  
7 feedbacks across the United States. *Science (80- )*. 352, 825–828. doi:10.1126/science.aaa7185.
- 8 Twardosz, R., and Kossowska-Cezak, U. (2013). Exceptionally hot summers in Central and Eastern Europe (1951-  
9 2010). *Theor. Appl. Climatol.* 112, 617–628. doi:10.1007/s00704-012-0757-0.
- 10 Udall, B., and Overpeck, J. (2017). The twenty-first century Colorado River hot drought and implications for the future.  
11 *Water Resour. Res.* 53, 2404–2418. doi:10.1002/2016WR019638.
- 12 Uhe, P., Otto, F. E. L., Haustein, K., van Oldenborgh, G. J., King, A. D., Wallom, D. C. H., et al. (2016). Comparison  
13 of methods: Attributing the 2014 record European temperatures to human influences. *Geophys. Res. Lett.* 43,  
14 8685–8693. doi:10.1002/2016GL069568.
- 15 Uhe, P., Sjoukje, P., Sarah, K., Kasturi, S., Joyce, K., Emmah, M., et al. (2017). Attributing drivers of the 2016 Kenyan  
16 drought. *Int. J. Climatol.* 38, e554–e568. doi:10.1002/joc.5389.
- 17 Ukkola, A. M., De Kauwe, M. G., Roderick, M. L., Abramowitz, G., and Pitman, A. J. (2020). Robust Future Changes  
18 in Meteorological Drought in CMIP6 Projections Despite Uncertainty in Precipitation. *Geophys. Res. Lett.* 47,  
19 e2020GL087820. doi:10.1029/2020GL087820.
- 20 Ukkola, A. M., Pitman, A. J., De Kauwe, M. G., Abramowitz, G., Herger, N., Evans, J. P., et al. (2018). Evaluating  
21 CMIP5 model agreement for multiple drought metrics. *J. Hydrometeorol.* 19, 969–988. doi:10.1175/JHM-D-17-  
22 0099.1.
- 23 Ukkola, A. M., Prentice, I. C., Keenan, T. F., Van Dijk, A. I. J. M., Viney, N. R., Myneni, R. B., et al. (2016). Reduced  
24 streamflow in water-stressed climates consistent with CO2 effects on vegetation. *Nat. Clim. Chang.* 6, 75–78.  
25 doi:10.1038/nclimate2831.
- 26 Underwood, B. S., Guido, Z., Gudipudi, P., and Feinberg, Y. (2017). Increased costs to US pavement infrastructure  
27 from future temperature rise. *Nat. Clim. Chang.* 7, 704–707. doi:10.1038/nclimate3390.
- 28 UNFCCC (2015). “Decision 1/CP.21: Adoption of the Paris Agreement,” in *Report of the Conference of the Parties on*  
29 *its twenty-first session, held in Paris from 30 November to 13 December 2015. Addendum: Part two: Action taken*  
30 *by the Conference of the Parties at its twenty-first session FCCC/CP/2015/10/Add.1.* (United Nations Framework  
31 Convention on Climate Change (UNFCCC)), 1–36. Available at: <https://unfccc.int/documents/9097>.
- 32 Utsumi, N., Kim, H., Kanae, S., and Oki, T. (2017). Relative contributions of weather systems to mean and extreme  
33 global precipitation. *J. Geophys. Res.* 122, 152–167. doi:10.1002/2016JD025222.
- 34 Valiya Veetil, A., and Mishra, A. K. (2020). Multiscale hydrological drought analysis: Role of climate, catchment and  
35 morphological variables and associated thresholds. *J. Hydrol.* 582. doi:10.1016/j.jhydrol.2019.124533.
- 36 Valverde, M. C., and Marengo, J. A. (2014). Extreme Rainfall Indices in the Hydrographic Basins of Brazil. *Open J.*  
37 *Mod. Hydrol.* 04, 10–26. doi:10.4236/ojmh.2014.41002.
- 38 van den Besselaar, E. J. M., Klein Tank, A. M. G., and Buishand, T. A. (2013). Trends in European precipitation  
39 extremes over 1951–2010. *Int. J. Climatol.* 33, 2682–2689. doi:10.1002/joc.3619.
- 40 Van Den Hurk, B., Best, M., Dirmeyer, P., Pitman, A., Polcher, J., and Santanello, J. (2011). Acceleration of land  
41 surface model development over a decade of glass. *Bull. Am. Meteorol. Soc.* 92, 1593–1600. doi:10.1175/BAMS-  
42 D-11-00007.1.
- 43 Van Den Hurk, B., Kim, H., Krinner, G., Seneviratne, S. I., Derksen, C., Oki, T., et al. (2016). LS3MIP (v1.0)  
44 contribution to CMIP6: The Land Surface, Snow and Soil moisture Model Intercomparison Project - Aims, setup  
45 and expected outcome. *Geosci. Model Dev.* 9, 2809–2832. doi:10.5194/gmd-9-2809-2016.
- 46 van den Hurk, B., Meijgaard, E. van, Valk, P. de, Heeringen, K.-J. van, and Gooijer, J. (2015). Analysis of a  
47 compounding surge and precipitation event in the Netherlands. *Environ. Res. Lett.* 10. doi:10.1088/1748-  
48 9326/10/3/035001.
- 49 Van Der Linden, E. C., Haarsma, R. J., and Van Der Schrier, G. (2019). Impact of climate model resolution on soil  
50 moisture projections in central-western Europe. *Hydrol. Earth Syst. Sci.* 23, 191–206. doi:10.5194/hess-23-191-  
51 2019.
- 52 van der Schrier, G., Barichivich, J., Briffa, K. R., and Jones, P. D. (2013). A scPDSI-based global data set of dry and  
53 wet spells for 1901-2009. *J. Geophys. Res. Atmos.* 118, 4025–4048. doi:10.1002/jgrd.50355.
- 54 van der Schrier, G., Rasmijn, L. M., Barkmeijer, J., Sterl, A., and Hazeleger, W. (2018). The 2010 Pakistan floods in a  
55 future climate. *Clim. Change* 148, 205–218. doi:10.1007/s10584-018-2173-7.
- 56 van der Wiel, K., Kapnick, S. B., van Oldenborgh, G. J., Whan, K., Philip, S., Vecchi, G. A., et al. (2017). Rapid  
57 attribution of the August 2016 flood-inducing extreme precipitation in south Louisiana to climate change. *Hydrol.*  
58 *Earth Syst. Sci.* 21, 897–921. doi:10.5194/hess-21-897-2017.
- 59 van Dijk, A. I. J. M., Beck, H. E., Crosbie, R. S., de Jeu, R. A. M., Liu, Y. Y., Podger, G. M., et al. (2013). The  
60 Millennium Drought in southeast Australia (2001–2009): Natural and human causes and implications for water  
61 resources, ecosystems, economy, and society. *Water Resour. Res.* 49, 1040–1057. doi:10.1002/wrcr.20123.

- 1 Van Huijgevoort, M. H. J., Hazenberg, P., Van Lanen, H. A. J., Teuling, A. J., Clark, D. B., Folwell, S., et al. (2013).  
2 Global multimodel analysis of drought in runoff for the second half of the twentieth century. *J. Hydrometeorol.*  
3 14, 1535–1552. doi:10.1175/JHM-D-12-0186.1.
- 4 Van Lanen, H. A. J., Laaha, G., Kingston, D. G., Gauster, T., Ionita, M., Vidal, J.-P., et al. (2016). Hydrology needed to  
5 manage droughts: the 2015 European case. *Hydrol. Process.* 30, 3097–3104. doi:10.1002/hyp.10838.
- 6 Van Lanen, H. A. J., Wanders, N., Tallaksen, L. M., and Van Loon, A. F. (2013). Hydrological drought across the  
7 world: Impact of climate and physical catchment structure. *Hydrol. Earth Syst. Sci.* 17, 1715–1732.  
8 doi:10.5194/hess-17-1715-2013.
- 9 Van Loon, A. F. (2015). Hydrological drought explained. *Wiley Interdiscip. Rev. Water* 2, 359–392.  
10 doi:10.1002/wat2.1085.
- 11 Van Loon, A. F., and Laaha, G. (2015). Hydrological drought severity explained by climate and catchment  
12 characteristics. *J. Hydrol.* 526, 3–14. doi:10.1016/j.jhydrol.2014.10.059.
- 13 Van Loon, A. F., Stahl, K., Di Baldassarre, G., Clark, J., Rangelcroft, S., Wanders, N., et al. (2016). Drought in a  
14 human-modified world: Reframing drought definitions, understanding, and analysis approaches. *Hydrol. Earth*  
15 *Syst. Sci.* 20, 3631–3650. doi:10.5194/hess-20-3631-2016.
- 16 Van Loon, A. F., and Van Lanen, H. A. J. (2012). A process-based typology of hydrological drought. *Hydrol. Earth*  
17 *Syst. Sci.* 16, 1915–1946. doi:10.5194/hess-16-1915-2012.
- 18 van Oldenborgh, G. J., Krikkken, F., Lewis, S., Leach, N. J., Lehner, F., Saunders, K. R., et al. (2021). Attribution of the  
19 Australian bushfire risk to anthropogenic climate change. *Nat. Hazards Earth Syst. Sci.* 21, 941–960.  
20 doi:10.5194/nhess-21-941-2021.
- 21 van Oldenborgh, G. J., Mitchell-Larson, E., Vecchi, G. A., de Vries, H., Vautard, R., and Otto, F. (2019). Cold waves  
22 are getting milder in the northern midlatitudes. *Environ. Res. Lett.* 14, 114004. doi:10.1088/1748-9326/ab4867.
- 23 van Oldenborgh, G. J., Otto, F. E. L., Hausteine, K., and AchutaRao, K. (2016). The Heavy Precipitation Event of  
24 December 2015 in Chennai, India. *Bull. Am. Meteorol. Soc.* 97, S87–S91. doi:10.1175/BAMS-D-16-0129.1.
- 25 van Oldenborgh, G. J., Philip, S., Kew, S., van Weele, M., Uhe, P., Otto, F., et al. (2018). Extreme heat in India and  
26 anthropogenic climate change. *Nat. Hazards Earth Syst. Sci.* 18, 365–381. doi:10.5194/nhess-18-365-2018.
- 27 van Oldenborgh, G. J., van der Wiel, K., Sebastian, A., Singh, R., Arrighi, J., Otto, F., et al. (2017). Attribution of  
28 extreme rainfall from Hurricane Harvey, August 2017. *Environ. Res. Lett.* 12, 124009. doi:10.1088/1748-  
29 9326/aa9ef2.
- 30 van Vuuren, D. P., Stehfest, E., den Elzen, M. G. J., Kram, T., van Vliet, J., Deetman, S., et al. (2011). RCP2.6:  
31 Exploring the possibility to keep global mean temperature increase below 2°C. *Clim. Change* 109, 95–116.  
32 doi:10.1007/s10584-011-0152-3.
- 33 Vanden Broucke, S., Wouters, H., Demuzere, M., and van Lipzig, N. P. M. (2019). The influence of convection-  
34 permitting regional climate modeling on future projections of extreme precipitation: dependency on topography  
35 and timescale. *Clim. Dyn.* 52, 5303–5324. doi:10.1007/s00382-018-4454-2.
- 36 Varela, V., Vlachogiannis, D., Sfetsos, A., Karozis, S., Politi, N., and Giroud, F. (2019). Projection of forest fire danger  
37 due to climate change in the French Mediterranean region. *Sustain.* 11. doi:10.3390/su11164284.
- 38 Varino, F., Arbogast, P., Joly, B., Riviere, G., Fandeur, M. L., Bovy, H., et al. (2018). Northern Hemisphere  
39 extratropical winter cyclones variability over the 20th century derived from ERA-20C reanalysis. *Clim. Dyn.*  
40 doi:10.1007/s00382-018-4176-5.
- 41 Vautard, R., Aalst, M., Boucher, O., Drouin, A., Hausteine, K., Kreienkamp, F., et al. (2020a). Human contribution to  
42 the record-breaking June and July 2019 heat waves in Western Europe. *Environ. Res. Lett.* doi:10.1088/1748-  
43 9326/aba3d4.
- 44 Vautard, R., Cattiaux, J., Yiou, P., Thépaut, J.-N., and Ciais, P. (2010). Northern Hemisphere atmospheric stilling partly  
45 attributed to an increase in surface roughness. *Nat. Geosci.* 3, 756–761. doi:10.1038/ngeo979.
- 46 Vautard, R., Gobiet, A., Jacob, D., Belda, M., Colette, A., Déqué, M., et al. (2013). The simulation of European heat  
47 waves from an ensemble of regional climate models within the EURO-CORDEX project. *Clim. Dyn.* 41, 2555–  
48 2575. doi:10.1007/s00382-013-1714-z.
- 49 Vautard, R., Kadyrov, N., Iles, C., Boberg, F., Buonomo, E., Bülow, K., et al. (2020b). Evaluation of the large EURO-  
50 CORDEX regional climate model ensemble. *J. Geophys. Res. Atmos.* doi:10.1029/2019JD032344.
- 51 Vautard, R., van Oldenborgh, G. J., Otto, F. E. L., Yiou, P., de Vries, H., van Meijgaard, E., et al. (2019). Human  
52 influence on European winter wind storms such as those of January 2018. *Earth Syst. Dynam.* 10, 271–286.  
53 doi:10.5194/esd-10-271-2019.
- 54 Vautard, R., Yiou, P., D’Andrea, F., de Noblet, N., Viovy, N., Cassou, C., et al. (2007). Summertime European heat and  
55 drought waves induced by wintertime Mediterranean rainfall deficit. *Geophys. Res. Lett.* 34.  
56 doi:10.1029/2006GL028001.
- 57 Vautard, R., Yiou, P., Otto, F., Stott, P., Christidis, N., Oldenborgh, G. J. van, et al. (2016). Attribution of human-  
58 induced dynamical and thermodynamical contributions in extreme weather events. *Environ. Res. Lett.* 11, 114009.  
59 doi:10.1088/1748-9326/11/11/114009.
- 60 Vautard, R., Yiou, P., van Oldenborgh, G.-J., Lenderink, G., Thao, S., Ribes, A., et al. (2015). Extreme Fall 2014  
61 Precipitation in the Cévennes Mountains. *Bull. Am. Meteorol. Soc.* 96, S56–S60. doi:10.1175/BAMS-D-15-

- 1 00088.1.
- 2 Veale, L., and Endfield, G. H. (2016). Situating 1816, the ‘year without summer’, in the UK. *Geogr. J.* 182, 318–330.
- 3 doi:10.1111/geoj.12191.
- 4 Vecchi, G. A., Delworth, T. L., Murakami, H., Underwood, S. D., Wittenberg, A. T., Zeng, F., et al. (2019). Tropical
- 5 cyclone sensitivities to CO<sub>2</sub> doubling: roles of atmospheric resolution, synoptic variability and background
- 6 climate changes. *Clim. Dyn.* doi:10.1007/s00382-019-04913-y.
- 7 Vecchi, G. A., and Soden, B. J. (2007). Increased tropical Atlantic wind shear in model projections of global warming.
- 8 *Geophys. Res. Lett.* 34. doi:10.1029/2006GL028905.
- 9 Velázquez, J. A., Schmid, J., Ricard, S., Muerth, M. J., Gauvin St-Denis, B., Minville, M., et al. (2013). An ensemble
- 10 approach to assess hydrological models’ contribution to uncertainties in the analysis of climate change impact on
- 11 water resources. *Hydrol. Earth Syst. Sci.* 17, 565–578. doi:10.5194/hess-17-565-2013.
- 12 Veldkamp, T. I. E., Wada, Y., Aerts, J. C. J. H., Döll, P., Gosling, S. N. N., Liu, J., et al. (2017). Water scarcity hotspots
- 13 travel downstream due to human interventions in the 20th and 21st century. *Nat. Commun.* 8, 15697.
- 14 doi:10.1038/ncomms15697.
- 15 Vetter, T., Reinhardt, J., Flörke, M., van Griensven, A., Hattermann, F., Huang, S., et al. (2017). Evaluation of sources
- 16 of uncertainty in projected hydrological changes under climate change in 12 large-scale river basins. *Clim.*
- 17 *Change* 141, 419–433. doi:10.1007/s10584-016-1794-y.
- 18 Vicente-Serrano, S. M., Domínguez-Castro, F., McVicar, T. R., Tomas-Burguera, M., Peña-Gallardo, M., Noguera, I.,
- 19 et al. (2020a). Global characterization of hydrological and meteorological droughts under future climate change:
- 20 The importance of timescales, vegetation-CO<sub>2</sub> feedbacks and changes to distribution functions. *Int. J. Climatol.*
- 21 40, 2557–2567. doi:10.1002/joc.6350.
- 22 Vicente-Serrano, S. M., Lopez-Moreno, J.-I., Beguería, S., Lorenzo-Lacruz, J., Sanchez-Lorenzo, A., García-Ruiz, J.
- 23 M., et al. (2014). Evidence of increasing drought severity caused by temperature rise in southern Europe. *Environ.*
- 24 *Res. Lett.* 9, 044001. doi:10.1088/1748-9326/9/4/044001.
- 25 Vicente-Serrano, S. M., McVicar, T. R., Miralles, D. G., Yang, Y., and Tomas-Burguera, M. (2020b). Unraveling the
- 26 influence of atmospheric evaporative demand on drought and its response to climate change. *WIREs Clim. Chang.*
- 27 11, e632. doi:10.1002/wcc.632.
- 28 Vicente-Serrano, S. M., Peña-Gallardo, M., Hannaford, J., Murphy, C., Lorenzo-Lacruz, J., Dominguez-Castro, F., et al.
- 29 (2019). Climate, Irrigation, and Land Cover Change Explain Streamflow Trends in Countries Bordering the
- 30 Northeast Atlantic. *Geophys. Res. Lett.* 46, 10821–10833. doi:10.1029/2019GL084084.
- 31 Vicente-Serrano, S. M., Quiring, S. M., Peña-Gallardo, M., Yuan, S., and Domínguez-Castro, F. (2020c). A review of
- 32 environmental droughts: Increased risk under global warming? *Earth-Science Rev.* 201, 102953.
- 33 doi:10.1016/j.earscirev.2019.102953.
- 34 Vicente-Serrano, S. M., Van der Schrier, G., Beguería, S., Azorin-Molina, C., Lopez-Moreno, J.-I. J.-I., der Schrier, G.,
- 35 et al. (2015). Contribution of precipitation and reference evapotranspiration to drought indices under different
- 36 climates. *J. Hydrol.* 526, 42–54. doi:10.1016/j.jhydrol.2014.11.025.
- 37 Vicente-Serrano, S. M., Zabalza-Martínez, J., Borràs, G., López-Moreno, J. I., Pla, E., Pascual, D., et al. (2017). Effect
- 38 of reservoirs on streamflow and river regimes in a heavily regulated river basin of Northeast Spain. *Catena* 149.
- 39 doi:10.1016/j.catena.2016.03.042.
- 40 Vicente-Serrano, S. M., Domínguez-Castro, F., Murphy, C., Hannaford, J., Reig, F., Peña-Angulo, D., et al. (2021).
- 41 Long-term variability and trends in meteorological droughts in Western Europe (1851–2018). *Int. J. Climatol.* 41,
- 42 E690–E717. doi:10.1002/joc.6719.
- 43 Vichot-Llano, A., Martínez-Castro, D., Bezanilla-Morlot, A., Centella-Artola, A., and Giorgi, F. (2021). Projected
- 44 changes in precipitation and temperature regimes and extremes over the Caribbean and Central America using a
- 45 multiparameter ensemble of RegCM4. *Int. J. Climatol.* 41, 1328–1350. doi:10.1002/joc.6811.
- 46 Vidal, J.-P., Hingray, B., Magand, C., Sauquet, E., and Ducharne, A. (2016). Hierarchy of climate and hydrological
- 47 uncertainties in transient low-flow projections. *Hydrol. Earth Syst. Sci.* 20, 3651–3672. doi:10.5194/hess-20-
- 48 3651-2016.
- 49 Vidale, P. L., Hodges, K., Vannièrè, B., Davini, P., Roberts, M. J., Strommen, K., et al. (2021). Impact of stochastic
- 50 physics and model resolution on the simulation of Tropical Cyclones in climate GCMs. *J. Clim.*, 1–85.
- 51 doi:10.1175/JCLI-D-20-0507.1.
- 52 Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tømmervik, H., Luks, B., Schuler, T. V., et al. (2016). Changes in
- 53 Winter Warming Events in the Nordic Arctic Region. *J. Clim.* 29, 6223–6244. doi:10.1175/JCLI-D-15-0763.1.
- 54 Villafuerte, M. Q., and Matsumoto, J. (2015). Significant Influences of Global Mean Temperature and ENSO on
- 55 Extreme Rainfall in Southeast Asia. *J. Clim.* 28, 1905–1919. doi:10.1175/JCLI-D-14-00531.1.
- 56 Villarini, G., Smith, J. A., and Vecchi, G. A. (2012). Changing Frequency of Heavy Rainfall over the Central United
- 57 States. *J. Clim.* 26, 351–357. doi:10.1175/JCLI-D-12-00043.1.
- 58 Vimont, D. J., and Kossin, J. P. (2007). The Atlantic Meridional Mode and hurricane activity. *Geophys. Res. Lett.* 34.
- 59 doi:10.1029/2007GL029683.
- 60 Vincent, L. A., Aguilar, E., Saindou, M., Hassane, A. F., Jumaux, G., Roy, D., et al. (2011). Observed trends in indices
- 61 of daily and extreme temperature and precipitation for the countries of the western Indian Ocean, 1961–2008. *J.*

- 1 *Geophys. Res.* 116, D10108. doi:10.1029/2010JD015303.
- 2 Vincent, L. A., Zhang, X., Mekis, É., Wan, H., and Bush, E. J. (2018). Changes in Canada's Climate: Trends in Indices  
3 Based on Daily Temperature and Precipitation Data. *Atmosphere-Ocean* 56, 332–349.  
4 doi:10.1080/07055900.2018.1514579.
- 5 Vogel, M. M., Hauser, M., and Seneviratne, S. I. (2020a). Projected changes in hot, dry and wet extreme events'  
6 clusters in CMIP6 multi-model ensemble. *Environ. Res. Lett.* 15, 094021. doi:10.1088/1748-9326/ab90a7.
- 7 Vogel, M. M., Orth, R., Cheruy, F., Hagemann, S., Lorenz, R., van den Hurk, B. J. J. M., et al. (2017). Regional  
8 amplification of projected changes in extreme temperatures strongly controlled by soil moisture-temperature  
9 feedbacks. *Geophys. Res. Lett.* 44, 1511–1519. doi:10.1002/2016GL071235.
- 10 Vogel, M. M., Zscheischler, J., Fischer, E. M., and Seneviratne, S. I. (2020b). Development of Future Heatwaves for  
11 Different Hazard Thresholds. *J. Geophys. Res. Atmos.* 125. doi:10.1029/2019JD032070.
- 12 Vogel, M. M., Zscheischler, J., and Seneviratne, S. I. (2018). Varying soil moisture-atmosphere feedbacks explain  
13 divergent temperature extremes and precipitation projections in central Europe. *Earth Syst. Dyn.* 9, 1107–1125.  
14 doi:10.5194/esd-9-1107-2018.
- 15 Vogel, M. M., Zscheischler, J., Wartenburger, R., Dee, D., and Seneviratne, S. I. (2019). Concurrent 2018 Hot  
16 Extremes Across Northern Hemisphere Due to Human-Induced Climate Change. *Earth's Futur.* 7,  
17 2019EF001189. doi:10.1029/2019EF001189.
- 18 Volosciuk, C., Maraun, D., Semenov, V. A., Tilinina, N., Gulev, S. K., and Latif, M. (2016). Rising Mediterranean Sea  
19 Surface Temperatures Amplify Extreme Summer Precipitation in Central Europe. *Sci. Rep.* 6, 32450.  
20 doi:10.1038/srep32450.
- 21 Vose, R. S., Easterling, D. R., Kunkel, K. E., LeGrande, A. N., and Wehner, M. F. (2017). "Temperature Changes in  
22 the United States," in *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, eds. D. J.  
23 Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock (Washington, DC,  
24 USA: U.S. Global Change Research Program), 185–206. doi:10.7930/J0N29V45.
- 25 Wada, Y., van Beek, L. P. H., Wanders, N., and Bierkens, M. F. P. (2013). Human water consumption intensifies  
26 hydrological drought worldwide. *Environ. Res. Lett.* 8, 034036. doi:10.1088/1748-9326/8/3/034036.
- 27 Wahl, T., Jain, S., Bender, J., Meyers, S. D., and Luther, M. E. (2015). Increasing risk of compound flooding from  
28 storm surge and rainfall for major US cities. *Nat. Clim. Chang.* 5, 1093. doi:10.1038/nclimate2736.
- 29 Waliser, D., and Guan, B. (2017). Extreme winds and precipitation during landfall of atmospheric rivers. *Nat. Geosci.*  
30 10, 179. doi:10.1038/ngeo2894.
- 31 Walsh, J. E., Ballinger, T. J., Euskirchen, E. S., Hanna, E., Mård, J., Overland, J. E., et al. (2020). Extreme weather and  
32 climate events in northern areas: A review. *Earth-Science Rev.* 209, 103324.  
33 doi:10.1016/j.earscirev.2020.103324.
- 34 Walsh, K. J. E. E., Camargo, S. J., Vecchi, G. A., Daloz, A. S., Elsner, J., Emanuel, K., et al. (2015). Hurricanes and  
35 climate: The U.S. Clivar working group on hurricanes. *Bull. Am. Meteorol. Soc.* 96, 997–1017.  
36 doi:10.1175/BAMS-D-13-00242.1.
- 37 Walsh, K. J. E., McBride, J. L., Klotzbach, P. J., Balachandran, S., Camargo, S. J., Holland, G., et al. (2016). Tropical  
38 cyclones and climate change. *Wiley Interdiscip. Rev. Clim. Chang.* 7, 65–89. doi:10.1002/wcc.371.
- 39 Wan, H., Zhang, X., and Zwiers, F. (2019). Human influence on Canadian temperatures. *Clim. Dyn.* 52, 479–494.  
40 doi:10.1007/s00382-018-4145-z.
- 41 Wanders, N., and Van Lanen, H. A. J. (2015). Future discharge drought across climate regions around the world  
42 modelled with a synthetic hydrological modelling approach forced by three general circulation models. *Nat.*  
43 *Hazards Earth Syst. Sci.* 15, 487–504. doi:10.5194/nhess-15-487-2015.
- 44 Wanders, N., and Wada, Y. (2015). Human and climate impacts on the 21st century hydrological drought. *J. Hydrol.*  
45 526, 208–220. doi:10.1016/j.jhydrol.2014.10.047.
- 46 Wang, H., Chen, Y., Xun, S., Lai, D., Fan, Y., and Li, Z. (2013a). Changes in daily climate extremes in the arid area of  
47 northwestern China. *Theor. Appl. Climatol.* 112, 15–28. doi:10.1007/s00704-012-0698-7.
- 48 Wang, H., and Schubert, S. (2014). Causes of the extreme dry conditions over California during early 2013 [in  
49 "Explaining Extreme Events of 2013 from a Climate Perspective"]. *Bull. Amer. Meteor. Soc.* 95, S7–S11.
- 50 Wang, K., Dickinson, R. E., and Liang, S. (2012). Global atmospheric evaporative demand over land from 1973 to  
51 2008. *J. Clim.* 25, 8353–8361. doi:10.1175/JCLI-D-11-00492.1.
- 52 Wang, P., Tang, J., Sun, X., Liu, J., and Juan, F. (2018a). Spatiotemporal characteristics of heat waves over China in  
53 regional climate simulations within the CORDEX-EA project. *Clim. Dyn.* doi:10.1007/s00382-018-4167-6.
- 54 Wang, P., Wu, X., Hao, Y., Wu, C., and Zhang, J. (2020). Is Southwest China drying or wetting? Spatiotemporal  
55 patterns and potential causes. *Theor. Appl. Climatol.* 139, 1–15. doi:10.1007/s00704-019-02935-4.
- 56 Wang, S.-Y. S., Zhao, L., Yoon, J.-H., Klotzbach, P., and Gillies, R. R. (2018b). Quantitative attribution of climate  
57 effects on Hurricane Harvey's extreme rainfall in Texas. *Environ. Res. Lett.* 13, 54014. doi:10.1088/1748-  
58 9326/aabb85.
- 59 Wang, W., Lee, X., Xiao, W., Liu, S., Schultz, N., Wang, Y., et al. (2018c). Global lake evaporation accelerated by  
60 changes in surface energy allocation in a warmer climate. *Nat. Geosci.* 11, 410–414. doi:10.1038/s41561-018-  
61 0114-8.

- 1 Wang, W., Zhou, W., Li, Y., Wang, X., and Wang, D. (2015). Statistical modeling and CMIP5 simulations of hot spell  
2 changes in China. *Clim. Dyn.* 44, 2859–2872. doi:10.1007/s00382-014-2287-1.
- 3 Wang, X., Jiang, D., and Lang, X. (2017a). Future extreme climate changes linked to global warming intensity. *Sci.*  
4 *Bull.* 62, 1673–1680. doi:10.1016/j.scib.2017.11.004.
- 5 Wang, X. L., Feng, Y., Chan, R., and Isaac, V. (2016). Inter-comparison of extra-tropical cyclone activity in nine  
6 reanalysis datasets. *Atmos. Res.* doi:10.1016/j.atmosres.2016.06.010.
- 7 Wang, X. L., Feng, Y., Compo, G. P., Swail, V. R., Zwiers, F. W., Allan, R. J., et al. (2013b). Trends and low  
8 frequency variability of extra-tropical cyclone activity in the ensemble of twentieth century reanalysis. *Clim. Dyn.*  
9 40, 2775–2800. doi:10.1007/s00382-012-1450-9.
- 10 Wang, X. L., Trewin, B., Feng, Y., and Jones, D. (2013c). Historical changes in Australian temperature extremes as  
11 inferred from extreme value distribution analysis. *Geophys. Res. Lett.* 40, 573–578. doi:10.1002/grl.50132.
- 12 Wang, Y., Lee, K.-H., Lin, Y., Levy, M., and Zhang, R. (2014). Distinct effects of anthropogenic aerosols on tropical  
13 cyclones. *Nat. Clim. Chang.* 4, 368. doi:10.1038/nclimate2144.
- 14 Wang, Y. W., and Yang, Y. H. (2014). China’s dimming and brightening: Evidence, causes and hydrological  
15 implications. *Ann. Geophys.* 32, 41–55. doi:10.5194/angeo-32-41-2014.
- 16 Wang, Y., Zhou, B., Qin, D., Wu, J., Gao, R., and Song, L. (2017b). Changes in mean and extreme temperature and  
17 precipitation over the arid region of northwestern China: Observation and projection. *Adv. Atmos. Sci.* 34, 289–  
18 305. doi:10.1007/s00376-016-6160-5.
- 19 Wang, Z., Jiang, Y., Wan, H., Yan, J., and Zhang, X. (2017c). Detection and Attribution of Changes in Extreme  
20 Temperatures at Regional Scale. *J. Clim.* 30, 7035–7047. doi:10.1175/JCLI-D-15-0835.1.
- 21 Wang, Z., Lin, L., Zhang, X., Zhang, H., Liu, L., and Xu, Y. (2017d). Scenario dependence of future changes in climate  
22 extremes under 1.5 °C and 2 °C global warming. *Sci. Rep.* 7, 46432. doi:10.1038/srep46432.
- 23 Ward, P. J., Couasnon, A., Eilander, D., Haigh, I. D., Hendry, A., Muis, S., et al. (2018). Dependence between high sea-  
24 level and high river discharge increases flood hazard in global deltas and estuaries. *Environ. Res. Lett.* 13,  
25 084012. doi:10.1088/1748-9326/aad400.
- 26 Wartenburger, R., Hirschi, M., Donat, M. G., Greve, P., Pitman, A. J., and Seneviratne, S. I. (2017). Changes in  
27 regional climate extremes as a function of global mean temperature: an interactive plotting framework. *Geosci.*  
28 *Model Dev. Discuss.*, 1–30. doi:10.5194/gmd-2017-33.
- 29 Wasko, C., and Nathan, R. (2019). Influence of changes in rainfall and soil moisture on trends in flooding. *J. Hydrol.*  
30 575, 432–441. doi:https://doi.org/10.1016/j.jhydrol.2019.05.054.
- 31 Wasko, C., and Sharma, A. (2017). Global assessment of flood and storm extremes with increased temperatures. *Sci.*  
32 *Rep.* 7, 7945. doi:10.1038/s41598-017-08481-1.
- 33 Watterson, I. G., J. Bathols, and Heady, C. (2014). What influences the skill of climate models over the continents?  
34 *Bull. Am. Meteorol. Soc.* 95, 689–700. doi:10.1175/BAMS-D-12-00136.1.
- 35 Weber, T., Haensler, A., Rechid, D., Pfeifer, S., Eggert, B., and Jacob, D. (2018). Analyzing Regional Climate Change  
36 in Africa in a 1.5, 2, and 3°C Global Warming World. *Earth’s Futur.* 6, 643–655. doi:10.1002/2017EF000714.
- 37 Wehner, M. F. (2020). Characterization of long period return values of extreme daily temperature and precipitation in  
38 the CMIP6 models: Part 2, projections of future change. *Weather Clim. Extrem.* 30, 100284.  
39 doi:10.1016/j.wace.2020.100284.
- 40 Wehner, M. F., Arnold, J. R., Knutson, T., Kunkel, K. E., and LeGrande, A. N. (2017). “Droughts, Floods, and  
41 Wildfires,” in *Climate Science Special Report: Fourth National Climate Assessment, Volume I*, eds. D. J.  
42 Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart, and T. K. Maycock (Washington, DC,  
43 USA: U.S. Global Change Research Program), 231–256. doi:10.7930/J0CJ8BNN.
- 44 Wehner, M. F., Reed, K. A., Li, F., Prabhat, Bacmeister, J., Chen, C.-T., et al. (2014). The effect of horizontal  
45 resolution on simulation quality in the Community Atmospheric Model, CAM5.1. *J. Adv. Model. Earth Syst.* 6,  
46 980–997. doi:10.1002/2013MS000276.
- 47 Wehner, M. F., Reed, K. A., Loring, B., Stone, D., and Krishnan, H. (2018a). Changes in tropical cyclones under  
48 stabilized 1.5 and 2.0 °C global warming scenarios as simulated by the Community Atmospheric Model under the  
49 HAPPI protocols. *Earth Syst. Dyn.* 9, 187–195. doi:10.5194/esd-9-187-2018.
- 50 Wehner, M. F., Zarzycki, C., and Patricola, C. (2019). “Estimating the Human Influence on Tropical Cyclone Intensity  
51 as the Climate Changes,” in *Hurricane Risk*, eds. J. Collins and K. Walsh (Cham, Switzerland: Springer), 235–  
52 260. doi:10.1007/978-3-030-02402-4\_12.
- 53 Wehner, M., Gleckler, P., and Lee, J. (2020). Characterization of long period return values of extreme daily temperature  
54 and precipitation in the CMIP6 models: Part 1, model evaluation. *Weather Clim. Extrem.* 30, 100283.  
55 doi:10.1016/j.wace.2020.100283.
- 56 Wehner, M., Prabhat, Reed, K. A., Stone, D., Collins, W. D., and Bacmeister, J. (2015). Resolution Dependence of  
57 Future Tropical Cyclone Projections of CAM5.1 in the U.S. CLIVAR Hurricane Working Group Idealized  
58 Configurations. *J. Clim.* 28, 3905–3925. doi:10.1175/JCLI-D-14-00311.1.
- 59 Wehner, M., Stone, D., Krishnan, H., AchutaRao, K., and Castillo, F. (2016). The Deadly Combination of Heat and  
60 Humidity in India and Pakistan in Summer 2015. *Bull. Am. Meteorol. Soc.* 97, S81–S86. doi:10.1175/BAMS-D-  
61 16-0145.1.

- 1 Wehner, M., Stone, D., Mitchell, D., Shiogama, H., Fischer, E., Graff, L. S., et al. (2018b). Changes in extremely hot  
2 days under stabilized 1.5 and 2.0°C global warming scenarios as simulated by the HAPPI multi-model ensemble.  
3 *Earth Syst. Dyn.* 9, 299–311. doi:10.5194/esd-9-299-2018.
- 4 Wehner, M., Stone, D., Shiogama, H., Wolski, P., Ciavarella, A., Christidis, N., et al. (2018c). Early 21st century  
5 anthropogenic changes in extremely hot days as simulated by the C20C+ detection and attribution multi-model  
6 ensemble. *Weather Clim. Extrem.* 20, 1–8. doi:10.1016/j.wace.2018.03.001.
- 7 Wehrli, K., Guillod, B. P., Hauser, M., Leclair, M., and Seneviratne, S. I. (2019). Identifying Key Driving Processes of  
8 Major Recent Heat Waves. *J. Geophys. Res. Atmos.* 124, 11746–11765. doi:10.1029/2019JD030635.
- 9 Wehrli, K., Hauser, M., and Seneviratne, S. I. (2020). Storylines of the 2018 Northern Hemisphere heatwave at pre-  
10 industrial and higher global warming levels. *Earth Syst. Dyn.* 11, 855–873. doi:10.5194/esd-11-855-2020.
- 11 Weldon, D., and Reason, C. J. C. (2014). Variability of rainfall characteristics over the South Coast region of South  
12 Africa. *Theor. Appl. Climatol.* 115, 177–185. doi:10.1007/s00704-013-0882-4.
- 13 Wen, Q. H., Zhang, X., Xu, Y., and Wang, B. (2013). Detecting human influence on extreme temperatures in China.  
14 *Geophys. Res. Lett.* 40, 1171–1176. doi:10.1002/grl.50285.
- 15 Wester, P., Mishra, A., Mukherji, A., and Shrestha, A. B. eds. (2019). *The Hindu Kush Himalaya Assessment:  
16 Mountains, Climate Change, Sustainability and People*. Cham, Switzerland: Springer doi:10.1007/978-3-319-  
17 92288-1.
- 18 Westra, S., Alexander, L. V., and Zwiers, F. W. (2013). Global Increasing Trends in Annual Maximum Daily  
19 Precipitation. *J. Clim.* 26, 3904–3918. doi:10.1175/JCLI-D-12-00502.1.
- 20 Westra, S., Fowler, H. J., Evans, J. P., Alexander, L. V., Berg, P., Johnson, F., et al. (2014). Future changes to the  
21 intensity and frequency of short-duration extreme rainfall. *Rev. Geophys.* 52, 522–555.  
22 doi:10.1002/2014RG000464.
- 23 Westra, S., White, C. J., and Kiem, A. S. (2016). Introduction to the special issue: historical and projected climatic  
24 changes to Australian natural hazards. *Clim. Change* 139, 1–19. doi:10.1007/s10584-016-1826-7.
- 25 Wetter, O., and Pfister, C. (2013). An underestimated record breaking event – why summer 1540 was likely warmer  
26 than 2003. 41–56. doi:10.5194/cp-9-41-2013.
- 27 Wetter, O., Pfister, C., Werner, J. P., Zorita, E., Wagner, S., Seneviratne, S. I., et al. (2014). The year-long  
28 unprecedented European heat and drought of 1540 – a worst case. *Clim. Change* 125, 349–363.  
29 doi:10.1007/s10584-014-1184-2.
- 30 Wever, N. (2012). Quantifying trends in surface roughness and the effect on surface wind speed observations. *J.  
31 Geophys. Res. Atmos.* 117, 1–14. doi:10.1029/2011JD017118.
- 32 Whan, K., Alexander, L. V., Imielska, A., Mcgree, S., Jones, D., Ene, E., et al. (2014). Trends and variability of  
33 temperature extremes in the tropical Western Pacific. *Int. J. Climatol.* 34, 2585–2603. doi:10.1002/joc.3861.
- 34 Whan, K., Zscheischler, J., Orth, R., Shongwe, M., Rahimi, M., Asare, E. O., et al. (2015). Impact of soil moisture on  
35 extreme maximum temperatures in Europe. *Weather Clim. Extrem.* 9, 57–67. doi:10.1016/j.wace.2015.05.001.
- 36 Whan, K., and Zwiers, F. (2016). Evaluation of extreme rainfall and temperature over North America in CanRCM4 and  
37 CRCM5. *Clim. Dyn.* 46, 3821–3843. doi:10.1007/s00382-015-2807-7.
- 38 Wickham, J. D., Wade, T. G., and Riitters, K. H. (2013). Empirical analysis of the influence of forest extent on annual  
39 and seasonal surface temperatures for the continental United States. *Glob. Ecol. Biogeogr.* 22, 620–629.  
40 doi:10.1111/geb.12013.
- 41 Wilcox, L. J., Yiou, P., Hauser, M., Lott, F. C., van Oldenborgh, G. J., Colfescu, I., et al. (2018). Multiple perspectives  
42 on the attribution of the extreme European summer of 2012 to climate change. *Clim. Dyn.* 50, 3537–3555.  
43 doi:10.1007/s00382-017-3822-7.
- 44 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., et al. (2005). From Dimming to Brightening:  
45 Decadal Changes in Solar Radiation at Earth’s Surface. *Science (80- )*. 308, 847–850.  
46 doi:10.1126/science.1103215.
- 47 Wilhelm, B., Ballesteros Cánovas, J. A., Macdonald, N., Toonen, W. H. J., Baker, V., Barriendos, M., et al. (2019).  
48 Interpreting historical, botanical, and geological evidence to aid preparations for future floods. *Wiley Interdiscip.  
49 Rev. Water* 6, e1318. doi:10.1002/wat2.1318.
- 50 Wilhite, D. A., and Pulwarty, R. S. (2017). “Drought as Hazard: Understanding the Natural and Social Context,” in  
51 *Drought and Water Crises: Integrating Science, Management, and Policy (2nd Edition)* (Boca Raton, FL, USA:  
52 CRC Press), 3–22. doi:10.1201/b22009.
- 53 Willems, P. (2013). Multidecadal oscillatory behaviour of rainfall extremes in Europe. *Clim. Change* 120, 931–944.  
54 doi:10.1007/s10584-013-0837-x.
- 55 Willett, K. M., Dunn, R. J. H., Thorne, P. W., Bell, S., de Podesta, M., Parker, D. E., et al. (2014). HadISDH land  
56 surface multi-variable humidity and temperature record for climate monitoring. *Clim. Past* 10, 1983–2006.  
57 doi:10.5194/cp-10-1983-2014.
- 58 Williams, A. P., Abatzoglou, J. T., Gershunov, A., Guzman-Morales, J., Bishop, D. A., Balch, J. K., et al. (2019).  
59 Observed Impacts of Anthropogenic Climate Change on Wildfire in California. *Earth’s Futur.* 7, 892–910.  
60 doi:10.1029/2019EF001210.
- 61 Williams, A. P., Allen, C. D., Macalady, A. K., Griffin, D., Woodhouse, C. A., Meko, D. M., et al. (2013). Temperature

- 1 as a potent driver of regional forest drought stress and tree mortality. *Nat. Clim. Chang.* 3, 292–297.  
2 doi:10.1038/nclimate1693.
- 3 Williams, A. P., Cook, E. R., Smerdon, J. E., Cook, B. I., Abatzoglou, J. T., Bolles, K., et al. (2020). Large contribution  
4 from anthropogenic warming to an emerging North American megadrought. *Science* (80- ). 368, 314–318.  
5 doi:10.1126/science.aaz9600.
- 6 Williams, A. P., Seager, R., Abatzoglou, J. T., Cook, B. I., Smerdon, J. E., and Cook, E. R. (2015). Contribution of  
7 anthropogenic warming to California drought during 2012–2014. *Geophys. Res. Lett.* 42, 6819–6828.  
8 doi:10.1002/2015GL064924.
- 9 Williams, A. P., Seager, R., Berkelhammer, M., Macalady, A. K., Crimmins, M. A., Swetnam, T. W., et al. (2014).  
10 Causes and Implications of Extreme Atmospheric Moisture Demand during the Record-Breaking 2011 Wildfire  
11 Season in the Southwestern United States. *J. Appl. Meteorol. Climatol.* 53, 2671–2684. doi:10.1175/JAMC-D-14-  
12 0053.1.
- 13 Willison, J., Robinson, W. A., and Lackmann, G. M. (2013). The Importance of Resolving Mesoscale Latent Heating in  
14 the North Atlantic Storm Track. *J. Atmos. Sci.* 70, 2234–2250. doi:10.1175/JAS-D-12-0226.1.
- 15 Wing, A. A., Camargo, S. J., Sobel, A. H., Kim, D., Moon, Y., Murakami, H., et al. (2019). Moist Static Energy Budget  
16 Analysis of Tropical Cyclone Intensification in High-Resolution Climate Models. *J. Clim.* 32, 6071–6095.  
17 doi:10.1175/JCLI-D-18-0599.1.
- 18 Winter, H. C., Brown, S. J., and Tawn, J. A. (2017). Characterising the changing behaviour of heatwaves with climate  
19 change. *Dyn. Stat. Clim. Syst.*, dzw006. doi:10.1093/climsys/dzw006.
- 20 Woldemeskel, F., and Sharma, A. (2016). Should flood regimes change in a warming climate? The role of antecedent  
21 moisture conditions. *Geophys. Res. Lett.* 43, 7556–7563. doi:10.1002/2016GL069448.
- 22 Woldemichael, A., Abel, T., Hossain, F., Pielke, R., and Beltrán-Przekurat, A. (2012). Understanding the impact of dam-  
23 triggered land use/land cover change on the modification of extreme precipitation. *Water Resour. Res.* 48.  
24 doi:10.1029/2011WR011684.
- 25 Wolski, P., Stone, D., Tadross, M., Wehner, M., and Hewitson, B. (2014). Attribution of floods in the Okavango basin,  
26 Southern Africa. *J. Hydrol.* 511, 350–358. doi:10.1016/j.jhydrol.2014.01.055.
- 27 Wolter, K., Eischeid, J. K., Quan, X.-W., Chase, T. n., Hoerling, M., Dole, R. M., et al. (2015). How Unusual was the  
28 Cold Winter of 2013/14 in the Upper Midwest? *Bull. Am. Meteorol. Soc.* 96, S10–S14. doi:10.1175/BAMS-D-15-  
29 00126.1.
- 30 Wood, R. R., and Ludwig, R. (2020). Analyzing Internal Variability and Forced Response of Subdaily and Daily  
31 Extreme Precipitation Over Europe. *Geophys. Res. Lett.* 47, e2020GL089300. doi:10.1029/2020GL089300.
- 32 Woodhouse, C. A., and Wise, E. K. (2020). The changing relationship between the upper and lower Missouri River  
33 basins during drought. *Int. J. Climatol.* 40, 5011–5028. doi:10.1002/joc.6502.
- 34 Woodward, C., Shulmeister, J., Larsen, J., Jacobsen, G. E., and Zawadzki, A. (2014). The hydrological legacy of  
35 deforestation on global wetlands. *Science* (80- ). 346, 844–847. doi:10.1126/science.1260510.
- 36 Woollings, T., Barriopedro, D., Methven, J., Son, S. W., Martius, O., Harvey, B., et al. (2018). Blocking and its  
37 Response to Climate Change. *Curr. Clim. Chang. Reports* 4, 287–300. doi:10.1007/s40641-018-0108-z.
- 38 Wu, J., Liu, Z., Yao, H., Chen, X., Chen, X., Zheng, Y., et al. (2018a). Impacts of reservoir operations on multi-scale  
39 correlations between hydrological drought and meteorological drought. *J. Hydrol.* 563, 726–736.  
40 doi:10.1016/j.jhydrol.2018.06.053.
- 41 Wu, J., Xu, Y., and Gao, X.-J. (2017). Projected changes in mean and extreme climates over Hindu Kush Himalayan  
42 region by 21 CMIP5 models. *Adv. Clim. Chang. Res.* 8, 176–184. doi:10.1016/j.accre.2017.03.001.
- 43 Wu, M., Nikulin, G., Kjellström, E., Belušić, D., Jones, C., and Lindstedt, D. (2020a). The impact of regional climate  
44 model formulation and resolution on simulated precipitation in Africa. *Earth Syst. Dyn.* 11, 377–394.  
45 doi:10.5194/esd-11-377-2020.
- 46 Wu, S.-Y. (2015). Changing characteristics of precipitation for the contiguous United States. *Clim. Change* 132, 677–  
47 692. doi:10.1007/s10584-015-1453-8.
- 48 Wu, W., McInnes, K., O’Grady, J., Hoeke, R., Leonard, M., and Westra, S. (2018b). Mapping Dependence Between  
49 Extreme Rainfall and Storm Surge. *J. Geophys. Res. Ocean.* 123, 2461–2474. doi:10.1002/2017JC013472.
- 50 Wu, X., Hao, Z., Tang, Q., Singh, V. P., Zhang, X., and Hao, F. (2021). Projected increase in compound dry and hot  
51 events over global land areas. *Int. J. Climatol.* 41, 393–403. doi:10.1002/joc.6626.
- 52 Wu, Y., and Polvani, L. M. (2017). Recent Trends in Extreme Precipitation and Temperature over Southeastern South  
53 America: The Dominant Role of Stratospheric Ozone Depletion in the CESM Large Ensemble. *J. Clim.* 30, 6433–  
54 6441. doi:10.1175/JCLI-D-17-0124.1.
- 55 Wu, Z., Yu, L., Du, Z., Zhang, H., Fan, X., and Lei, T. (2020b). Recent changes in the drought of China from 1960 to  
56 2014. *Int. J. Climatol.* 40, 3281–3296. doi:10.1002/joc.6397.
- 57 Wuebbles, D., Meehl, G., Hayhoe, K., Karl, T. R., Kunkel, K., Santer, B., et al. (2014). CMIP5 Climate Model  
58 Analyses: Climate Extremes in the United States. *Bull. Am. Meteorol. Soc.* 95, 571–583. doi:10.1175/BAMS-D-  
59 12-00172.1.
- 60 Wurbs, R. A., and Ayala, R. A. (2014). Reservoir evaporation in Texas, USA. *J. Hydrol.* 510, 1–9.  
61 doi:10.1016/j.jhydrol.2013.12.011.



- 1 Xia, Y., Ek, M. B., Wu, Y., Ford, T., and Quiring, S. M. (2015). Comparison of NLDAS-2 Simulated and NASMD  
2 Observed Daily Soil Moisture. Part I: Comparison and Analysis. *J. Hydrometeorol.* 16, 1962–1980.  
3 doi:10.1175/JHM-D-14-0096.1.
- 4 Xia, Y., Sheffield, J., Ek, M. B., Dong, J., Chaney, N., Wei, H., et al. (2014). Evaluation of multi-model simulated soil  
5 moisture in NLDAS-2. *J. Hydrol.* 512, 107–125. doi:10.1016/j.jhydrol.2014.02.027.
- 6 Xiao, C., Wu, P., Zhang, L., and Song, L. (2016). Robust increase in extreme summer rainfall intensity during the past  
7 four decades observed in China. *Sci. Rep.* 6, 38506. doi:10.1038/srep38506.
- 8 Xiao, K., Griffis, T. J., Baker, J. M., Bolstad, P. V., Erickson, M. D., Lee, X., et al. (2018a). Evaporation from a  
9 temperate closed-basin lake and its impact on present, past, and future water level. *J. Hydrol.* 561, 59–75.  
10 doi:10.1016/j.jhydrol.2018.03.059.
- 11 Xiao, M., Udall, B., and Lettenmaier, D. P. (2018b). On the Causes of Declining Colorado River Streamflows. *Water  
12 Resour. Res.* 54, 6739–6756. doi:10.1029/2018WR023153.
- 13 Xie, W., Zhou, B., You, Q., Zhang, Y., and Ullah, S. (2020). Observed changes in heat waves with different severities  
14 in China during 1961–2015. *Theor. Appl. Climatol.* 141, 1529–1540. doi:10.1007/s00704-020-03285-2.
- 15 Xie, X., Liang, S., Yao, Y., Jia, K., Meng, S., and Li, J. (2015). Detection and attribution of changes in hydrological  
16 cycle over the Three-North region of China: Climate change versus afforestation effect. *Agric. For. Meteorol.*  
17 203, 74–87. doi:10.1016/j.agrformet.2015.01.003.
- 18 Xin, X., Wu, T., Zhang, J., Yao, J., and Fang, Y. (2020). Comparison of CMIP6 and CMIP5 simulations of  
19 precipitation in China and the East Asian summer monsoon. *Int. J. Climatol.* doi:10.1002/joc.6590.
- 20 Xu, L., Chen, N., and Zhang, X. (2019a). Global drought trends under 1.5 and 2 °C warming. *Int. J. Climatol.* 39,  
21 2375–2385. doi:10.1002/joc.5958.
- 22 Xu, R., Yu, P., Abramson, M. J., Johnston, F. H., Samet, J. M., Bell, M. L., et al. (2020). Wildfires, Global Climate  
23 Change, and Human Health. *N. Engl. J. Med.* 383, 2173–2181. doi:10.1056/NEJMs2028985.
- 24 Xu, W., Yuan, W., Dong, W., Xia, J., Liu, D., and Chen, Y. (2013). A meta-analysis of the response of soil moisture to  
25 experimental warming. *Environ. Res. Lett.* 8, 044027. doi:10.1088/1748-9326/8/4/044027.
- 26 Xu, Y., Wu, J., Shi, Y., Zhou, B., Li, R., and Wu, J. (2016a). Change in Extreme Climate Events over China Based on  
27 CMIP5 Change in Extreme Climate Events over China Based on CMIP5. *Atmos. Ocean. Sci. Lett.* 8, 185–192.  
28 doi:10.3878/AOSL20150006.
- 29 Xu, Y., Zhang, X., Wang, X., Hao, Z., Singh, V. P., and Hao, F. (2019b). Propagation from meteorological drought to  
30 hydrological drought under the impact of human activities: A case study in northern China. *J. Hydrol.* 579.  
31 doi:10.1016/j.jhydrol.2019.124147.
- 32 Xu, Y., Zhou, B. T., Wu, J., Han, Z. Y., Zhang, Y. X., and Wu, J. (2017). Asian climate change under 1.5–4 °C  
33 warming targets. *Adv. Clim. Chang. Res.* 8, 99–107. doi:10.1016/j.accre.2017.05.004.
- 34 Xu, Z., Jiang, Y., Jia, B., and Zhou, G. (2016b). Elevated-CO2 Response of Stomata and Its Dependence on  
35 Environmental Factors. *Front. Plant Sci.* 7, 657. doi:10.3389/fpls.2016.00657.
- 36 Yamada, Y., Kodama, C., Satoh, M., Nakano, M., Nasuno, T., and Sugi, M. (2019). High-Resolution Ensemble  
37 Simulations of Intense Tropical Cyclones and Their Internal Variability During the El Niños of 1997 and 2015.  
38 *Geophys. Res. Lett.* 46, 7592–7601. doi:10.1029/2019GL082086.
- 39 Yamada, Y., Kodama, C., Satoh, M., Sugi, M., Roberts, M. J., Mizuta, R., et al. (2021). Evaluation of the contribution  
40 of tropical cyclone seeds to changes in tropical cyclone frequency due to global warming in high-resolution multi-  
41 model ensemble simulations. *Prog. Earth Planet. Sci.* 8, 11. doi:10.1186/s40645-020-00397-1.
- 42 Yamada, Y., Oouchi, K., Satoh, M., Tomita, H., and Yanase, W. (2010). Projection of changes in tropical cyclone  
43 activity and cloud height due to greenhouse warming: global cloud-system-resolving approach. *Geophys Res Lett*  
44 37.
- 45 Yamada, Y., Satoh, M., Sugi, M., Kodama, C., Noda, A. T., Nakano, M., et al. (2017). Response of Tropical Cyclone  
46 Activity and Structure to Global Warming in a High-Resolution Global Nonhydrostatic Model. *J. Clim.* 30, 9703–  
47 9724. doi:10.1175/JCLI-D-17-0068.1.
- 48 Yamaguchi, M., Chan, J. C. L., Moon, I.-J., Yoshida, K., and Mizuta, R. (2020). Global warming changes tropical  
49 cyclone translation speed. *Nat. Commun.* 11, 47. doi:10.1038/s41467-019-13902-y.
- 50 Yamaguchi, M., and Maeda, S. (2020a). Increase in the Number of Tropical Cyclones Approaching Tokyo since 1980.  
51 *J. Meteorol. Soc. Japan. Ser. II* 98, 775–786. doi:10.2151/jmsj.2020-039.
- 52 Yamaguchi, M., and Maeda, S. (2020b). Slowdown of typhoon translation speeds in mid-latitudes in September  
53 influenced by the Pacific Decadal Oscillation and global warming. *J. Meteor. Soc. Japan* 98.  
54 doi:10.2151/jmsj.2020-068.
- 55 Yanase, W., Satoh, M., Taniguchi, H., and Fujinami, H. (2012). Seasonal and intraseasonal modulation of tropical  
56 cyclogenesis environment over the bay of bengal during the extended summer monsoon. *J. Clim.* 25, 2914–2930.  
57 doi:10.1175/JCLI-D-11-00208.1.
- 58 Yang, C., Li, L., and Xu, J. (2018a). Changing temperature extremes based on CMIP5 output via semi-parametric  
59 quantile regression approach. *Int. J. Climatol.* 38, 3736–3748. doi:10.1002/joc.5524.
- 60 Yang, J. A., Kim, S., Mori, N., and Mase, H. (2018b). Assessment of long-term impact of storm surges around the  
61 Korean Peninsula based on a large ensemble of climate projections. *Coast. Eng.* 142, 1–8.

- 1 doi:10.1016/j.coastaleng.2018.09.008.
- 2 Yang, M.-Z., Zhong, P. A., Li, J.-Y., Chen, J., Wang, W.-Z., Zhu, F.-L., et al. (2018c). Spatial and temporal  
3 characteristics of pan evaporation in the Huaihe river basin during 1951–2015. *Appl. Ecol. Environ. Res.* 16,  
4 7635–7655. doi:10.15666/aecer/1606\_76357655.
- 5 Yang, S.-H., Kang, N.-Y., Elsner, J. B., and Chun, Y. (2018d). Influence of Global Warming on Western North Pacific  
6 Tropical Cyclone Intensities during 2015. *J. Clim.* 31, 919–925. doi:10.1175/JCLI-D-17-0143.1.
- 7 Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R., and Donohue, R. J. (2019). Hydrologic implications of  
8 vegetation response to elevated CO<sub>2</sub> in climate projections. *Nat. Clim. Chang.* 9, 44–48. doi:10.1038/s41558-  
9 018-0361-0.
- 10 Yang, Y., Zhang, S., McVicar, T. R., Beck, H. E., Zhang, Y., and Liu, B. (2018e). Disconnection Between Trends of  
11 Atmospheric Drying and Continental Runoff. *Water Resour. Res.* 54, 4700–4713. doi:10.1029/2018WR022593.
- 12 Yang, Y., Zhang, S., Roderick, M. L., McVicar, T. R., Yang, D., Liu, W., et al. (2020). Comparing Palmer Drought  
13 Severity Index drought assessments using the traditional offline approach with direct climate model outputs.  
14 *Hydrol. Earth Syst. Sci.* 24, 2921–2930. doi:10.5194/hess-24-2921-2020.
- 15 Yao, J., Chen, Y., Chen, J., Zhao, Y., Tuoliewubieke, D., Li, J., et al. (2020). Intensification of extreme precipitation in  
16 arid Central Asia. *J. Hydrol.*, 125760. doi:10.1016/j.jhydrol.2020.125760.
- 17 Yao, Y., Luo, D., Dai, A., and Simmonds, I. (2017). Increased Quasi Stationarity and Persistence of Winter Ural  
18 Blocking and Eurasian Extreme Cold Events in Response to Arctic Warming. Part I: Insights from Observational  
19 Analyses. *J. Clim.* 30, 3549–3568. doi:10.1175/JCLI-D-16-0261.1.
- 20 Yatagai, A., Minami, K., Masuda, M., and Sueto, N. (2019). Development of Intensive APHRODITE Hourly  
21 Precipitation Data for Assessment of the Moisture Transport That Caused Heavy Precipitation Events. *SOLA*  
22 15A, 43–48. doi:10.2151/sola.15A-008.
- 23 Ye, Z., and Li, Z. (2017). Spatiotemporal variability and trends of extreme precipitation in the Huaihe river basin, a  
24 climatic transitional zone in East China. *Adv. Meteorol.* 2017. doi:10.1155/2017/3197435.
- 25 Yettella, V., and Kay, J. E. (2017). How will precipitation change in extratropical cyclones as the planet warms?  
26 Insights from a large initial condition climate model ensemble. *Clim. Dyn.* 49, 1765–1781. doi:10.1007/s00382-  
27 016-3410-2.
- 28 Yin, H., Donat, M. G., Alexander, L. V., and Sun, Y. (2015). Multi-dataset comparison of gridded observed temperature  
29 and precipitation extremes over China. *Int. J. Climatol.* 35, 2809–2827. doi:10.1002/joc.4174.
- 30 Yin, H., and Sun, Y. (2018). Detection of Anthropogenic Influence on Fixed Threshold Indices of Extreme  
31 Temperature. *J. Clim.* 31, 6341–6352. doi:10.1175/JCLI-D-17-0853.1.
- 32 Yin, H., Sun, Y., and Donat, M. G. (2019). Changes in temperature extremes on the Tibetan Plateau and their  
33 attribution. *Environ. Res. Lett.* 14, 124015. doi:10.1088/1748-9326/ab503c.
- 34 Yin, H., Sun, Y., Wan, H., Zhang, X., and Lu, C. (2017). Detection of anthropogenic influence on the intensity of  
35 extreme temperatures in China. *Int. J. Climatol.* 37, 1229–1237. doi:10.1002/joc.4771.
- 36 Yiou, P., Jézéquel, A., Naveau, P., Otto, F. E. L., Vautard, R., and Vrac, M. (2017). A statistical framework for  
37 conditional extreme event attribution. *Adv. Stat. Climatol. Meteorol. Oceanogr.* 3, 17–31. doi:10.5194/ascmo-3-  
38 17-2017.
- 39 Yokoyama, C., Tsuji, H., and Takayabu, Y. N. (2020). The Effects of an Upper-Tropospheric Trough on the Heavy  
40 Rainfall Event in July 2018 over Japan. *J. Meteorol. Soc. Japan. Ser. II* 98, 235–255. doi:10.2151/jmsj.2020-013.
- 41 Yoshida, K., Sugi, M., Mizuta, R., Murakami, H., and Ishii, M. (2017). Future Changes in Tropical Cyclone Activity in  
42 High-Resolution Large-Ensemble Simulations. *Geophys. Res. Lett.* 44, 9910–9917. doi:10.1002/2017GL075058.
- 43 You, Q., Jiang, Z., Kong, L., Wu, Z., Bo, Y., Kang, S., et al. (2017). A comparison of heat wave climatologies and  
44 trends in China based on multiple definitions. *Clim. Dyn.* 48, 3975–3989. doi:10.1007/s00382-016-3315-0.
- 45 Young, I. R., Vinoth, J., Zieger, S., and Babanin, A. V. (2012). Investigation of trends in extreme value wave height  
46 and wind speed. *J. Geophys. Res. Ocean.* 117, 1–13. doi:10.1029/2011JC007753.
- 47 Yu, B., Lin, H., Kharin, V. V., and Wang, X. L. (2019a). Interannual variability of North American winter temperature  
48 extremes and its associated circulation anomalies in observations and CMIP5 simulations. *J. Clim.*, JCLI-D-19-  
49 0404.1. doi:10.1175/JCLI-D-19-0404.1.
- 50 Yu, B., Lin, H., and Soulard, N. (2019b). A Comparison of North American Surface Temperature and Temperature  
51 Extreme Anomalies in Association with Various Atmospheric Teleconnection Patterns. *Atmos.* 10.  
52 doi:10.3390/atmos10040172.
- 53 Yu, B., Lin, H., Wu, Z. W., and Merryfield, W. J. (2018). The Asian–Bering–North American teleconnection:  
54 seasonality, maintenance, and climate impact on North America. *Clim. Dyn.* 50, 2023–2038. doi:10.1007/s00382-  
55 017-3734-6.
- 56 Yu, E., Sun, J., Chen, H., and Xiang, W. (2015). Evaluation of a high-resolution historical simulation over China:  
57 climatology and extremes. *Clim. Dyn.* 45, 2013–2031. doi:10.1007/s00382-014-2452-6.
- 58 Yu, M., Li, Q., Hayes, M. J., Svoboda, M. D., and Heim, R. R. (2014). Are droughts becoming more frequent or severe  
59 in China based on the Standardized Precipitation Evapotranspiration Index: 1951–2010? *Int. J. Climatol.* 34, 545–  
60 558. doi:10.1002/joc.3701.
- 61 Yu, R., and Zhai, P. (2020). Changes in compound drought and hot extreme events in summer over populated eastern

- 1 China. *Weather Clim. Extrem.* 30, 100295. doi:10.1016/j.wace.2020.100295.
- 2 Yuan, S., and Quiring, S. M. (2017). Evaluation of soil moisture in CMIP5 simulations over the contiguous United  
3 States using in situ and satellite observations. *Hydrol. Earth Syst. Sci.* 21, 2203–2218. doi:10.5194/hess-21-2203-  
4 2017.
- 5 Yuan, S., Quiring, S. M., and Leason, Z. T. (2021). Historical Changes in Surface Soil Moisture Over the Contiguous  
6 United States: An Assessment of CMIP6. *Geophys. Res. Lett.* 48, 1–9. doi:10.1029/2020GL089991.
- 7 Yuan, W., Zheng, Y., Piao, S., Ciais, P., Lombardozzi, D., Wang, Y., et al. (2019). Increased atmospheric vapor  
8 pressure deficit reduces global vegetation growth. *Sci. Adv.* 5, eaax1396. doi:10.1126/sciadv.aax1396.
- 9 Yuan, X., Wang, L., and Wood, E. F. (2018a). Anthropogenic Intensification of Southern African Flash Droughts as  
10 Exemplified by the 2015/16 Season [in “Explaining Extreme Events of 2016 from a Climate Perspective”]. *Bull.*  
11 *Am. Meteorol. Soc.* 99, S86–S90. doi:10.1175/BAMS-D-17-0077.1.
- 12 Yuan, X., Wang, S., and Hu, Z.-Z. (2018b). Do Climate Change and El Niño Increase Likelihood of Yangtze River  
13 Extreme Rainfall? [in “Explaining Extreme Events of 2016 from a Climate Perspective”]. *Bull. Am. Meteorol.*  
14 *Soc.* 99, S113–S117. doi:10.1175/BAMS-D-17-0089.1.
- 15 Zaherpour, J., Gosling, S. N., Mount, N., Schmied, H. M., Veldkamp, T. I. E., Dankers, R., et al. (2018). Worldwide  
16 evaluation of mean and extreme runoff from six global-scale hydrological models that account for human  
17 impacts. *Environ. Res. Lett.* 13, 065015. doi:10.1088/1748-9326/aac547.
- 18 Zahid, M., Blender, R., Lucarini, V., and Bramati, M. C. (2017). Return levels of temperature extremes in southern  
19 Pakistan. *Earth Syst. Dyn.* 8, 1263–1278. doi:10.5194/esd-8-1263-2017.
- 20 Zahid, M., and Rasul, G. (2012). Changing trends of thermal extremes in Pakistan. *Clim. Change.* doi:10.1007/s10584-  
21 011-0390-4.
- 22 Zampieri, M., Ceglar, A., Dentener, F., and Toreti, A. (2017). Wheat yield loss attributable to heat waves, drought and  
23 water excess at the global, national and subnational scales. *Environ. Res. Lett.* 12, 064008. doi:10.1088/1748-  
24 9326/aa723b.
- 25 Zampieri, M., D’Andrea, F., Vautard, R., Ciais, P., de Noblet-Ducoudré, N., and Yiou, P. (2009). Hot European  
26 Summers and the Role of Soil Moisture in the Propagation of Mediterranean Drought. *J. Clim.* 22, 4747–4758.  
27 doi:10.1175/2009JCLI2568.1.
- 28 Zappa, G., Hawcroft, M. K., Shaffrey, L., Black, E., and Brayshaw, D. J. (2015). Extratropical cyclones and the  
29 projected decline of winter Mediterranean precipitation in the CMIP5 models. *Clim. Dyn.* doi:10.1007/s00382-  
30 014-2426-8.
- 31 Zappa, G., Shaffrey, L. C., and Hodges, K. I. (2013a). The Ability of CMIP5 Models to Simulate North Atlantic  
32 Extratropical Cyclones. *J. Clim.* 26, 5379–5396. doi:10.1175/JCLI-D-12-00501.1.
- 33 Zappa, G., Shaffrey, L. C., Hodges, K. I., Sansom, P. G., and Stephenson, D. B. (2013b). A multimodel assessment of  
34 future projections of north atlantic and european extratropical cyclones in the CMIP5 climate models. *J. Clim.* 26,  
35 5846–5862. doi:10.1175/JCLI-D-12-00573.1.
- 36 Zappa, G., and Shepherd, T. G. (2017). Storylines of Atmospheric Circulation Change for European Regional Climate  
37 Impact Assessment. *J. Clim.* 30, 6561–6577. doi:10.1175/JCLI-D-16-0807.1.
- 38 Zarzycki, C. M. (2016). Tropical Cyclone Intensity Errors Associated with Lack of Two-Way Ocean Coupling in High-  
39 Resolution Global Simulations. *J. Clim.* 29, 8589–8610. doi:10.1175/JCLI-D-16-0273.1.
- 40 Zarzycki, C. M. (2018). Projecting Changes in Societally Impactful Northeastern U.S. Snowstorms. *Geophys. Res. Lett.*  
41 45, 12,067–12,075. doi:10.1029/2018GL079820.
- 42 Zarzycki, C. M., and Jablonowski, C. (2015). Experimental Tropical Cyclone Forecasts Using a Variable-Resolution  
43 Global Model. *Mon. Weather Rev.* 143, 4012–4037. doi:10.1175/MWR-D-15-0159.1.
- 44 Zarzycki, C. M., Jablonowski, C., and Taylor, M. A. (2014). Using Variable-Resolution Meshes to Model Tropical  
45 Cyclones in the Community Atmosphere Model. *Mon. Weather Rev.* 142, 1221–1239. doi:10.1175/MWR-D-13-  
46 00179.1.
- 47 Zarzycki, C. M., and Ullrich, P. A. (2017). Assessing sensitivities in algorithmic detection of tropical cyclones in  
48 climate data. *Geophys. Res. Lett.* 44, 1141–1149. doi:10.1002/2016GL071606.
- 49 Zeder, J., and Fischer, E. M. (2020). Observed extreme precipitation trends and scaling in Central Europe. *Weather*  
50 *Clim. Extrem.* 29, 100266. doi:10.1016/j.wace.2020.100266.
- 51 Zeleke, T. T., Giorgi, F., Diro, G. T., and Zaitchik, B. F. (2017). Trend and periodicity of drought over Ethiopia. *Int. J.*  
52 *Climatol.* 37, 4733–4748. doi:10.1002/joc.5122.
- 53 Zhai, J., Mondal, S. K., Fischer, T., Wang, Y., Su, B., Huang, J., et al. (2020a). Future drought characteristics through a  
54 multi-model ensemble from CMIP6 over South Asia. *Atmos. Res.* 246. doi:10.1016/j.atmosres.2020.105111.
- 55 Zhai, R., Tao, F., Lall, U., Fu, B., Elliott, J., and Jägermeyr, J. (2020b). Larger Drought and Flood Hazards and Adverse  
56 Impacts on Population and Economic Productivity Under 2.0 than 1.5°C Warming. *Earth’s Futur.* 8.  
57 doi:10.1029/2019EF001398.
- 58 Zhan, R., Chen, B., and Ding, Y. (2018). Impacts of SST anomalies in the Indian-Pacific basin on Northwest Pacific  
59 tropical cyclone activities during three super El Niño years. *J. Oceanol. Limnol.* 36, 20–32. doi:10.1007/s00343-  
60 018-6321-8.
- 61 Zhan, R., and Wang, Y. (2017). Weak Tropical Cyclones Dominate the Poleward Migration of the Annual Mean

- 1 Location of Lifetime Maximum Intensity of Northwest Pacific Tropical Cyclones since 1980. *J. Clim.* 30, 6873–  
2 6882. doi:10.1175/JCLI-D-17-0019.1.
- 3 Zhan, Y. J., Ren, G. Y., Shrestha, A. B., Rajbhandari, R., Ren, Y. Y., Sanjay, J., et al. (2017). Changes in extreme  
4 precipitation events over the Hindu Kush Himalayan region during 1961–2012. *Adv. Clim. Chang. Res.* 8, 166–  
5 175. doi:10.1016/j.accre.2017.08.002.
- 6 Zhang, C., Liu, F., and Shen, Y. (2018a). Attribution analysis of changing pan evaporation in the Qinghai-Tibetan  
7 Plateau, China: Pan Evaporation in the Qinghai-Tibetan Plateau. *Int. J. Climatol.* 38. doi:10.1002/joc.5431.
- 8 Zhang, C., and Wang, Y. (2018). Why is the simulated climatology of tropical cyclones so sensitive to the choice of  
9 cumulus parameterization scheme in the WRF model? *Clim. Dyn.* 51, 3613–3633. doi:10.1007/s00382-018-4099-  
10 1.
- 11 Zhang, D., Zhang, Q., Qiu, J., Bai, P., Liang, K., and Li, X. (2018b). Intensification of hydrological drought due to  
12 human activity in the middle reaches of the Yangtze River, China. *Sci. Total Environ.* 637–638, 1432–1442.  
13 doi:10.1016/j.scitotenv.2018.05.121.
- 14 Zhang, G., Murakami, H., Knutson, T. R., Mizuta, R., and Yoshida, K. (2020a). Tropical cyclone motion in a changing  
15 climate. *Sci. Adv.* 6, eaaz7610. doi:10.1126/sciadv.aaz7610.
- 16 Zhang, J., Chen, H., and Zhang, Q. (2019a). Extreme drought in the recent two decades in northern China resulting  
17 from Eurasian warming. *Clim. Dyn.* 52, 2885–2902. doi:10.1007/s00382-018-4312-2.
- 18 Zhang, L., Zhou, T., Chen, X., Wu, P., Christidis, N., and Lott, F. C. (2020b). The late spring drought of 2018 in South  
19 China. *Bull. Am. Meteorol. Soc.* 101, S59–S64. doi:10.1175/BAMS-D-19-0202.1.
- 20 Zhang, M., Chen, Y., Shen, Y., and Li, B. (2019b). Tracking climate change in Central Asia through temperature and  
21 precipitation extremes. *J. Geogr. Sci.* 29, 3–28. doi:10.1007/s11442-019-1581-6.
- 22 Zhang, M., Chen, Y., Shen, Y., and Li, Y. (2017a). Changes of precipitation extremes in arid Central Asia. *Quat. Int.*  
23 436, 16–27. doi:10.1016/j.quaint.2016.12.024.
- 24 Zhang, P., Jeong, J. H., Yoon, J. H., Kim, H., Simon Wang, S. Y., Linderholm, H. W., et al. (2020c). Abrupt shift to  
25 hotter and drier climate over inner East Asia beyond the tipping point. *Science (80-. )*. 370, 1095–1099.  
26 doi:10.1126/science.abb3368.
- 27 Zhang, P., Ren, G., Xu, Y., Wang, X. L., Qin, Y., Sun, X., et al. (2019c). Observed Changes in Extreme Temperature  
28 over the Global Land Based on a Newly Developed Station Daily Dataset. *J. Clim.* 32, 8489–8509.  
29 doi:10.1175/JCLI-D-18-0733.1.
- 30 Zhang, Q., Gu, X., Singh, V. P., Kong, D., and Chen, X. (2015a). Spatiotemporal behavior of floods and droughts and  
31 their impacts on agriculture in China. *Glob. Planet. Change* 131, 63–72. doi:10.1016/j.gloplacha.2015.05.007.
- 32 Zhang, Q., Gu, X., Singh, V. P., Xiao, M., and Xu, C.-Y. (2015b). Flood frequency under the influence of trends in the  
33 Pearl River basin, China: Changing patterns, causes and implications. *Hydrol. Process.* 29, 1406–1417.  
34 doi:10.1002/hyp.10278.
- 35 Zhang, Q., Ni, X., and Zhang, F. (2017b). Decreasing trend in severe weather occurrence over China during the past 50  
36 years. *Sci. Rep.* 7, 42310. doi:10.1038/srep42310.
- 37 Zhang, R., Su, F., Jiang, Z., Gao, X., Guo, D., Ni, J., et al. (2015c). An overview of projected climate and  
38 environmental changes across the Tibetan Plateau in the 21st century. *Chinese Sci. Bull.* doi:10.1360/N972014-  
39 01296.
- 40 Zhang, W., Li, W., Zhu, L., Ma, Y., Yang, L., Lott, F. C., et al. (2020d). Anthropogenic Influence on 2018 Summer  
41 Persistent Heavy Rainfall in Central Western China. *Bull. Am. Meteorol. Soc.* 101, S65–S70. doi:10.1175/BAMS-  
42 D-19-0147.1.
- 43 Zhang, W., and Luo, D. (2019). A Nonlinear Theory of Atmospheric Blocking: An Application to Greenland Blocking  
44 Changes Linked to Winter Arctic Sea Ice Loss. *J. Atmos. Sci.* 77, 723–751. doi:10.1175/JAS-D-19-0198.1.
- 45 Zhang, W., Vecchi, G. A., Murakami, H., Delworth, T. L., Paffendorf, K., Jia, L., et al. (2016a). Influences of Natural  
46 Variability and Anthropogenic Forcing on the Extreme 2015 Accumulated Cyclone Energy in the Western North  
47 Pacific. *Bull. Am. Meteorol. Soc.* 97, S131–S135. doi:10.1175/BAMS-D-16-0146.1.
- 48 Zhang, W., Vecchi, G. A., Murakami, H., Villarini, G., and Jia, L. (2016b). The Pacific meridional mode and the  
49 occurrence of tropical Cyclones in the western North Pacific. *J. Clim.* 29, 381–398. doi:10.1175/JCLI-D-15-  
50 0282.1.
- 51 Zhang, W., Villarini, G., Scoccimarro, E., and Napolitano, F. (2020e). Examining the precipitation associated with  
52 medicanes in the high-resolution ERA-5 reanalysis data. *Int. J. Climatol.* n/a. doi:10.1002/joc.6669.
- 53 Zhang, W., Villarini, G., Vecchi, G. A., and Smith, J. A. (2018c). Urbanization exacerbated the rainfall and flooding  
54 caused by hurricane Harvey in Houston. *Nature* 563, 384–388. doi:10.1038/s41586-018-0676-z.
- 55 Zhang, W., and Zhou, T. (2019). Significant Increases in Extreme Precipitation and the Associations with Global  
56 Warming over the Global Land Monsoon Regions. *J. Clim.* 32, 8465–8488. doi:10.1175/JCLI-D-18-0662.1.
- 57 Zhang, X., Flato, G., Kirchmeier-Young, M., Vincent, L., Wan, H., Wang, X., et al. (2019d). “Changes in Temperature  
58 and Precipitation Across Canada,” in *Canada’s Changing Climate Report*, eds. E. Bush and D. S. Lemmen  
59 (Ottawa, ON, Canada: Government of Canada), 112–193. Available at: <https://changingclimate.ca/CCCR2019/>.
- 60 Zhang, X. S., Amirthanathan, G. E., Bari, M. A., Laugesen, R. M., Shin, D., Kent, D. M., et al. (2016c). How  
61 streamflow has changed across Australia since the 1950s: evidence from the network of hydrologic reference

- 1 stations. *Hydrol. Earth Syst. Sci.* 20, 3947–3965. doi:10.5194/hess-20-3947-2016.
- 2 Zhang, X. S., Amirthanathan, G. E., Bari, M. A., Laugesen, R. M., Shin, D., Kent, D. M., et al. (2016d). How  
3 streamflow has changed across Australia since the 1950s: evidence from the network of hydrologic reference  
4 stations. *Hydrol. Earth Syst. Sci.* 20, 3947–3965. doi:10.5194/hess-20-3947-2016.
- 5 Zhang, X., Wan, H., Zwiers, F. W., Hegerl, G. C., and Min, S.-K. (2013). Attributing intensification of precipitation  
6 extremes to human influence. *Geophys. Res. Lett.* 40, 5252–5257. doi:10.1002/grl.51010.
- 7 Zhang, X., Zwiers, F. W., Li, G., Wan, H., and Cannon, A. J. (2017c). Complexity in estimating past and future extreme  
8 short-duration rainfall. *Nat. Geosci.* 10, 255. doi:10.1038/ngeo2911.
- 9 Zhang, Y., Fan, J., Logan, T., Li, Z., and Homeyer, C. R. (2019e). Wildfire Impact on Environmental Thermodynamics  
10 and Severe Convective Storms. *Geophys. Res. Lett.* 46, 10082–10093. doi:10.1029/2019GL084534.
- 11 Zhang, Y., Pang, X., Xia, J., Shao, Q., Yu, E., Zhao, T., et al. (2019f). Regional Patterns of Extreme Precipitation and  
12 Urban Signatures in Metropolitan Areas. *J. Geophys. Res. Atmos.* 124, 641–663. doi:10.1029/2018JD029718.
- 13 Zhang, Y., Wang, H., Sun, J., and Drange, H. (2010). Changes in the tropical cyclone genesis potential index over the  
14 western north pacific in the SRES A2 scenario. *Adv. Atmos. Sci.* 27, 1246–1258. doi:10.1007/s00376-010-9096-1.
- 15 Zhang, Z., and Colle, B. A. (2017). Changes in Extratropical Cyclone Precipitation and Associated Processes during the  
16 Twenty-First Century over Eastern North America and the Western Atlantic Using a Cyclone-Relative Approach.  
17 *J. Clim.* 30, 8633–8656. doi:10.1175/JCLI-D-16-0906.1.
- 18 Zhang, Z., Ralph, F. M., and Zheng, M. (2019g). The Relationship Between Extratropical Cyclone Strength and  
19 Atmospheric River Intensity and Position. *Geophys. Res. Lett.* 46, 1814–1823. doi:10.1029/2018GL079071.
- 20 Zhang, Z., Wang, K., Chen, D., Li, J., and Dickinson, R. (2019h). Increase in Surface Friction Dominates the Observed  
21 Surface Wind Speed Decline during 1973–2014 in the Northern Hemisphere Lands. *J. Clim.* 32, 7421–7435.  
22 doi:10.1175/JCLI-D-18-0691.1.
- 23 Zhao, C., Brissette, F., Chen, J., and Martel, J. L. J.-L. (2020). Frequency change of future extreme summer  
24 meteorological and hydrological droughts over North America. *J. Hydrol.* 584, 124316.  
25 doi:10.1016/j.jhydrol.2019.124316.
- 26 Zhao, C., Lin, Y., Wu, F., Wang, Y. Y., Li, Z., Rosenfeld, D., et al. (2018). Enlarging Rainfall Area of Tropical  
27 Cyclones by Atmospheric Aerosols. *Geophys. Res. Lett.* 45, 8604–8611. doi:10.1029/2018GL079427.
- 28 Zhao, K., and Jackson, R. B. (2014). Biophysical forcings of land-use changes from potential forestry activities in  
29 North America. *Ecol. Monogr.* 84, 329–353. doi:10.1890/12-1705.1.
- 30 Zhao, M., and Held, I. M. (2011). TC-Permitting GCM Simulations of Hurricane Frequency Response to Sea Surface  
31 Temperature Anomalies Projected for the Late-Twenty-First Century. *J. Clim.* 25, 2995–3009. doi:10.1175/JCLI-  
32 D-11-00313.1.
- 33 Zhao, M., Held, I. M., and Lin, S.-J. (2012). Some Counterintuitive Dependencies of Tropical Cyclone Frequency on  
34 Parameters in a GCM. *J. Atmos. Sci.* 69, 2272–2283. doi:10.1175/JAS-D-11-0238.1.
- 35 Zhao, M., Held, I. M., Lin, S. J., and Vecchi, G. A. (2009). Simulations of global hurricane climatology, interannual  
36 variability, and response to global warming using a 50-km resolution GCM. *J. Clim.* 22, 6653–6678.  
37 doi:10.1175/2009JCLI3049.1.
- 38 Zhao, S., Deng, Y., and Black, R. X. (2016). Warm Season Dry Spells in the Central and Eastern United States:  
39 Diverging Skill in Climate Model Representation. *J. Clim.* 29, 5617–5624. doi:10.1175/JCLI-D-16-0321.1.
- 40 Zhao, T., and Dai, A. (2015). The magnitude and causes of global drought changes in the twenty-first century under a  
41 low-moderate emissions scenario. *J. Clim.* 28, 4490–4512. doi:10.1175/JCLI-D-14-00363.1.
- 42 Zhao, T., and Dai, A. (2017). Uncertainties in historical changes and future projections of drought. Part II: model-  
43 simulated historical and future drought changes. *Clim. Change* 144, 535–548. doi:10.1007/s10584-016-1742-x.
- 44 Zhao, Y., Ducharne, A., Sultan, B., Braconnot, P., and Vautard, R. (2015). Estimating heat stress from climate-based  
45 indicators: present-day biases and future spreads in the CMIP5 global climate model ensemble. *Environ. Res.*  
46 *Lett.* 10, 084013. doi:10.1088/1748-9326/10/8/084013.
- 47 Zheng, C., Zhang, R., Shi, W., Li, X., and Chen, X. (2017). Trends in significant wave height and surface wind speed in  
48 the China Seas between 1988 and 2011. *J. Ocean Univ. China* 16, 717–726. doi:10.1007/s11802-017-3213-z.
- 49 Zheng, F., Westra, S., and Sisson, S. A. (2013). Quantifying the dependence between extreme rainfall and storm surge  
50 in the coastal zone. *J. Hydrol.* 505, 172–187. doi:10.1016/j.jhydrol.2013.09.054.
- 51 Zheng, H., Chiew, F. H. S., Potter, N. J., and Kirono, D. G. C. (2019). Projections of water futures for Australia: An  
52 update. in *23rd International Congress on Modelling and Simulation – Supporting Evidence-Based Decision*  
53 *Making: The Role of Modelling and Simulation, MODSIM 2019*, 1000–1006.  
54 doi:10.36334/modsim.2019.k7.zhengh.
- 55 Zhou, B., Wen, Q. H., Xu, Y., Song, L., and Zhang, X. (2014). Projected changes in temperature and precipitation  
56 extremes in China by the CMIP5 multimodel ensembles. *J. Clim.* 27, 6591–6611. doi:10.1175/JCLI-D-13-  
57 00761.1.
- 58 Zhou, B., Xu, Y., Wu, J., Dong, S., and Shi, Y. (2016a). Changes in temperature and precipitation extreme indices over  
59 China: Analysis of a high-resolution grid dataset. *Int. J. Climatol.* 36, 1051–1066. doi:10.1002/joc.4400.
- 60 Zhou, C., Chen, D., Wang, K., Dai, A., and Qi, D. (2020). Conditional Attribution of the 2018 Summer Extreme Heat  
61 over Northeast China: Roles of Urbanization, Global Warming, and Warming-Induced Circulation Changes. *Bull.*

- 1 *Am. Meteorol. Soc.* 101, S71–S76. doi:10.1175/BAMS-D-19-0197.1.
- 2 Zhou, C., Wang, K., and Qi, D. (2018). Attribution of the July 2016 Extreme Precipitation Event Over China's Wuhang  
3 [in "Explaining Extreme Events of 2016 from a Climate Perspective"]. *Bull. Am. Meteorol. Soc.* 99, S107–S111.  
4 doi:10.1175/BAMS-D-17-0090.1.
- 5 Zhou, C., Wang, K., Qi, D., and Tan, J. (2019a). Attribution of a Record-Breaking Heatwave Event in Summer 2017  
6 over the Yangtze River Delta [in "Explaining Extreme Events of 2017 from a Climate Perspective"]. *Bull. Am.*  
7 *Meteorol. Soc.* 100, S97–S103. doi:10.1175/BAMS-D-18-0134.1.
- 8 Zhou, S., Williams, A. P., Berg, A. M., Cook, B. I., Zhang, Y., Hagemann, S., et al. (2019b). Land–atmosphere  
9 feedbacks exacerbate concurrent soil drought and atmospheric aridity. *Proc. Natl. Acad. Sci.* 116, 18848–18853.  
10 doi:10.1073/pnas.1904955116.
- 11 Zhou, S., Williams, A. P., Lintner, B. R., Berg, A. M., Zhang, Y., Keenan, T. F., et al. (2021). Soil moisture–  
12 atmosphere feedbacks mitigate declining water availability in drylands. *Nat. Clim. Chang.* 11, 38–44.  
13 doi:10.1038/s41558-020-00945-z.
- 14 Zhou, T., Song, F., Lin, R., Chen, X., and Chen, X. (2013). The 2012 north China floods: explaining an extreme rainfall  
15 event in the context of a longer-term drying tendency [in "Explaining Extreme Events of 2012 from a Climate  
16 Perspective"]. *Bull. Am. Meteorol. Soc.* 94, S49–S52. doi:10.1175/BAMS-D-13-00085.1.
- 17 Zhou, W., Tang, J., Wang, X., Wang, S., Niu, X., and Wang, Y. (2016b). Evaluation of regional climate simulations  
18 over the CORDEX-EA-II domain using the COSMO-CLM model. *Asia-Pacific J. Atmos. Sci.* 52, 107–127.  
19 doi:10.1007/s13143-016-0013-0.
- 20 Zhu, S., Ge, F., Fan, Y., Zhang, L., Sielmann, F., Fraedrich, K., et al. (2020). Conspicuous temperature extremes over  
21 Southeast Asia: seasonal variations under 1.5 °C and 2 °C global warming. *Clim. Change* 160, 343–360.  
22 doi:10.1007/s10584-019-02640-1.
- 23 Zieger, S., Babanin, A. V., and Young, I. R. (2014). Changes in ocean surface wind with a focus on trends in regional  
24 and monthly mean values. *Deep. Res. Part I Oceanogr. Res. Pap.* 86, 56–67. doi:10.1016/j.dsr.2014.01.004.
- 25 Zipser, E. J., Cecil, D. J., Liu, C., Nesbitt, S. W., and Yorty, D. P. (2006). Where are the most: Intense thunderstorms on  
26 Earth? *Bull. Am. Meteorol. Soc.* 87, 1057–1071. doi:10.1175/BAMS-87-8-1057.
- 27 Zittis, G. (2017). Observed rainfall trends and precipitation uncertainty in the vicinity of the Mediterranean, Middle  
28 East and North Africa. *Theor. Appl. Climatol.* doi:10.1007/s00704-017-2333-0.
- 29 Zolina, O., Simmer, C., Belyaev, K., Gulev, S. K., and Koltermann, P. (2013). Changes in the duration of European wet  
30 and dry spells during the last 60 years. *J. Clim.* 26, 2022–2047. doi:10.1175/JCLI-D-11-00498.1.
- 31 Zollo, A. L., Rillo, V., Bucchignani, E., Montesarchio, M., and Mercogliano, P. (2016). Extreme temperature and  
32 precipitation events over Italy: assessment of high-resolution simulations with COSMO-CLM and future  
33 scenarios. *Int. J. Climatol.* 36, 987–1004. doi:10.1002/joc.4401.
- 34 Zscheischler, J., and Fischer, E. M. (2020). The record-breaking compound hot and dry 2018 growing season in  
35 Germany. *Weather Clim. Extrem.* 29, 100270. doi:10.1016/j.wace.2020.100270.
- 36 Zscheischler, J., Fischer, E. M., and Lange, S. (2018a). The effect of bias adjustment on impact modeling. *Earth Syst.*  
37 *Dyn. Discuss.*, 1–17. doi:10.5194/esd-2018-68.
- 38 Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., et al. (2020). A typology of  
39 compound weather and climate events. *Nat. Rev. Earth Environ.* 1, 333–347. doi:10.1038/s43017-020-0060-z.
- 40 Zscheischler, J., and Seneviratne, S. I. (2017). Dependence of drivers affects risks associated with compound events.  
41 *Sci. Adv.* 3, e1700263. doi:10.1126/sciadv.1700263.
- 42 Zscheischler, J., Westra, S., van den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., et al. (2018b). Future  
43 climate risk from compound events. *Nat. Clim. Chang.* 8, 469–477. doi:10.1038/s41558-018-0156-3.
- 44 Zulkafli, Z., Buytaert, W., Manz, B., Rosas, C. V., Willems, P., Lavado-Casimiro, W., et al. (2016). Projected increases  
45 in the annual flood pulse of the Western Amazon. *Environ. Res. Lett.* 11, 014013. doi:10.1088/1748-  
46 9326/11/1/014013.
- 47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

1  
2  
3  
4  
5  
6  
7  
8**Appendix 11.A****[START TABLE 11.A.1 HERE]****Table 11.A.1:** Common drought metrics, associated drought types, drought indices, general description and associated references

Drought metric	Associated drought type	Drought indices	Comments	Representative references
Precipitation deficit	Referred to as “meteorological drought”	Standardized Precipitation Index (SPI), Consecutive Dry Days (CDD), Precipitation deciles and percentiles.	SPI is defined for given time scales in order to identify precipitation deficits over different periods. The SPI shows flexibility to account for different time scales by summing precipitation over $k$ months, termed accumulation periods. CDD is usually based on daily precipitation records. Dry-spell length is another commonly used term. The number of dry days (NDD) is also used in some publications.	(Donat et al., 2013a; Orłowsky and Seneviratne, 2013; Sillmann et al., 2013a; Spinoni et al., 2014; Kingston et al., 2015; Stage et al., 2017; Coppola et al., 2021b)
Excess atmospheric evaporative demand (AED)	Driver for agricultural and ecological drought, together with precipitation through its impact on evapotranspiration and vegetation stress under soil moisture deficits	Potential evaporation anomalies, Evaporative Demand Drought Index (EDDI).	AED can be measured locally by means of evaporation pans. Physically-based models (e.g., Penman-Monteith) using all aerodynamic and radiative drivers from observations produce robust estimates of the the observed magnitude and variability of the evaporative demand. On the contrary, empirical estimates based on air temperature are affected by more uncertainties (Section 11.6.1.2), especially when applied to climate change projections. AED is an upper bound for actual evapotranspiration (ET) but also induces additional vegetation stress under dry conditions (Section 11.6.1.2).	(Hobbins et al., 2012, 2016; Sheffield et al., 2012; Wang et al., 2012; McEvoy et al., 2016; Roberts et al., 2018; Stephens et al., 2018; Sun et al., 2018c; Vicente-Serrano et al., 2020b)
Soil moisture deficits	Usually referred to as “agricultural drought”. Also relevant for ecological droughts.	Soil moisture anomalies (SMA), Standardized Soil Moisture Index (SSMI)	Networks of ground-based soil moisture measurements are available in different regions, but are very sparse and cover very short periods. Surface soil moisture can be monitored from satellites, but only since the 1980s at the earliest. Physically-based land surface models retrieve soil moisture using meteorological variables (precipitation, radiation, wind, temperature, humidity) as input.	(Dorigo et al., 2011, 2015, 2017; Seneviratne et al., 2013; Orłowsky and Seneviratne, 2013; AghaKouchak, 2014; Sohrabi et al., 2015; Zhao and Dai, 2015; Stillman et al., 2016; Yuan and Quiring, 2017; Berg and Sheffield, 2018; Hanel et al., 2018; Samaniego et al., 2018; Seager et al., 2019; Ford and Quiring, 2019; Moravec et al., 2019)
Streamflow and surface water deficits	Usually referred to as “hydrological drought”	SRI (Standardized Runoff Index), SSI (Standardized Streamflow Index), threshold level methods, SGI (Standardized Groundwater Index)	Usually based on monthly records of hydrological variables (e.g., streamflow, groundwater, reservoir storages), although daily streamflow is also used using threshold level methods. Observational data is available but not in all regions.	(Bloomfield and Marchant, 2013; Van Lanen et al., 2013; Wada et al., 2013; Forzieri et al., 2014; Prudhomme et al., 2014; Schewe et al., 2014; Van Loon, 2015; Van Loon and Laaha, 2015; Gosling et al., 2017)

<p>Atmospheri-based drought indices</p>	<p>Metrics of drought severity based on meteorological variables, combining precipitation and AED as drivers.</p>	<p>Standardized-Precipitation Evapotranspiration Index (SPEI), Palmer Drought Severity Index (PDSI)</p>	<p>These drought indices are generated using precipitation and AED. The quality of the outputs depend on the method used to determine the AED. They are widely used for drought monitoring and early warning. These indices are not intended to be a soil moisture or water-balance proxy.</p>	<p>(Dai, 2013; Beguería et al., 2014; Cook et al., 2014a; Mitchell et al., 2014; Stagge et al., 2015; Vicente-Serrano et al., 2015; Dai et al., 2018; Mukherjee et al., 2018b; Yang et al., 2020)</p>
---	---	---	--	---


1  
2  
3

**[END TABLE 11.A.1 HERE]**



1 **Figures**

2



3

4


5

6

7


**Figure 11.1:** Chapter 11 visual abstract of contents.

1  
2




3 **Figure 11.2:** Time series of observed temperature anomalies for global average annual mean temperature (black), land  
4 average annual mean temperature (green), land average annual hottest daily maximum temperature (TXx,  
5 purple), and land average annual coldest daily minimum temperature (TNn, blue). Global and land mean  
6 temperature anomalies are relative to their 1850-1900 means based on the multi-product mean annual  
7 time series assessed in Section 2.3.1.1.3 (see text for references). TXx and TNn anomalies are relative to  
8 their respective 1961-1990 means and are based on the HadEX3 dataset (Dunn et al., 2020) using values  
9 for grid boxes with at least 90% temporal completeness over 1961-2018. Further details on data sources  
10 and processing are available in the chapter data table (Table 11.SM.9).  
11  
12  
13

Scaling of regional annual hottest temperature (TXx)




1  
 2 **Figure 11.3:** Regional mean changes in annual hottest daily maximum temperature (TXx) for AR6 land regions and  
 3 the global land, against changes in global mean surface air temperature (GSAT) as simulated by CMIP6  
 4 models under different forcing scenarios SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. (a)  
 5 shows individual models from the CMIP6 ensemble (grey), the multi-model median under three selected  
 6 SSPs (colours), and the multi-model median (black). (b) to (l) show the multi-model-median for the  
 7 pooled data for individual AR6 regions. Numbers in parantheses indicate the linear scaling between  
 8 regional TXx and GSAT. The black line indicates the 1:1 reference scaling between TXx and GSAT. See  
 9 Atlas.1.3.2 for the definition of regions. For details on the methods see Supplementary Material 11.SM.2.

**Overview of observed changes for cold, hot, and wet extremes and their potential human contribution.**



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15


**Figure 11.4:** Overview of observed changes for cold, hot, and wet extremes and their potential human contribution. Shown are the direction of change and the confidence in 1) the observed changes in how cold and hot as well as wet extremes have already changed across the world and 2) in the contribution of whether human-induced climate change contributed in causing to these changes (attribution). In each region changes in extremes are indicated by colour (orange – increase in the type of extreme, blue – decrease, both colours – there are changes of opposing direction within the region the signal depends on the exact event definition, grey – there are no changes observed, and no fill – the data/evidence is too sparse to make an assessment). The squares and dots next to the symbol indicate the level of confidence for observing the trend and the human contribution, respectively. The more black dots/squares the higher the level of confidence. The information on this figure is based on regional assessment of the literature on observed trends, detection and attribution and event attribution in section 11.9.



1  
2

**Box 11.1, Figure 1:** Multi-model (CMIP5) mean fractional changes (in % per degree of warming) for (a) annual maximum precipitation (Rx1day), (b) changes in Rx1day due to the thermodynamic contribution and (c) changes in Rx1day due to the dynamic contribution estimated as the difference between the total changes and the thermodynamic contribution. Changes were derived from a linear regression for the period 1950–2100. Uncertainty is represented using the simple approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models ( $n=22$ ) agree on sign of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box Atlas 1. A detailed description of the estimation of dynamic and thermodynamic contributions is given in Pfahl et al. (2017). Adapted from (Pfahl et al., 2017), originally published in Nature Climate Change/ Springer Nature. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).


15



1  
2  
3  
4  
5


**Figure 11.5:** Confidence and likelihood of past changes and projected future changes at 2°C of global warming on the global scale. The information in this figure is based on Tables 11.1 and 11.2.

1  
2



3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

**Figure 11.6:** Projected changes in the frequency of extreme temperature events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1851-1900 baseline. Extreme temperatures are defined as the maximum daily temperatures that were exceeded on average once during a 10-year period (10-year event, blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land and the AR6 regions. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the frequency changes across the multi model ensemble, and the whiskers extend to the 90% uncertainty range. The dotted line indicates no change in frequency. The results are based on the multi-model ensemble from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Adapted from (Li et al., 2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

**Figure 11.7:** Projected changes in the frequency of extreme precipitation events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1951-1990 baseline. Extreme precipitation is defined as the maximum daily precipitation (Rx1day) that was exceeded on average once during a 10-year period (10-year event, blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land and the AR6 regions. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the frequency changes across the multi model ensemble, and the whiskers extend to the 90% uncertainty range. The dotted line indicates no change in frequency. The results are based on the multi-model ensemble from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Adapted from (Li et al., 2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).




1  
2

(a) Annual hottest temperature (TXx)

CMIP6	0.1	0.1	0.2	0.5	0.7	0.8	0.7	0.7	0.7	0.6	0.5	0.3	0.3	0.4	0.6	0.2	0.4	0.3	0.3	0.8	0.9	0.9	0.4
CMIP5	0.1	0.1	0.2	0.5	0.5	0.5	0.4	0.7	0.4	0.4	0.4	0.2	0.3	0.4	0.9	0.3	0.4	0.3	0.3	1.0	0.7	0.7	0.4
	glob.	oc.	land	GIC	NEC	CNA	ENA	NWN	WNA	NCA	SCA	CAR	NWS	SAM	SSA	SWS	SES	NSA	NES	NEU	CEU	EEU	MED
CMIP6	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.7	0.7	0.8	0.8	0.4	0.5	0.6	0.2	0.6	0.4	0.4	0.4	0.6	0.7	0.8	1.1
CMIP5	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.6	0.7	0.8	0.8	0.3	0.5	0.6	0.2	0.3	0.2	0.4	0.5	0.7	0.7	0.6	1.0
	WAF	SAH	NEAF	CEAF	SWAF	SEAF	CAF	RAR	RFE	ESB	WSB	WCA	TIB	EAS	ARP	SAS	SEA	NAU	CAU	SAU	NZ	EAN	WAN

(b) Annual maximum daily precipitation (Rx1day)


CMIP6	0.3	0.4	0.4	0.8	0.7	1.4	1.0	0.8	1.7	2.0	2.2	2.8	0.9	1.3	2.1	2.7	0.8	1.2	1.2	0.8	1.0	1.2	2.0
CMIP5	0.3	0.4	0.4	0.7	0.7	1.4	1.0	0.8	1.5	1.9	2.0	2.7	0.8	1.1	2.2	2.2	0.9	0.9	1.3	0.9	1.0	1.2	2.1
	glob.	oc.	land	GIC	NEC	CNA	ENA	NWN	WNA	NCA	SCA	CAR	NWS	SAM	SSA	SWS	SES	NSA	NES	NEU	CEU	EEU	MED
CMIP6	0.9	1.5	1.1	0.8	1.9	1.2	0.6	0.9	0.9	1.0	0.9	1.3	1.0	1.1	2.6	0.9	0.8	2.1	2.2	2.6	1.7	0.6	0.9
CMIP5	1.3	2.7	1.3	0.8	1.7	1.3	0.7	0.8	1.0	0.9	1.1	1.8	1.0	0.9	2.6	1.1	0.8	1.9	1.9	2.5	1.9	0.7	1.0
	WAF	SAH	NEAF	CEAF	SWAF	SEAF	CAF	RAR	RFE	ESB	WSB	WCA	TIB	EAS	ARP	SAS	SEA	NAU	CAU	SAU	NZ	EAN	WAN



**Figure 11.8:** Global and regional-scale emergence of changes in temperature (a) and precipitation (b) extremes for the globe (glob.), global oceans (oc.), global lands (land), and the AR6 regions. Colours indicate the multi-model mean global warming level at which the difference in 20-year means of the annual maximum daily maximum temperature (TXx) and the annual maximum daily precipitation (Rx1day) become significantly different from their respective mean values during the 1851–1900 base period. Results are based on simulations from the CMIP5 and CMIP6 multi-model ensembles. See Atlas.1.3.2 for the definition of regions. Adapted from Seneviratne and Hauser, 2020) under the terms of the Creative Commons Attribution license.

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15


1




2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13

**Cross-Chapter Box 11.1, Figure 1:** Schematic representation of relationship between emission scenarios, global warming levels (GWLs), regional climate responses, and impacts. The illustration shows the implied uncertainty problem associated with differentiating between 1.5, 2°C, and other GWLs. Focusing on GWL raises questions associated with emissions pathways to get to these temperatures (scenarios), as well as questions associated with regional climate responses and the associated impacts at the corresponding GWL (the impacts question). Adapted from (James, Washington, Schlessner, Rogelj, & Conway, 2017) and (Rogelj, 2013) under the terms of the Creative Commons Attribution license.

### Annual mean precipitation response at 2°C




### Annual mean temperature response at 2°C



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24


**Cross-Chapter Box 11.1, Figure 2:** (a-c) CMIP6 multi-model mean precipitation change at 2°C GWL (20-yr mean) in three different SSP scenarios relative to 1850-1900. All models reaching the corresponding GWL in the corresponding scenario are averaged. The number of models averaged across is shown at the top right of the panel. The maps for the other two SSP scenarios SSP1-1.9 (five models only) and SSP3-7.0 (not shown) are consistent. (d-f) Same as (a-c) but for annual mean temperature. (g) Annual mean temperature change at 2°C in CMIP6 models with high warming rate reaching the GWL in the corresponding scenario before the earliest year of the assessed very likely range (section 4.3.4) (h) Climate response at 2°C GWL across all SSP1-1.9, SSP2-2.6, SSP2-4.5, SSP3-7.0 and SSP5-8.5 in all other models not shown in (g). The good agreement of (g) and (h) demonstrate that the mean temperature response at 2°C is not sensitive to the rate of warming and thereby the GSAT warming of the respective models in 2081-2100. Uncertainty is represented using the advanced approach: No overlay indicates regions with robust signal, where  $\geq 66\%$  of models show change greater than variability threshold and  $\geq 80\%$  of all models agree on sign of change; diagonal lines indicate regions with no change or no robust signal, where  $< 66\%$  of models show a change greater than the variability threshold; crossed lines indicate regions with conflicting signal, where  $\geq 66\%$  of models show change greater than variability threshold and  $< 80\%$  of all models agree on sign of change. For more information on the advanced approach, please refer to the Cross-Chapter Box Atlas.1.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10

**Cross-Chapter Box 11.1, Figure. 3:** Illustration of the AR6 GWL sampling approach to derive the timing and the response at a given GWL for the case of CMIP6 data. For the mapping of scenarios/time slices into GWLs for CMIP6, please refer to Table 4.2. Respective numbers for the CMIP6 multi-model experiment are provided in the Chapter 11 Supplementary Material (11.SM.1). Note that the time frames used to derived the GWL time slices can also include different number of years (e.g. 30 years for some analyses).


1  
2  
3



4  
5  
6  
7  
8  
9  
10  
11  
12  
13

**Figure 11.9:** Linear trends over 1960–2018 in the annual maximum daily maximum temperature (TXx, a), the annual minimum daily minimum temperature (TNn, b), and the annual number of days when daily maximum temperature exceeds its 90<sup>th</sup> percentile from a base period of 1961–1990 (TX90p, c), based on the HadEX3 data set (Dunn et al., 2020). Linear trends are calculated only for grid points with at least 66% of the annual values over the period and which extend to at least 2009. Areas without sufficient data are shown in grey. No overlay indicates regions where the trends are significant at  $p = 0.1$  level. Crosses indicate regions where trends are not significant. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).


1



2  
3  
4  
5  
6  
7  
8  
9  
10

**Figure 11.10:** Multi-model mean bias in temperature extremes (°C) for the period 1979-2014, calculated as the difference between the CMIP6 multi-model mean and the average of observations from the values available in HadEX3 for (a) the annual hottest temperature (TXx) and (b) the annual coldest temperature (TNn). Areas without sufficient data are shown in grey. Adapted from Wehner et al. (2020) under the terms of the Creative Commons Attribution license. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).


1  
2



3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

**Figure 11.11:** Projected changes in (a-c) annual maximum temperature (TXx) and (d-f) annual minimum temperature (TNn) at 1.5°C, 2°C, and 4°C of global warming compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-model ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The numbers in the top right indicate the number of simulations included. Uncertainty is represented using the simple approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models agree on sign of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1




2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

**Figure 11.12:** Projected changes in the intensity of extreme temperature events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1851-1900 baseline. Extreme temperature events are defined as the daily maximum temperatures (TXx) that were exceeded on average once during a 10-year period (10-year event, blue) and that once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the intensity changes across the multi-model ensemble, and the whiskers extend to the 90% uncertainty range. The results are based on the multi-model ensemble from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Based on (Li et al., 2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).




1  
2



3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16


**Figure 11.13:** Signs and significance of the observed trends in annual maximum daily precipitation (Rx1day) during 1950–2018 at 8345 stations with sufficient data. (a) Percentage of stations with statistically significant trends in Rx1day; green dots show positive trends and brown dots negative trends. Box-and-whisker plots indicate the expected percentage of stations with significant trends due to chance estimated from 1000 bootstrap realizations under a no-trend null hypothesis. The boxes mark the median, 25<sup>th</sup> percentile, and 75<sup>th</sup> percentile. The upper and lower whiskers show the 97.5<sup>th</sup> and the 2.5<sup>th</sup> percentiles, respectively. Maps of stations with positive (b) and negative (c) trends. The light color indicates stations with non-significant trends and the dark color stations with significant trends. Significance is determined by a two-tailed test conducted at the 5% level. Adapted from Sun et al. (2020). © American Meteorological Society. Used with permission. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1  
2



3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13


**Figure 11.14:** Multi-model mean bias in annual maximum daily precipitation (Rx1day, %) for the period 1979-2014, calculated as the difference between the CMIP6 multi-model mean and the average of available observational or reanalysis products including (a) ERA5, (b) HadEX3, and (c) and REGEN. Bias is expressed as the percent error relative to the long-term mean of the respective observational data products. Brown indicates that models are too dry, while green indicates that they are too wet. Areas without sufficient observational data are shown in grey. Adapted from Wehner et al. (2020) under the terms of the Creative Commons Attribution license. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13

**Figure 11.15:** Projected changes in the intensity of extreme precipitation events under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1851-1900 baseline. Extreme precipitation events are defined as the daily precipitation (Rx1day) that was exceeded on average once during a 10-year period (10-year event, blue) and once during a 50-year period (50-year event, orange) during the 1851-1900 base period. Results are shown for the global land. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the intensity changes across the multi model median, and the whiskers extend to the 90% uncertainty range. The results are based on the multi-model ensemble estimated from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Based on Li et al. (2020a). Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).


1



2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14

**Figure 11.16:** Projected changes in annual maximum daily precipitation at (a) 1.5°C, (b) 2°C, and (c) 4°C of global warming compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-model ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The numbers on the top right indicate the number of simulations included. Uncertainty is represented using the simple approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models agree on sign of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1  
2  
3




4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14

**Figure 11.17:** Observed linear trend for (a) consecutive dry days (CDD) during 1960-2018, (b) standardized precipitation index (SPI) and (c) standardized precipitation-evapotranspiration index (SPEI) during 1951-2016. CDD data are from the HadEx3 dataset (Dunn et al., 2020), trend calculation of CDD as in Figure 11.9. Drought severity is estimated using 12-month SPI (SPI-12) and 12-month SPEI (SPEI-12). SPI and SPEI datasets are from Spinoni et al. (2019). The threshold to identify drought episodes was set at -1 SPI/SPEI units. Areas without sufficient data are shown in grey. No overlay indicates regions where the trends are significant at  $p = 0.1$  level. Crosses indicate regions where trends are not significant. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1  
2  
3


Changes in 10-year soil moisture drought in drying regions



4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**Figure 11.18:** Projected changes in the intensity (a) and frequency (b) of drought under 1°C, 1.5°C, 2°C, 3°C, and 4°C global warming levels relative to the 1850-1900 baseline. Summaries are computed for the AR6 regions in which there is at least medium confidence in increase in agriculture/ ecological drought at the 2°C warming level (“drying regions”), including W. North-America, C. North-America, N. Central-America, S. Central-America, N. South-America, N. E. South-America, South-American-Monsoon, S.W.South-America, S.South-America, West & Central-Europe, Mediterranean, W.Southern-Africa, E.Southern-Africa, Madagascar, E.Australia, S.Australia (c). A drought event is defined as a 10-year drought event whose annual mean soil moisture was below its 10th percentile from the 1850-1900 base period. For each box plot, the horizontal line and the box represent the median and central 66% uncertainty range, respectively, of the frequency or the intensity changes across the multi-model ensemble, and the whiskers extend to the 90% uncertainty range. The line of zero in (a) indicates no change in intensity, while the line of one in (b) indicates no change in frequency. The results are based on the multi-model ensemble estimated from simulations of global climate models contributing to the sixth phase of the Coupled Model Intercomparison Project (CMIP6) under different SSP forcing scenarios. Intensity changes in (a) are expressed as standard deviations of the interannual variability in the period 1850-1900 of the corresponding model. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources and processing are available in the chapter data table (Table 11.SM.9).

1  
2  
3




4  
5  
6  
7  
8  
9  
10  
11

**Figure 11.19:** Projected changes in (a-c) the number of consecutive dry days (CDD), (d-f) annual mean soil moisture over the total column, and (g-l) the frequency and intensity of one-in-ten year soil moisture drought for the June-to-August and December-to-February seasons at 1.5°C, 2°C, and 4°C of global warming compared to the 1851-1900 baseline. Results are based on simulations from the CMIP6 multi-model ensemble under the SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. The numbers in the top right indicate the number of simulations included. Uncertainty is represented using the simple

1 approach: no overlay indicates regions with high model agreement, where  $\geq 80\%$  of models agree on sign  
2 of change; diagonal lines indicate regions with low model agreement, where  $< 80\%$  of models agree on  
3 sign of change. For more information on the simple approach, please refer to the Cross-Chapter Box  
4 Atlas 1. For details on the methods see Supplementary Material 11.SM.2. Further details on data sources  
5 and processing are available in the chapter data table (Table 11.SM.9).  
6  
7




1



**Figure 11.20:** Summary schematic of past and projected changes in tropical cyclone (TC), extratropical cyclone (ETC), atmospheric river (AR), and severe convective storm (SCS) behaviour. Global changes (blue shading) from top to bottom: 1) Increased mean and maximum rain-rates in TCs, ETCs, and ARs [past (*low confidence* due to lack of reliable data) & projected (*high confidence*)]. 2) Increased proportion of stronger TCs [past (*medium confidence*) & projected (*high confidence*)]. 3) Decrease or no change in global frequency of TC genesis [past (*low confidence* due to lack of reliable data) & projected (*medium confidence*)]. 4) Increased and decreased ETC wind-speed, depending on the region, as storm-tracks change [past (*low confidence* due to lack of reliable data) & projected (*medium confidence*)]. Regional changes, from left to right: 1) Poleward TC migration in the western North Pacific and subsequent changes in TC exposure [past (*medium confidence*) & projected (*medium confidence*)]. 2) Slowdown of TC forward translation speed over the contiguous US and subsequent increase in TC rainfall [past (*medium confidence*) & projected (*low confidence* due to lack of directed studies)]. 3) Increase in mean and maximum SCS rain-rate and increase in springtime SCS frequency and season length over the contiguous US [past (*low confidence* due to lack of reliable data) & projected (*medium confidence*)].

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31




1  
2  
3  
4  
5  
6  
7  
8  
9  
10

**Box 11.4, Figure 1:** Analysis of the percentage of land area affected by temperature extremes larger than two (orange) or three (blue) standard deviations in June-July-August (JJA) between 30°N and 80°N using a normalization. The more appropriate estimate is the corrected normalization. These panels show for both estimates a substantial increase in the overall land area affected by very high hot extremes since 1990 onward. Adapted from Sippel et al. (2015).

1

July 2018




2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

**Box 11.4, Figure 2:** Meteorological conditions in July 2018. The color shading shows the monthly mean near-surface air temperature anomaly with respect to 1981 to 2010. Contour lines indicate the geopotential height in m, highlighted are the isolines on 12'000 m and 12'300 m, which indicate the approximate positions of the polar-front jet and subtropical jet, respectively. The light blue-green ellipse shows the approximate extent of the strong precipitation event that occurred at the beginning of July in the region of Japan and Korea. All data is from the global ECMWF Reanalysis v5 (ERA5, Hersbach et al., 2020).

**FAQ 11.1: How will changes in climate extremes compare with changes in climate averages?**

The direction and magnitude of future changes in climate extremes and averages depend on the variable considered.



Future **changes in temperature** averages and extremes will be **similar**

Future **changes in precipitation** averages and extremes can be **very different**

1  
2

**FAQ 11.1, Figure 1: Global maps of future changes in surface temperature (top panels) and precipitation (bottom panels) for long-term average (left) and extreme conditions (right).** All changes were estimated using the CMIP6 ensemble mean for a scenario with a global warming of 4°C relative to 1850-1900 temperatures. Average surface temperatures refer to the warmest three-month season (summer in mid- to high-latitudes) and extreme temperature refer to the hottest day in a year. Precipitation changes, which can include both rainfall and snowfall changes, are normalized by 1850-1900 values and shown in percentage; extreme precipitation refers to the largest daily rainfall in a year.

3

### FAQ 11.2: Will climate change cause unprecedented extremes?

Yes, in a changing climate, extreme events may be unprecedented when they occur with...



**Larger magnitude**



**Increased frequency**



**New locations**



**Different timing**




**New combinations (compound)**

1  
2  
3

**FAQ 11.2, Figure 1:** New types of unprecedented extremes that will occur as a result of climate change.

### FAQ 11.3: Climate change and extreme events

Extreme events have become more probable and more intense. Many of these changes can be attributed to human influence on the climate.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

**FAQ 11.3, Figure 1: Changes in climate result in changes in the magnitude and probability of extremes.**

Example of how temperature extremes differ between a climate with pre-industrial greenhouse gases (shown in blue) and the current climate (shown in orange) for a representative region. The horizontal axis shows the range of extreme temperatures, while the vertical axis shows the annual chance of each temperature event’s occurrence. Moving towards the right indicates increasingly hotter extremes that are more rare (less probable). For hot extremes, an extreme event of a particular temperature in the pre-industrial climate would be more probable (vertical arrow) in the current climate. An event of a certain probability in the pre-industrial climate would be warmer (horizontal arrow) in the current climate. While the climate under greenhouse gases at the pre-industrial level experiences a range of hot extremes, such events are hotter and more frequent in the current climate.