

Variational data assimilation for morphodynamic model parameter estimation

Conference or Workshop Item

Presentation

Presentation slides

Smith, P. J. ORCID: <https://orcid.org/0000-0003-4570-4127>,
Dance, S. L. ORCID: <https://orcid.org/0000-0003-1690-3338>,
Baines, M. J., Nichols, N. K. ORCID: <https://orcid.org/0000-0003-1133-5220> and Scott, T. R. (2009) Variational data assimilation for morphodynamic model parameter estimation. In: Centre for Australian Weather and Climate Research (CAWCR) Workshop on Ensemble Prediction and Data Assimilation, February 2009, Australian Bureau of Meteorology, Melbourne. Available at <https://centaur.reading.ac.uk/1705/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: http://www.cawcr.gov.au/events/cawcr_da_ep3/abstracts.htm#PSmith

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Variational data assimilation for morphodynamic model parameter estimation

Department of Mathematics, University of Reading:

Polly Smith*, Sarah Dance, Mike Baines, Nancy Nichols,

Environmental Systems Science Centre, University of Reading:

Tania Scott

*email: p.j.smith@reading.ac.uk

This project is funded under the Natural Environment Research Council (NERC) Flood Risk From Extreme Events (FREE) programme, with additional funding provided by the Environment Agency as part of the CASE (Co-operative Awards in Science and Engineering) scheme. Thanks also to HR Wallingford for visits and useful discussions.

Outline

- Background
 - what is morphodynamic modelling?
 - why do we need morphodynamic models?
- Data assimilation and parameter estimation
 - how can data assimilation provide us with better estimates of uncertain model parameters?
 - what is state augmentation?
 - how do we model the error covariances?
 - Application to a simple 1D morphodynamic model
 - does it work?
- Summary

Terminology

- **Bathymetry** - the underwater equivalent to topography
 - coastal bathymetry is dynamic and evolves with time
 - water action erodes, transports, and deposits sediment, which changes the bathymetry, which alters the water action, and so on
- **Morphodynamics** - the study of the evolution of the bathymetry in response to the flow induced sediment transport
- **Morphodynamic prediction**
 - why?
 - how?



Kent channel


1960s

1970s

1997


18km

- Channel movement
 - impacts on habitats in the bay
 - affects access to ports
 - has implications for flooding during storm events



Scale
0 1 2 3
kilometres

Picture courtesy of Nigel Cross,
Lancaster City Council



Morecambe
Bay

Kent channel


1960s

1970s

1997

18km

- Channel movement
 - impacts on habitats in the bay
 - affects access to ports
 - has implications for flooding during storm events



Scale
0 1 2 3
kilometres

Picture courtesy of Nigel Cross,
Lancaster City Council

Morphodynamic modelling

- Operational coastal flood forecasting is limited near-shore by lack of knowledge of evolving bathymetry
 - but it is impractical to continually monitor large coastal areas
- Modelling is difficult
 - longer term changes are driven by shorter term processes
 - uncertainty in initial conditions and parameters
- An alternative approach is to use data assimilation
 - improved estimates of current model bathymetry
 - improved predictions of future bathymetry
 - better value for money for investments in observations & model development



Parameter estimation

- Model equations depend on parameters
 - exact values are unknown
 - inaccurate parameter values can lead to growth of model error
 - affects predictive ability of the model
- How do we estimate these values *a priori*?
 - theoretical values
 - calibration

or ...

- data assimilation
 - choose parameters based on observations
 - state augmentation: model parameters are estimated alongside the model state

3D Var cost function

- Measures the distance of the solution from the background and observations weighted by the inverse of their errors

$$\tilde{J}(\mathbf{w}) = (\mathbf{w} - \mathbf{w}^b)^T \tilde{\mathbf{B}}^{-1} (\mathbf{w} - \mathbf{w}^b) + (\mathbf{y} - \tilde{\mathbf{h}}(\mathbf{w}))^T \mathbf{R}^{-1} (\mathbf{y} - \tilde{\mathbf{h}}(\mathbf{w}))$$

background term

observation term

$\mathbf{w} = \begin{pmatrix} \mathbf{z} \\ \mathbf{p} \end{pmatrix}$ is the augmented model state vector and $\tilde{\mathbf{B}}$ and \mathbf{R} are the covariance matrices of the background and observation errors.

Notation:

- \mathbf{z} model state vector
- \mathbf{p} vector of parameters
- \mathbf{w}^b background state
- \mathbf{y} vector of observations
- $\tilde{\mathbf{h}}$ augmented observation operator

Background error covariance

Augmented B matrix:

$$\tilde{\mathbf{B}} = \begin{pmatrix} \mathbf{B}_{zz} & \mathbf{B}_{zp} \\ (\mathbf{B}_{zp})^T & \mathbf{B}_{pp} \end{pmatrix}.$$

If we assume errors are unbiased

$\mathbf{B}_{zz} = E(\boldsymbol{\varepsilon}_b \boldsymbol{\varepsilon}_b^T)$ state covariance

$\mathbf{B}_{pp} = E(\boldsymbol{\varepsilon}_p \boldsymbol{\varepsilon}_p^T)$ parameter covariance

$\mathbf{B}_{zp} = E(\boldsymbol{\varepsilon}_b \boldsymbol{\varepsilon}_p^T)$ state parameter cross covariances

where $\boldsymbol{\varepsilon}_b = \mathbf{z}^b - \mathbf{z}^t$ and $\boldsymbol{\varepsilon}_p = \mathbf{p}^b - \mathbf{p}^t$

Augmented gain matrix

$$\begin{aligned}\tilde{\mathbf{K}} &= \tilde{\mathbf{B}}\tilde{\mathbf{H}}^T \left[\tilde{\mathbf{H}}\tilde{\mathbf{B}}\tilde{\mathbf{H}}^T + \mathbf{R} \right]^{-1} \\ &= \begin{pmatrix} \mathbf{B}_{zz}\mathbf{H}^T \\ \mathbf{B}_{zp}^T\mathbf{H}^T \end{pmatrix} \left[\mathbf{H}\mathbf{B}_{zz}\mathbf{H}^T + \mathbf{R} \right]^{-1} \\ &\stackrel{\text{def}}{=} \begin{pmatrix} \mathbf{K}_z \\ \mathbf{K}_p \end{pmatrix}\end{aligned}$$

$$\begin{aligned}\mathbf{z}^a &= \mathbf{z}^b + \mathbf{K}_z(\mathbf{y} - \mathbf{h}(\mathbf{z}^b)) \\ \mathbf{p}^a &= \mathbf{p}^b + \mathbf{K}_p(\mathbf{y} - \mathbf{h}(\mathbf{z}^b))\end{aligned}$$

Simple 1D model

Based on the sediment conservation equation

$$\frac{\partial z}{\partial t} = - \left(\frac{1}{1 - \varepsilon} \right) \frac{\partial q}{\partial x}$$

where $z(x,t)$ is the bathymetry, t is time, q is the sediment transport rate in the x direction and ε is the sediment porosity.

For the sediment transport rate we use the power law

$$q = Au^n$$

where $u(x,t)$ is the depth averaged current and **A and n are parameters whose values need to be set**


If we assume that water flux (F) and height (H) are constant

$$F = u(H - z)$$

we can rewrite the sediment conservation equation as

$$\frac{\partial z}{\partial t} + a(z, H, F, \varepsilon, A, n) \frac{\partial z}{\partial x} = 0$$

where $a(z, H, F, \varepsilon, A, n)$ is the advection velocity or bed celerity.





Can we use data assimilation to estimate the parameters A and n ?

State-parameter cross covariances

The Extended Kalman filter

Augmented system model:

$$\mathbf{w}_{k+1} = \begin{pmatrix} \mathbf{z}_{k+1} \\ \mathbf{p}_{k+1} \end{pmatrix} = \begin{pmatrix} \mathbf{f}(\mathbf{z}_k, \mathbf{p}_k) \\ \mathbf{p}_k \end{pmatrix} = \tilde{\mathbf{f}}(\mathbf{w}_k)$$

State forecast:

$$\mathbf{w}_{k+1}^f = \tilde{\mathbf{f}}_k(\mathbf{w}_k^a)$$

Error covariance forecast:

$$\mathbf{P}_{k+1}^f = \mathbf{F}_k \mathbf{P}_k^a \mathbf{F}_k^T$$

Tangent linear model:

$$\mathbf{F}_k = \left. \frac{\partial \tilde{\mathbf{f}}}{\partial \mathbf{w}} \right|_{\mathbf{w}_k^a} = \left(\begin{array}{cc} \frac{\partial \mathbf{f}(\mathbf{z}, \mathbf{p})}{\partial \mathbf{z}} & \frac{\partial \mathbf{f}(\mathbf{z}, \mathbf{p})}{\partial \mathbf{p}} \\ \mathbf{0} & \mathbf{I} \end{array} \right) \bigg|_{\mathbf{z}_k^a, \mathbf{p}_k^a} = \left(\begin{array}{cc} \mathbf{M}_k & \mathbf{N}_k \\ \mathbf{0} & \mathbf{I} \end{array} \right)$$

Error covariance forecast:

$$\mathbf{P}_{k+1}^f = \left(\begin{array}{cc} \mathbf{M}_k \mathbf{P}_{\mathbf{z}\mathbf{z}_k}^a \mathbf{M}_k^T + \mathbf{N}_k \mathbf{P}_{\mathbf{p}\mathbf{p}_k}^a \mathbf{N}_k^T & \mathbf{N}_k \mathbf{P}_{\mathbf{p}\mathbf{p}_k}^a \\ \mathbf{P}_{\mathbf{p}\mathbf{p}_k}^a \mathbf{N}_k^T & \mathbf{P}_{\mathbf{p}\mathbf{p}_k}^a \end{array} \right)$$

Hybrid approach



$$\tilde{\mathbf{B}}_k = \begin{pmatrix} \mathbf{B}_{zz} & \mathbf{N}_k \mathbf{B}_{pp} \\ \mathbf{B}_{pp} \mathbf{N}_k^T & \mathbf{B}_{pp} \end{pmatrix}$$



State-parameter cross covariances for our simple model are given by



$$\begin{aligned} \mathbf{B}_{zpk} &= \mathbf{N}_k \mathbf{B}_{pp} \\ &= \begin{pmatrix} \frac{\partial \mathbf{f}_k}{\partial A} & \frac{\partial \mathbf{f}_k}{\partial n} \end{pmatrix} \begin{pmatrix} \sigma_A^2 & \sigma_{An} \\ \sigma_{An} & \sigma_n^2 \end{pmatrix} \\ &= \begin{pmatrix} \sigma_A^2 \frac{\partial \mathbf{f}_k}{\partial A} + \sigma_{An} \frac{\partial \mathbf{f}_k}{\partial n} & \sigma_n^2 \frac{\partial \mathbf{f}_k}{\partial n} + \sigma_{An} \frac{\partial \mathbf{f}_k}{\partial A} \end{pmatrix} \end{aligned}$$



Model setup



- Assume perfect model and observations
- Identical twin experiments
 - reference solution generated using Gaussian initial data and parameter values $A = 0.002 \text{ ms}^{-1}$ and $n = 3.4$
- Use incorrect model inputs
 - inaccurate initial bathymetry
 - inaccurate parameter estimates
- 3D Var algorithm is applied sequentially
 - observations taken at fixed grid points & assimilated every hour
 - the cost function is minimized iteratively using a quasi-Newton descent algorithm
- Covariances
 - \mathbf{B}_{zz} fixed
 - \mathbf{B}_{zp} time varying











Summary

- Presented a novel approach to model parameter estimation using data assimilation
 - demonstrated the technique using a simple morphodynamic model
- Results are very encouraging
 - scheme is capable of recovering near-perfect parameter values
 - improves model performance
- What next ...?
 - can our scheme be successfully applied to more complex models?
 - can we determine an optimal observing network?

Morecambe Bay at sunset



Simple Models of Changing Bathymetry with Data Assimilation

P.J Smith, M.J. Baines, S.L. Dance, N.K. Nichols and T.R. Scott

Department of Mathematics, University of Reading

Numerical Analysis Report 10/2007*

Data Assimilation for Parameter Estimation with Application to a Simple Morphodynamic Model

P.J Smith, M.J. Baines, S.L. Dance, N.K. Nichols and T.R. Scott

Department of Mathematics, University of Reading

Mathematics Report 2/2008*

Variational data assimilation for parameter estimation: application to a simple morphodynamic model

P.J Smith, M.J. Baines, S.L. Dance, N.K. Nichols and T.R. Scott

Submitted to Ocean Dynamics PECS 2008 Special Issue*

*available from <http://www.reading.ac.uk/maths/research/>