

# *The role of the pod in seed development: strategies for manipulating yield*

Article

Accepted Version

Bennett, E. J., Roberts, J. A. and Wagstaff, C. ORCID:  
<https://orcid.org/0000-0001-9400-8641> (2011) The role of the  
pod in seed development: strategies for manipulating yield.  
New Phytologist, 190 (4). pp. 838-853. ISSN 0028-646X doi:  
<https://doi.org/10.1111/j.1469-8137.2011.03714.x> Available at  
<https://centaur.reading.ac.uk/26411/>

It is advisable to refer to the publisher's version if you intend to cite from the  
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/j.1469-8137.2011.03714.x>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law,  
including copyright law. Copyright and IPR is retained by the creators or other  
copyright holders. Terms and conditions for use of this material are defined in  
the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online



## The Role of the Pod in Seed Development: Strategies for Manipulating Yield

Journal:	<i>New Phytologist</i>
Manuscript ID:	NPH-TR-2011-11339.R1
Manuscript Type:	TR - Commissioned Material - Tansley Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Bennett, Emma; University of Reading, Department of Food and Nutritional Sciences Roberts, Jeremy; Nottingham University, School of Biosciences Wagstaff, Carol; University of Reading, Department of Food and Nutritional Sciences, Centre for Food Security,
Key Words:	pod, seed, silique, senescence, resource allocation, yield, Arabidopsis, Brassica

SCHOLARONE™  
Manuscripts

1 *Tansley Review*

2 **The Role of the Pod in Seed Development: Strategies for Manipulating Yield**

3 Emma J Bennett<sup>1</sup>, Jeremy A Roberts<sup>2</sup>, and Carol Wagstaff<sup>1</sup>

4 <sup>1</sup> Department of Food and Nutritional Sciences, University of Reading, PO Box 226, Whiteknights, Reading,  
5 RG6 6AP, UK; <sup>2</sup> Plant and Crop Sciences Division, School of Biosciences, University of Nottingham, Sutton  
6 Bonington Campus, Loughborough, Leicestershire, LE12 5RD, UK

7  
8 *Author for correspondence:*

9 *Carol Wagstaff*

10 *Tel: +44 118 378 5362*

11 *Email: [c.wagstaff@reading.ac.uk](mailto:c.wagstaff@reading.ac.uk)*

12

13

14 **Key words:** pod, seed, silique, senescence, resource allocation, yield, nutrition, *Brassica*,

15 *Arabidopsis*

16

17

18

## 19 **The Role of the Pod in Seed Development: Strategies for Manipulating Yield**

### 20 **Summary**

21           Pods play a key role in encapsulating the developing seeds and protecting them from  
22 pests and pathogens. In addition to this protective function it has been shown that the  
23 photosynthetically active pod wall contributes assimilates and nutrients to fuel seed growth.  
24 Recent work has revealed that signals originating from the pod may also act to co-ordinate  
25 grain filling and regulate the reallocation of reserves from damaged seeds to those that have  
26 retained viability. In this review we consider the evidence that pods can regulate seed growth  
27 and maturation, particularly in members of the Brassicaceae family, and explore how the  
28 timing and duration of pod development might be manipulated to enhance either the quantity  
29 of crop yield or its nutritional properties.

30

### 31 **I. Introduction**

32           A pod, or silique as it is known in the Brassicaceae, is a photosynthetically active  
33 organ that encloses the seeds during their development. Once seed maturation is complete  
34 the bivalve pod splits longitudinally and its contents are released. In this review we will focus  
35 our discussions primarily on members of the Brassicaceae family and seek to demonstrate  
36 that the function of the pod extends far beyond simply safeguarding the maturing seeds. In  
37 this review we will refer to siliques by the more generic term pod, as much of the data  
38 generated from the study of Brassicaceous species has wider relevance across other families in  
39 the plant kingdom that reproduce through the formation of pods which enclose their seeds.  
40 Whilst pods are not essential for an individual's existence they play a paramount role in the  
41 survival of a species and could therefore be considered as one of the most important organs of  
42 a plant. In light of this it is perhaps surprising that relatively little research has been carried

43 out on pod growth and development and how this might be manipulated to enhance crop  
44 yield.

45 Seeds constitute an important source of dietary protein due to their high  
46 concentrations of seed storage proteins (SSPs) which can help alleviate malnutrition under  
47 circumstances when the consumption of animal protein is low. In the Brassicaceae family the  
48 main SSPs can be classified into either 12S globulins or 2S albumins, precursors of which are  
49 synthesised on the rough endoplasmic reticulum and, after maturation, reside in protein  
50 storage vacuoles within the seed (Herman & Larkins, 1999). Despite their nutritional  
51 benefits, seeds are deficient in some essential amino acids (see Mandal & Mandal, 2000 for a  
52 review) and are therefore incapable of completely alleviating protein malnutrition without  
53 supplementation. Hence yield enhancement is not only concerned with net increases in  
54 marketable produce but also in strategies to improve nutritional quality. The latter term  
55 encompasses many different aspects of seed composition but with regard to protein it  
56 represents the correlation between the amino acid profile of a seed and a balanced diet as  
57 recommended by the World Health Organisation (WHO) (Mandal & Mandal, 2000). Recent  
58 figures from WHO estimate that whilst the number of underweight children has fallen to 16%  
59 of the global population a simultaneous increase in inhabitants means that this still represents  
60 104 million undernourished children worldwide (World Health Organisation, 2010). In stark  
61 contrast, a global obesity epidemic is also occurring and as of the year 2000 WHO estimated  
62 that 300 million adults worldwide were obese (World Health Organization, 2000), a larger  
63 group than those considered to be undernourished. Intriguingly, there is evidence that one  
64 approach to help alleviate obesity might be to improve SSP levels and seed nutritional  
65 quality as high protein diets have been shown to assist in reducing and maintaining a healthy  
66 body mass (Claessens *et al.*, 2009).

67           The increasing acknowledgement that food security is a growing global problem was  
68 highlighted by Ban Ki-moon, speaking at a UN summit on solving the world's food crisis in  
69 2008, when he predicted that there would need to be a 50% increase in global food  
70 production by 2030. Thus, a rising world population (predicted to reach around 9 billion by  
71 2050), against a backdrop of climate change, makes the need to optimise yield of paramount  
72 importance. Maximising the efficiency of crop growth is one way global food demands can  
73 be met, and with a growing body of evidence that the pod directly influences seed  
74 composition it is bringing our ability to manipulate pod growth and development to the  
75 forefront of yield enhancement strategies.

76

## 77 **II. Pod Structure and Development**

78           The Brassicaceae family contains more than 3000 species and these produce non-  
79 fleshy fruit in the form of a silique that emerges from the gynoecium following ovule  
80 fertilisation. Recent transcript profiling analyses has provided evidence to support the  
81 assertion that the pod wall represents a modified leaf (Ma *et al.*, 2005; Scutt *et al.*, 2006;  
82 Wagstaff *et al.*, 2009). In the model *Brassica* species *Arabidopsis* the pod wall (pericarp) is  
83 composed of two fused carpels that undergo cell expansion between fertilisation and  
84 maturity, causing the pod to elongate about seven times its initial length between fertilisation  
85 and maturity (Sessions & Zambryski, 1995; Louvet *et al.*, 2006) . In contrast, the pods from  
86 members of the Fabaceae family are formed from a single carpel so, whilst the fruit from  
87 both families is commonly referred to as a pod, the term silique is reserved for members of  
88 the Brassicaceae family.

### 89 *2.1. The Pod Wall Structure*

90           The pericarp has been classified into three functional cell layers; the exocarp,  
91 mesocarp and endocarp, which are all characteristic components of fruit cell walls. The

92 exocarp comprises a single celled epidermal layer that is populated with stomata to facilitate  
93 gaseous exchange. The mesocarp is composed of layers of chlorenchyma cells that are rich in  
94 chloroplasts (Sessions & Zambryski, 1995). Finally, the endocarp consists of two dissimilar  
95 cell layers, a surface layer (ena) made up of large thin walled cells and an inner layer (enb)  
96 formed from small tightly packed cells as a result of several anticlinal cell divisions (Spence  
97 *et al.*, 1996). The silique wall is not entirely uniform and a narrow dehiscence zone (DZ),  
98 approximately two cell layers in width, spans the length of the silique between the valve and  
99 the replum (for review see Ferrandiz *et al.*, 1999; Ferrandiz, 2002). Such pericarp  
100 differentiation is necessary to target cellular degradation to the middle lamella between DZ  
101 cells, thus allowing the pod to shatter and release its mature seeds (Meakin & Roberts, 1990).  
102 Ultrastructural analysis of mature green silique walls has revealed developmental patterns  
103 associated with the onset of senescence, as they contain fewer thylakoids per granum than  
104 would normally be observed in leaf plastids (Wagstaff *et al.*, 2009). This feature, often  
105 associated with the structural reorganisation of thylakoid membranes, accompanies reduced  
106 PSI and PSII activity during senescence (Prakash *et al.*, 2001). Additionally, the decrease in  
107 chlorophyll in the pod wall precedes the decline observed in seeds which remain  
108 photosynthetic for longer (Wagstaff *et al.*, 2009); together these ultrastructural and  
109 physiological features may be required for optimisation of photosynthate accumulation in  
110 seeds. The consequences of such structural changes may be to enable a greater percentage of  
111 incident light to reach the seeds. In a crop such as oilseed rape (*Brassica napus*) this could  
112 potentially enhance its yield as ATP and NADPH are predicted to be required for the  
113 biosynthesis of seed storage products such as lipids (Fuhrmann *et al.*, 1994; Aach & Heise,  
114 1998; Schwender *et al.*, 2004; Goffman *et al.*, 2005). Given that only 20-30% of the incident  
115 light passes through the silique wall, and the spectral quality also changes in favour of the far  
116 red (FR) wavelengths, oilseed rape seeds develop within a shaded environment. These FR



117 wavelengths may also trigger the induction of seed dormancy, which is a common feature of  
118 freshly harvested seed from this crop, since such wavelengths are known to inhibit seed  
119 germination (Borthwick *et al.*, 1951). Using similar strategies to shade leaves, seeds have a  
120 lower chlorophyll a/b ratio to enable them to capture a greater amount of the available light  
121 (Eastmond *et al.*, 1996; King *et al.*, 1998) and it is reasonable to assume that selection  
122 pressures would have favoured a temporal separation of the loss of photosynthetic capacity  
123 between pods and seeds. One advantage of having a temporal division in photosynthetic  
124 maximums is that a greater amount of incident light is capable of reaching the developing  
125 seeds which can enhance ATP production leading to increased oil synthesis (Ruuska *et al.*,  
126 2002). High levels of anthocyanins are contained within the testa of *M. truncatula* seeds,  
127 which not only give the seeds colour (Abirached-Darmency *et al.*, 2005) but potentially  
128 impart protection against photo-oxidative stress; for a review of anthocyanin function see  
129 Archetti *et al.* (2009). As the silique wall begins the process of senescence and chlorophyll  
130 catabolism the enclosed seeds, which have matured within a shaded environment, are slowly  
131 exposed to higher levels of incident light. Consequentially the anthocyanins in the seed testa  
132 may afford some protection against increased UVB exposure and associated build up of  
133 reactive oxygen species.

## 134 2.2. Pod Wall Development

135 Despite being characteristically viewed merely as a protective organ, for instance the  
136 pod morphology of alfalfa has been shown to be instrumental in safeguarding seeds against  
137 chalcid wasps (Small & Brookes, 1984), the role of the pod appears to alter during the  
138 course of development. For instance, transcriptional profiling of the pod wall at different  
139 developmental stages has revealed that the observed changes in pod anatomy and chlorophyll  
140 levels throughout pod maturation correlated with alterations in transcription factor expression  
141 patterns (Wagstaff *et al.*, 2009). Indeed there is a strong association between pod

142 development and seed size (Pechan & Morgan, 1985) which has prompted the suggestion that  
143 pod length could be used as an indication of crop yield (Diepenbrock, 2000).

144

### 145 **III. The Pod as a Sink in Plant Resource Allocation**

146 The allocation of resources to developing siliques remains poorly understood, with the  
147 majority of research concentrating on nutrient remobilisation out of senescing leaves.

148 Nevertheless, it has been established that at anthesis the pod becomes a resource sink capable  
149 of storing remobilised nitrogen (N) and carbon (C) for utilisation upon germination (Harvey,

150 1973; BuchananWollaston, 1997; Diepenbrock, 2000; Rossato *et al.*, 2001; Schiltz *et al.*,

151 2005). This recycling of nutrients is essential for producing seeds that contain high

152 concentrations of storage compounds such as proteins, lipids and starch, which is why the

153 phloem remains functional throughout senescence (Feller & Fischer, 1994). Unlike other

154 plant organs, such as taproots, the pod is considered a sink throughout development (Rossato

155 *et al.*, 2001) although in practice the pod wall contributes assimilates to the developing

156 embryo during the final stages of seed maturation (Rochat & Boutin, 1991), for example,

157 20% of the N accumulated in pea seeds has been shown to be remobilised from the adjacent

158 pod wall (Schiltz *et al.*, 2005).

#### 159 *3.1. Nitrogen Uptake*

160 There is a considerable body of evidence to support the view that resources from

161 vegetative parts of the plant are remobilised into the pod (Harvey, 1973; Flinn *et al.*, 1977;

162 Schiltz *et al.*, 2005). <sup>15</sup>N labelling experiments have demonstrated that about 48% of the N

163 cycling through oilseed rape ends up in mature pods (Rossato *et al.*, 2001) and it is presumed

164 that most of this originates from vegetative tissues, since little N uptake occurs during

165 flowering and pod development. This observation indicates that N fertilisation after flowering

166 would only have minimal effects on plant yield. Our understanding of the pathways involved

167 in remobilising N from the leaves to developing pods are superficial, however, recent  
168 insights into the metabolic role of the enzyme pyruvate orthophosphate dikinase (PPDK) in  
169 metabolism may have shed some light on this problem. PPDK interconverts pyruvate and  
170 phosphoenolpyruvate and is central to photosynthesis in C4 plants, but it is also up-regulated  
171 during leaf senescence of C3 plants where it functions in a pathway that generates the  
172 transport amino acid glutamine, which is then loaded into the phloem. Over-expressing  
173 cytosolic PPDK results in more efficient amino acid transport and hence N remobilisation  
174 from the leaves is accelerated during senescence, leading to increases in Arabidopsis seed  
175 weight and N content, but not seed number (Taylor *et al.*, 2010). One possible explanation for  
176 these observations is that an extended growth period and larger rosette size might provide a  
177 greater initial resource pool in the vegetative tissues from which to reallocate storage  
178 compounds. However, while elevating the soil N content has been shown to enhance crop  
179 yield by increasing the number of pods per plant such an approach does not impact upon pod  
180 or seed weight (Allen & Morgan, 1972). This implies that the reproductive strategy of oilseed  
181 rape is to direct assimilates into additional pods when resources are plentiful rather than to  
182 produce seeds containing a greater concentration of storage products (Gammelvind *et al.*,  
183 1996). Combining the increased N remobilisation efficiency of PPDK over-expressers with  
184 higher soil N might substantially enhance yield in terms of both pod quantity and quality.

### 185 3.2. Leaf Senescence

186 Leaf senescence is a highly co-ordinated process that enables maximum recovery and  
187 remobilisation of nutrients from the leaves. At the onset of Arabidopsis leaf senescence there  
188 is an increase in the transcription of genes such as the ABC, sugar, peptide, amino acid and  
189 cation transporters in addition to the potential mobilisation of sulphur released upon protein  
190 degradation (Buchanan-Wollaston & Ainsworth, 1997). The start of leaf senescence is also  
191 accompanied by a concomitant increase in pod CO<sub>2</sub> metabolism indicating that this organ has

192 an elevated rate of photosynthesis (Gammelvind *et al.*, 1996; Robinson & Hill, 1999),  
193 potentially to enable nutrient uptake into the pod. As pods are photosynthetic organs, capable  
194 of generating reducing energy and ATP, their exact sink requirements are still a matter of  
195 debate. In oilseed rape, Allen and Morgan (1972) predicted that pods were capable of  
196 supporting their own growth, but subsequent examination of different *Brassica* species  
197 indicates that the photoassimilate contribution by the pod wall to developing seeds might be  
198 species specific (Ramana & Ghildiyal, 1997). The mechanisms by which resources are  
199 allocated into individual seeds is unknown for, whilst *Arabidopsis* fills its seeds in a uniform,  
200 co-ordinated manner, other species such as peas have larger and heavier seeds in the middle  
201 of their pods compared to those at the distal and proximal extremes of the pod (Harvey,  
202 1973). A fuller understanding of the source-sink relationship could prove to be crucial in  
203 improving crop yield as the sinks compete for the available resources.

### 204 3.3. Leaf-Pod Push-Pull Export and Import System

205 In some species the import of resources into the developing seeds is closely correlated  
206 with the capacity of leaves to export assimilates (Wittenbach & Vernon, 1983). However, in  
207 *Arabidopsis* the development of reproductive structures only minimally influences leaf  
208 senescence with organ age having a far greater effect. Selective pod removal had almost no  
209 impact on individual leaf senescence in *Arabidopsis*, but overall plant longevity was  
210 increased by 20-50 days according to Nooden & Penney (2001), but in our hands the rosette  
211 leaves remained green if all but the main inflorescence was removed (Figure 1). Whole plant  
212 senescence can be delayed through the removal of seeds from both pea and soybean pods  
213 (Lockhart & Gottschall, 1961; Lindoo & Nooden, 1977) and it has been proposed that the pea  
214 seed coat determines sink strength (Rochat & Boutin, 1992). Further work implies that this  
215 'pulling power' is coordinated with the breakdown of leaf storage products (Taylor *et al.*,  
216 2010) and it is likely that the pod sink strength is not a fixed entity but instead co-ordinated

217 throughout development to balance the ‘needs’ of the seeds, for instance N remobilisation  
218 decreases during the later stages of seed filling (Schiltz *et al.*, 2005). The plastic nature of  
219 whole plant resource allocation can be observed in many of the soybean de-podding  
220 experiments performed by Nooden and co-workers (Lindoo & Nooden, 1976; Lindoo &  
221 Nooden, 1977; Nooden *et al.*, 1978; Nooden & Murray, 1982) and from studies examining  
222 sterile mutants (Nooden & Penney, 2001). The number of pods also has the capacity to affect  
223 leaf photosynthesis, for instance selective pod removal in soybeans leads to reduced rates of  
224 CO<sub>2</sub> exchange within these plants, probably due to stomatal closure, a consequence of  
225 increased photoassimilate accumulation within the leaves brought about by having fewer  
226 sinks to export resources to (Setter & Brun, 1980; Wittenbach & Vernon, 1983). This  
227 indicates the presence of a dynamic feedback loop in which the pods signal their resource  
228 requirements to the leaves, causing the remobilisation of photoassimilates relative to the  
229 signal strength received (Figure 1). Hence, when fewer pods are present in de-podded plants  
230 the leaves temporarily halt photoassimilate production in response to an accumulation of  
231 carbohydrate in the leaves and an absence of ‘pull’ from the pods. An alternative view is that  
232 the sinks do not pull in resources but instead ‘free load’ by altering the conductance of  
233 plasmodesmata at the phloem-sink interface to affect the rate of nutrient unloading, matching  
234 this to their resource requirements (see Lalonde *et al.*, 2003 for a review).

235         Maximum remobilisation capacity is critical for R selected species, such as  
236 *Arabidopsis*, whose reproductive strategy is to produce viable seeds as quickly as possible. A  
237 potential response to such a life history trait is that *Arabidopsis* determines its seed set based  
238 upon the nutritional supply during the reproductive stage, rather than it being predetermined  
239 by growth and development during the vegetative phase, a trait associated with weeds  
240 growing in unpredictable environments (Schulze *et al.*, 1990). Such a trait could be desirable  
241 for commercial *Brassica* and legume crops to improve seed set.

242

243 **IV. Resource Transport into the Seeds via the Pod**244 *4.1. Transport from the pod wall*

245           The ability to manipulate resource partitioning and assimilate transport into the seeds  
246 and pods could help maximise overall yield (Wardlaw, 1990). Studies using detached pods  
247 have shown that the photosynthetic tissues of the pod wall are capable of generating 60% of  
248 seed assimilates (reviewed in Diepenbrock, 2000), although it must be noted that *in vitro* pod  
249 growth results in a decrease in internal pod O<sub>2</sub> concentrations compared to growth *in vivo*,  
250 which can alter the amount of storage compound within a seed (Musgrave *et al.*, 2008). As  
251 development progresses there is an increase in the compounds exported from the pod wall  
252 into the seeds. During this period the pod wall efficiently remobilises any accumulated N into  
253 the seeds such that upon harvest 80% of the total shoot N in oilseed rape has been relocated  
254 into the seeds. Seeds are capable of receiving the majority of their amino acids through the  
255 phloem-mediated pathway (Okumoto *et al.*, 2002), which can come from the pod wall as well  
256 as vegetative organs. Such remobilisation decreases during the later stages of development  
257 (Schjoerring *et al.*, 1995; Schiltz *et al.*, 2005), potentially due to the absence of sucrose  
258 synthase (SUS) activity in the pod wall and funiculus (Fallahi *et al.*, 2008) which is predicted  
259 to provide energy for phloem loading and unloading of solutes at this site (Fallahi *et al.*,  
260 2008). Whether photoassimilates generated during pod wall photosynthesis can be re-  
261 allocated to other pods is not yet clear, but such a mechanism could prove valuable given that  
262 pods higher up in the canopy receive a greater amount of incident light and are thus capable  
263 of increased photosynthetic rates whilst being largely exempt from the problems of self  
264 shading.

265 *4.2. Phloem unloading at the pod*

266           The import of resources into the pod is primarily concerned with phloem unloading,  
267 but this area has received little attention compared with phloem loading at the source site  
268 (Patrick, 1997), which is partly due to the great diversity between different sink types. Seeds  
269 are well adapted for the uptake of photoassimilates translocated from the pod wall. In  
270 *Medicago truncatula* the micropylar region of the seed coat, a small opening in the outer  
271 epidermis of the ovule located at one end of the seed, contains a vascular system believed to  
272 be instrumental in nutrient transport into the developing seed (Abirached-Darmency *et al.*,  
273 2005). Photoassimilates enter the funiculus which leads to the vascular bundle in the seed  
274 coat where unloading can occur (Van Dongen, 2003; Stadler *et al.*, 2005).

275 Whilst there is a comprehensive understanding of how resources are transported via the  
276 xylem and phloem to the pod junction subsequent steps describing the mechanism of transfer  
277 from the pod petiole to the seed-funiculus are still poorly defined. The current notion is that  
278 transport initially occurs through the symplastic pathway using plasmodesmata and is driven  
279 by simple diffusion and or bulk flow. However solutes must be subsequently translocated via  
280 apoplastic pathways to move between seed and pod tissues, plus the presence of a selectively  
281 permeable apoplastic pathway connecting the maternal tissue and phloem helps prevent  
282 nutrient loss. In addition, an apoplastic pathway can run in parallel to the symplastic pathway  
283 between the phloem and maternal tissues, but the exact details surrounding resource transport  
284 into the seeds are still under review (Patrick & Offler, 2001; Lalonde *et al.*, 2003).

285           It is predicted that the concentration of storage products within the seeds helps  
286 regulate the efficiency with which resources are transported around the plant (Schulze *et al.*,  
287 1994). One such mechanism of regulating supply and demand between the phloem and pod  
288 involves the reduction of apoplastic sucrose levels within sink tissues (reviewed in Patrick,  
289 1997), a further indication that the pods act as sinks to pull resources in from the surrounding

290 tissues. This flow of solutes is driven along the phloem by passive transport caused by  
291 differences in turgor pressure between the source and sink organs (Patrick, 1997).

#### 292 4.3. Nutrient transporters

293 Numerous nutrient transporters are located within the funiculus and at the base of the  
294 pod in the pedicel. One of the many Arabidopsis sulphate transporters, *SULTR2-1*, controls  
295 the translocation of sulphur into seeds and is potentially capable of regulating the import of  
296 this element into seed storage proteins, (Awazuhara *et al.*, 2005). Regulating sulphate uptake  
297 into pods can directly impact upon both seed quality and yield yet *SULTR2-1* mRNA levels  
298 do not alter regardless of the sulphur concentration that plants are grown in, highlighting the  
299 fundamental importance of this transporter in maintaining an import system (Awazuhara *et*  
300 *al.*, 2005).

301 Since only small amounts of nitrate are directly translocated from the roots into the  
302 seeds (Chopin *et al.*, 2007), by itself nitrate is unlikely to contribute much N towards seed  
303 nutrition but instead it is predicted to serve as a signalling molecule, or to alter the osmotic  
304 balance during the early stages of seed filling (McIntyre, 1997; Chopin *et al.*, 2007), although  
305 during periods of nitrogen deficiency it was postulated that nitrates might have a greater role  
306 to play in enhancing seed nutrition (Fan *et al.*, 2009). As the Arabidopsis amino acid  
307 transporter *AAP8*, which has a similar expression pattern to *SULTR2-1* (Awazuhara *et al.*,  
308 2005), is present in the funiculus and pod vascular tissue it is predicted to be responsible for  
309 enabling the import of organic nitrogen into the seeds (Okumoto *et al.*, 2002), leading to the  
310 hypothesis that amino acids enter the pod vascular tissue and are transported through the  
311 funiculus where they are imported into seeds at the micropylar region (Awazuhara *et al.*,  
312 2005).

#### 313 4.4. Impact of seed wounding



314           Although seeds clearly function as a sink for assimilates and nutrients during  
315 development the impact of seed wounding or abortion on transport processes has received  
316 little attention. All seeds within a pod are connected to the vascular trace by the funiculus. In  
317 *Arabidopsis*, the response regulator gene *ARR22* has been shown to be expressed specifically  
318 at the junction between the funiculus and chalazal tissues (Gattolin *et al.*, 2006; Horak *et al.*,  
319 2008). Despite chalazal tissues important role in supplying nutrients to the developing seeds,  
320 silencing of *ARR22* does not give rise to a morphologically detectable phenotype (Gattolin *et*  
321 *al.*, 2006; Horak *et al.*, 2008), although ectopically expressing the gene results in the  
322 generation of extremely dwarfed plants. If seeds from Pro<sub>ARR22</sub>::GUS plants are punctured  
323 then expression of the reporter is rapidly up-regulated in the chalazal region indicating that the  
324 *ARR22* transcript accumulates as a result of injury to the seed (Gattolin *et al.*, 2006). Recent  
325 research has revealed that the wounding of *Arabidopsis* pods is accompanied by rapid  
326 changes in transcript profile, causing mRNAs encoding seed storage proteins to dramatically  
327 decline, whereas those encoding proteins involved in proteolysis substantially increase,  
328 suggesting that damaged seeds may initiate a resource remobilisation programme (Naomab  
329 and Roberts unpublished). Intriguingly these changes do not occur in plants where the *ARR22*  
330 gene has been silenced, indicating that this response regulator protein plays an important role  
331 in signalling the presence of tissue damage within a seed. The demonstration that wounding  
332 causes an increase in the expression of genes involved in protein breakdown suggests that  
333 there may be a mechanism by which aborted seeds could redistribute their assimilates to  
334 support the development of those that will progress to maturation, indicating that one possible  
335 role for *ARR22* is to modulate assimilate partitioning into seeds contained within a pod. The  
336 precise mechanism that co-ordinates seed filling within an individual pod is unknown and,  
337 whilst in a weed such as *Arabidopsis* this is uniform, the trait may have been lost in some  
338 domesticated crops where seed size within an individual pod can vary considerably. The

339 spatial and temporal expression of *ARR22* make it a strong candidate for having a role in the  
340 co-ordination process and by manipulating its expression it might be possible to extend or  
341 reduce the seed filling period.

342

## 343 **V. Pod Senescence and Dehiscence**

### 344 *5.1. Method of Seed Dispersal*

345 At a plant level, uncoordinated pod senescence and dehiscence is advantageous and  
346 limits the seed loss that can occur if ripe pods shatter during temporarily adverse  
347 environmental conditions that subsequently inhibit germination of the next generation of  
348 plants. However, in a commercial setting, premature and uncoordinated pod shattering results  
349 in substantial pre-harvest losses and therefore significantly reduces net yield. For instance up  
350 to 20% of oilseed is lost per annum due to premature pod shatter whilst in Birdsfoot trefoil  
351 (*Lotus corniculatus* L.) this can be as high as 50% during adverse weather conditions  
352 (MacLeod, 1981). Such events can also impede subsequent crop growth due to the emergence  
353 of volunteer plants in the following growing season. Hence preventing premature pod shatter  
354 would instantly increase net crop yield, which, aside from the economic implications, would  
355 undoubtedly contribute to a viable solution of sustaining an escalating world population. Due  
356 to its economic importance, *Brassica napus* is a popular candidate species for investigating  
357 methods aimed at preventing or delaying pod shatter. Nevertheless, the development of a  
358 genotype that fails to shatter may not provide an ideal solution to this problem as this could  
359 compromise the postharvest processing chain due to difficulties in removing seeds from the  
360 pods without damage. A more amenable approach might be to delay or suspend the final  
361 stages of pod development after seed maturation to prevent individual pods shattering until  
362 all the pods have fully developed, at which time an internal or external stimulus could be  
363 applied to co-ordinate pod shattering across the crop.

### 364 5.2.1 Dehiscence

365 The pod wall of Brassicaceae family members is typically composed of two valves  
366 connected by a replum. In between these lignified cell types is a narrow band of valve margin  
367 cells that forms the dehiscence zone (DZ) and remains a non-lignified separation layer (SL)  
368 throughout pod development (Ferrandiz, 2002). Pod shattering is either initiated at the base of  
369 the pod where the pedicel meets the replum, as in oilseed rape (Morgan *et al.*, 2000), or at the  
370 pod tip and will continue along the DZ until the valves have completely separated (Davies &  
371 Bruce, 1997). A highly co-ordinated sequence of cellular and molecular events are required  
372 to bring about the dissolution of the middle lamella between cells of the DZ, and separation is  
373 precipitated in part by water loss from the pod wall, causing the valve cells to shrink and  
374 creating the tension necessary to pull them apart (Meakin & Roberts, 1990; Liljegren *et al.*,  
375 2004).

### 376 5.2.2 Genetic dissection of dehiscence

377 Much of the research undertaken in this area has focused on the differentiation and  
378 development of the DZ in the model species *Arabidopsis*. Formation of this non-lignified  
379 region is controlled by several MADS-Box genes which are capable of not only negatively  
380 regulating each other's expression but also acting independently in the valve, valve margin  
381 and replum cell layers. Expression of the functionally redundant *SHATTERPROOF 1*  
382 (*SHP1*) and *SHATTERPROOF 2* (*SHP2*) genes specifies the DZ and promotes lignification of  
383 adjacent cells (Flanagan *et al.*, 1996; Liljegren *et al.*, 2000). To confine the DZ to the valve  
384 margin *FRUITFULL* (*FUL*) represses *INDEHISCENT* (*IND*), *ALCATRAZ* (*ALC*) *SHP1* and  
385 *SHP2* expression in the adjacent valve cells as well as preventing lignification of the DZ (Gu  
386 *et al.*, 1998; Ferrándiz, 2000; Rajani & Sundaresan, 2001; Liljegren *et al.*, 2004). The  
387 transcription factor *SPATULA* (*SPT*) might also function in pod dehiscence since its  
388 expression is identical to that of *FUL* from mid pod development leading the authors to

389 propose that these two genes share regulatory roles (Heisler *et al.*, 2001), although more  
390 recent work from the same group has suggested that *SPT* is regulated by *IND* (Groszmann *et*  
391 *al.*, 2010). *SHP1* and *SHP2* transcriptionally activate *ALC* expression, which is required for  
392 establishing the SL between the valve margin and replum (Rajani & Sundaresan, 2001). At  
393 the onset of the pod shatter process, pectin in the SL cell walls is degraded by hydrolytic  
394 enzymes, such as ADPG1 and ADPG2 polygalacturonase (PG) enzymes which, in  
395 combination with increased pod wall tension, enables dehiscence to proceed (Ogawa *et al.*,  
396 2009). In addition to the PG enzymes it is predicted that a cyclic nucleotide-gated ion  
397 channel, *AtCNGC2*, might be involved in regulating programmed cell death within the DZ  
398 cells in Arabidopsis (Köhler *et al.*, 2001). The *IND* gene also functions downstream of the  
399 SHP transcription factors and is similarly required for differentiation of the valve margin  
400 cells, as well as lignification of the adjacent valve and replum cell layers (Liljegren *et al.*,  
401 2004). However *IND* also appears to be transcriptionally activated by factors other than *SHP1*  
402 and *SHP2*, since low expression levels can be detected in the valves in the *shp/shp2/ful* triple  
403 mutant (Ferrándiz, 2000; Liljegren *et al.*, 2004). *IND* forms a self regulating network and is  
404 also required for the expression of *ADPG1* in the DZ (Ogawa *et al.*, 2009); thus if *IND* is not  
405 expressed to define the valve margins then the PG enzyme which would breakdown this cell  
406 layer also fails to be produced. All five transcription factors involved in patterning the  
407 silique: *IND*, *ALC*, *SHP1*, *SHP2*, and *FUL* are required for lignification of the valve layer and  
408 hence seed dispersal (Liljegren *et al.*, 2004). Furthermore, these MADS-box genes are  
409 predicted to be repressed by *REPLUMLESS* (*RPL*) which functions to specify the replum cell  
410 layer adjacent to the valve margin and hence maintain the DZ at the valve margin (Roeder *et*  
411 *al.*, 2003).

412 Mutating the transcription factors involved in specifying the Arabidopsis DZ can  
413 create an indehiscence phenotype, for instance in the *FUL* gain of function (Ferrándiz, 2000),

414 *shp1shp2* double knockout (Liljegren *et al.*, 2000), *IND* loss of function (Liljegren *et al.*,  
415 2004) and *ALC* loss of function (Rajani & Sundaresan, 2001) plants the DZ is prevented  
416 from forming properly, demonstrating that the genetic manipulation of orthologues in  
417 *Brassica* species might be a suitable strategy for controlling pod shatter. However, there are  
418 potential limitations to this strategy as ectopic expression of the *FUL* gene in *B. juncea* has  
419 been revealed to make pods resistant to threshing (Østergaard *et al.*, 2006). A more recent  
420 approach showed that it is possible to fine-tune the severity of the shatter phenotype through  
421 inducing point mutations in *Brassica* orthologues of the *IND* gene and selecting those  
422 variants with a commercially useful degree of valve margin disruption (Girin *et al.*, 2010).  
423 Whilst wheat has been cultivated for thousands of years, the large scale commercialisation of  
424 oilseed rape is a more recent development and, within the genetic diversity of material that  
425 exists, some cultivars exhibit a greater resistance to pod shatter than commercial varieties  
426 (Morgan *et al.*, 1998). This diversity provides an extensive genetic pool that can be  
427 investigated and should aid the identification of better mechanisms for successfully  
428 controlling pod dehiscence.

### 429 5.3. Seed abscission

430 For seed dispersal to take place, not only does the pod have to ‘unzip’, but the seed  
431 must also detach from the funiculus. Like dehiscence, the regulation of seed abscission is a  
432 highly co-ordinated event culminating in wall dissolution at the hilum. Work on *Arabidopsis*  
433 has shown that *HECATE3* (*HEC3*), which directs the expression of *ADPG1* in the seed  
434 abscission zone, and *SEEDSTICK* (*STK*) are required for normal seed shedding (Pinyopich *et*  
435 *al.*, 2003; Ogawa *et al.*, 2009).. This aspect of plant development provides a further avenue  
436 for manipulating and potentially co-ordinating crop yield.

### 437 5.4. Hormonal regulation of dehiscence and seed abscission

438 A role for the plant hormones ethylene and auxin in regulating the timing of  
439 abscission has been documented extensively, with ethylene promoting and auxin inhibiting  
440 the process (Sexton & Roberts, 1982). Although a peak in ethylene production has been  
441 shown to precede dehiscence, exposure to the gas does not hasten pod shatter (Meakin &  
442 Roberts, 1990). Changes in auxin levels during pod development have also been identified  
443 but it is not clear to what extent the hormone regulates the dehiscence process (Johnson-  
444 Flanagan & Spencer, 1994; Chauvaux *et al.*, 1997; Child *et al.*, 1998). A recent publication  
445 (Sorefan *et al.*, 2009) demonstrated that local changes in pod auxin concentration are crucial  
446 for the differentiation of the DZ. The local auxin minimum generated at the valve margins  
447 seems to be produced by *IND* which acts to regulate auxin transport and as such increasing  
448 indole-3-acetic acid (IAA) levels at the valve margin leads to the development of an  
449 indehiscent phenotype due to the absence of this cell layer (Sorefan *et al.*, 2009). A recent  
450 publication by Arnaud *et al.* (2010) showed that gibberellin is a direct target, and is  
451 absolutely required, for the correct functioning of *IND*. The same authors concluded that *ALC*  
452 interacts directly with DELLA repressors, which antagonize *ALC* function but are  
453 destabilized by gibberellin. Taken together, these findings show that the gibberellin/DELLA  
454 pathway has a key role in patterning the Arabidopsis fruit and its eventual dehiscence.

455 Seeds are a major source of ethylene synthesis but their climateric production of the  
456 gas may only accelerate the onset of senescence rather than promote dehiscence *per se* (John  
457 *et al.*, 1995). This is highlighted by examining parthenocarpic pods which also produce a  
458 peak in ethylene and undergo shatter, albeit at a delayed rate, indicating that the pod wall has  
459 the capacity to produce the gas (Meakin & Roberts, 1990; Child *et al.*, 1998). The  
460 Arabidopsis protein AtTRP1, an orthologue of a tomato protein which interacts with the  
461 tomato ethylene receptors LeETR1 and NR, is highly expressed in the seed abscission zone

462 (Lin *et al.*, 2009). This observation suggests a possible role for AtTRP1 in regulating seed  
463 shedding which could be tested in lines where the gene is silenced or over-expressed.

464

## 465 **VI. The Role of Plant Phytohormones in Pod Development**

### 466 *6.1.1 The influence of salicylic acid on seed yield*

467 Plant phytohormones function in many aspects of development including cell  
468 differentiation, elongation, pattern formation and coping with abiotic and biotic stresses, all  
469 of which help maintain a high reproductive capacity. Depending on the tissue location and  
470 developmental stage phytohormones can either act synergistically or antagonistically towards  
471 one other which, in addition to pleiotropic effects, make their roles in pod and seed  
472 development difficult to discern.

473 Phytohormones can directly influence yield, for instance decreasing salicylic acid  
474 (SA) levels in Arabidopsis *NahG* transgenic lines and *sid2* mutants increases both the number  
475 of seeds per pod and the number of pods per plant, the latter resulting from an enhanced  
476 branching phenotype (Abreu & Munné-Bosch, 2009). Such physiological alterations were  
477 also correlated with a change in seed composition whereby N, vitamin E and pro-vitamin A  
478 content were enriched. This increase is a likely consequence of the late flowering and delayed  
479 senescent phenotype associated with SA deficient plants, thus enabling a longer period for  
480 resource translocation into the developing seeds (Martinez *et al.*, 2004; Abreu & Munné-  
481 Bosch, 2009). Such findings complement other studies which demonstrate that the  
482 constitutive overproduction of SA reduces seed yield (Mauch *et al.*, 2001). A decrease in  
483 seed weight was however reported in plants with reduced SA abundance (Abreu & Munné-  
484 Bosch, 2009) and, since germination potential was never measured, it is unknown how these  
485 altered ratios of nutritional compounds in the seed affect viability. Despite the implications  
486 for improving crop yield the effects of manipulating SA levels are not fully understood and,

487 in light of the fact that exogenous application increased seed yield in a grass species (Joaquin  
488 *et al.*, 2007), in contrast to the findings in *Arabidopsis*, this highlights a potentially species  
489 specific role for SA. In addition to this SA has a fundamental role in plant defence against  
490 microbial pathogen attack (reviewed in Vlot *et al.*, 2009) and environmental stresses  
491 (reviewed in Horváth *et al.*, 2007), hence complete KOs are unlikely to be commercially  
492 viable. Nevertheless, SA knockout lines that are regulated by a pod-specific promoter might  
493 extend the pod developmental period and allow more resource reallocation into the seeds,  
494 without compromising innate resistance strategies.

#### 495 6.1.2 Ethylene mutants

496 Since the discovery of ethylene as a biologically active and readily diffusible plant  
497 growth regulator (Neljubov, 1901) it has been associated with many processes including, but  
498 not restricted to, seed germination, growth, timing of organ senescence, fruit ripening and  
499 abscission (Abeles *et al.*, 1992). Ethylene can temporally and spatially regulate numerous  
500 aspects of plant development, with fleshy and dehiscent fruits becoming more competent to  
501 respond to ethylene ripening signals as they age (Joaquin *et al.*, 2007). For instance a burst in  
502 seed ethylene production correlates with the onset of pod dehiscence (Oeller *et al.*, 1991),  
503 highlighting the importance of ethylene in regulating the timing of developmental events,  
504 even if it is not necessarily inducing such responses. The interaction between different  
505 hormone pathways remains largely undiscovered, although ethylene is currently known to  
506 assist in plant responses to JA, SA, auxin ABA and cytokinin signalling and together they  
507 play an important role in responding to biotic and abiotic stresses. This has led to  
508 considerable effort being invested in uncovering the ethylene response pathway (for reviews  
509 see Ecker & Stepanova, 2000; Guo & Ecker, 2004) with recent studies focusing on the  
510 mechanisms of sensing and reacting to ethylene signals through a family of cell surface  
511 receptors. *Arabidopsis* has five ethylene receptors (encoded by *ERS1*, *ERS2*, *ETR1*, *ETR2* and



512 *EIN4*) and mutations conferring dominant ethylene insensitivity all occur in the hydrophobic  
513 regions of the N-terminal ethylene sensor domain (Bleecker *et al.*, 1988; Hua *et al.*, 1995;  
514 Ecker *et al.*, 1998), implying that there are only a limited number of genetic locations in  
515 which mutations are capable of causing ethylene insensitivity (Bleecker *et al.*, 1998). The fact  
516 that ethylene receptor mutants have subtly altered phenotypes and encode distinct proteins  
517 implies a functional specificity for the different receptors. This view was upheld by Zhou *et*  
518 *al.* (2007) who argued that they are not functionally redundant but, as previously suggested,  
519 may mediate the response of more than one signal (Bleecker *et al.*, 1998). For instance a link  
520 between glucose sensitivity and the ethylene pathway (Zhou *et al.*, 1998) has been made  
521 since glucose acts to decrease *EIN3* levels (Yanagisawa *et al.*, 2003), whilst in the monocot  
522 rice it has been shown that reduced expression of the ethylene receptor *ETR2* can increase  
523 thousand grain weight by up to 4% through altering starch acclimation and increasing sugar  
524 translocation into the filling grains (Wuriyanghan *et al.*, 2009). The ethylene insensitive  
525 receptor mutant *etr1-1* demonstrates a retarded leaf senescence phenotype which corresponds  
526 to a delay in the expression of other senescence associated genes (SAGs). However, this  
527 extended visual longevity does not correlate with functionality of the photosynthetic  
528 apparatus, so *etr1-1* leaves have entered into the senescence programme despite retaining  
529 higher chlorophyll levels for longer (Grbic & Bleecker, 1995), potentially indicating that  
530 *etr1-1* mutants are unlikely to positively affect seed yield.

531 Examination of microarray data for Arabidopsis (Table 1; developmental data from  
532 Wagstaff *et al.*, 2009; wound response data taken from Naomab, 2008) revealed that the only  
533 ethylene receptor showing at least a doubling of transcript levels during developmental pod  
534 wall senescence was *ETR2* which increased 2.8-fold from mature green to yellow senescent  
535 pod walls. This gene did not produce any signal on the wounded tissue arrays, indicating that  
536 it does not have a role in the wound response at the transcriptional level; the same could be

537 said for *EIN4* but to a lesser extent. *ERS1* and *ERS2* were up-regulated 1.9-fold and 1.5-fold  
538 respectively in wild type pods 90min after wounding the intact pods multiple times with a  
539 pin, although only *ERS1* showed any developmental response, perhaps indicating that there is  
540 a segregation of ethylene receptors with respect to the signals they respond to. Most *ACC*  
541 *Synthase (ACS)* and *ACC Oxidase (ACO)* genes present in the Arabidopsis genome did not  
542 produce a signal on the microarrays, indicating that the process of ethylene biosynthesis is  
543 regulated at the post-transcriptional level. The exception was *ACO4* which was 19-fold up-  
544 regulated during developmental senescence and 11.4-fold increased 90 minutes after  
545 wounding. In contrast, *ACS2* does not appear to change during developmental senescence but  
546 it was 4.7-fold induced by the wound signal. Wounding also increased expression of genes  
547 encoding sugar transporters/signalling molecules, although these did not change during  
548 senescence. The glucose transporter *GPT1*, the hexokinase glucose sensor *HXK1*, *SUC2*,  
549 *SUC3*, *SUC4* and *SUC5* were all up-regulated by wounding. Of these *SUC2* increases the  
550 most (8-fold) within the 90 minute response period. It would appear therefore, that  
551 developmentally programmed resource allocation is regulated slightly differently to resource  
552 re-allocation that occurs after an unexpected event such as wounding which will compromise  
553 the viability of the seeds within that pod. Ethylene appears to have a stronger association with  
554 the wound response, despite its traditional links with the senescence process in other plant  
555 organs, and genes encoding sugar transporters within the pod only appear to be up-regulated  
556 after wounding, indicating that they may be more involved with resource export than import.  
557

## 558 6.2. A role for ABA in seed development

559 Abscisic acid (ABA) is traditionally associated with stress responses, and  
560 consequentially growth retardation, but in the absence of such environmental insults it is  
561 required for normal seed maturation and is able to promote cellular growth, including in the

562 pod (Cheng *et al.*, 2002). In wheat the ABA:ethylene ratio affects the rate of grain filling and,  
563 since this is quite sensitive, imposing a small stress such as mild drought amplifies ABA  
564 levels within wheat grains and correlates with an increase in grain filling (Yang *et al.*, 2006).  
565 Similar results have been observed in oilseed rape (*B. napus*) and *Medicago truncatula* where  
566 raised ABA levels induced by osmotic stress stimulated a higher production of SSP  
567 transcripts and accumulation of free amino acids respectively (Wilén *et al.*, 1990; Planchet *et*  
568 *al.*, 2010). Plant sensitivity to ABA is controlled by *ABI3*, which in turn is required for the  
569 accumulation of SSPs within the seed (Nambara *et al.*, 1992). Whilst the oilseed rape  
570 experiments described above were performed on excised embryos, it does suggest that a  
571 controlled application of ABA to the pods may increase the abundance of SSPs in the seeds  
572 without the need to implement a water stress.

### 573 6.3 A role for other phytohormones

574 Gibberellins (GAs) are another class of phytohormone that have numerous functions  
575 within the plant including helping to break seed dormancy, regulating plant growth and floral  
576 induction. In *Arabidopsis* normal pod development requires GA levels to be kept within a  
577 confined range, as increased concentrations result in fewer seeds per pod (Rieu *et al.*, 2008)  
578 and a decrease in pod wall length and weight (Srinivasan & Morgan, 1996). Correlations  
579 between GA and cytokinin levels also appear crucial for regulating pod wall growth  
580 (Srinivasan & Morgan, 1996). Fertilisation triggers an auxin-mediated promotion of GA  
581 synthesis specifically in the ovule which is then transported to the valves where GA targets  
582 DELLA proteins for degradation and therefore releases the repression of fruit growth seen in  
583 unfertilised pods (Marti *et al.*, 2007; Dorcey *et al.*, 2009).

584 Despite not being classed as a hormone, glucose also helps to regulate phytohormone  
585 levels and it is capable of functioning like a hormonal signalling molecule by indicating the  
586 plant's nutrient status (Arenas-Huertero *et al.*, 2000; Cheng *et al.*, 2002; Rolland *et al.*, 2002).

587 For instance, there are interactions between sugar and nitrogen signalling that can affect the  
588 carbon-nitrogen balance, (Sheen *et al.*, 1999), indicating that the capacity of a plant to sense  
589 changes in the glucose concentration within individual organs can regulate phytohormone  
590 production and consequentially mediate the source-sink nutrient balance (Cheng *et al.*, 2002).  
591 This theory is further supported by the observation that the Arabidopsis glucose  
592 insensitive/ABA-deficient mutant *gin1/aba2* has smaller pods than wild type and  
593 consequentially produces far more aborted embryos per pod than wild type, although any  
594 mature seeds are the normal size (Cheng *et al.*, 2002).

### 595 6.3. Altering the developmental period

596 Cytokinins help to regulate the timing of senescence and, since their levels fall at the  
597 onset of this process, exogenous application can delay senescence (Nooden *et al.*, 1979). This  
598 knowledge has enabled leaf senescence to be postponed by attaching a promoter from the  
599 senescence specific gene *SAG12* to the gene encoding isopentyl transferase (*IPT*), which  
600 catalyses cytokinin biosynthesis, generating auto-inhibition of senescence through the  
601 maintenance of pre-senescence cytokinin levels. In tobacco this prolonged the flowering  
602 period and photosynthetic lifespan which together resulted in a 50% increase in dry weight  
603 and seed yield, although it was not reported whether this also affected seed composition (Gan  
604 & Amasino, 1995). However, the *SAG12:IPT* construct in wheat only resulted in delayed  
605 senescence and not an increase in seed yield which the authors suggested was due to  
606 interference by the construct with the normally extremely rapid relocation of resources from  
607 senescing leaves to reproductive sinks (Sýkorová *et al.*, 2008). The tight correlation between  
608 developmental period and seed yield raises the possibility that a similar system of auto-  
609 regulation of senescence could be implemented to co-ordinate pod development, at least in  
610 dicotyledonous plants. Theoretically, delaying senescence and extending the photosynthetic  
611 period would increase the potential for seed filling and prevent the onset of dehiscence. The

612 process of senescence and dehiscence could subsequently be coordinated across the whole  
613 plant if the inhibition provided by cytokinin could be turned off in a controlled manner, for  
614 example by using an inducible promoter system.

615

## 616 **VII. Siliqua biosynthesis of compounds for the seed**

### 617 *7.1. Siliqua and seed photosynthesis*

618 Since the onset of leaf senescence occurs prior to the last pod forming, and before  
619 seed fill is complete, embryos have to rely upon pod or seed wall and stem photosynthesis to  
620 generate the remainder of their photoassimilates required for viable seed production.

621 Enclosure within a pod limits the photosynthetic capacity of the seed itself; in contrast the  
622 pod has a photosynthetic potential far greater than that of a leaf if assessed on the assimilate  
623 produced per unit of chlorophyll basis (King *et al.*, 1997). Carbon photosynthates stored  
624 within the pod wall are thought to be remobilised to developing seeds as a decrease in hexose  
625 levels corresponds with a concomitant increase in seed growth (King *et al.*, 1997).

626 Additionally *de novo* starch synthesis within oilseeds is presumed to be insufficient to  
627 account for the final oil levels observed, indicating the importance of translocating  
628 carbohydrates, such as sucrose and hexose, across the pod wall. To enable seeds to generate  
629 some of their own photoassimilates the pod wall in oilseed rape has a sclerenchyma cell layer  
630 nearest to the inner pod cavity that is predicted to act as a barrier to gas diffusion and  
631 therefore aid a build-up of CO<sub>2</sub> around the seeds (King *et al.*, 1998). Developing seeds are  
632 capable of fixing this CO<sub>2</sub> and consequentially generating energy for the synthesis of seed  
633 storage products, although predictions suggest that the quantity of CO<sub>2</sub> is not enough to  
634 sustain photosynthesis in the seed itself. In the pea pod, for example, it has been calculated  
635 that respiration accounts for the loss of more C than is incorporated into the fruit during the  
636 photosynthetic period. In the second half of seed development the pod is only capable of

637 producing about 10% of the carbon required by the seed (Flinn *et al.*, 1977) most likely due  
638 to onset of chlorophyll catabolism in the pod wall, but this reinforces the absolute necessity to  
639 re-allocate resources around the plant.

#### 640 *7.2.1 Seed storage proteins*

641 SSPs accumulate within both protein storage vacuoles and the endoplasmic reticulum  
642 (Crofts *et al.*, 2004). During the seed filling phase the import of amino acids into the embryo  
643 only occurs via the phloem and requires the amino acid co-transporter located within the pod  
644 vascular system that is encoded by *AAP2* (Hirner *et al.*, 1998). *AAP2* might therefore have an  
645 important role in transporting amino acids from vegetative plant organs and the pod wall into  
646 the seed. Should it emerge that SSP transcripts found in the pod wall (Wagstaff *et al.*, 2009)  
647 are transcribed into proteins which are subsequently translocated to the seeds *AAP2* might be  
648 involved in their transport. This leads to the possibility that targeting pod wall transcripts  
649 could increase the concentration of nutritionally essential sulphur proteins normally lacking  
650 within oil seeds.

#### 651 *7.2.2 Synthesising molecules in the pod wall*

652 A microarray analysis in *B. napus* revealed that seeds express genes encoding many  
653 essential storage compounds (Yu *et al.*, 2010). This finding does not automatically signify  
654 that all these genes were translated into proteins *in situ* but it does indicate that the seed does  
655 not necessarily need to import them from the pod wall. This needs to be weighed against the  
656 observation that, despite seeds containing the enzymes for synthesising some of their own  
657 compounds such as glucosinolates, they are still produced in the pod wall and translocated  
658 into the developing embryos (Bilsborrow *et al.*, 1993; Zhao *et al.*, 1993; Du & Halkier,  
659 1998). The reasons for a plant utilising this strategy is unclear; energetically it would be more  
660 efficient for the seeds to synthesise their own storage compounds so the factors that  
661 determine whether the seeds, pod wall, or both, translate these remain to be elucidated.

### 662 7.3. Lipids and oils

663 At higher light intensities seeds are capable of synthesising more fatty acids,  
664 indicating that an increased rate of photosynthesis might be responsible for this (Schwender  
665 *et al.*, 2004; Goffman *et al.*, 2005). Therefore it follows that either making the pod walls  
666 thinner to improve light penetration, or providing the seeds with more energy, could improve  
667 oil synthesis. This topic is laced with controversy, for whilst Eastmond *et al.* (1996) believes  
668 that the low level of incident light reaching a seed is insufficient to generate enough reducing  
669 power in the form of NADPH for lipid biosynthesis, two independent studies (Willms *et al.*,  
670 1999; Schwender *et al.*, 2004) refute this claim. Instead both support the notion that green  
671 oilseeds are well adapted to low light levels and as such can produce enough energy for fatty  
672 acid synthesis. Regardless of who is correct, the implication of this is that an increased oil  
673 content could be achieved by having seeds and or pod walls which remained  
674 photosynthetically active for longer, potentially by utilising the *SAG12:IPT* auto-regulation  
675 system discussed above or a stay green phenotype which remains functionally photosynthetic  
676 and has delayed senescence.

### 677 7.4. Translocation of molecules from the pod wall into the developing seeds

678 The transport of molecules from the pod wall into the seeds represents a centripetal  
679 mode of transport towards the inner integuments of the pod presumably via the single entry  
680 point at the base of the pod in the vascular system. Seeds of the legume species *Medicago*  
681 *truncatula* are highly specialised for the importation of nutrients with their micropylar region  
682 containing a vascular system organised into tracheids (Abirached-Darmency *et al.*, 2005).  
683 The presence of sucrose synthase (SUS) in phloem associated companion cells (Fallahi *et al.*,  
684 2008) when the siliques are fully mature supports the previously identified role of SUS in  
685 phloem loading/unloading (Martin *et al.*, 1993; Nolte & Koch, 1993). It is predicted that SUS  
686 could be vital for transporting assimilates generated in the pod wall into developing seeds via

687 the pod wall phloem (Fallahi *et al.*, 2008) especially since the SUS protein exhibits a spatial  
688 change throughout development. Initially SUS is highly expressed within the pod wall and  
689 funiculus, but by the later stages of development it is only found in the embryo and aleurone  
690 layer of the seed (Fallahi *et al.*, 2008), indicating a translocation of assimilates from the pod  
691 wall into the seeds via the phloem tissue.

692

### 693 **VIII. Conclusion**

694 This review has highlighted the contributions that a pod can make to the development  
695 of the encapsulated seed and identified strategies for manipulating resource allocation  
696 (summarised in Figure 2). In addition to providing protection from biotic and abiotic stresses  
697 it is evident that the photosynthetically active pod can contribute assimilates and nutrients  
698 that are subsequently imported into the developing seeds and a ‘push-pull’ model is proposed  
699 where the strength of the sink exerted by the seeds determines the degree to which resources  
700 are remobilised from other parts of the plant. Transcript profiling of the pod wall during  
701 development has revealed that the tissue shares features in common with leaf material and it  
702 is possible that shared events take place during pod and leaf senescence. Manipulation of the  
703 timing of pod senescence may make it possible to enhance the duration of grain filling and  
704 increase protein, carbohydrate or lipid content into the developing seed. In weed species, such  
705 as *Arabidopsis*, the partitioning of assimilates is highly co-ordinated within a pod so that at  
706 dehiscence all seeds are at an equivalent stage of maturation; although the timing of  
707 dehiscence across the plant, or a population of plants, is generally uncoordinated so that pods  
708 release their contents over a long period of time. In domesticated crops, such as peas and  
709 beans, grain filling may be less well orchestrated within a pod or an ear, and some seeds may  
710 act a stronger sinks than others. In contrast, coordination across the whole plant, and between  
711 plants in a monoculture, is extremely good as a consequence of the strong selection imposed



712 by man for uniformity as crops have been domesticated. The mechanism that co-ordinates  
713 assimilate import into seeds positioned at different sites in a pod is unclear. However, recent  
714 transcriptional analyses in *Arabidopsis* has identified a response regulator, *ARR22*, expressed  
715 within the micropylar tissues, that plays a key role in regulating the response of seeds to  
716 wounding and could contribute to regulating the assimilate import/export. Further 'omic  
717 analyses of pod tissues, particularly those of the pod wall, will assist in dissecting the  
718 contribution of the pod to the development of the seed. Armed with this information it  
719 should be possible to devise strategies to manipulate pod development so that we can not  
720 only enhance seed yield but also, and perhaps even more importantly, its nutritional value.

721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747

## References

- Aach H, Heise K. 1998.** On the compartmentation of triacylglycerol synthesis in developing seeds of *Brassica napus*. *Botanica Acta* **111**: 123-129.
- Abeles F, Morgan P, Saltveit M. 1992.** *Ethylene in Plant Biology*: Academic Press San Diego (USA).
- Abirached-Darmency M, Abdel-gawwad MR, Conejero G, Verdeil JL, Thompson R. 2005.** *In situ* expression of two storage protein genes in relation to histo-differentiation at mid-embryogenesis in *Medicago truncatula* and *Pisum sativum* seeds. *Journal of Experimental Botany* **56**: 2019-2028.
- Abreu ME, Munné-Bosch S. 2009.** Salicylic acid deficiency in *NahG* transgenic lines and *sid2* mutants increases seed yield in the annual plant *Arabidopsis thaliana*. *Journal of Experimental Botany* **60**: 1261-1271.
- Allen EJ, Morgan DG. 1972.** A quantitative analysis of the effects of nitrogen on the growth, development and yield of oilseed rape. *The Journal of Agricultural Science* **78**: 315-324.
- Archetti M, Döring TF, Hagen SB, Hughes NM, Leather SR, Lee DW, Lev-Yadun S, Manetas Y, Ougham HJ, Schaberg PG, Thomas H. 2009.** Unravelling the evolution of autumn colours: an interdisciplinary approach. *Trends in Ecology & Evolution* **24**: 166-173.
- Arenas-Huertero F, Arroyo A, Zhou L, Sheen J, Leo P. 2000.** Analysis of *Arabidopsis* glucose insensitive mutants , *gin5* and *gin6* , reveals a central role of the plant hormone ABA in the regulation of plant vegetative development by sugar. *Genes & Development* **14**: 2085-2096.
- Arnaud N, Girin T, Sorefan K, Fuentes S, Wood TA, Lawrenson T, Sablowski R, Østergaard L. 2010.** Gibberellins control fruit patterning in *Arabidopsis thaliana*. *Genes and Development* **24**: 2127-2132.
- Awazuhara M, Fujiwara T, Hayashi H, Watanabe-Takahashi A, Takahashi H, Saito K. 2005.** The function of SULTR2;1 sulfate transporter during seed development in *Arabidopsis thaliana*. *Physiologia Plantarum* **125**: 95-105.

- 748 **Bilsborrow PE, Evans EJ, Murray F, Zhao FJ. 1993.** Glucosinolate changes in developing pods of  
 749 single and double low varieties of autumn-sown oilseed rape (*B. napus*). *Annals of Applied*  
 750 *Biology* **122**: 135-143.
- 751 **Bleecker AB, Esch JJ, Hall AE, Rodríguez FI, Binder BM. 1998.** The ethylene-receptor family  
 752 from *Arabidopsis*: structure and function. *Philosophical Transactions of the Royal Society of*  
 753 *London. Series B, Biological Sciences* **353**: 1405-1412.
- 754 **Bleecker AB, Estelle MA, Somerville C, Kende H. 1988.** Insensitivity to ethylene conferred by a  
 755 dominant mutation in *Arabidopsis thaliana*. *Science* **241**: 1086-1089.
- 756 **Borthwick HA, Hendricks SB, Parker MW, Toole EH, Toole VK. 1951.** A reversible  
 757 photoreaction controlling seed germination. *Proceedings of the National Academy of Sciences*  
 758 **38**: 662-666.
- 759 **Buchanan-Wollaston V, Ainsworth C. 1997.** Leaf senescence in *Brassica napus*: cloning of  
 760 senescence related genes by subtractive hybridisation. *Plant Molecular Biology* **33**: 821-834.
- 761 **Buchanan-Wollaston V. 1997.** The molecular biology of leaf senescence. *Journal of Experimental*  
 762 *Botany* **48**: 181-199.
- 763 **Chauvaux N, Child R, John K, Ulvskov P, Borkhardt B, Prinsen E, Van Onckelen HA. 1997.**  
 764 The role of auxin in cell separation in the dehiscence zone of oilseed rape pods. *Journal of*  
 765 *Experimental Botany* **48**: 1423-1429.
- 766 **Cheng W-H, Endo A, Zhou L, Penney J, Chen H-C, Arroyo A, Leon P, Nambara E, Asami T,**  
 767 **Seo M, Koshiba T, Sheen J. 2002.** A unique short-chain dehydrogenase / reductase in  
 768 *Arabidopsis* glucose signaling and abscisic acid biosynthesis and function. *The Plant Cell* **14**:  
 769 2723-2743.
- 770 **Child RD, Chauvaux N, John K, Ulvskov P, Van Onckelen HA. 1998.** Ethylene biosynthesis in  
 771 oilseed rape pods in relation to pod shatter. *Journal of Experimental Botany* **49**: 829-838.
- 772 **Chopin F, Orsel M, Dorbe MF, Chardon F, Truong HN, Miller AJ, Krapp A, Daniel-Vedele F.**  
 773 **2007.** The *Arabidopsis* ATNRT2.7 nitrate transporter controls nitrate content in seeds. *Plant*  
 774 *Cell* **19**: 1590-1602.

- 775 **Claessens M, van Baak MA, Monsheimer S, Saris WHM. 2009.** The effect of a low fat, high  
776 protein or high carbohydrate *ad libitum* diet on weight loss maintenance and metabolic risk  
777 factors. *International Journal of Obesity* **33**: 296-304.
- 778 **Crofts AJ, Washida H, Okita TW, Ogawa M, Kumamaru T. 2004.** Targeting of proteins to  
779 endoplasmic reticulum-derived compartments in plants. The importance of RNA localization.  
780 *Plant Physiology* **136**: 3414-3419.
- 781 **Davies GC, Bruce DM. 1997.** Fracture mechanics of oilseed rape pods. *Journal of Materials Science*  
782 **32**: 5895-5899.
- 783 **Diepenbrock W. 2000.** Yield analysis of winter oilseed rape (*Brassica napus* L.): a review. *Field*  
784 *Crops Research* **67**: 35-49.
- 785 **Dorcey E, Urbez C, Blázquez MA, Carbonell J, Perez-Amador MA. 2009.** Fertilization-dependent  
786 auxin response in ovules triggers fruit development through the modulation of gibberellin  
787 metabolism in Arabidopsis. *Plant Journal* **58**: 318-332.
- 788 **Du LC, Halkier BA. 1998.** Biosynthesis of glucosinolates in the developing silique walls and seeds  
789 of *Sinapis alba*. *Phytochemistry* **48**: 1145-1150.
- 790 **Eastmond P, Koláčá L, Rawsthorne S. 1996.** Photosynthesis by developing embryos of oilseed rape  
791 (*Brassica napus* L.). *Journal of Experimental Botany* **47**: 1763-1769.
- 792 **Ecker JR, Hua J, Sakai H, Nourizadeh S, Chen QG, Bleecker AB, Meyerowitz EM. 1998.** *EIN4*  
793 and *ERS2* are members of the putative ethylene receptor gene family in Arabidopsis. *The*  
794 *Plant Cell* **10**: 1321.
- 795 **Ecker JR, Stepanova AN. 2000.** Ethylene signaling: from mutants to molecules. *Current Opinion in*  
796 *Plant Biology* **3**: 353-360.
- 797 **Fallahi H, Scofield GN, Badger MR, Chow WS, Furbank RT, Ruan Y-L. 2008.** Localization of  
798 sucrose synthase in developing seed and siliques of *Arabidopsis thaliana* reveals diverse roles  
799 for SUS during development. *Journal of Experimental Botany* **59**: 3283-3295.
- 800 **Fan S-C, Lin C-S, Hsu P-K, Lin S-H, Tsay Y-F. 2009.** The Arabidopsis nitrate transporter NRT1.7,  
801 expressed in phloem, is responsible for source-to-sink remobilization of nitrate. *The Plant*  
802 *Cell* **21**: 2750-2761.

- 803 **Feller U, Fischer A. 1994.** Nitrogen metabolism in senescing leaves. *Critical Reviews in Plant*  
804 *Science* **13**: 241-273.
- 805 **Ferrándiz C. 2002.** Regulation of fruit dehiscence in Arabidopsis. *Journal of Experimental Botany*  
806 **53**: 2031-2038.
- 807 **Ferrándiz C. 2000.** Negative regulation of the *SHATTERPROOF* Genes by *FRUITFULL* during  
808 Arabidopsis fruit development. *Science* **289**: 436-438.
- 809 **Ferrándiz C, Pelaz S, Yanofsky MF. 1999.** Control of carpel and fruit development in Arabidopsis.  
810 *Annual Review of Biochemistry* **68**: 321-354.
- 811 **Flanagan CA, Hu Y, Ma H 1996.** Specific regulation of the *AGL1* MADS-box gene suggests  
812 regulatory functions in Arabidopsis gynoecium and ovule development. *The Plant Journal*.  
813 **10**: 343-353.
- 814 **Flinn AM, Atkins Ca, Pate JS. 1977.** Significance of photosynthetic and respiratory exchanges in  
815 the carbon economy of the developing pea fruit. *Plant Physiology* **60**: 412-418.
- 816 **Fuhrmann J, Johnen T, Heise KP. 1994.** Compartmentation of fatty acid metabolism in zygotic  
817 rape embryos. *Journal of Plant Physiology* **143**: 565-569.
- 818 **Gammelvind L, Schjoerring J, Mogensen V, Jensen C, Bock J. 1996.** Photosynthesis in leaves and  
819 siliques of winter oilseed rape (*Brassica napus* L). *Plant and Soil* **186**: 227-236.
- 820 **Gan S, Amasino RM. 1995.** Inhibition of leaf senescence by autoregulated production of cytokinin.  
821 *Science* **270**: 1986-1988.
- 822 **Gattolin S, Alandete-Saez M, Elliott K, Gonzalez-Carranza Z, Naomab E, Powell C, Roberts**  
823 **JA. 2006.** Spatial and temporal expression of the response regulators *ARR22* and *ARR24* in  
824 *Arabidopsis thaliana*. *Journal of Experimental Botany* **57**: 4225-4233.
- 825 **Girin T, Stephenson P, Goldsack CM, Kempin SA, Perez A, Pires N, Sparrow PA, Wood TA,**  
826 **Yanofsky MF, Ostergaard L. 2010.** Brassicaceae *INDEHISCENT* genes specify valve  
827 margin cell fate and repress replum formation. *Plant Journal* **63**: 329-338.
- 828 **Goffman FD, Alonso AP, Schwender J, Shachar-Hill Y, Ohlrogge JB. 2005.** Light enables a very  
829 high efficiency of carbon storage in developing embryos of rapeseed. *Plant Physiology* **138**:  
830 2269-2279.

- 831 **Grbic V, Bleecker AB. 1995.** Ethylene regulates the timing of leaf senescence in Arabidopsis. *Plant*  
832 *Journal* **8**: 595-602.
- 833 **Groszmann M, Bylstra Y, Lampugnani ER, Smyth DR. 2010.** Regulation of tissue-specific  
834 expression of *SPATULA*, a bHLH gene involved in carpel development, seedling germination,  
835 and lateral organ growth in Arabidopsis. *Journal of Experimental Botany* **61**: 1495-1508.
- 836 **Gu Q, Ferrándiz C, Yanofsky MF, Martienssen R. 1998.** The *FRUITFULL* MADS-box gene  
837 mediates cell differentiation during Arabidopsis fruit development. *Development* **125**: 1509-  
838 1517.
- 839 **Guo H, Ecker JR. 2004.** The ethylene signaling pathway: new insights. *Current Opinion in Plant*  
840 *Biology* **7**: 40-49.
- 841 **Harvey D. 1973.** The Translocation of <sup>14</sup>C-photosynthate in *Pisum sativum* L. *Annals of Botany* **37**:  
842 787-794.
- 843 **Heisler MG, Atkinson a, Bylstra YH, Walsh R, Smyth DR. 2001.** *SPATULA*, a gene that controls  
844 development of carpel margin tissues in Arabidopsis, encodes a bHLH protein. *Development*  
845 **128**: 1089-1098.
- 846 **Herman EM, Larkins BA. 1999.** Protein storage bodies and vacuoles. *Plant Cell* **11**: 601-613.
- 847 **Hirner B, Fischer WN, Rentsch D, Kwart M, Frommer WB. 1998.** Developmental control of  
848 H<sup>+</sup>/amino acid permease gene expression during seed development of Arabidopsis. *Plant*  
849 *Journal* **14**: 535-544.
- 850 **Hoch Wa, Zeldin EL, McCown BH. 2001.** Physiological significance of anthocyanins during  
851 autumnal leaf senescence. *Tree Physiology* **21**: 1-8.
- 852 **Horak J, Grefen C, Berendzen KW, Hahn A, Stierhof YD, Stadelhofer B, Stahl M, Koncz C,**  
853 **Harter K. 2008.** The *Arabidopsis thaliana* response regulator *ARR22* is a putative AHP  
854 phospho-histidine phosphatase expressed in the chalaza of developing seeds. *BMC Plant*  
855 *Biology* **8**: 77.
- 856 **Horváth E, Szalai G, Janda T. 2007.** Induction of abiotic stress tolerance by salicylic acid signaling.  
857 *Journal of Plant Growth Regulation* **26**: 290-300.

- 858 **Hua J, Chang C, Sun Q, Meyerowitz EM. 1995.** Ethylene insensitivity conferred by Arabidopsis  
859 *ERS* gene. *Science* **269**: 1712-1714.
- 860 **Joaquin TBM, Trejo C, Hernandez-Garay A, Perez PJ, Garcia ADG, Quero CAR. 2007.** Effects  
861 of ethephon, salicylic acid and cidef-4 on the yield and quality of guinea grass seed. *Tropical*  
862 *Grasslands* **41**: 55-60.
- 863 **John I, Drake R, Farrell A, Cooper W, Lee P, Horton P, Grierson D. 1995.** Delayed leaf  
864 senescence in ethylene deficient acc-oxidase antisense tomato plants - molecular and  
865 physiological analysis. *Plant Journal* **7**: 483-490.
- 866 **Johnson-Flanagan AM, Spencer MS. 1994.** Ethylene production during development of mustard  
867 (*Brassica juncea*) and canola (*Brassica napus*) seed. *Plant Physiology* **106**: 601-606.
- 868 **King SP, Badger MR, Furbank RT. 1998.** CO<sub>2</sub> refixation characteristics of developing canola seeds  
869 and silique wall. *Australian Journal of Plant Physiology* **25**: 377-386.
- 870 **King SP, Lunn JE, Furbank RT. 1997.** Carbohydrate content and enzyme metabolism in developing  
871 canola siliques. *Plant Physiology* **114**: 153-160.
- 872 **Köhler C, Merkle T, Roby R, Neuhaus G. 2001.** Developmentally regulated expression of a cyclic  
873 nucleotide-gated ion channel from Arabidopsis indicates its involvement in programmed cell  
874 death. *Planta* **213**: 327-332.
- 875 **Lalonde S, Tegeder M, Throne-Holst M, Frommer WB, Patrick JW. 2003.** Phloem loading and  
876 unloading of sugars and amino acids. *Plant, Cell and Environment* **26**: 37-56.
- 877 **Liljegren SJ, Ditta GS, Eshed HY, Savidge B, Bowman JL, Yanofsky MF. 2000.**  
878 *SHATTERPROOF* MADS-box genes control seed dispersal in Arabidopsis. *Nature* **404**: 766-  
879 770.
- 880 **Liljegren SJ, Roeder AHK, Kempin Sa, Gremski K, Østergaard L, Guimil S, Reyes DK,**  
881 **Yanofsky MF. 2004.** Control of fruit patterning in Arabidopsis by *INDEHISCENT*. *Cell* **116**:  
882 843-853.
- 883 **Lin ZF, Ho CW, Grierson D. 2009.** *AtTRP1* encodes a novel TPR protein that interacts with the  
884 ethylene receptor ERS1 and modulates development in Arabidopsis. *Journal of Experimental*  
885 *Botany* **60**: 3697-3714.

- 886 **Lindoo SJ, Nooden LD. 1976.** The interrelation of fruit development and leaf senescence in ' Anoka '  
887 soybeans. *Science* **137**: 218-223.
- 888 **Lindoo SJ, Nooden LD. 1977.** On the behavior of the senescence signal in Anoka soybeans'.  
889 *Experimental Biology*: 1136-1140.
- 890 **Lockhart Ja, Gottschall V. 1961.** Fruit-induced & apical senescence in *Pisum sativum* L. *Plant*  
891 *Physiology* **36**: 389-398.
- 892 **Louvet R, Cavel E, Gutierrez L, Guenin S, Roger D, Gillet F, Guerineau F, Pelloux J. 2006.**  
893 Comprehensive expression profiling of the pectin methylesterase gene family during silique  
894 development in *Arabidopsis thaliana*. *Planta* **224**: 782-791.
- 895 **Ma L, Sun N, Liu X, Jiao Y, Zhao H, Deng XW. 2005.** Organ-specific expression of Arabidopsis  
896 genome during development. *Plant Physiology* **138**: 80-91.
- 897 **MacLeod J. 1981.** Harvesting in oilseed rape. *Oilseed Rape Book. A manual for growers, farmers*  
898 *and advisors (Green, C., ed.). Cambridge: Cambridge Agricultural Publishing*: 107-120.
- 899 **Mandal S, Mandal RK. 2000.** Seed storage proteins and approaches for improvement of their  
900 nutritional quality by genetic engineering. *Current Science* **79**: 576-589.
- 901 **Marti C, Orzaez D, Ellul P, Moreno V, Carbonell J & Granell A. 2007.** Silencing of DELLA  
902 induces facultative parthenocarpy in tomato fruits. *Plant Journal*. **52**: 865-876.
- 903 **Martin T, Frommer WB, Salanoubat M, Willmitzer L. 1993.** Expression of an Arabidopsis  
904 sucrose synthase gene indicates a role in metabolization of sucrose both during phloem  
905 loading and sink organs. *The Plant Journal* **4**: 367-377.
- 906 **Martinez C, Pons E, Prats G, Leon J. 2004.** Salicylic acid regulates flowering time and links  
907 defence responses and reproductive development. *The Plant Journal* **37**: 209-217.
- 908 **Mauch F, Mauch-Mani B, Gaille C, Kull B, Haas D, Reimmann C. 2001.** Manipulation of  
909 salicylate content in *Arabidopsis thaliana* by the expression of an engineered bacterial  
910 salicylate synthase. *The Plant Journal* **25**: 67-77.
- 911 **McIntyre G. 1997.** The Role of nitrate in the osmotic and nutritional control of plant development.  
912 *Australian Journal of Plant Physiology* **24**: 103-118.



- 913 **Meakin PJ, Roberts JA. 1990.** Dehiscence of fruit in oilseed rape (*Brassica napus* L.). *Journal of*  
 914 *Experimental Botany* **41**: 1003-1011.
- 915 **Morgan CL, Bruce DM, Child R, Ladbroke ZL, Arthur AE. 1998.** Genetic variation for pod  
 916 shatter resistance among lines of oilseed rape developed from synthetic *B. napus*. *Field Crops*  
 917 *Research* **58**: 153-165.
- 918 **Morgan CL, Ladbroke ZL, Bruce DM, Child R, Arthur AE. 2000.** Breeding oilseed rape for pod  
 919 shattering resistance. *Journal of Agricultural Science* **135**: 347-359.
- 920 **Musgrave M, Allen J, Blasiak J, Tuominen L, Kuang A. 2008.** *In vitro* seed maturation in *Brassica*  
 921 *rapa* L.: Relationship of silique atmosphere to storage reserve deposition. *Environmental and*  
 922 *Experimental Botany* **62**: 247-253.
- 923 **Nambara E, Naito S, McCourt P. 1992.** A mutant of *Arabidopsis* which is defective in seed  
 924 development and storage protein accumulation is a new *ABI3* allele. *The Plant Journal* **2**:  
 925 435-441.
- 926 **Naomab E. 2008.** The role of response regulators during *Arabidopsis* pod development. *University of*  
 927 *Nottingham Thesis*.
- 928 **Neljubov D. 1901.** Über die horizontale nutation der stengel von *Pisum sativum* und einiger anderer.  
 929 *Pflanzen Beitrage und Botanik Zentralblatt*. **10**: 128-139
- 930 **Nolte KD, Koch KE. 1993.** Companion-cell specific localization of sucrose synthase in zones of  
 931 phloem loading and unloading. *Plant Physiology* **101**: 899-905.
- 932 **Nooden LD, Kahanak GM, Okatan Y. 1979.** Prevention of monocarpic senescence in soybeans  
 933 with auxin and cytokinin: an antidote for self-destruction. *Science* **206**: 841-843.
- 934 **Nooden LD, Murray BJ. 1982.** Transmission of the monocarpic senescence signal via the xylem in  
 935 soybean. *Plant Physiology* **69**: 754-756.
- 936 **Nooden LD, Penney JP. 2001.** Correlative controls of senescence and plant death in *Arabidopsis*  
 937 *thaliana* (Brassicaceae). *Journal of Experimental Botany* **52**: 2151-2159.
- 938 **Nooden LD, Rupp DC, Derman BD. 1978.** Seraration of seed development from monocarpic  
 939 senescence in soybeans. *Nature* **271**: 354-357.

- 940 **Oeller PW, Min-wong L, Taylor LP, Pike DA, Athanasios T. 1991.** Reversible Inhibition of  
941 Tomato antisense RNA. *Science* **437**: 5-7.
- 942 **Ogawa M, Kay P, Wilson S, Swain SM. 2009.** Arabidopsis dehiscence zone polygalacturonase 1  
943 (*ADPG1*), *ADPG2*, and *QUARTET2* are polygalacturonases required for cell separation  
944 during reproductive development in Arabidopsis. *The Plant Cell* **21**: 216-233.
- 945 **Okumoto S, Schmidt R, Tegeder M, Fischer WN, Rentsch D, Frommer WB, Koch W. 2002.**  
946 High affinity amino acid transporters specifically expressed in xylem parenchyma and  
947 developing seeds of Arabidopsis. *Journal of Biological Chemistry* **277**: 45338-45346.
- 948 **Østergaard L, Kempin Sa, Bies D, Klee HJ, Yanofsky MF. 2006.** Pod shatter-resistant Brassica  
949 fruit produced by ectopic expression of the *FRUITFULL* gene. *Plant Biotechnology Journal*  
950 **4**: 45-51.
- 951 **Patrick JW. 1997.** Phloem unloading: sieve element unloading and post-sieve element transport.  
952 *Annual review of Plant Physiology and Plant Molecular Biology* **48**: 191-222.
- 953 **Patrick JW, Offler CE. 2001.** Compartmentation of transport and transfer events in developing  
954 seeds. *Journal of Experimental Botany* **52**: 551-564.
- 955 **Pechan Pa, Morgan DG. 1985.** Defoliation and its effects on pod and seed development in oil seed  
956 rape (*Brassica napus* L.). *Journal of Experimental Botany* **36**: 458-468.
- 957 **Pinyopich A, Ditta GS, Savidge B, Liljegren SJ, Baumann E, Wisman E, Yanofsky MF. 2003.**  
958 Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature*  
959 **424**: 85-88.
- 960 **Planchet E, Rannou1 O, Ricoult C, Boutet-Mercey S, Maia-Grondard A, Limami AM. 2010.**  
961 Nitrogen metabolism responses to water deficit act through both abscisic acid (ABA) -  
962 dependent and independent pathways in *Medicago truncatula* during post-germination.  
963 *Journal of Experimental Botany*: doi: 10.1093/jxb/erq294 online early.
- 964 **Prakash JS, Baig Ma, Mohanty P. 2001.** Senescence induced structural reorganization of thylakoid  
965 membranes in *Cucumis sativus* cotyledons; LHC II involvement in reorganization of  
966 thylakoid membranes. *Photosynthesis Research* **68**: 153-161.

- 967 **Rajani S, Sundaresan V. 2001.** The Arabidopsis myc / bHLH gene *ALCATRAZ* enables cell  
968 separation in fruit dehiscence. *Current Biology* **11**: 1914-1922.
- 969 **Ramana S, Ghildiyal MC. 1997.** Contribution of leaf photosynthesis towards seed yield in Brassica  
970 species. *Journal of Agronomy and Crop Science* **178**: 185-187.
- 971 **Rieu I, Eriksson S, Powers SJ, Gong F, Griffiths J, Woolley L, Benloch R, Nilsson O, Thomas  
972 SG, Hedden P, Phillips AL. 2008.** Genetic analysis reveals that C19 -GA 2-oxidation is a  
973 major gibberellin inactivation pathway in Arabidopsis. *The Plant Cell* **20**: 2420-2436.
- 974 **Robinson CK, Hill SA. 1999.** Altered resource allocation during seed development in Arabidopsis  
975 caused by the *abi3* mutation. *Plant Cell and Environment* **22**: 117-123.
- 976 **Rochat C, Boutin J-P. 1991.** Metabolism of phloem-borne amino acids in maternal tissues of fruit of  
977 nodulated or nitrate-fed pea plants (*Pisum sativum* L ). **42**: 207-214.
- 978 **Rochat C, Boutin J-P. 1992.** Temporary storage compounds and sucrose-starch metabolism in seed  
979 coats during pea seed development (*Pisum sativum*). *Physiologia Plantarum* **85**: 567-572.
- 980 **Roeder AHK, Ferra C, Yanofsky MF. 2003.** The role of the REPLUMLESS homeodomain protein  
981 in patterning the Arabidopsis fruit. *Current Biology* **13**: 1630-1635.
- 982 **Rolland F, Moore B, Sheen J. 2002.** Sugar sensing and signaling in plants. *Plant Cell* **14**: 185-205.
- 983 **Rossato L, Lainé P, Ourry a. 2001.** Nitrogen storage and remobilization in *Brassica napus* L. during  
984 the growth cycle: nitrogen fluxes within the plant and changes in soluble protein patterns.  
985 *Journal of Experimental Botany* **52**: 1655-1663.
- 986 **Ruuska SA, Girke T, Benning C, Ohlrogge JB. 2002.** Contrapuntal networks of gene expression  
987 during Arabidopsis seed filling. *Plant Cell* **14**: 1191-1206.
- 988 **Schiltz S, Munier-Jolain N, Judy C, Burstin J, Salon C. 2005.** Dynamics of exogenous nitrogen  
989 partitioning and nitrogen remobilization from vegetative organs in pea revealed by <sup>15</sup>n in  
990 vivo labeling throughout seed filling. *Plant Physiology* **137**: 1463-1473.
- 991 **Schjoerring J, Bock J, Gammelvind L, Jensen C, Mogensen V. 1995.** Nitrogen incorporation and  
992 remobilization in different shoot components of field-grown winter oilseed rape (*Brassica  
993 napus* L) as affected by rate of nitrogen application and irrigation. *Plant and Soil* **177**: 255-  
994 264.

- 995 **Schulze W, Schulze E, Stadler J, Heilmeier H, Mooney HA. 1990.** Growth and reproduction of  
996 *Arabidopsis thaliana* in relation to storage of starch and nitrate in the wild-type and in starch-  
997 deficient and nitrate-uptake-deficient mutants. *Environment*: 795-809.
- 998 **Schulze W, Schulze E, Stadler J, Heilmeier H, Stitt M, Mooney H. 1994.** Growth and  
999 Reproduction of *Arabidopsis thaliana* in relation to storage of starch and nitrate in the wild-  
1000 type and in starch deficient and nitrate uptake deficient mutants. *Plant Cell and Environment*  
1001 **17**: 795-809.
- 1002 **Schwender J, Ruuska SA, Ohlrogge JB. 2004.** The capacity of green oilseeds to utilize  
1003 photosynthesis to drive biosynthetic processes. *Plant Physiology* **136**: 2700-2709.
- 1004 **Scutt CP, Vinauger-Douard M, Fourquin C, Finet C, Dumas C. 2006.** An evolutionary  
1005 perspective on the regulation of carpel development. *Journal of Experimental Botany* **57**:  
1006 2143-2152
- 1007 **Sessions RA, Zambryski PC. 1995.** *Arabidopsis* gynoecium structure in the wild type and in *ettin*  
1008 mutants. *Development* **1532**: 1519-1532.
- 1009 **Setter TL, Brun WA. 1980.** Stomatal closure and photosynthetic inhibition in soybean leaves  
1010 induced by petiole girdling and pod removal. *Plant Physiology* **65**: 884-887.
- 1011 **Sexton R, Roberts JA. 1982.** Cell Biology of Abscission. *Annual Review of Plant Pathology* **33**:  
1012 133-162.
- 1013 **Sheen J, Zhou L, Jang JC. 1999.** Sugars as signaling molecules. *Current Opinion in Plant Biology*  
1014 **2**: 410-418.
- 1015 **Sýkorová B, Kurešová G, Daskalova S, Trčková M, Hoyerová K, Raimanová I, Motyka V,**  
1016 **Trávníčková A, Elliott MC, Kamínek M. 2008.** Senescence-induced ectopic expression of  
1017 the *A. tumefaciens ipt* gene in wheat delays leaf senescence, increases cytokinin content,  
1018 nitrate influx, and nitrate reductase activity, but does not affect grain yield. *Journal of*  
1019 *Experimental Botany* **59**: 377-387.
- 1020 **Small E, Brookes B. 1984.** Coiling of alfalfa pods in relation to resistance against seed chalcids:  
1021 additional observations. *Canadian Journal of Plant Science* **64**: 659-665

- 1022 **Sorefan K, Girin T, Liljegren SJ, Ljung K, Robles P, Galvan-Ampudia CS, Offringa R, Friml J,**  
 1023 **Yanofsky MF, Ostergaard L. 2009.** A regulated auxin minimum is required for seed  
 1024 dispersal in *Arabidopsis*. *Nature* **459**: 583-586.
- 1025 **Spence J, Vercher Y, Gates P, Harris N. 1996.** Pod shatter in *Arabidopsis thaliana*, *Brassica napus*  
 1026 and *B. juncea*. *Journal of Microscopy* **181**: 195-203.
- 1027 **Srinivasan A, Morgan DG. 1996.** Growth and development of the pod wall in spring rape (*Brassica*  
 1028 *napus*) as related to the presence of seeds and exogenous phytohormones. *The Journal of*  
 1029 *Agricultural Science* **127**: 487-500.
- 1030 **Stadler R, Lauterbach C, Sauer N. 2005.** Cell-to-cell movement of green fluorescent protein reveals  
 1031 post-phloem transport in the outer integument and identifies symplastic domains in  
 1032 *Arabidopsis* seeds and embryos. *Plant Physiology* **139**: 701-712.
- 1033 **Taylor L, Nunes-Nesi A, Parsley K, Leiss A, Leach G, Coates S, Wingler A, Fernie AR, Hibberd**  
 1034 **JM. 2010.** Cytosolic pyruvate, orthophosphate dikinase functions in nitrogen remobilization  
 1035 during leaf senescence and limits individual seed growth and nitrogen content. *The Plant*  
 1036 *Journal*: 641-652.
- 1037 **Van Dongen JT. 2003.** Structure of the Developing Pea Seed Coat and the Post-phloem Transport  
 1038 Pathway of Nutrients. *Annals of Botany* **91**: 729-737.
- 1039 **Vlot aC, Dempsey DMA, Klessig DF. 2009.** Salicylic Acid, a multifaceted hormone to combat  
 1040 disease. *Annual Review of Phytopathology* **47**: 177-206.
- 1041 **Wagstaff C, Yang TJ, Stead AD, Buchanan-Wollaston V, Roberts JA. 2009.** A molecular and  
 1042 structural characterization of senescing *Arabidopsis* siliques and comparison of transcriptional  
 1043 profiles with senescing petals and leaves. *Plant Journal* **57**: 690-705.
- 1044 **Wardlaw B. 1990.** The control of carbon partitioning in plants. *New Phytologist* **116**: 341-381.
- 1045 **Wilén RW, Mandel RM, Pharis RP, Holbrook LA, Moloney MM. 1990.** Effects of abscisic acid  
 1046 and high osmoticum on storage protein gene expression in microspore embryos of *Brassica*  
 1047 *napus*. *Plant Physiology* **94**: 875-881.
- 1048 **Willms J, Salon C, Layzell D. 1999.** Evidence for light-stimulated fatty acid synthesis in soybean  
 1049 fruit. *Plant Physiology* **120**: 1117-1128.

- 1050 **Wittenbach, Vernon A. 1983.** Effect of pod removal on leaf photosynthesis and soluble protein  
 1051 composition of field-grown soybeans. *Plant Physiology* **73**: 121-124.
- 1052 **Wuriyangan H, Zhang B, Cao W-H, Ma B, Lei G, Liu Y-F, Wei W, Wu H-J, Chen L-J, Chen**  
 1053 **H-W, Cao Y-R, He S-J, Zhang W-K, Wang X-J, Chen S-Y, Zhang J-S. 2009.** The  
 1054 ethylene receptor ETR2 delays floral transition and affects starch accumulation in rice. *The*  
 1055 *Plant Cell* **21**: 1473-1494.
- 1056 **Yanagisawa S, Yoo S-d, Sheen J. 2003.** Differential regulation of *EIN3* stability by glucose and  
 1057 ethylene signalling in plants. *Nature* **425**: 521-525.
- 1058 **Yang J, Zhang J, Liu K, Wang Z, Liu L. 2006.** Abscisic acid and ethylene interact in wheat grains  
 1059 in response to soil drying during grain filling. *New Phytologist* **171**: 293-303.
- 1060 **Yu B, Gruber M, Khachatourians GG, Hegedus DD, Hannoufa A. 2010.** Gene expression  
 1061 profiling of developing *Brassica napus* seed in relation to changes in major storage  
 1062 compounds. *Plant Science* **178**: 381-389
- 1063 **Zhao F, Bilsborrow PE, Evans EJ, Syers JK. 1993.** Sulphur turnover in the developing pods of  
 1064 single and double low varieties of oilseed rape (*Brassica napus* L). *Journal of the Science of*  
 1065 *Food and Agriculture* **62**: 111-119.
- 1066 **Zhou L, Jang JC, Jones TL, Sheen J. 1998.** Glucose and ethylene signal transduction crosstalk  
 1067 revealed by an Arabidopsis glucose-insensitive mutant. *Proceedings of the National Academy*  
 1068 *of Sciences of the United States of America* **95**: 10294-10299.
- 1069 **Zhou X, Liu Q, Xie F, Wen C-K. 2007.** RTE1 is a Golgi-associated and ETR1-dependent negative  
 1070 regulator of ethylene responses. *Plant Physiology* **145**: 75-86.
- 1071

**Table 1.** Expression of genes involved in ethylene and sugar transport/signalling during developmental pod senescence (green to yellow pods) and wound response (0-90min after wounding)<sup>1</sup>.

Function	Gene	AGI	Fold change in senescent pods	Fold change in wounded pods
Ethylene receptors	ETR1	At1g66340	0.94	1.50
	ETR2	At3g23150	2.83	NP
	ERS1	At2g40940	1.70	1.99
	ERS2	At1g04310	1.01	1.55

	EIN4	At3g04580	1.93	0.86
Ethylene biosynthesis	ACS2	At1g01480	NP	4.73
	ACS4	At2g22810	NP	NP
	ACS9	At3g49700	NP	NP
	ACS11	At4g08040	NP	NP
	ACO4	At1g05010	19.80	11.39
Sugar transport	GPT1	At5g54800	0.84	2.17
	GPT2	At1g61800	0.18	NP
	GLT1	At5g16150	1.12	1.11
	SUC2	At1g22710	0.92	8.31
	SUC3	At2g02860	1.24	1.52
	SUC4	At1g09960	0.77	3.54
	SUC5	At1g71890	0.10	1.33
	HXK1	At4g29130	0.84	2.45

<sup>1</sup>Shaded values indicate a fold increase greater than 1.5; NP indicates no signal present on array.

## Figure legends

Figure 1. Push-Pull model of resource allocation. Vegetative green organs such as leaves and stems produce assimilates which are pushed into the central pool of resources for that plant. Photosynthetic activity of the pod also contributes to the resource pool early in development, but it becomes a sink during senescence and seed maturation. A negative feedback loop, hypothesised to be mediated by an unknown signal originating from the immature pod, prevents early remobilisation of resources away from photosynthetic organs which is only broken as the pull from the maturing seeds becomes strong enough to initiate remobilisation from the central pool. The strength of the pull is proportional to the number of maturing pods on the plant; hence selective pod removal prevents senescence of the rosette leaves as the number of sinks is reduced.

Figure 2. Ways to manipulate yield. Blue boxes indicate targets strategies for yield manipulation; green boxes indicate the tools that could be used; pink boxes show the consequence of manipulation for each strategy. Coloured edges of blue boxes are linked with lines of the same colour. References in brackets are listed below and in full within the reference section of the main manuscript. (1) Allen & Morgan, 1972; (2) Taylor et al., 2010; (3) Abreu & Munné-Bosch, 2009; (4) Wuriyanghan et al., 2009; (5) Yang et al., 2006; (6) Wilen et al., 1990; (7) Maia-Grondard & Limami, 2010; (8) Gan & Amasino, 1995; (9) Sorefan et al., 2009.






Figure 1. Push-Pull model of resource allocation. Vegetative green organs such as leaves and stems produce assimilates which are pushed into the central pool of resources for that plant. Photosynthetic activity of the pod also contributes to the resource pool early in development, but it becomes a sink during senescence and seed maturation. A negative feedback loop, hypothesised to be mediated by an unknown signal originating from the immature pod, prevents early remobilisation of resources away from photosynthetic organs which is only broken as the pull from the maturing seeds becomes strong enough to initiate remobilisation from the central pool. The strength of the pull is proportional to the number of maturing pods on the plant; hence selective pod removal prevents senescence of the rosette leaves as the number of sinks is reduced.

34x15mm (600 x 600 DPI)

Review




Figure 2. Ways to manipulate yield. Blue boxes indicate targets strategies for yield manipulation; green boxes indicate the tools that could be used; pink boxes show the consequence of manipulation for each strategy. Coloured edges of blue boxes are linked with lines of the same colour. References in brackets are listed below and in full within the reference section of the main manuscript. (1) Allen & Morgan, 1972; (2) Taylor et al., 2010; (3) Abreu & Munné-Bosch, 2009; (4) Wuriyangan et al., 2009; (5) Yang et al., 2006; (6) Wilen et al., 1990; (7) Maia-Grondard & Limami, 2010; (8) Gan & Amasino, 1995; (9) Sorefan et al., 2009.

41x25mm (600 x 600 DPI)