

Recent progress in understanding and projecting regional and global mean sea-level change

Article

Accepted Version

Clark, P. U., Church, J. A., Gregory, J. M. ORCID:
<https://orcid.org/0000-0003-1296-8644> and Payne, A. J. (2015)
Recent progress in understanding and projecting regional and
global mean sea-level change. *Current Climate Change
Reports*, 1 (4). pp. 224-246. ISSN 2198-6061 doi:
<https://doi.org/10.1007/s40641-015-0024-4> Available at
<https://centaur.reading.ac.uk/44088/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s40641-015-0024-4>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Recent Progress in Understanding and Projecting Regional and Global Mean**
2 **Sea-Level Change**

3
4
5 Peter U. Clark¹, John A. Church², and Jonathan M. Gregory^{3,4}, and Anthony J. Payne⁵

6
7 ¹College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR
8 97330, USA

9
10 ²CSIRO Oceans and Atmosphere Flagship, Hobart, Tasmania 7000, Australia

11
12 ³NCAS-Climate, University of Reading, Reading RG6 6BB, UK

13
14 ⁴Met Office Hadley Centre, Exeter EX1 3PB, UK

15
16 ⁵School of Geographical Sciences, University of Bristol, BS8 1SS, UK

17 Abstract

18 Considerable progress has been made in understanding present and future regional and
19 global sea level in the two years since publication of the Fifth Assessment Report (AR5) of the
20 Intergovernmental Panel on Climate Change. Here we evaluate how the new results affect the
21 AR5's assessment of (i) historical sea-level rise, including attribution of that rise and
22 implications for the sea-level budget, (ii) projections of the components and of total global mean
23 sea level (GMSL), and (iii) projections of regional variability and emergence of the
24 anthropogenic signal. In each of these cases, new work largely provides additional evidence in
25 support of the AR5 assessment, providing greater confidence in those findings. Recent analyses
26 confirm the 20th century sea-level rise, with some analyses showing a slightly smaller rate before
27 1990 and some a slightly larger value than reported in the AR5. There is now more evidence of
28 an acceleration in the rate of rise. Ongoing ocean heat uptake and associated thermal expansion
29 have continued since 2000, and are consistent with ocean thermal expansion reported in the AR5.
30 A significant amount of heat is being stored deeper in the water column, with a larger rate of heat
31 uptake since 2000 compared to the previous decades and with the largest storage in the Southern
32 Ocean. The first formal detection studies for ocean thermal expansion and glacier mass loss since
33 the AR5 have confirmed the AR5 finding of a significant anthropogenic contribution to sea-level
34 rise over the last 50 years. New projections of glacier loss from two regions suggest smaller
35 contributions to GMSL rise from these regions than in studies assessed by the AR5; additional
36 regional studies are required to further assess whether there are broader implications of these
37 results. Mass loss from the Greenland Ice Sheet, primarily as a result of increased surface
38 melting, and from the Antarctic Ice Sheet, primarily as a result of increased ice discharge, has
39 accelerated. The largest estimates of acceleration in mass loss from the two ice sheets for 2003-

40 2013 equal or exceed the acceleration of GMSL rise calculated from the satellite altimeter sea-
41 level record over the longer period of 1993-2014. However, when increased mass
42 gain in land water storage and parts of East Antarctica, and decreased mass
43 loss from glaciers in Alaska and some other regions, are taken into account,
44 the net acceleration in the ocean mass gain is consistent with the satellite
45 altimeter record. New studies suggest that a marine ice-sheet instability (MISI) may have been
46 initiated in parts of the West Antarctic Ice Sheet (WAIS), but that it will affect only a limited
47 number of ice streams in the 21st century. New projections of mass loss from the Greenland and
48 Antarctic Ice Sheets by 2100, including a contribution from parts of WAIS undergoing unstable
49 retreat, suggest a contribution that falls largely within the *likely* range (i.e., two-thirds
50 probability) of the AR5. These new results increase confidence in the AR5 *likely* range,
51 indicating that there is a greater probability that sea-level rise by 2100 will lie in this range with a
52 corresponding decrease in the likelihood of an additional contribution of several tens of
53 centimeters above the *likely* range. In view of the comparatively limited state of knowledge and
54 understanding of rapid ice-sheet dynamics, we continue to think that it is not yet possible to
55 make reliable quantitative estimates of future GMSL rise outside the *likely* range. Projections of
56 21st-century GMSL rise published since the AR5 depend on results from expert elicitation, but
57 we have low confidence in conclusions based on these approaches. New work on regional
58 projections and emergence of the anthropogenic signal suggests that the two commonly predicted
59 features of future regional sea-level change (the increasing tilt across the Antarctic Circumpolar
60 Current and the dipole in the North Atlantic) are related to regional changes in wind stress and
61 surface heat flux. Moreover, it is expected that sea-level change in response to anthropogenic
62 forcing, particularly in regions of relatively low unforced variability such as the low-latitude

63 Atlantic, will be detectable over most of the ocean by 2040. The east-west contrast of sea-level
64 trends in the Pacific observed since the early 1990s cannot be satisfactorily accounted for by
65 climate models, nor yet definitively attributed either to unforced variability or forced climate
66 change.

67 **1. Introduction**

68 Understanding and projecting regional and global mean sea-level change is of critical
69 importance to assessing socio-economic impacts and for planning for adaptation in the highly
70 populated low-lying coastal zones of the world. Today, about 10 per cent of the world's
71 population (>600 million people) and about 65% of the world's cities with populations of greater
72 than 5 million are located at elevations less than 10 m above sea level (McGranahan et al., 2007),
73 and ~150 million people live within one meter of the high-tide level (Lichter et al., 2011). Over
74 the last four decades, rapid migration towards the coast and development has significantly
75 increased exposure of populations and assets to extreme sea-level events, with about 270 million
76 people and US\$13 trillion worth of assets being exposed to the such events in 2010 (Jongman et
77 al., 2012). Continued population growth, economic development, and urbanization, combined
78 with additional sea-level rise and associated increase in frequency of extreme sea-level events,
79 will further increase the risk and impacts in coastal zones (Wong et al., 2014).

80 Planning for and adapting to sea-level change requires an assessment of the expected
81 magnitude of change and its uncertainties. Working Group I (WGI) of the Fifth Assessment
82 Report (AR5) of the Intergovernmental Panel on Climate Change (IPCC) assessed sea-level
83 change from several perspectives: (1) past sea-level change with an emphasis on previous warm
84 intervals, the past millennium, and over the instrumental period (since ~1700), (2) contributions
85 to global mean sea-level (GMSL) rise during the instrumental period, (3) projections of GMSL
86 change during the 21st century and over the longer term (to 2500), (4) projections of 21st century
87 regional sea-level change, and (5) projections of 21st century sea-level extremes and waves
88 (Church et al., 2013a).

89 The calibrated uncertainty language of the IPCC uses terms to indicate the assessed
90 likelihood of an outcome or a result. When referring to the IPCC assessment of likelihood of an
91 outcome, we follow their convention in italicizing the terms. For example, *likely* means that there
92 is probability of two-thirds or more that the outcome may lie within the *likely* range whereas *very*
93 *likely* means a probability of at least 90% within the range. Uncertainty is quantified using 90%
94 uncertainty intervals unless otherwise stated. The 90% uncertainty interval, reported in square
95 brackets (e.g., 0.21 [0.16 to 0.26] m), is expected to have a 90% likelihood of covering the value
96 that is being estimated; it is only specific to the *very likely* range when identified as such.

97 The AR5 concluded that 21st-century global mean surface air temperature (SAT) change
98 is *likely* (i.e., two-thirds probability, NB not *very likely*) to lie within the 5-95% range (i.e. from
99 the 5th to the 95th percentile) of the CMIP5 model projections; this is a measure of the spread
100 resulting from different choices of structure and parameters in the models (Collins et al., 2013).
101 Accordingly, the AR5 interpreted the 5-95% range of model results as the *likely* range for each
102 of the GMSL rise contributions that is projected on the basis of CMIP5 results (thermohaline,
103 glaciers, ice-sheet surface mass balance) (Church et al., 2013a), and for consistency, the model
104 5-95% range was also interpreted as *likely* for projected contributions from rapid ice-sheet
105 dynamics and land water storage. As an example, reporting a *likely* range of projected sea-level
106 rise of 0.2 to 0.8 m means that it is *likely* (i.e., at least 66% probability) that sea level will lie
107 within the 0.2 to 0.8 m uncertainty range, and a probability of 33% or less that it will lie outside
108 that range (not necessarily symmetrically distributed). The AR5 was not able to assess a *very*
109 *likely* range for projected GMSL rise because there was (and still is) no assessment available of
110 the *very likely* range for global mean SAT change, and because the probability of ice-sheet
111 dynamical changes that would give rise to greater values could not be robustly quantified.

112 The AR5 assessment of sea-level change made three important advances beyond the
113 Fourth Assessment Report (AR4) (Meehl et al., 2007). Firstly, the AR5 demonstrated that, when
114 an allowance for potential ice-sheet contributions is included, the observed GMSL rise is
115 consistent with the sum of the estimated contributions since 1900, and that models and
116 observations are consistent regarding the contributions from thermal expansion and glaciers over
117 the last 50 years. The budget was closed for 1993–2010 (corresponding to the period of
118 continuous satellite observations of sea level and ice sheets), and for 1971–2010 (with reasonable
119 estimates of ice-sheet contributions). These findings imply improved physical understanding of
120 the causes of past GMSL change, and greater confidence in the reliability of models for making
121 projections.

122 Secondly, the AR5 included future rapid changes in ice-sheet dynamics in its sea-level
123 projections. This could not be done in the AR4 because there were no existing models and
124 insufficient scientific understanding of the accelerations in ice-sheet outflow that had only
125 recently been observed (Meehl et al., 2007). Accordingly, the AR4 did not provide a best
126 estimate or *likely* range of 21st-century sea-level change. By the time of the AR5, understanding
127 and modelling of these contributions from ice-sheet dynamics had developed sufficiently to
128 allow an assessment of a *likely* increase of global mean sea level for each of the four
129 Representative Concentration Pathways (RCPs) of future atmospheric composition
130 (Meinshausen et al., 2011; Moss et al., 2010) used by WG1 for climate projections (for example,
131 0.52 to 0.98 m by 2100 for RCP8.5) (Church et al., 2013a). The AR5 could not exclude the
132 possibility of higher sea levels, but concluded that only the collapse of the marine-based sections
133 of the Antarctic ice sheet, if initiated, could cause GMSL to rise substantially higher (estimated
134 at several tenths of a meter) than the *likely* range in the 21st century (Church et al., 2013a).

135 However, significant uncertainties, particularly related to the dynamical Antarctic ice-sheet
136 contribution, remain.

137 Thirdly, the inclusion of the effect of rapid ice-sheet dynamical change meant that the
138 AR5, unlike the AR4, was able to make projections of the regional distribution of sea-level
139 change. This led to the conclusion that, by the end of the 21st century, it is *very likely* that
140 regional sea-level rise will be positive over about 95% of the world ocean, and that about 70% of
141 the global coastlines are projected to experience a relative sea-level change within 20% of the
142 GMSL change (Church et al., 2013a).

143 The AR5 projections of 21st century sea-level change have been criticized for taking a
144 “moderate line” (Kerr, 2013), for being “conservative” (Rahmstorf, 2013), and for being
145 “misleading” (Mooney, 2014). As explained by Church et al. (2013c), such criticisms are based
146 in part on a misunderstanding of how the results are reported, particularly with regard to the
147 IPCC calibrated uncertainty language and on whether the results were given for 2081-2100 or for
148 2100. For example, under the highest scenario considered, the AR5 reported a *likely* range (with
149 two-thirds probability) of 0.45 to 0.82 m for 2081-2100 and 0.52 to 0.98 m for 2100, but Kerr
150 (2013) reported the latter as “a worst case of 1 meter.”

151 In the two and a half years since March 2013, the cutoff date for literature assessed by the
152 AR5, there has been considerable progress in understanding several of the key issues on sea-level
153 change discussed by the AR5. In this review we summarize the main findings by the AR5 and
154 discuss how literature published since the cutoff date compares to the AR5 assessment, including
155 whether any modification of that assessment is required, and with particular attention to
156 projections beyond the *likely* range.

157 **2. Historical sea level**

158 *2a. The tide-gauge record (~1770-2015)*

159 The AR5 (Church et al., 2013a; Rhein et al., 2013) concluded that the trend in GMSL for
160 the 1900 to 2010 was $1.7 \pm 0.2 \text{ mm yr}^{-1}$ (1.5 mm yr^{-1} from 1901 to 1990), and accelerated during
161 the 20th century in the presence of multi-decadal variability, with estimates that ranged from
162 $0.000 [-0.002 \text{ to } 0.002] \text{ mm yr}^{-2}$ to $0.013 [0.007 \text{ to } 0.019] \text{ mm yr}^{-2}$. New and longer tide-gauge
163 records have become available since the AR5. Updates of two of the GMSL reconstructions used
164 in the AR5 give $1.77 \pm 0.28 \text{ mm yr}^{-1}$ (Wenzel and Schroter, 2014) and $1.9 \pm 0.3 \text{ mm yr}^{-1}$ since 1900
165 (and $3.1 \pm 0.6 \text{ mm yr}^{-1}$ for 1993 to 2009) (Jevrejeva et al., 2014b), in the upper half of the AR5
166 range. Hay et al. (2015) considered the “fingerprints” of mass loss from glaciers and ice sheets in
167 their reconstruction of GMSL. These effects on regional sea level are due to the change in the
168 geopotential field (jointly determined by gravitation and Earth rotation) and the elastic response
169 of the lithosphere, both practically instantaneous and caused by the geographical redistribution of
170 mass on the Earth’s surface when mass is transferred from land into the ocean. As shorthand, we
171 later (section 8) refer to these effects as “GeLi.” By this method, they estimated a trend of $1.2 \pm$
172 0.2 mm yr^{-1} for 1901 to 1990, which overlaps the AR5 range of $1.5 \pm 0.2 \text{ mm yr}^{-1}$ for the
173 corresponding period, but is lower primarily because their estimate has very little sea-level
174 change during 1950-1970. Including the GeLi fingerprints is an advance over assuming
175 geographically uniform sea-level contributions from land ice (as in Church and White (2011)),
176 but estimating the magnitude of the weight factors for the many fingerprints used by Hay et al. in
177 the presence of large regional decadal variability is challenging given the time-varying and
178 incomplete distribution of observations. Hamlington and Thompson (2015) drew attention to the
179 tide-gauges from the Arctic, Alaska and Japan that are included in Hay et al. (2015) but not the
180 other reconstructions. The Arctic gauges have negative trends for 1950-1970, and high-latitude

181 gauges in general have the most uncertain glacial isostatic adjustment (Jevrejeva et al., 2014b).
182 However, Hay (personal communication) reports that their results are essentially unchanged if
183 they use the same tide-gauge data set as Church and White (2011).

184 Becker et al. (2014), Bos et al. (2014), Beenstock et al. (2014), and Dangendorf et al.
185 (2015) suggested that long-term variability in local and global mean sea level has resulted in
186 previous studies underestimating uncertainties. Using refined uncertainty estimates, Becker et al.
187 and Dangendorf et al. estimated a larger uncertainty, due to unforced and naturally forced
188 variability, and thus argued the *minimum* long-term anthropogenic GMSL trend since 1900 is 1
189 mm yr⁻¹ and 0.6 mm yr⁻¹, respectively (both at the 99% confidence level). Since the variability
190 can be of either sign, these studies also imply the possibility that the anthropogenic GMSL trend
191 could be larger than observed, and partially offset by variability. Beenstock et al. (2015) argued
192 that tide-gauges with trends that were not significantly different to zero should be excluded from
193 the global mean, that the longer tide-gauge time series are from locations with larger rates of sea-
194 level rise, and therefore that GMSL estimates reported in the AR5 are biased high. We disagree
195 with the first argument, which could be made to disprove the existence of any global mean trend
196 that is partly obscured by local variability, whereas averaging over many locations reduces that
197 variability. The second argument is doubtful because it assumes constant regional patterns of
198 forced sea-level change through the century.

199 A number of different techniques have been used to estimate acceleration of *local* relative
200 sea level (for example, see Visser et al. (2015)), but the results are controversial (see for example
201 Kenigson and Han (2014); Piecuch and Ponte (2015)). Haigh et al. (2014) demonstrated the
202 difficulty of estimating accelerations from local tide-gauge observations, the necessity for long
203 time series (Douglas (1992) argued that almost 50 years was required), and the importance of

204 removing unforced and naturally forced variability.

205 There are now more estimates of acceleration of GMSL over the 19th to 20th century and
206 these are generally larger than those available at the time of the AR5, ranging from $0.0042 \pm$
207 $0.0092 \text{ mm yr}^{-2}$ (Wenzel and Schroter, 2014) to $0.02 \pm 0.01 \text{ mm yr}^{-2}$ (Jevrejeva et al., 2014b)
208 (see Cahill et al. (2015); Hay et al. (2015); Hogarth (2014); Olivieri and Spada (2013); Spada et
209 al. (2015) for intermediate estimates). Jorda (2014) estimated that at least 2.2 mm yr^{-1} of the
210 recent sea-level trend estimated from altimetry cannot be attributed to unforced multidecadal
211 variability, implying that the change in trend between 1900-1990 and the altimeter period is at
212 least partly forced.

213 In summary, recent analyses confirm the 20th century sea-level rise, with some analyses
214 showing a slightly smaller rate before 1990 and some a slightly larger value than reported in the
215 AR5. There is now more evidence of an acceleration in the rate of rise.

216 *2b. The satellite altimeter period*

217 According to the AR5, the rate of sea-level rise measured by the
218 TOPEX/POSEIDON/Jason1/2 satellite altimeter missions over 1993 to 2012 was $3.2 \pm 0.4 \text{ mm}$
219 yr^{-1} , with interannual variability that was related to climate variability, particularly the El Niño-
220 Southern Oscillation phenomenon. Cazenave et al. (2014) found a deceleration from the first to
221 the second decade of the altimeter record, but demonstrated that there was no significant
222 reduction of the underlying rate of sea-level rise over this two-decade period if the effects of
223 interannual climate variation on the storage of water on land (particularly in Australia; Fasullo et
224 al. (2013)) and thermal expansion are excluded. Confirming the importance of interannual
225 variability, Yi et al. (2015) have shown that since the La Niña event of 2010, the rate of sea-level
226 rise has been substantially larger than the 1993-2015 average as a result of a decrease of water

227 stored on land, increased ocean thermal expansion, and faster loss of mass from the ice sheets
228 (particularly Greenland).

229 As a result of a major and ongoing effort to improve the quality of the altimeter records
230 (Ablain et al., 2015), the ERS-1/ERS-2/Envisat GMSL trend over 1993 to 2010 was revised
231 upward from $1.59 \pm 0.5 \text{ mm yr}^{-1}$ to $2.36 \pm 0.5 \text{ mm yr}^{-1}$, compared with their TOPEX/Jason
232 estimate of $2.98 \pm 0.4 \text{ mm yr}^{-1}$ over the same period. Watson et al. (2015) examined the quality
233 of the TOPEX/Jason1/2 altimeter record by a careful comparison with sea level measured by
234 coastal and island tide gauges. They found small but significant trends in the sea-level biases,
235 mostly in the first six years (the TOPEX satellite) of the record. After correction for these biases,
236 they estimated a corrected GMSL trend over 1993 to mid-2014 of $2.6 \text{ to } 2.9 \pm 0.4 \text{ mm yr}^{-1}$
237 (dependent on the vertical land motion correction adopted), and found an acceleration of $0.041 \pm$
238 0.058 mm yr^{-2} , in contrast to the deceleration reported by Cazenave et al. (2014). (The
239 acceleration is not significantly different from zero, but it is significantly different from the
240 deceleration over the same period of $-0.057 \pm 0.058 \text{ mm yr}^{-2}$ if these bias drifts were not
241 corrected.) Reprocessing of the TOPEX record is currently underway and should shed light on
242 the validity or otherwise of these corrections and revised GMSL estimates.

243 Analysis of 13 years (2002-2015) of data from GRACE, Global Navigation Satellite
244 System, satellite laser ranging and the Ocean Circulation and Climate of the ocean bottom
245 pressure (Wu and Heflin, 2015) give an acceleration of global mass (non-steric) component of
246 $0.04 \pm 0.09 \text{ mm yr}^{-2}$, similar to the above satellite altimeter estimate (for latitudes less than 65°)
247 but with larger uncertainty estimates. This ocean mass is the balance between accelerated mass
248 loss from the Greenland and West Antarctic Ice Sheets and increased mass gain in land water
249 storage and parts of East Antarctica, and decreased mass loss from glaciers in Alaska and some

250 other regions.

251 The recent analyses confirm that the rate of sea-level rise since 1993 is larger than prior
252 to 1990, with suggestions of a slightly smaller rate than reported in the AR5 and with a small
253 (but not statistically significant) acceleration.

254 **3. Sea-level contributions**

255 *3a. Steric sea-level change*

256 The AR5 estimated rates of thermal expansion of 0.8 [0.5 to 1.1] mm yr⁻¹ for 1971 to
257 2010 and 1.1 [0.8 to 1.4] mm yr⁻¹ for 1993 to 2010. Some of the studies discussed in the AR5
258 (e.g. Lyman et al. (2010)), but not all (e.g. (Church et al., 2011b; Church et al., 2013d)), reported
259 a sharp spike in ocean heat uptake in the early 2000s followed by a slowing of the rate of heat
260 uptake (and thus ocean thermal expansion). Since then, Abraham et al. (2013) published a major
261 review on the evolving observing system, the reduction of XBT biases, and estimates of heat
262 content and thermosteric sea-level trends over different periods, generally confirming the AR5
263 assessment of trends since 1971 and 1993. These estimates are dependent on a range of
264 uncertainties, including ocean climatologies used, vertical resolution (Cheng and Zhu, 2014a;
265 Cheng and Zhu, 2014b) and mapping techniques (Chang et al., 2014).

266 The apparent surface warming “hiatus” has prompted many studies, including a focus on
267 ocean heat uptake, but with less attention specifically addressing the related thermosteric sea-
268 level change. Balmaseda et al. (2013) and Chen and Tung (2014) demonstrated an ongoing ocean
269 heat uptake during the hiatus, but with a greater accumulation deeper in the water column,
270 between the depth of most previous upper ocean estimates (300 m and 700 m) and 2000 m.
271 Balmaseda et al. (2013) demonstrated a clearer response to volcanic eruptions than earlier studies
272 (e.g. Domingues et al. (2008)) and a greater rate of heat uptake after 2000 than in the 1990s.

273 Chen and Tung (2014) argued that most of the heat uptake occurred in the North Atlantic and in
274 the Southern Ocean whereas Nieves et al. (2015) argued for the importance of heat uptake in the
275 100 m to 300 m layer of the Indian and Pacific Oceans.

276 Ocean heat content changes are directly related to top of the atmosphere net radiation
277 fluxes (Palmer and McNeall, 2014). Allan et al. (2014) used atmospheric reanalysis and 20th
278 century simulations to extend the CERES satellite radiation observations of the Earth's global
279 energy balance (anchored to the estimates of ocean energy uptake; Loeb et al. (2012)) back to the
280 1980s. Smith et al. (2015) extended these series back to 1960 and demonstrated an increasing
281 ocean heat uptake, the impact of volcanic eruptions, and that the spike in ocean heat uptake and
282 subsequent decrease in the rate of ocean warming after 2000 in some studies (see above) was
283 probably an artifact of errors in XBT bias corrections and/or incomplete ocean coverage.
284 Wunsch and Heimbach (2014) used a sophisticated data assimilation technique to infer a deep
285 ocean cooling, in direct contrast to the direct observations of Purkey and Johnson (2010). The
286 reason for this difference in the Wunsch and Heimbach results is unclear but could relate to the
287 omission of the geothermal heat flux from the ocean seafloor (approximately equivalent to the
288 difference between their results to the direct observations) or the dominance of upper ocean
289 observations over the sparse deep observations in their analysis.

290 Extended altimeter observations of sea-surface height, GRACE observations of ocean
291 mass, and Argo observations of upper ocean thermal expansion have shown an approximate
292 closure of the sea-level budget (see Leuliette (2015) for a review). Von Shuckmann et al. (2014)
293 and Dieng et al. (2015) argued that ocean heat-uptake estimates were biased low because of
294 inadequate coverage and mapping, particularly in the region of the tropical Asian archipelago.

295 Uncertainties in the sea-level budget are too large for the deep-ocean contribution to be inferred
296 (von Schuckmann et al. 2014; Llovel et al. 2014; Dieng et al. 2015).

297 Durack et al. (2014) found that, in the CMIP3 and CMIP5 climate model simulations,
298 60% of the heat uptake occurred in the Southern Hemisphere. In contrast, the observational
299 estimates ranged from about 35% (Levitus et al., 2012) to about 50% (Domingues et al. 2008),
300 suggesting that historical ocean heat content estimates may be biased low by various amounts
301 because of lack of data in the Southern Hemisphere. Based on Argo data from 2006 to 2014,
302 Roemmich et al. (2015) found an even larger ratio of Southern to Northern Hemisphere ocean
303 heat uptake (67 to 98%) possibly as a result of greater negative aerosol forcing in the Northern
304 Hemisphere and/or ocean heat-uptake processes. This larger ratio raises concern about the use of
305 the model results for adjusting the observational estimates, as suggested by Durack et al. (2014).
306 One of the mapping techniques used in Roemmich et al. (2015; reduced space optimal
307 interpolation) also directly addressed the mapping deficiencies in the tropical Asian archipelago
308 identified in von Schuckmann et al. (2014) and indicates that over the 2006 to 2014 period, the
309 thermal expansion in the upper 2,000 m was $0.8 \pm 0.5 \text{ mm yr}^{-1}$ (Didier Monselesan, personal
310 communication), and $0.9 \pm 0.5 \text{ mm yr}^{-1}$ for the full depth ocean (using the Purkey and Johnson
311 2010 abyssal ocean estimates), close to the AR5 estimate for 1993-2010.

312 Halosteric trends over the historical period from 1950 to 2010 are important regionally in
313 reinforcing thermosteric trends in the Pacific and counteracting them in the Atlantic (generally of
314 order 25% of thermosteric trends) (Durack et al., 2015) but are not important for the global ocean
315 steric change. Purkey et al. (2014) demonstrated agreement between regional ocean-mass trends
316 determined from GRACE data and the difference between altimeter sea-level observations and
317 ocean steric sea-level change.

318 In summary, recent analyses clearly indicate an ongoing ocean heat uptake and associated
319 thermal expansion since 2000, with a significant amount of heat being stored deeper in the water
320 column and with the largest storage in the Southern Ocean. Quantitatively, the new results are
321 consistent with ocean thermal expansion reported in the AR5, and with a larger rate of heat
322 uptake since 2000 compared to the previous decades.

323 *3b. Glacier mass loss*

324 Glacier mass loss was a major contributor to sea-level rise during the 20th century. The
325 AR5 estimated sea-level contributions from glaciers of 0.69 [0.61 to 0.77] mm yr⁻¹ for 1901-
326 1990, 0.68 [0.31 to 1.05] mm yr⁻¹ for 1971-2010 and 0.86 [0.49 to 1.23] mm yr⁻¹ for 1993-2010.
327 The estimates diverge most before 1950 and after 2003. Extensions and corrections to the glacier
328 inventory, combined with a significantly increased number of geodetic mass-balance and glacier-
329 length measurements, have led to convergence of the various estimates such that they now agree
330 within uncertainties over all the periods considered (Marzeion et al., 2015) (Marzeion et al.,
331 2015). There have been only small changes in the average of these estimates, and values are
332 consistent with those reported in the AR5, increasing confidence in those estimates.

333 *3c. Greenland Ice Sheet*

334 The AR5 found that the Greenland ice sheet has lost mass over the last two decades and
335 that the rate of loss has increased (Vaughan et al., 2013). Mass loss was about equally partitioned
336 between changes in surface mass balance (SMB, snow accumulation minus runoff) and increased
337 discharge into the ocean as icebergs, with the former dominating in southwest, central-north, and
338 northeast sectors and the latter in southeast and central-west sectors. While on average SMB has
339 become progressively less positive over the last two decades, there have been considerable
340 spatial and temporal variations in rates of discharge. Observations indicated that the contribution

341 to GMSL had *very likely* increased from 0.09 [−0.02 to 0.20] mm yr^{−1} for 1992–2001 to 0.59
342 [0.43 to 0.76] mm yr^{−1} for 2002–2011, with a total of $\sim 8.0 \pm 1.4$ mm from 1992 to 2012,
343 implying an acceleration of ~ 0.05 [0.02 to 0.08] mm yr^{−2}. These numbers include the
344 contribution from glaciers peripheral to the ice sheet, because some methods used to measure
345 mass loss are unable to separate them from the ice sheet.

346 Several studies published since the AR5 using various remote-sensing techniques support
347 the AR5 conclusions. Based on laser altimetry measurements from IceSat, Csatho et al. (2014)
348 estimated a contribution from the Greenland ice sheet to GMSL of 0.68 mm yr^{−1} for 2003–2009.
349 Based on gravity measurements from GRACE, Velicogna et al. (2014) found mass loss across
350 much of the ice sheet for the 2003–2013 period, and reported a contribution to GMSL rise of 0.77
351 ± 0.16 mm yr^{−1}, with an acceleration of 0.069 ± 0.003 mm yr^{−2}. The largest losses occurred in the
352 southeastern and northwestern sectors of the ice sheet (70% combined), while the southwestern
353 sector experienced the greatest acceleration in mass loss. SMB accounted for $\sim 67\%$ of the total
354 mass loss during this period. Using Operation IceBridge ice-thickness measurements to better
355 constrain discharge rates for 178 marine-terminating outlet glaciers, Enderlin et al. (2014) found
356 that 15 glaciers accounted for 77% of the total mass loss by discharge since 2000 and only four
357 glaciers accounted for 50% of the total. The contributions of SMB and discharge were nearly
358 equal until ~ 2006 , but the contribution of discharge has since decreased, with a corresponding
359 increase in mass loss due to SMB changes, which has reached 84% of the total since 2009.
360 Enderlin et al. (2014) estimated that the contribution to GMSL increased from 0.43 ± 0.09 mm
361 yr^{−1} over 2000–2005 to 0.73 ± 0.05 mm yr^{−1} over 2005–2009 and 1.05 ± 0.14 mm yr^{−1} over
362 2009–2012, with an acceleration of 0.08 ± 0.02 mm yr^{−2}, giving a total contribution of $8.23 \pm$
363 0.93 mm over 2000–2012. Wouters et al. (2013) and Wu and Helfin (2015) cautioned that these

364 calculated accelerations are for short periods and are affected by longer term climate variability.
365 Based on radar altimetry measurements from CryoSat-2 for the three-year period starting in
366 January 2011, Helm et al. (2014) confirmed the recent high contribution to GMSL (1.04 ± 0.07
367 mm yr^{-1}), with greatest elevation changes on the western, southeastern, and northeastern
368 margins. Based on data from several measurement platforms, Khan et al. (2014) also
369 documented recent pronounced thinning of the northeast Greenland ice stream. Van Angelen et
370 al. (2014) emphasized the importance of the observed persistent negative anomalies in the
371 summertime North Atlantic oscillation (NAO) index for the recent changes in the observed
372 SMB, which exceeded the CMIP5 simulations, supporting the AR5 conclusion that internally
373 generated regional climate variability has been the dominant cause of recent negative SMB
374 (Church et al., 2013a). The CMIP5 models do not consistently project negative anomalies in the
375 summertime in the NAO index and a return of the NAO to more positive values may lead to a
376 partial recovery in the SMB changes.

377 In summary, recent publications indicate an acceleration of mass loss from the Greenland
378 ice sheet, primarily as a result of increased surface melting. The reported contributions and the
379 computed accelerations are at the upper end of the *likely* range assessed in the AR5.

380 *3d. Antarctic Ice Sheet*

381 In the near-absence of surface melting and runoff, Antarctica's mass budget is dominated
382 by snow accumulation and ice discharge across the grounding line into floating ice shelves. In
383 the AR5 assessment, the Antarctic ice sheet (including the peripheral glaciers) was losing mass
384 and *likely* contributed $0.27 [0.16 \text{ to } 0.37] \text{ mm yr}^{-1}$ to GMSL over 1993–2010, and $0.41 [0.20 \text{ to}$
385 $0.61] \text{ mm yr}^{-1}$ over 2005–2010, suggesting an acceleration of $\sim 0.01 [-0.02 \text{ to } 0.05] \text{ mm yr}^{-2}$
386 (Vaughan et al., 2013). The acceleration has been caused by an increase in discharge in the

387 Antarctic Peninsula and the Amundsen Sea sector of West Antarctica, and was somewhat offset
388 by a mass gain over East Antarctica due to increased snowfall.

389 Papers published since 2013 largely corroborate the AR5 assessment. Based on GRACE
390 measurements for the period 2003-2013, Velicogna et al. (2014) estimated that the ice sheet has
391 contributed to GMSL rise at a rate of $0.18 \pm 0.12 \text{ mm yr}^{-1}$ with an acceleration of 0.03 ± 0.01
392 mm yr^{-2} ; greatest rates of mass loss are from the Amundsen Sea area (equivalent to 0.32 ± 0.02
393 mm yr^{-1}), which accounts for 94% of the mass loss from West Antarctica, and the Antarctic
394 Peninsula, with loss from both areas being dominated by dynamics. Dronning Maud Land in East
395 Antarctica experienced a mass gain accounting for a fall in GMSL of $0.17 \pm 0.02 \text{ mm yr}^{-1}$ since
396 2008 (Lenaerts et al., 2013). Harig and Simons (2015) used techniques to increase the spatial
397 resolution of the GRACE data, thus better resolving regional variations, and arrived at similar
398 estimates. As noted above, Wouters et al. (2013) and Wu and Helfin (2015) cautioned that these
399 calculated accelerations are for short periods and are affected by longer term climate variability.

400 Measurements of recent elevation changes over the Antarctic ice sheet from CryoSat-2
401 show a contribution to GMSL rise of $0.45 \pm 0.14 \text{ mm yr}^{-1}$ since 2010, 75% of which is derived
402 from the Amundsen Sea sector of West Antarctica (McMillan et al., 2014), and a contribution of
403 $0.35 \pm 0.23 \text{ mm yr}^{-1}$ since 2011 (Helm et al., 2014). Sutterly et al. (2014) compared four
404 independent estimates of the mass balance of the Amundsen Sea area, identified by all studies as
405 the primary region of Antarctica currently experiencing mass loss. The four methods agree in
406 terms of mass loss and acceleration in loss at the regional scale. The contribution to GMSL rise
407 was $0.23 \pm 0.01 \text{ mm yr}^{-1}$ over 1992–2013 and $0.28 \pm 0.03 \text{ mm yr}^{-1}$ over 2003–2011.

408 Rignot et al. (2014) showed that the grounding lines of glaciers draining the Amundsen
409 Sea sector of West Antarctica retreated by up to 35 km from 1992 to 2011. These rapid retreats

410 have proceeded along regions where the bed slopes downwards away from the coast, the
411 configuration which is potentially subject to the marine ice-sheet instability. Upstream of the
412 2011 grounding line positions, there are no major bed obstacles that would prevent irreversible
413 retreat in this sector of the West Antarctic ice sheet (Rignot et al., 2014). Ice-sheet model
414 experiments suggest that current retreat of PIG (Favier et al., 2014) and Thwaites Glacier (TG)
415 (Joughin et al., 2014) may be irreversible.

416 The AR5 concluded that the changes in the Amundsen Sea region are due to high ocean
417 heat flux causing thinning of ice shelves and grounded ice and grounding line retreat. Schmidtke
418 et al. (2014) documented warming at the bottom of the Amundsen and Bellingshausen seas that is
419 linked to increased heat content and to a shoaling of the mid-depth temperature maximum over
420 the continental slope, allowing warmer, saltier Circumpolar Deep Water (CDW) access to the
421 shelf in recent years. Paolo et al. (2015) found that thinning of Antarctic ice shelves has
422 increased since 1994, being most intense in the Amundsen and Bellingshausen regions. The
423 average thickness of ice shelves has decreased by 5-8% in less than two decades, with some
424 shelves thinning at rates that imply complete loss in less than 100 years. Greatest thinning has
425 occurred near the deep grounding lines, consistent with ocean forcing from increased flux of
426 CDW across the continental shelf.

427 In summary, recent publications support the AR5 assessment that mass loss from the
428 Antarctic ice sheet is accelerating, with most of that loss coming from the Amundsen Sea sector
429 of the WAIS. Observations and model simulations since the AR5 suggest that this acceleration
430 may be associated with a marine ice-sheet instability that may have been initiated in parts of the
431 WAIS, and that this may have been triggered by increased flux of CDW across the continental
432 shelf. There are also indications of increased accumulation in East Antarctica.

433 When the Antarctic and Greenland ice sheets are taken together, the largest estimates of
434 the acceleration in their contribution to sea level (range from 0.112 to 0.140 mm yr⁻²) equal or
435 exceed the acceleration in GMSLR calculated from the satellite altimeter sea-level record (0.099
436 mm yr⁻²) (Watson et al., 2015), although we note that the ice-sheet estimates (2003-2013) are
437 only for the latter half of the satellite altimeter period (1993 to mid-2014), during which time the
438 sea-level acceleration may be larger. Wu and Heflin (2015) also reported increased mass gain in
439 land water storage and parts of East Antarctica, and decreased mass loss from glaciers in Alaska
440 and some other regions such that the acceleration in the ocean mass gain is consistent with the
441 satellite altimeter record.

442 **4. The sea-level budget**

443 Sea-level estimates from satellite altimetry, GRACE observations of ocean mass, and
444 Argo observations of thermal expansion of the upper 2000 m of the ocean combine to close the
445 sea-level budget since 2005 within uncertainties (see recent updates by Leuliette (2015) and Yi et
446 al. (2015)). There have not yet been any new comprehensive attempts to close the budget since
447 1993 with the revised satellite altimeter data, nor have there have been any new assessments of
448 the budget of GMSL rise before the satellite altimeter period. Hay et al. (2015) suggested that
449 their lower rate of 20th-century GMSL rise is more consistent than earlier reconstructions with
450 the sum of contributions. We note that there are several possible combinations of the
451 contributions presented by Gregory et al. (2013) that could match the reconstruction of Hay et al.
452 Moreover, their estimated rate of sea-level rise prior to 1950 is slightly larger than that of Church
453 and White (2011). The small changes in the average of the updated estimates of the glacier
454 contribution from Marzion et al. (2015) are unlikely to have a major impact on closure of the
455 sea-level budget for any of the periods considered.

456 **5. Attribution of sea-level change**

457 Based on understanding of physical processes and results from climate models, the AR5
458 assessed the causes of observed changes in the primary components that contribute to GMSL
459 change (Bindoff et al., 2013). Advances in observations and understanding of changes in global
460 ocean heat content since the AR4 led them to conclude that it is *very likely* that there has been a
461 substantial contribution from anthropogenic forcing since the 1970s, which has thus contributed
462 to thermosteric sea-level rise. For land ice, Bindoff et al. (2013) concluded that it is *likely* that
463 anthropogenic forcing has contributed to mass loss of glaciers since the 1960s and surface
464 melting of the Greenland ice sheet since 1993. Based on the assessment of the components,
465 Bindoff et al. (2013) concluded that it is *very likely* that there is a substantial contribution from
466 anthropogenic forcing to GMSL rise since the 1970s. On the other hand, there was too little
467 evidence available to attribute the causes of the observed mass loss from the Antarctic ice sheet
468 since 1993.

469 Since the AR5, several new studies have quantified the contribution of anthropogenic
470 forcing to changes in the thermosteric and glacier sea-level components, which have been the
471 largest contributors to GMSL rise since 1900 (Church et al., 2013a; Gregory et al., 2013).
472 Marcos and Amores (2014) compared the global thermosteric rise derived from observations of
473 ocean temperature between 0-700 m to results derived from the CMIP5 experiments associated
474 with “natural” (solar and volcanic) and “historical-only” forcings, where the latter include
475 forcing from anthropogenic greenhouse gases and aerosols. For each of the two sets of
476 experiments, they used a signal-to-noise maximizing empirical orthogonal function analysis to
477 separate the signal that is a result of the forcing from the internal variability of the system. This
478 forced response was projected on to the observations to compute the fraction of variability

479 associated with the forced response. On this basis, they find that 87% (range of 72 to 100%) of
480 the observed thermosteric sea-level rise since 1970 is associated with anthropogenic forcing.

481 Slangen et al. (2014b) compared observed estimates of thermosteric sea-level rise from
482 1957 to 2005 over the full depth of the ocean to CMIP5 experiments forced by five different
483 scenarios, including the natural and historical forcing used by Marcos and Amores (2014) as well
484 as anthropogenic-only, greenhouse gas-only, and anthropogenic aerosol-only forcings (Figure 1).
485 They find that the best agreement between the observed and modeled thermosteric sea-level
486 change is with the experiments that include all anthropogenic forcings, indicating a substantial
487 anthropogenic influence on the observed period of sea-level change. The addition of natural
488 forcing that accounts for externally forced variability in the historical experiment improves the
489 agreement on decadal time scales. Based on multiple regression analyses of observed
490 temperature changes onto the simulated responses to the various forcings, Slangen et al. (2014b)
491 derived the scaling factors (β_{est}) by which the simulated responses must be multiplied to obtain
492 the best fit to the observations. These indicate that the modeled anthropogenic response
493 ($\beta_{\text{est}}=1.08 \pm 0.13$) and a reduced modeled natural-only response ($\beta_{\text{est}}=0.70 \pm 0.30$) fit the
494 observations best, similar to what was found in attribution studies of global mean surface
495 temperature (Bindoff et al., 2013).

496 Marzeion et al. (2014a) simulated changes in global glacier SMB since 1851 using a
497 calibrated glacier model forced by the climatology from the natural and historical CMIP5
498 experiments and compared these with observations of global glacier SMB since 1960. Although
499 the natural forcing causes negative SMB over the full modeled period, the simulated SMB is too
500 positive compared to the observations since 1991. In contrast, the SMB simulated by the
501 historical forcing is consistent with the observations for the full period, accounting for an

502 increasing percentage of the total mass loss that reaches $69 \pm 24\%$ since 1991. The additional
503 contribution to GMSL from the anthropogenic influence is ~ 35 mm, with most of this coming in
504 recent decades.

505 In summary, the first formal detection studies for ocean thermal expansion and glacier
506 mass loss since the AR5 was published have confirmed the AR5 finding of a significant
507 anthropogenic contribution to sea-level rise over the last 50 years.

508 **6. Glacier and ice-sheet projections**

509 *6a. Glaciers*

510 The AR5 projected 21st century changes in global glacier SMB (except for Antarctic
511 glaciers) from CMIP5 global mean surface air-temperature projections using a parameterized
512 scheme derived from the results of four published glacier models, with each glacier model given
513 equal weight in the projections (Church et al., 2013b). Each of the glacier models includes
514 detailed treatments of SMB and the evolution of glacier hypsometry. Projected glacier mass
515 losses for 2100 outside Antarctica suggest a *likely* contribution of 0.05 to 0.17 m of sea-level
516 equivalent (SLE) under RCP2.6, corresponding to loss of 15 to 55% of present glacier volume,
517 and 0.10 to 0.26 m SLE under RCP8.5, or 35 to 85% (Church et al., 2013a).

518 Two new studies have examined regional changes in glacier mass over the 21st century.
519 Clarke et al. (2015) modeled glacier mass loss by 2100 for western Canada. They used AOGCM
520 projections for the four RCP scenarios to force a regional glacier model that includes SMB and,
521 for the first time, a detailed physics-based model of ice dynamics that better represents changes
522 in glacier hypsometry. Sensitivity tests of the model excluding ice dynamics show that ice-
523 volume loss is systematically underestimated. Model projections for 2100 suggest that, relative
524 to 2005, western Canada glaciers will lose $\sim 60\%$ of their volume for RCP2.6 and $\sim 80\%$ for

525 RCP8.5, as compared to model results from Marzeion et al. (2012), which suggest ~80% mass
526 loss for RCP2.6 and 100% for RCP8.5, amounting to a difference of <1 mm GMSL. Marzeion et
527 al. (2014b) noted that other 21st century projections similarly showed relatively small differences
528 in glacier mass loss under large differences in forcing scenarios, and concluded that these small
529 differences are because the 21st century response is dominated by the response to climate change
530 during the previous century (i.e., committed glacier loss to previous warming) as well as the
531 effect of hypsometric changes that reduce glacier sensitivity to the forcing as the glaciers
532 decrease in size and retreat to higher elevations.

533 Lang et al. (2015) used a regional climate model coupled to a surface energy balance
534 model to calculate changes in SMB of glaciers in Svalbard under RCP8.5. Their modeled domain
535 is based on a fixed topography, and thus does not capture changes in glacier elevation and
536 hypsometry that can significantly influence rates of mass loss (Marzeion et al., 2014b). By 2100,
537 they find that the accumulation area of all glacierized areas in Svalbard will disappear, with a
538 contribution of ~7 mm SLE corresponding to a loss of 31% of present volume. This contribution
539 is ~25-50% of previous estimates (Marzeion et al., 2012; Radic et al., 2014), with the large
540 differences suggested to reflect the lower resolution climate models and more simplified SMB
541 models used in the earlier studies.

542 In summary, new studies of glacier loss from two regions suggest contributions to GMSL
543 rise that is less than studies assessed by the AR5 that included simulations from these regions
544 (e.g., Marzeion et al., 2012). Because the AR5 did not provide regional estimates, it is not
545 possible to make direct comparisons to these regional studies. Additional regional studies are
546 required to further assess whether there are broader implications of these results.

547 *6b. Greenland Ice Sheet*

548 The Greenland ice sheet loses mass through a combination of changes in SMB and
549 dynamical changes associated with discharge into the ocean, including enhanced basal sliding,
550 and the interaction between SMB and ice flow. The AR5 included both of these components in
551 projections for the 21st century. Projections of SMB were made based on a relation between mass
552 loss and temperature derived from regional climate modeling forced with CMIP5 AOGCMs
553 (Fettweis et al., 2013), with additional allowances for methodological uncertainties and changes
554 in ice-sheet topography (Church et al., 2013b). Projections of rapid change in discharge were
555 based largely on flowline modeling of four of the larger outlet glaciers, with results scaled up by
556 ~5 times because these glaciers drain ~20% of the ice sheet by area (Nick et al., 2013), giving
557 *likely* ranges of 0.02 to 0.09 m by 2100 for RCP8.5, and 0.01 to 0.06 m for the three RCPs of
558 lower forcing, for which there was insufficient information to assess scenario dependence
559 (Church et al., 2013b). *Likely* ranges for projections of the total contributions to GMSL rise by
560 2100 relative to 1986–2005 range from 0.04 to 0.12 m for RCP2.6 to 0.09 to 0.28 m for RCP8.5
561 (Church et al., 2013b). In general, the contributions from SMB and dynamics were assessed as
562 nearly equal for each scenario except RCP8.5, where SMB is greater.

563 Since the AR5, several studies that documented large spatial and temporal variability in
564 discharge of outlet glaciers (Section 3c) suggest that a simple extrapolation and scaling of a few
565 of the largest glaciers for projections may be inaccurate. Csatho et al. (2014) found that mass loss
566 is not proportional to drainage basin area, and that the majority of Greenland mass loss during
567 the 2003–2009 period (~80%) was due to thinning of small to moderately sized drainage basins
568 rather than the four large glaciers modeled by Nick et al. (2013). In contrast, Enderlin et al.
569 (2014) found that these four glaciers represented 42% of the ice-sheet discharge change from
570 2000 to 2012 as opposed to the 20% expected from an assumption of proportionality with

571 drainage area. Based on assumptions about regional variations in discharge through the 21st
572 century, Enderlin et al. (2014) estimated a contribution to GMSL rise of 80 mm, similar to the
573 upper limit of the AR5 *likely* range for the RCP8.5 scenario (85 mm).

574 A number of studies have documented rapid drainage of supraglacial lakes to the base of
575 the ice sheet through conduits, confirming earlier studies that suggested that such drainage
576 increases the rate of short-term ice motion by basal sliding (Joughin et al., 2008; Zwally et al.,
577 2002). Leeson et al. (2015) showed that the area over which supraglacial lakes are distributed
578 will increase by ~50% by 2060 under RCP8.5. However, studies since the AR5 support the
579 conclusion that enhanced basal lubrication makes an insignificant contribution to the *likely* range
580 of sea-level rise over the 21st century. Some studies suggested that this occurs through the
581 subglacial drainage system becoming more efficient with increased discharge, thus reducing
582 basal sliding (Sole et al., 2013; Sundal et al., 2011). Mayaud et al. (2014) forced a subglacial
583 hydrological model of the western ice-sheet margin in the Paakitsoq region with surface runoff
584 that was derived from climate model output for the RCP2.6, 4.5 and 8.5 scenarios. They showed
585 that as runoff increases in response to a warmer climate, the subglacial drainage system
586 transitions from a less efficient network with associated higher basal sliding to a more efficient
587 network with an associated reduction in basal sliding. Shannon et al. (2013) used four ice-sheet
588 models forced by a regional climate model under the A1B scenario to examine the contributions
589 of changes in SMB and of outflow associated with enhanced basal sliding. Enhanced basal
590 sliding was represented by a set of parameterizations that encompasses the range of observations
591 of annual runoff and increase in basal ice flow. Experiments for changes in SMB only find a
592 contribution to GMSL rise by 2100 ranging from 47.5 mm to 61 mm, lying within the AR5 *likely*
593 range of 30 to 150 mm for the same scenario (Church et al., 2013b). Shannon et al. (2013) found

594 that the contribution from basal sliding remains below ~5% of the contribution from SMB.

595 The AR5 assessed the positive feedback between SMB and ice-sheet height as
596 contributing 0 to 15% of SMB change over the course of the 21st century. Based on climate
597 projections under the A1B scenario used to force five ice-sheet models, Edwards et al. (2014)
598 found that the SMB-elevation feedback accounts for an additional 1.8%-6.9% (95% confidence
599 interval) sea-level rise by 2100, in agreement with the AR5.

600 Fürst et al. (2015) forced an ice-sheet model that accounts for basal lubrication and ocean
601 thermal forcing of ice discharge with reanalysis data over the period 2005-2010. The consequent
602 mass loss, of which ~40% is from ice discharge, contributes 0.62 mm yr^{-1} to GMSL rise, similar
603 to observations (Section 3c), giving confidence in the ice-sheet model. With forcing from an
604 ensemble of ten CMIP5 AOGCMs under the four RCP scenarios, Fürst et al. (2015) found the
605 contributions to GMSL rise by 2100 relative to 2000 range from $0.04 \pm 0.02 \text{ m}$ (1 std. dev.) for
606 RCP2.6 to $0.10 \pm 0.03 \text{ m}$ for RCP8.5, which are in good agreement with the AR5 *likely* ranges.
607 Of this mass loss, the contribution from SMB dominates, with discharge decreasing by 2100 as
608 the margin thins and retreats from the coast; enhanced basal lubrication accounts for less than
609 1% of the discharge.

610 In summary, projections of Greenland mass loss support the AR5 assessment that
611 enhanced basal lubrication will make an insignificant contribution sea-level rise over the 21st
612 century. Projections of mass loss from discharge and SMB fall within the AR5 *likely* ranges, but
613 differ from the AR5 in suggesting a greater contribution from SMB, with decreasing discharge
614 associated with the ice-sheet margin retreating from the coast.

615 *6c. Antarctic Ice Sheet*

616 The AR5 projected the contribution to GMSL from the Antarctic ice sheet for changes in
617 SMB (primarily changes in snow accumulation) and outflow across the grounding line to
618 floating ice shelves. In the latter case, mass is subsequently lost in roughly equal amounts by
619 calving and marine melt from the lower ice surface (Rignot et al. (2013). Projections of the
620 combined (SMB and outflow) contribution to GMSL by 2100 relative to 1986-2005 were
621 assessed to fall in a *likely* range of -0.04 to 0.16 m for RCP2.6 and -0.08 to 0.14 m for RCP8.5
622 (Church et al., 2013b). Projections of the outflow contribution to GMSL by 2100 were assessed
623 to fall in a *likely* range of -0.02 to 0.185 m, with no scenario dependence specified due to
624 insufficient information. The assessed contribution from a potential marine ice-sheet instability
625 (MISI) (section 3d) lay outside the *likely* range of the outflow contribution and was characterized
626 by a magnitude of an additional several tens of centimeters (Church et al., 2013a).

627 Studies published since the AR5 can be divided into three groups: those for Pine Island
628 Glacier (PIG) and the other glaciers of the Amundsen Sea Embayment (ASE); studies of the
629 whole of the West Antarctic ice sheet (WAIS) and individual East Antarctic glaciers; and studies
630 of the whole of Antarctica.

631 Favier et al. (2014) reported a comparison of three models of the PIG's evolution over 50
632 years in response to a step change in melt starting from a situation similar to the present day.
633 They obtained a three-to-six fold increase in the rate of mass loss compared to the present-day
634 observations with GMSL contributions in the range of 9 to 25 mm over 50 years. Further
635 experiments in which the initial change in melt is removed suggested that PIG's current retreat
636 may be irreversible. In similar experiments, Seroussi et al. (2014) found GMSL contributions of
637 up to 20 mm in 50 years. Both papers are consistent with the AR5 assessment that PIG's end-of-
638 the-century contribution can be characterized by centimeters of GMSL rise, and lie well within

639 the AR5 assessment of the *likely* contribution from Antarctic rapid ice dynamical change (Table
640 1) (Church et al., 2013a).

641
642 **Table 1: Projections of mass loss by rapid ice-sheet dynamics from the Antarctic ice sheet**
643 **for the 21st century**

Reference	Region of ice sheet	Mass loss (m)	Period
AR5	all of Antarctica	-0.020 to 0.185	1996-2100
Favier et al. (2014)	PIG	0.009 to 0.025	2000-2050
Seroussi et al. (2014)	PIG	up to 0.020	2000-2050
Joughin et al. (2014)	TG	up to 0.025	2000-2100
Gong et al. (2014)	Lambert	up to 0.009	2000-2100
Sun et al. (2014)	Totten	up to 0.020	2000-2100
Cornford et al. (2015)	WAIS	up to 0.200	2000-2100
Levermann et al. (2014)	all of Antarctica (RCP2.6)	0.02 to 0.140	2000-2100
Levermann et al. (2014)	all of Antarctica (RCP8.5)	0.04 to 0.210	2000-2100

644
645 At the time of the AR5, there was no evidence of grounding-line retreat for PIG's larger
646 neighbor Thwaites Glacier (TG). Subsequent research has shown, however, that retreat in other
647 parts of the ASE (most notably TG) is just as pronounced as for PIG (Rignot et al., 2014).
648 Joughin et al. (2014) simulated the consequences of this retreat and found a moderate
649 contribution to GMSL over a century of $<0.25 \text{ mm yr}^{-1}$ (similar to PIG's assessed contribution)
650 with larger rates ($>1 \text{ mm yr}^{-1}$) predicted as the grounding line retreats on to deeper bedrock after
651 200 to 900 years.

652 Cornford et al. (2015) simulated the response of the WAIS to both atmospheric and
653 oceanic forcing derived from regional models that were forced, in turn, by climate models for the

654 A1B and E1 (strong mitigation) scenarios (Figure 2). In particular, the oceanic forcing includes
655 the influx of warm water under the Filchner-Ronne ice shelf reported by Hellmer et al. (2012),
656 although this only occurs towards the end of the current century. Substantial mass loss is
657 therefore dominated by the ASE and contributes up to 50 mm GMSL by 2100, including a
658 contribution from the retreat of TG. Increased outflow is compensated by an increase in SMB,
659 particularly for the Filchner-Ronne sector. This compensation is, however, weaker in the E1
660 scenario, producing the surprising result that the contribution to GMSL is larger in this
661 mitigation scenario than in A1B. This results from the contributions from outflow (oceanic
662 forcing) and SMB being independent, with the latter responding to greenhouse forcing. The
663 compensation term (additional outflow between 0 and 35% of SMB) employed by the AR5 may
664 therefore over estimate this effect and hence exaggerate the contribution to GMSL rise. Finally,
665 more extreme scenarios, in which melt rate is increased across all ice shelves from 1980 (at odds
666 with the regional modeling results and observations), produce contributions of up to 200 mm by
667 2100 (Table 1). In more idealized modeling, Wright et al. (2014) identify Institute and Moller ice
668 streams draining into the Filchner-Ronne ice shelf as close to the threshold for retreat, albeit on a
669 millennial timescale, with associated contributions of ~ 0.14 m to GMSL over this timescale.

670 Gong et al (2014) used a similar methodology to Cornford et al. (2015) for the Lambert
671 Glacier and found that increased outflow is more than compensated by increased SMB, and that
672 only loss of the Amery ice shelf (for instance by large-scale crevassing, see below) leads to
673 GMSL rise (up to 9 mm by 2100 depending on the area lost). Sun et al. (2014) also used similar
674 methods and obtain a contribution from Totten Glacier, East Antarctica, of no more than 20 mm
675 by the end of the century.

676 Pollard et al. (2015) investigated the consequences of crevasse-based loss of the large ice
677 shelves followed (in their model) by the collapse of the ice cliffs left at grounding lines, resulting
678 in large contributions to GMSL of the order of ten meters per millennium (10 mm yr^{-1}). Kuipers
679 Munneke et al. (2014) used a model of firn densification to assess the probability of meltwater
680 ponding on Antarctica's ice shelves in the future, which may then lead to their destabilization
681 through large-scale crevassing such as caused the collapse of the Larsen B ice shelf (MacAyeal
682 et al., 2003). However, they found this effect would not be important outside of the Antarctic
683 Peninsula during the present century, in accord with the assessment by the AR5.

684 Levermann et al. (2014) developed a probabilistic approach to projecting the contribution
685 of Antarctica to GMSL by employing linear response functions based on the response of three
686 ice-sheet models from the SEARISE exercise (Nowicki et al., 2013) to an idealized step increase
687 in marine melt. A scaling is developed between global mean temperature change and ocean
688 temperatures around Antarctica based on the CMIP5 ensemble, which combined with the
689 response functions forms a probabilistic framework for making projections. The approach
690 successfully reproduces the observed contribution between 1992 and 2001, provided a suitable
691 delay between global mean and regional ocean temperatures is used. With this delay, 66%-
692 probability ranges of end-of-the-century GMSL change are 0.02-0.14 m for RCP2.6 and 0.04-
693 0.21 m for RCP8.5. These ranges are broadly similar to the AR5 assessed *likely* range (-0.02 to
694 0.185 m), for which scenario dependence was not assessed, but the upper end for RCP8.5 is
695 slightly higher. A potential contributing factor to the slightly larger range may be the retreat
696 patterns displayed the SEARISE models, which in some cases are fairly extensive.

697 In summary, new studies suggest that a MISI may have been initiated in parts of the
698 WAIS, but that only a few WAIS ice streams will experience MISI during the 21st century. New

699 projections of mass loss from the Greenland and Antarctic Ice Sheets by 2100, including a MISI
700 contribution from parts of WAIS, suggest a contribution that falls largely within the *likely* range
701 (i.e., two-thirds probability) of the AR5. These new results increase confidence in the AR5 *likely*
702 range, indicating that there is a greater probability that sea-level rise by 2100 will fall in that
703 range with a corresponding decrease in the likelihood of an additional contribution of several
704 tens of centimeters above the *likely* range.

705 **7. Projections of global mean sea-level rise by 2100**

706 The AR5 assessed two categories of models used for projections of 21st-century sea-level
707 rise: process-based models that simulate the underlying processes and interactions, and semi-
708 empirical models (SEMs) that are based on statistical relationships between observed GMSL and
709 global mean temperature or total radiative forcing (RF) (Church et al., 2013a). The AR5 assigned
710 medium confidence to process-based projections, due to (i) understanding of the modeled
711 physical processes, (ii) the consistency of the models with wider physical understanding of those
712 processes as elements of the climate system, (iii) the consistency of modeled and observed
713 contributions, and (iv) the consistency of observed and modeled GMSL. As discussed at length
714 in the AR5, SEM projections are based on the assumption that future sea-level change will have
715 the same relationship to global mean temperature change or RF as it had in the period of
716 calibration. This assumption implies uncertainty that is difficult to quantify. Consequently there
717 is low agreement and no consensus in the scientific community about their reliability, giving one
718 reason for which they were assigned low confidence by the AR5. The other is that, in nearly
719 every case, SEMs projected a substantially higher sea level by 2100 than the process-based
720 models, but no satisfactory physical explanation was available for this. To date there have been
721 no new projections of sea level using revised estimates of sea level to train semi-empirical

722 models. Orlic and Parasic (2015) analyzed three variants of semi-empirical models and showed
723 that the projections are similar through the middle of the 21st century but diverge after that and
724 urge caution in their use beyond the 21st century.

725 On the basis of their higher assessed confidence, the AR5 projections for the 21st century
726 are based on process-based models (Church et al., 2013a). For RCP2.6, the *likely* range of sea-
727 level rise by 2100 is 0.28 to 0.61 m. For RCP8.5, the *likely* range by 2100 is 0.52 to 0.98 m with
728 a rate during 2081–2100 of 8 to 16 mm yr⁻¹. Church et al. (2013a) concluded that “there is
729 currently insufficient evidence to evaluate the probability of specific levels above the assessed
730 *likely* range” (p. 1140), but that sea levels substantially higher than the *likely* range would only
731 occur in the 21st century if the marine-based sections of the Antarctic ice sheet were to collapse.
732 The potential contribution could not be precisely quantified, but they determined with medium
733 confidence that “this additional contribution would not exceed several tenths of a meter of sea-
734 level rise during the 21st century” (p. 1140).

735 Three studies since the AR5 have made projections for 21st-century GMSL rise based on
736 some level of expert elicitation. Horton et al. (2014) presented projections of GMSL rise based
737 on a survey involving 90 individuals with some experience in publishing on sea level. The
738 respondents estimated 66% confidence intervals for GMSL rise by 2100, with respect to the
739 time-mean of 1986 to 2005. Their median estimate of the *likely* range was 0.7 to 1.2 m for
740 scenario RCP8.5; they noted that their range lies above the AR5 *likely* range of 0.52 to 0.98 m.
741 They also reported a *very likely* range of 0.5 to 1.5 m for RCP8.5. In commenting on Horton et
742 al., Gregory et al. (2014) suggested that some respondents might have assumed an extensive
743 Antarctic marine ice-sheet instability to lie within the *likely* range of possibilities. According to
744 the AR5, a MISI is *unlikely* to occur, but if initiated it could add up to several tenths of a metre to

745 GMSL rise during the 21st century, making the AR5 and Horton et al. consistent. However, as
746 discussed in section 6c, the several detailed ice-sheet modeling studies that have been published
747 since the AR5 suggest a contribution from Antarctic ice-sheet rapid dynamical change, including
748 the possibility of MISI, that falls largely within the AR5 *likely* range, with only one estimate
749 being higher, by 0.025 m, or one-tenth of the difference (0.22 m) between the upper estimates of
750 the *likely* range by the AR5 and Horton et al.

751 Kopp et al. (2014) combined results from process-based models and expert elicitation to
752 derive projections of 21st-century GMSL rise from each of the main components, which were
753 then summed to derive the total rise. For the ice sheets, they reconciled the scenario differences
754 between the AR5 projections and those from the expert elicitation by using the AR5 to
755 characterize the median and *likely* ranges and expert elicitation to calibrate the shape of the tails.
756 They used results from a single glacier model (Marzeion et al., 2012), one of four considered by
757 the AR5, that by itself has a narrower *likely* range than in the AR5, results from a selection of
758 CMIP5 GCMs that give a higher and wider *likely* range than in the AR5, and a relationship
759 between changes in land water storage and population that was not considered by the AR5 with
760 the *likely* range falling within the AR5 *likely* range. Despite these differences, Kopp et al. (2014)
761 derive a similar median (0.79m) and *likely* range (0.60 to 1.00 m) for the total GMSL rise for the
762 RCP8.5 scenario as in the AR5. In addition, when including the results from the expert
763 elicitation, they estimated a *very likely* range of 0.52 to 1.21 m, and a *virtually certain* range
764 (99% probability) of 0.39 to 1.76 m.

765 Jevrejeva et al. (2014a) combined the uncertainty estimates from the AR5 results for
766 thermosteric, glacier, and land water storage for the RCP8.5 scenario with results from an
767 assessment on contributions from ice-sheet dynamics by 13 experts (Bamber and Aspinall, 2013)

768 to derive a probability distribution function (PDF) for 21st-century GMSL rise. The main
769 difference between the ice-sheet contributions lies in the shape assumed for the uncertainty
770 distribution. The AR5 assumed it to be uniform within the *likely* range from the AR5; Jevrejeva
771 et al. used the positively skewed distribution from the expert elicitation, which had a long tail
772 extending towards the high end, dominated by Antarctica and reflecting the wide spread and thus
773 lack of consensus among the expert estimates (Bamber and Aspinall, 2013). Consequently their
774 combined PDF for GMSL rise is positively skewed by the Antarctic contribution, and their
775 median and *likely* range are 0.79 [0.58 to 1.20] m, to be compared with 0.74 m [0.52 to 0.98] m
776 in the AR5. Jevrejeva et al. (2014a) report a *very likely* range of 0.45 to 1.80 m. These latter
777 values are significantly larger than Kopp et al. and Horton et al. and inconsistent with recent
778 results on simulation of the Antarctic ice sheet contributions using state of the art models
779 discussed in section 6c.

780 In summary, the three studies of projections of 21st-century GMSL rise published since
781 the AR5 all depend on results from expert elicitation. Jevrejeva et al. (2014a) and Kopp et al.
782 (2014) used them to help characterize the tails of the distribution about the *likely* range derived
783 from process-based models, providing a means to infer ranges for higher probabilities, whereas
784 Horton et al. (2014) relied entirely on them to characterize the *likely* range. We have low
785 confidence in conclusions based on expert elicitation because, as noted by Gregory et al. (2014),
786 the respondents are not asked to justify, and we cannot know, how they arrived at their
787 conclusions. Although the physical processes have operated in the past, to our knowledge, the
788 circumstances of anthropogenic forcing are entirely unprecedented in recent Earth history. In
789 these circumstances, intuition should not be trusted, because there is no relevant experience in
790 which it could have been trained to make the right decision, and we believe that the only rational

791 approach to quantitative assessment is by transparent reasoned analysis of available results. For
792 instance, Figure 2 of Horton et al. shows that several of the respondents placed the 83-percentile
793 for GMSLR by 2100 for RCP8.5 above 2.5 m, i.e. more than 1.5 m above the AR5 *likely* range,
794 the largest estimate being at about 6 m. No physically plausible scenarios for values above 2.5 m
795 have been published in the peer-reviewed literature; on the contrary, it has been argued that they
796 are physically untenable (Pfeffer et al., 2008). Furthermore, these projections are inconsistent
797 with recent results on simulation of the Antarctic ice sheet contributions using state-of-the-art
798 models (section 6).

799 **8. Regional projections and emergence time of the forced signal**

800 GMSLR due to the increase in the volume of the global ocean affects relative sea level
801 everywhere, but projected local RSL change differs from GMSLR because of various effects,
802 which we can put in five categories:

- 803 1. The ocean dynamic response (the change in the ocean dynamic topography) to the change in
804 its 3-D density field (arising from temperature and salinity change) and circulation, caused by the
805 effect of climate change on surface heat, freshwater and momentum (wind stress) fluxes. Many
806 studies consider the sum of ocean dynamic sea-level change and the contribution of global steric
807 change (thermal expansion) to GMSLR; we refer to this sum as “ocean climate change”.
- 808 2. The “GeLi” effects, or “fingerprints” (section 2b) of land-ice change, on the geoid and the
809 elastic response of the lithosphere.
- 810 3. Glacial isostatic adjustment (GIA), which is the ongoing change in the geopotential field and
811 deformation of the solid Earth due to the reduction of ice-sheet mass since the Last Glacial
812 Maximum ~20,000 years ago.

813 4. The redistribution of atmospheric mass, i.e. the change in mean sea-level pressure. This term
814 is shown in the AR5 and earlier to be small compared with other multidecadal changes.

815 5. Local tectonic effects, which are not discussed further here (see Kopp et al. (2015) for a
816 discussion).

817 The AR4 considered ocean dynamical change, but not GeLi effects, because it was not
818 able to make a projection of rapid ice-sheet dynamic change, and therefore could not produce
819 estimates of regional sea-level change due to all contributions. Subsequent advances (following
820 Katsman et al. (2011), Slangen et al. (2012), and Church et al. (2011a)) enabled this to be done
821 in the AR5, which therefore presented projections of RSL change including all the above effects
822 except tectonics, which were also excluded. A number of regional projections and projections at
823 coastal tide gauge sites, building on the AR4 and AR5 assessments and using the AR5 approach,
824 have also been completed (Carson et al., 2015b; CSIRO, 2014; Han et al., 2014; Hunter et al.,
825 2013; Kopp et al., 2014; Little et al., 2015; McInnes et al., 2015; Simpson et al., 2014; Slangen et
826 al., 2014a). Slangen et al. (2014a) also made RSL projections including all effects based on
827 CMIP5 simulations of ocean climate change, combined with somewhat different estimates for
828 the land-ice contributions from the AR5 assessment. They pointed out that the distribution of
829 RSL change is positively skewed (also seen in AR5, Fig 13.22) because the GeLi effect of land-
830 ice change is slightly positive in large low-latitude areas and strongly negative in small high-
831 latitude areas near to the ice sheets and glaciers.

832 As in the AR5, Slangen et al. (2014a) projected RSL change of less than the global mean
833 (by up to 50%) near West Antarctica and Greenland, due to GeLi effects of the ice sheets. On the
834 other hand, the AR5 and Slangen et al. (2014a) projected RSL rise exceeding the global mean
835 (by more than 20%) on the east coast of North America and in a roughly zonal band on the north

836 side of the Antarctic Circumpolar Current. These phenomena are due to ocean climate change.
837 They are commonly seen in AOGCMs and have previously been noted (e.g. AR4, Pardaens et al.
838 (2011)), but their magnitude differs greatly among models (Bouttes et al., 2012; Yin, 2012).
839 Bouttes and Gregory (2014) provided evidence that the model spread in projected changes in
840 ocean surface fluxes of momentum (wind stress), heat, and freshwater forcing all contribute to
841 the model spread of projected ocean dynamic SL change.

842 Bouttes et al. (2012) demonstrated that the north-south zonal dipole (positive ocean
843 dynamical sea-level rise to the north of the ACC and negative to the south) is largely a
844 thermosteric effect caused by and roughly proportional to the change in the mid-latitude westerly
845 winds in the Southern Ocean, which AOGCMs generally project to increase in magnitude and
846 shift southward, consequently shifting and tilting the isopycnals in the ocean. Frankcombe et al.
847 (2013) reproduced this effect in an eddy-permitting ocean model (1/4-degree resolution). They
848 pointed out that the same feature in RSL change can be seen in the altimeter trends since 1993
849 and could be due to anthropogenic influence (forced by greenhouse gases or ozone depletion) on
850 the Southern Annular Mode (Cai and Cowan, 2007; Thompson et al., 2011). It is possible that
851 the effect of eddy saturation in eddy-resolving models might reduce the effect, although Suzuki
852 and Ishii (2011) showed similar sea-level change in eddy-permitting and lower-resolution
853 models. In CMIP5 models, Bouttes and Gregory (2014) found that surface heat flux change also
854 contributes to producing the feature, which is apparent in CMIP5 historical simulations as well
855 as in projections, and in some regions it is already larger than simulated unforced trends of the
856 length of the altimeter period, consistent with it being a response to anthropogenic forcing
857 (Bilbao et al., 2015).

858 The enhanced RSL rise in the North Atlantic is also a part of dipole feature, with reduced
859 RSL rise to the south, though its geographical detail is model dependent (Bouttes et al., 2014;
860 Swingedouw et al., 2013). This feature has previously been associated with weakening of the
861 AMOC due to buoyancy forcing, and is caused by a change in surface heat flux (a reduction of
862 heat loss in the North Atlantic in a warming climate) and increase of surface freshwater flux in
863 CMIP5 models (Bouttes and Gregory, 2014), as in earlier models. RSL rise in the north of the
864 North Atlantic, whether due to heat or freshwater input, is mainly the direct result of the added
865 buoyancy (Bouttes et al., 2014; Swingedouw et al., 2013), and the consequent weakening of the
866 AMOC brings about a redistribution of heat which opposes the effect of the surface flux change
867 on RSL in most of the affected area, but reinforces it along the east coast of North America
868 (Bouttes et al., 2014), where the projected RSL rise is consequently particularly large (Yin et al.,
869 2009). Howard et al. (2014) assessed the magnitude of the effects of increased freshwater inflow
870 from ice-sheet mass loss on ocean dynamic sea-level change and the AMOC to be small
871 compared with the effects of changing surface fluxes from the atmosphere.

872 The dominant feature of observed RSL change in the altimeter period is the contrast of
873 rising sea level in the west and falling in the east in the Pacific, due to changes in wind stress
874 forcing (England et al., 2014; Griffies et al., 2014). It has been argued that unforced variability
875 associated with the Pacific Decadal Oscillation (PDO) (Hamlington et al., 2013; Hamlington et
876 al., 2014; Merrifield et al., 2012; Zhang and Church, 2012) contributes to this pattern, although
877 Frankcombe et al. (2015) found that the relationship between the sea-level pattern and the PDO
878 is not statistically robust in short observational records. Hamlington et al. (2014) removed the
879 influence of PDO statistically, and isolated a remaining sea-level pattern associated with
880 warming in the western tropical Pacific, which they suggested could be due to anthropogenic

881 warming of the tropical Indian Ocean. However, this residual sea-level pattern was somewhat
882 different to that found in Zhang and Church (2012) and may not be robust because of the short
883 available satellite altimeter record (about half a PDO cycle), and Palanisamy et al. (2015) also
884 disputed this interpretation. The east-west Pacific pattern (Figure 3) does not appear to be
885 anthropogenic in CMIP5 historical simulations, but on the other hand it is too large to be
886 consistent with unforced variability as simulated by the AOGCMs (Bilbao et al., 2015; Carson et
887 al., 2015a; Palanisamy et al., 2015). Consequently this remains an important phenomenon that
888 still requires an explanation. It could also be associated with some of the explanations advanced
889 for the “hiatus” of global warming (England et al., 2015; Kosaka and Xie, 2013; Meehl et al.,
890 2011).

891 Under a given future scenario, RSL change due to ocean climate change will be due to a
892 combination of forced response and unforced variability of the climate system. The former grows
893 with time, while the latter, although staying roughly constant (Little et al., 2015), may be
894 substantial. Hu and Deser (2013) showed that even by the middle of this century, unforced
895 variability leads to an uncertainty of a up to factor of two in coastal projections, the largest
896 spread (in proportional terms) being on the north Pacific and north Atlantic coasts, despite the
897 very small spread in projections of global mean thermal expansion. Bordbar et al. (2015) found
898 that even after 100 years, the spread in local projections due to unforced variability may be
899 comparable to global thermal expansion.

900 The “time of emergence” is when the forced signal becomes discernible because it is
901 sufficiently large compared with unforced variability (Hawkins and Sutton, 2012). Several recent
902 studies have evaluated the time of emergence of sea-level change, with various definitions, in the
903 CMIP5 dataset. For ocean dynamical sea-level change (which excludes GMSLR), the local

904 signal is not detectable in the majority of the ocean even after 100 years of anthropogenic forcing
905 (Bordbar et al., 2015; Lyu et al., 2014) (Figures 4, 5). It only emerges clearly in the Southern
906 Ocean near Antarctica, where projected sea-level change is markedly less than the global average
907 due to the north-south zonal dipole. A non-uniform pattern of change (i.e. contrasts between
908 dynamic sea-level change in different regions) should be detectable in only a few years, and
909 indeed already may be detected in the Southern Ocean (Bilbao et al., 2015).

910 If dynamical sea-level change and thermal expansion are taken together (i.e. considering
911 sea-level change due to ocean climate change), the signal of forced sea-level change relative to
912 the reference mean level of 1986-2005 emerges in about half of the ocean area by 2040 (Lyu et
913 al., 2014) (Figures 4, 5). Instead of using a reference level, a comparison of trends in projections
914 of forced sea-level change starting in 1990 with simulated unforced trends of the same length
915 shows a detectable signal by the early 2030s in half of the ocean area (Richter and Marzeion,
916 2014). In the MPI-ESM-LR AOGCM, chosen as an example, trends of 20 years starting in 2006
917 exceed one standard deviation of unforced variability in more than half of the area, and trends of
918 50 years in more than 90% (Carson et al., 2015a). With both techniques, the region of earliest
919 emergence is the low-latitude Atlantic, where unforced variability is particularly small (Bilbao et
920 al., 2015; Carson et al., 2015a; Little et al., 2015; Lyu et al., 2014; Richter and Marzeion, 2014).
921 Early emergence is also noted in some coastal areas, especially the Atlantic coast of North
922 America (Carson et al., 2015a; Richter and Marzeion, 2014). By contrast, in much the Southern
923 Ocean, where variability is high and the forced response is small, the signal of change may not
924 be detectable until late in the 21st century or afterwards (Bilbao et al., 2015; Lyu et al., 2014;
925 Richter and Marzeion, 2014). If RSL including all contributions to GMSLR (not just thermal

926 expansion) and GeLi effects is considered, the signal of forced change is discernible in about half
927 of the ocean area as early as 2020 (Lyu et al., 2014) (Figures 4, 5).

928 Bilbao et al. (2015) showed that in CMIP5 models the patterns of predicted sea-level
929 change due to ocean climate change (Figure 6) are stable in time and fairly independent of
930 scenario. Consequently accurate projections can be made by scaling a fixed pattern with a time-
931 dependent magnitude, as for surface air temperature, but ocean volume-mean temperature is
932 generally a slightly more accurate predictor than global mean surface air temperature for scaling
933 the pattern of sea-level change. Recent work has begun to explore the impact of individual
934 radiative forcings (greenhouse gases, aerosols and natural forcing) on historical sea-level change
935 (Slangen et al., in press).

936 In summary, there has been new work in three main areas since the AR5. First, analysis
937 of models indicates that the two commonly predicted features of future regional sea-level
938 change, namely the increasing tilt across the ACC and the dipole in the North Atlantic (both with
939 enhanced sea-level rise on their northern side), are related to regional changes in wind stress and
940 surface heat flux. Much remains to be understood about the geographical patterns and remote
941 influences of surface flux changes, and their effects on ocean circulation and interior transports,
942 in order to make confident regional projections. Second, it is expected that sea-level change in
943 response to anthropogenic forcing in regions of relatively low unforced variability, such as the
944 low-latitude Atlantic, will be detectable within 20 years or less. We note that an anthropogenic
945 influence has already been detected on the global mean (Section 5). Third, the east-west contrast
946 in the Pacific of sea-level trends observed since the early 1990s cannot be satisfactorily
947 accounted for by climate models, nor yet definitively attributed either to unforced variability
948 (such as the PDO) or forced climate change.

949 **9. Synthesis**

950 Recent analyses confirm the 20th-century sea-level rise, with some studies showing a
951 slightly smaller rate before 1990 and some a slightly larger value than reported in the AR5.
952 There is now clearly more evidence of an acceleration in the rate of rise from the 19th to the 20th
953 century (as reported in the AR5), during the 20th century, from the pre-1990 rate to the rate
954 during the altimeter period, and with a positive (but not significant) rate in the altimeter record.

955 Ongoing ocean heat uptake and associated thermal expansion have continued and
956 increased since 2000 (compared with previous decades), with a significant amount of heat being
957 stored deeper in the water column and with the largest storage in the Southern Ocean. New
958 estimates of global glacier mass loss indicate improved agreement between studies and thus
959 smaller uncertainties over the 20th century, but with little change in their average value. Two
960 regional studies suggest glacier contributions to GMSL rise that are less than assessed by the
961 AR5, but additional studies are required to assess the implications for the global estimates. The
962 acceleration of mass loss from Greenland, primarily as a result of increased surface melting is
963 continuing. Mass loss from the Antarctic ice sheet is also accelerating, with most of that loss
964 coming from discharge from the Amundsen Sea sector of the WAIS that is larger than the
965 estimated increase in accumulation in East Antarctica. Observations and model simulations
966 suggest that this acceleration may be associated with a marine ice-sheet instability that may have
967 been initiated in parts of the WAIS by an increased flux of CDW across the continental shelf.
968 When the Antarctic and Greenland ice sheets are taken together, the estimated acceleration is
969 larger than observed from the satellite altimeter sea-level record, including at the upper end
970 where their combined acceleration (range from 0.112 to 0.140 mm yr⁻²) exceeds that inferred
971 from analysis of the satellite altimeter sea-level record (0.099 mm yr⁻²) (Watson et al., 2015).

972 However, we note that the ice-sheet estimates (2003-2013) are only for the latter half of the
973 satellite altimeter period (1993 to mid-2014), during which time the sea-level acceleration may
974 be larger. Also increased mass gain in land water storage and parts of East Antarctica, and
975 decreased mass loss from glaciers in Alaska and some other regions such that the total
976 acceleration in the ocean mass gain is consistent with the satellite altimeter record. There have
977 not yet been any new comprehensive attempts to close the budget since 1900 or 1993, but it
978 would appear that following the methodology of Gregory et al. (2013), the sum of contributions
979 can explain the observed rise.

980 The first formal detection studies for ocean thermal expansion and glacier mass loss,
981 which were the two largest contributors to 20th-century GMSL rise, have confirmed the AR5
982 assessment of a significant anthropogenic contribution to sea-level rise over the last 50 years.

983 Projections of mass loss from Greenland ice-sheet discharge and SMB fall within the
984 AR5 *likely* ranges, but differ from the AR5 in suggesting a greater contribution from SMB, a
985 decreasing discharge as the ice-sheet margin retreats from the coast, and with an insignificant
986 contribution from enhanced basal lubrication. Post-AR5 studies of mass loss from the Greenland
987 and Antarctic Ice Sheets by 2100, including a MISI contribution from parts of WAIS, suggest a
988 contribution that falls largely within the *likely* range (i.e., two-thirds probability) of the AR5.

989 Three post-AR5 studies of projections of 21st-century GMSL derive a *very likely* range.
990 However, these projections are based on expert elicitations and we have low confidence in
991 deriving a *very likely* range in projections from such an approach.

992 Analyses of models indicate that the two commonly predicted features of future regional
993 sea-level change, namely the increasing tilt across the ACC and the dipole in the North Atlantic
994 (both with enhanced sea-level rise on their northern side), are related to regional changes in wind

995 stress and surface heat flux. However, much remains to be understood about the geographical
996 patterns and remote influences of surface flux changes, and their effects on ocean circulation and
997 sea level. In particular, the east-west contrast in the Pacific of sea-level trends observed since the
998 early 1990s cannot be satisfactorily accounted for by climate models, nor yet definitely attributed
999 either to unforced variability (such as the PDO) or forced climate change.

1000 In view of the comparatively limited state of knowledge and understanding of rapid ice-
1001 sheet dynamics, we continue to think that it is not yet possible to make reliable quantitative
1002 estimates of future GMSL rise outside the *likely* range. However, new ice-sheet modeling results
1003 increase confidence in the AR5 *likely* range, indicating that there is a greater probability that sea-
1004 level rise by 2100 will fall in that range with a corresponding decrease in the likelihood of an
1005 additional contribution of several tens of centimeters above the *likely* range.

1006 Despite these uncertainties, it is clear that the sea-level response to anthropogenic forcing
1007 will be detectable over most of the ocean by 2040, confirming the importance of sea-level
1008 change as a major issue that society will have to confront during the 21st century and beyond.
1009 The AR5 projections and the updates clearly indicate the rate of rise in 2100 is directly related to
1010 the future emissions and that avoiding a larger sea-level rise will require significant and urgent
1011 mitigation of greenhouse gas emissions. As emphasized by Lowe and Gregory (2010): “It is vital
1012 to continue to monitor sea level and its components and to develop a capability to make reliable
1013 projections. [A]s we cannot provide certainties, we must become better at explaining the
1014 uncertainties to decision makers. These uncertainties imply a need to keep open a range of
1015 adaptation [and mitigation] options and to be able to change the approach as the predictions
1016 become more robust” (p. 43).

1017 **Acknowledgments**

1018 PUC was supported by the U.S. National Science Foundation Antarctic Glaciology Program
1019 (ANT-1043517) and the Marine Geology and Geophysics Program (OCE-1335197). JAC was
1020 supported by the Australian Government Department of the Environment, the Bureau of
1021 Meteorology and CSIRO through the Australian Climate Change Science Programme. We thank
1022 two reviewers for their helpful comments.

1023 **Conflict of Interest Statement**

1024 On behalf of all authors, the corresponding author states that there is no conflict of
1025 interest.

1026 **Figure Captions**

1027
 1028 Figure 1. Thermosteric global mean sea-level change (m) with respect to 2005. CMIP5
 1029 multimodel mean (red) $\pm 2 \sigma$ (light grey), realization mean (blue), individual realizations (dark
 1030 grey). Numbers of models/realizations in brackets. Observations (black lines): Domingues (solid
 1031 + uncertainties), Levitus (dashes), Ishii (dots). **(a)** Internal variability, **(b)** natural forcing only,
 1032 **(c)** greenhouse gas forcing only, **(d)** aerosol forcing only, **(e)** anthropogenic forcing only, and **(f)**
 1033 all forcings combined. (From Slangen et al., 2104)

1034
 1035 Figure 2. Net change in volume above flotation in the WAIS over the course of the combined
 1036 experiment in of both atmospheric and oceanic forcing derived from regional models that were
 1037 forced, in turn, using climate models for the A1B and E1 (strong mitigation) scenarios. Only the
 1038 Amundsen Sea Embayment experiences a net loss (ΔV) in all of the combined experiments.
 1039 Nonetheless, the result is a net loss over West Antarctica as a whole. Note that Thwaites glacier
 1040 does not retreat in the combined anomaly experiments (which use the synthetic accumulation),
 1041 and the ASE could contribute an extra $9 \times 10^3 \text{ km}^3$ loss by 2100 and $40 \times 10^3 \text{ km}^3$ by 2200. (From
 1042 Cornford et al. 2015)

1043
 1044 Figure 3. Observed sea level change trends (mm yr^{-1}) from satellite altimetry between 1993–
 1045 2012. The hatching indicates trends that are significant (at the 5% level) with respect to at least
 1046 2/3 of CMIP5 pre-industrial control simulations.

1047
 1048 Figure 4. Multimodel ensemble median time-of-emergence (ToE) for regional sea-level change
 1049 under RCP8.5. Different change signals are used: **a**, ocean dynamic topography change (dynamic
 1050 sea level); **b**, ocean climate change (dynamic sea-level change plus global mean thermosteric
 1051 sea-level change); **c**, relative sea-level change (total sea-level change). Warm (cold) colours
 1052 represent rising (falling) sea level; light grey areas have no emergence before 2080; deep grey
 1053 colour means no agreement among models; white colour means no data coverage or over land.
 1054 (From Lyu et al. 2014)

1055
 1056 Figure 5. The cumulative fraction of the total area with the emergence of change signals before
 1057 the given time from the multimodel ensemble median patterns. Dynamic sea level (blue), ocean
 1058 climate change (dynamic sea level plus global mean thermosteric sea level, black), relative sea-
 1059 level (total sea level, red), surface air temperature (green). Dash-dot lines are for RCP4.5 and
 1060 solid lines for RCP8.5. (From Lyu et al. 2014)

1061
 1062 Figure 6. CMIP5 model ensemble mean and **b** standard deviation of the forced patterns of ocean
 1063 dynamic topography change ($\text{m}/^\circ\text{C}$) calculated using ocean volume mean temperature ($^\circ\text{C}$) as
 1064 predictor for the historical + RCP4.5 simulations between 1993–2099. (From Bilboa et al. 2015)

1065

1066 **References**

- 1067 Ablain, M., Cazenave, A., Larnicol, G., Balmaseda, M., Cipollini, P., Faugere, Y., Fernandes,
1068 M.J., Henry, O., Johannessen, J.A., Knudsen, P., Andersen, O., Legeais, J., Meyssignac,
1069 B., Picot, N., Roca, M., Rudenko, S., Scharffenberg, M.G., Stammer, D., Timms, G. and
1070 Benveniste, J., 2015. Improved sea level record over the satellite altimetry era
1071 (1993-2010) from the Climate Change Initiative project. *Ocean Science*, 11(1): 67-
1072 82.
- 1073 Abraham, J.P., Baringer, M., Bindoff, N.L., Boyer, T., Cheng, L.J., Church, J.A., Conroy, J.L.,
1074 Domingues, C.M., Fasullo, J.T., Gilson, J., Goni, G., Good, S.A., Gorman, J.M., Gouretski,
1075 V., Ishii, M., Johnson, G.C., Kizu, S., Lyman, J.M., Macdonald, A.M., Minkowycz, W.J.,
1076 Moffitt, S.E., Palmer, M.D., Piola, A.R., Reseghetti, F., Schuckmann, K., Trenberth, K.E.,
1077 Velicogna, I. and Willis, J.K., 2013. A Review of global ocean temperature
1078 observations: Implications for ocean heat content estimates and climate change.
1079 *Reviews of Geophysics*, 51(3): 450-483.
- 1080 Allan, R.P., Liu, C., Loeb, N.G., Palmer, M.D., Roberts, M., Smith, D. and Vidale, P.-L., 2014.
1081 Changes in global net radiative imbalance 1985–2012. *GEOPHYSICAL RESEARCH*
1082 *LETTERS*, 41(15): 5588-5597.
- 1083 Balmaseda, M.A., Trenberth, K.E. and Källén, E., 2013. Distinctive climate signals in
1084 reanalysis of global ocean heat content. *GEOPHYSICAL RESEARCH LETTERS*, 40(9):
1085 1754-1759.
- 1086 Bamber, J.L. and Aspinall, W.P., 2013. An expert judgement assessment of future sea level
1087 rise from the ice sheets. *Nature Climate Change*, 3: 424-427.
- 1088 Becker, M., Karpytchev, M. and Lennartz-Sassinek, S., 2014. Long-term sea level trends:
1089 Natural or anthropogenic? *Geophysical Research Letters*, 41: 5571-5580.
- 1090 Beenstock, M., Felsenstein, D., Frank, E. and Reingewertz, Y., 2015. Tide gauge location and
1091 the measurement of global sea level rise. *Environmental and Ecological Statistics*,
1092 22(1): 179-206.
- 1093 Bilbao, R.A.F., Gregory, J.M. and Bouttes, N., 2015. Analysis of the regional pattern of sea
1094 level change due to ocean dynamics and density change for 1993–2009 in
1095 observations and CMIP5 AOGCMs. *Climate Dynamics*.
- 1096 Bindoff, N.L., Stott, P.A., AchutaRao, K.M., Allen, M.R., Gillett, N., Gutzler, D., Hansingo, K.,
1097 Hegerl, G., Hu, Y., Jain, S., IMokhov, I.I., Overland, J., Perlwitz, J., Sebbari, R. and Zhang,
1098 X., 2013. Detection and attribution of climate change: from global to regional. In: T.F.
1099 Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V.
1100 Bex and P.M. Midgley (Editors), *Climate Change 2013: The Physical Science Basis*.
1101 Contribution of Working Group I to the Fifth Assessment Report of the
1102 Intergovernmental Panel on Climate Change. Cambridge University Press,
1103 Cambridge, U.K., pp. 867-952.
- 1104 Bordbar, M.H., Martin, T., Latif, M. and Park, W., 2015. Effects of long-term variability on
1105 projections of twenty-first-century dynamic sea level. *Nature Climate Change*, 5(4):
1106 343-347.
- 1107 Bos, M.S., S. D. P. Williams, S.D.P., Araujo, I.B. and Bastos, L., 2014. The effect of temporal
1108 correlated noise on the sea level rate and acceleration uncertainty. *Geophysical*
1109 *Journal International* 196: 1423-1430.

- 1110 Bouttes, N. and Gregory, J.M., 2014. Attribution of the spatial pattern of CO₂-forced sea
1111 level change to ocean surface flux changes. *Environmental Research Letters*, 9(3).
1112 Bouttes, N., Gregory, J.M., Kuhlbrodt, T. and Smith, R.S., 2014. The drivers of projected
1113 North Atlantic sea level change. *Climate Dynamics*, 43(5-6): 1531-1544.
1114 Bouttes, N., Gregory, J.M., Kuhlbrodt, T. and Suzuki, T., 2012. The effect of windstress
1115 change on future sea level change in the Southern Ocean. *Geophysical Research*
1116 *Letters*, 39.
- 1117 Cahill, N., Kemp, A.C., Horton, B.P. and Parnell, A.C., 2015. Modeling sea-level change using
1118 errors-in-variables integrated Gaussian processes *Annals of Applied Statistics*, 9(2):
1119 547-571.
- 1120 Cai, W.J. and Cowan, T., 2007. Trends in Southern Hemisphere circulation in IPCC AR4
1121 models over 1950-99: Ozone depletion versus greenhouse forcing. *Journal of*
1122 *Climate*, 20(4): 681-693.
- 1123 Carson, M., Kohl, A. and Stammer, D., 2015a. The Impact of Regional Multidecadal and
1124 Century-Scale Internal Climate Variability on Sea Level Trends in CMIP5 Models.
1125 *Journal of Climate*, 28(2): 853-861.
- 1126 Carson, M., Kohl, A., Stammer, D., Aimee Slangen, Katsman, C.C.A., van de Wal, R.S.W.,
1127 Church, J. and White, N., 2015b. Coastal sea level changes, observed and projected
1128 during the 20th and 21st century. *Climate Dynamics*, in press.
- 1129 Cazenave, A., Dieng, H.B., Meyssignac, B., von Schuckmann, K., Decharme, B. and Berthier, E.,
1130 2014. The rate of sea-level rise. *Nature Climate Change*, 4(5): 358-361.
- 1131 Chang, Y.-S., Vecchi, G.A., Rosati, A., Zhang, S. and Yang, X., 2014. Comparison of global
1132 objective analyzed T-S fields of the upper ocean for 2008–2011. *Journal of Marine*
1133 *Systems*, 137(0): 13-20.
- 1134 Chen, X. and Tung, K.-K., 2014. Varying planetary heat sink led to global-warming
1135 slowdown and acceleration. *Science*, 345(6199): 897-903.
- 1136 Cheng, L. and Zhu, J., 2014a. Influences of the Choice of Climatology on Ocean Heat Content
1137 Estimation. *Journal of Atmospheric and Oceanic Technology*, 32(2): 388-394.
- 1138 Cheng, L. and Zhu, J., 2014b. Uncertainties of the Ocean Heat Content Estimation Induced by
1139 Insufficient Vertical Resolution of Historical Ocean Subsurface Observations. *Journal*
1140 *of Atmospheric and Oceanic Technology*, 31(6): 1383-1396.
- 1141 Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A., Merrifield,
1142 M.A., Milne, G.A., Nerem, R.S., Nunn, P.D., Payne, A.J., Pfeffer, W.T., D., S. and
1143 Unnikrishnan, A.S., 2013a. Sea level change. In: T.F. Stocker, D. Qin, G.-K. Plattner, M.
1144 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (Editors),
1145 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
1146 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*
1147 *Cambridge University Press, Cambridge, UK, pp. 1137-1216.*
- 1148 Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A., Merrifield,
1149 M.A., Milne, G.A., Nerem, R.S., Nunn, P.D., Payne, A.J., Pfeffer, W.T., D., S. and
1150 Unnikrishnan, A.S., 2013b. Sea level change supplementary material. In: T.F. Stocker,
1151 D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and
1152 P.M. Midgley (Editors), *Climate Change 2013: The Physical Science Basis.*
1153 *Contribution of Working Group I to the Fifth Assessment Report of the*
1154 *Intergovernmental Panel on Climate Change. Cambridge University Press,*
1155 *Cambridge, U.K., pp. 13SM-1-13SM-8.*

- 1156 Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A., Merrifield,
1157 M.A., Milne, G.A., Nerem, R.S., Nunn, P.D., Payne, A.J., Pfeffer, W.T., Stammer, D. and
1158 Unnikrishnan, A.S., 2013c. Sea-level rise by 2100. *Science*, 342(6165): 1445-1445.
- 1159 Church, J.A., Gregory, J.M., White, N.J., Platten, S.M. and Mitrovica, J.X., 2011a. Understanding
1160 and Projecting Sea Level Change. *Oceanography*, 24(2): 130-143.
- 1161 Church, J.A. and White, N.J., 2011. Sea-level rise from the late 19th to the early 21st century.
1162 *Surveys in Geophysics*, 32: 585-602.
- 1163 Church, J.A., White, N.J., Konikow, L.F., Domingues, C.M., Cogley, J.G., Rignot, E., Gregory, J.M.,
1164 van den Broeke, M.R., Monaghan, A.J. and Velicogna, I., 2011b. Revisiting the Earth's
1165 sea-level and energy budgets from 1961 to 2008. *Geophysical Research Letters*, 38.
- 1166 Church, J.A., White, N.J., Konikow, L.F., Domingues, C.M., Cogley, J.G., Rignot, E., Gregory, J.M.,
1167 van den Broeke, M.R., Monaghan, A.J. and Velicogna, I., 2013d. Revisiting the Earth's
1168 sea-level and energy budgets from 1961 to 2008 (vol 38, L18601, 2011).
1169 *Geophysical Research Letters*, 40(15): 4066-4066.
- 1170 Clarke, G.K.C., Jarosch, A.H., Anslow, F.S., Radic, V. and Menounos, B., 2015. Projected
1171 deglaciation of western Canada in the twenty-first century. *Nature Geoscience*, 8(5):
1172 372-377.
- 1173 Collins, M., R. Knutti, J. Arblaster, J.-L. Dufresne, T. Fichefet, P. Friedlingstein, X. Gao and W.J.
1174 Gutowski, T.J., G. Krinner, M. Shongwe, C. Tebaldi, A.J. Weaver and M. Wehner, 2013.
1175 Long-term climate change: Projections, commitments and irreversibility. In: T.F.
1176 Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V.
1177 Bex and P.M. Midgley (Editors), *Climate Change 2013: The Physical Science Basis*.
1178 Contribution of Working Group I to the Fifth Assessment Report of the
1179 Intergovernmental Panel on Climate Change. Cambridge University Press,
1180 Cambridge, UK, pp. 1029-1136.
- 1181 Cornford, S.L., Martin, D.F., Payne, A.J., Ng, E.G., Le Brocq, A.M., Gladstone, R.M., Edwards,
1182 T.L., Shannon, S.R., Agosta, C., van den Broeke, M.R., Hellmer, H.H., Krinner, G.,
1183 Ligtenberg, S.R.M., Timmermann, R. and Vaughan, D.G., 2015. Century-scale
1184 simulations of the response of the West Antarctic Ice Sheet to a warming climate.
1185 *The Cryosphere Discuss.*, 9(2): 1887-1942.
- 1186 Csatho, B.M., Schenk, A.F., van der Veen, C.J., Babonis, G., Duncan, K., Rezvanbehbahani, S.,
1187 van den Broeke, M.R., Simonsen, S.B., Nagarajan, S. and van Angelen, J.H., 2014. Laser
1188 altimetry reveals complex pattern of Greenland Ice Sheet dynamics. *Proceedings of*
1189 *the National Academy of Sciences of the United States of America*, 111(52): 18478-
1190 18483.
- 1191 CSIRO, A.B.o.M.a., 2014. *Climate Variability, Extremes and Change in the Western Tropical*
1192 *Pacific: New Science and Updated Country Reports*. Pacific-Australia Climate Change
1193 Science and Adaptation Planning Program Technical Report, Australian Bureau of
1194 Meteorology and Commonwealth Scientific and Industrial Research Organisation,
1195 Melbourne, Australia.
- 1196 Dieng, H., Palanisamy, H., Cazenave, A., Meyssignac, B. and von Schuckmann, K., 2015. The
1197 Sea Level Budget Since 2003: Inference on the Deep Ocean Heat Content. *Surveys in*
1198 *Geophysics*, 36(2): 209-229.
- 1199 Domingues, C.M., Church, J.A., White, N.J., Gleckler, P.J., Wijffels, S.E., Barker, P.I.M. and Dunn,
1200 J.R., 2008. Improved estimates of upper-ocean warming and multi-decadal sea-level
1201 rise. *Nature*, 453(7198): 1090.

- 1202 Douglas, B.C., 1992. Global sea-level acceleration. *Journal of Geophysical Research-Oceans*,
1203 97: 12699-12706.
- 1204 Durack, P.J., Gleckler, P.J., Landerer, F.W. and Taylor, K.E., 2014. Quantifying underestimates
1205 of long-term upper-ocean warming. *Nature Clim. Change*, 4(11): 999-1005.
- 1206 Edwards, T.L., Fettweis, X., Gagliardini, O., Gillet-Chaulet, F., Goelzer, H., Gregory, J.M.,
1207 Hoffman, M., Huybrechts, P., Payne, A.J., Perego, M., Price, S., Quiquet, A. and Ritz, C.,
1208 2014. Effect of uncertainty in surface mass balance-elevation feedback on
1209 projections of the future sea level contribution of the Greenland ice sheet.
1210 *Cryosphere*, 8(1): 195-208.
- 1211 Enderlin, E.M., Howat, I.M., Jeong, S., Noh, M.J., van Angelen, J.H. and van den Broeke, M.R.,
1212 2014. An improved mass budget for the Greenland ice sheet. *Geophysical Research*
1213 *Letters*, 41(3): 866-872.
- 1214 England, M.H., Kajtar, J.B. and Maher, N., 2015. Robust warming projections despite the
1215 recent hiatus. *Nature Climate Change*, 5: 394-396.
- 1216 England, M.H., McGregor, S., Spence, P., Meehl, G.A., Timmermann, A., Cai, W.J., Sen Gupta, A.,
1217 McPhaden, M.J., Purich, A. and Santoso, A., 2014. Recent intensification of wind-
1218 driven circulation in the Pacific and the ongoing warming hiatus. *Nature Climate*
1219 *Change*, 4(3): 222-227.
- 1220 Fasullo, J.T., Boening, C., Landerer, F.W. and Nerem, R.S., 2013. Australia's unique influence
1221 on global sea level in 2010-2011. *Geophysical Research Letters*, 40(16): 4368-4373.
- 1222 Favier, L., Durand, G., Cornford, S.L., Gudmundsson, G.H., Gagliardini, O., Gillet-Chaulet, F.,
1223 Zwinger, T., Payne, A.J. and Le Brocq, A.M., 2014. Retreat of Pine Island Glacier
1224 controlled by marine ice-sheet instability. *Nature Climate Change*, 4(2): 117-121.
- 1225 Fettweis, X., Franco, B., Tedesco, M., van Angelen, J.H., Lenaerts, J.T.M., van den Broeke, M.R.
1226 and Gallee, H., 2013. Estimating Greenland ice sheet surface mass balance
1227 contribution to future sea level rise using the regional atmospheric model MAR. *The*
1228 *Cryosphere*, 7: 469-489.
- 1229 Frankcombe, L.M., McGregor, S. and England, M.H., 2015. Robustness of the modes of Indo-
1230 Pacific sea level variability. *Climate Dynamics*, 45(5-6): 1281-1298.
- 1231 Frankcombe, L.M., Spence, P., Hogg, A.M., England, M.H. and Griffies, S.M., 2013. Sea level
1232 changes forced by Southern Ocean winds. *Geophysical Research Letters*, 40(21):
1233 5710-5715.
- 1234 Fürst, J.J., Goelzer, H. and Huybrechts, P., 2015. Ice-dynamic projections of the Greenland ice
1235 sheet in response to atmospheric and oceanic warming. *The Cryosphere*, 9: 1039-
1236 1062.
- 1237 Gong, Y., Cornford, S.L. and Payne, A.J., 2014. Modelling the response of the Lambert
1238 Glacier-Amery Ice Shelf system, East Antarctica, to uncertain climate forcing over
1239 the 21st and 22nd centuries. *Cryosphere*, 8(3): 1057-1068.
- 1240 Gregory, J.M., Church, J.A., Clark, P.U., Payne, A.J., Merrifield, M.A., Nerem, R.S., Nunn, P.D.,
1241 Pfeffer, W.T. and Stammer, D., 2014. Comment on "Expert assessment of sea-level
1242 rise by AD 2100 and AD 2300", by Horton et al. (2014). *Quaternary Science Reviews*,
1243 97: 193-194.
- 1244 Gregory, J.M., White, N.J., Church, J.A., Bierkens, M.F.P., Box, J.E., van den Broeke, M.R.,
1245 Cogley, J.G., Fettweis, X., Hanna, E., Huybrechts, P., Konikow, L.F., Leclercq, P.W.,
1246 Marzeion, B., Oerlemans, J., Tamisiea, M.E. and Van de Wal, R.S.W., 2013. Twentieth-

- 1247 century global-mean sea level rise: Is the whole greater than the sum of the parts?
 1248 *Journal of Climate*, 26: 4476-4499.
- 1249 Griffies, S.M., Yin, J.J., Durack, P.J., Goddard, P., Bates, S.C., Behrens, E., Bentsen, M., Bi, D.H.,
 1250 Biastoch, A., Boning, C.W., Bozec, A., Chassignet, E., Danabasoglu, G., Danilov, S.,
 1251 Domingues, C.M., Drange, H., Farneti, R., Fernandez, E., Greatbatch, R.J., Holland,
 1252 D.M., Ilicak, M., Large, W.G., Lorbacher, K., Lu, J.H., Marsland, S.J., Mishra, A., Nurser,
 1253 A.J.G., Melia, D.S.Y., Palter, J.B., Samuels, B.L., Schroter, J., Schwarzkopf, F.U.,
 1254 Sidorenko, D., Treguier, A.M., Tseng, Y.H., Tsujino, H., Uotila, P., Valcke, S., Voldoire,
 1255 A., Wang, Q., Winton, M. and Zhang, X.B., 2014. An assessment of global and regional
 1256 sea level for years 1993-2007 in a suite of interannual CORE-II simulations. *Ocean*
 1257 *Modelling*, 78: 35-89.
- 1258 Haigh, I.D., Wahl, T., Rohling, E.J., Price, R.M., Pattiaratchi, C.B., Calafat, F.M. and Dangendorf,
 1259 S., 2014. Timescales for detecting a significant acceleration in sea level rise. *Nature*
 1260 *Communications*, 5.
- 1261 Hamlington, B.D., Leben, R.R., Strassburg, M.W., Nerem, R.S. and Kim, K.-Y., 2013.
 1262 Contribution of the Pacific decadal oscillation to global mean sea level trends.
 1263 *Geophysical Research Letters*, 40: 5171-5175.
- 1264 Hamlington, B.D., Strassburg, M.W., Leben, R.R., Han, W., Nerem, R.S. and Kim, K.Y., 2014.
 1265 Uncovering an anthropogenic sea-level rise signal in the Pacific Ocean. *Nature*
 1266 *Climate Change*, 4(9): 782-785.
- 1267 Hamlington, B.D. and Thompson, P.R., 2015. Considerations for estimating the 20th century
 1268 trend in global mean sea level. *Geophysical Research Letters*, 42: 4102-4109.
- 1269 Han, G., Ma, Z., Bao, H. and Slangen, A., 2014. Regional differences of relative sea level
 1270 changes in the Northwest Atlantic: Historical trends and future projections. *Journal*
 1271 *of Geophysical Research: Oceans*, 119(1): 156-164.
- 1272 Harig, C. and Simons, F.J., 2015. Accelerated West Antarctic ice mass loss continues to
 1273 outpace East Antarctic gains. *Earth and Planetary Science Letters*, 415: 134-141.
- 1274 Hawkins, E. and Sutton, R., 2012. Time of emergence of climate signals. *Geophysical*
 1275 *Research Letters*, 39.
- 1276 Hay, C.C., Morrow, E., Kopp, R.E. and Mitrovica, J.X., 2015. Probabilistic reanalysis of
 1277 twentieth-century sea-level rise. *Nature*, 517(7535): 481-+.
- 1278 Hellmer, H.H., Kauker, F., Timmermann, R., Determann, J. and Rae, J., 2012. Twenty-first-
 1279 century warming of a large Antarctic ice-shelf cavity by a redirected coastal current.
 1280 *Nature*, 485: 225-228.
- 1281 Helm, V., Humbert, A. and Miller, H., 2014. Elevation and elevation change of Greenland and
 1282 Antarctica derived from CryoSat-2. *Cryosphere*, 8(4): 1539-1559.
- 1283 Hogarth, P., 2014. Preliminary analysis of acceleration of sea level rise through the
 1284 twentieth century using extended tide gauge data sets (August 2014). *Journal of*
 1285 *Geophysical Research-Oceans*, 119(11): 7645-7659.
- 1286 Horton, B., Rahmstorf, S., Engelhart, S.E. and Kemp, A.C., 2014. Expert assessment of sea-
 1287 level rise by AD 2100 and AD 2300. *Quaternary Science Review*, 84: 1-6.
- 1288 Howard, T., Ridley, J., Pardaens, A.K., Hurkmans, R.T.W.L., Payne, A.J., Giesen, R.H., Lowe,
 1289 J.A., Bamber, J.L., Edwards, T.L. and Oerlemans, J., 2014. The land-ice contribution to
 1290 21st-century dynamic sea level rise. *Ocean Science*, 10(3): 485-500.
- 1291 Hu, A.X. and Deser, C., 2013. Uncertainty in future regional sea level rise due to internal
 1292 climate variability. *Geophysical Research Letters*, 40(11): 2768-2772.

- 1293 Hunter, J.R., Church, J.A., White, N.J. and Zhang, X., 2013. Towards a global regionally
1294 varying allowance for sea-level rise. *Ocean Engineering*, 71(0): 17-27.
- 1295 Jevrejeva, S., Grinsted, A. and Moore, J.C., 2014a. Upper limit for sea level projections by
1296 2100. *Environmental Research Letters*, 9(10).
- 1297 Jevrejeva, S., Moore, J.C., Grinsted, A., Matthews, A.P. and Spada, G., 2014b. Trends and
1298 acceleration in global and regional sea levels since 1807. *Global and Planetary
1299 Change*, 113: 11-22.
- 1300 Jongman, B., Ward, P.J. and Aerts, J.C.J.H., 2012. Global exposure to river and coastal
1301 flooding: long term trends and changes. *Global Environmental Change: Human and
1302 Policy Dimensions*, 22: 823-835.
- 1303 Jorda, G., 2014. Detection time for global and regional sea level trends and accelerations.
1304 *Journal of Geophysical Research-Oceans*, 119(10): 7164-7174.
- 1305 Joughin, I., Das, S.B., King, M.A., Smith, B.E., Howat, I.M. and Moon, T., 2008. Seasonal
1306 speedup along the western flank of the Greenland Ice Sheet. *Science*, 320(5877):
1307 781-783.
- 1308 Joughin, I., Smith, B.E. and Medley, B., 2014. Marine Ice Sheet Collapse Potentially Under
1309 Way for the Thwaites Glacier Basin, West Antarctica. *Science*, 344(6185): 735-738.
- 1310 Katsman, C.A., Sterl, A., Beersma, J.J., van den Brink, H.W., Church, J.A., Hazeleger, W., Kopp,
1311 R.E., Kroon, D., Kwadijk, J., Lammensen, R., Lowe, J., Oppenheimer, M., Plag, H.-P.,
1312 Ridley, J., von Storch, H., Vaughan, D.G., Vellinga, P., Vermeersen, L.L.A., van de Wal,
1313 R.S.W. and Weisse, R., 2011. Exploring high-end scenarios for local sea level rise to
1314 develop flood protection strategies for a low-lying delta—the Netherlands as an
1315 example. *Climatic Change*, 109: 617-645.
- 1316 Kenigson, J.S. and Han, W., 2014. Detecting and understanding the accelerated sea level rise
1317 along the east coast of the United States during recent decades. *Journal of
1318 Geophysical Research-Oceans*, 119(12): 8749-8766.
- 1319 Kerr, R.A., 2013. A stonger IPCC report. *Science*, 342: 43.
- 1320 Khan, S.A., Kjaer, K.H., Bevis, M., Bamber, J.L., Wahr, J., Kjeldsen, K.K., Bjork, A.A., Korsgaard,
1321 N.J., Stearns, L.A., van den Broeke, M.R., Liu, L., Larsen, N.K. and Muresan, I.S., 2014.
1322 Sustained mass loss of the northeast Greenland ice sheet triggered by regional
1323 warming. *Nature Climate Change*, 4(4): 292-299.
- 1324 Kopp, R.E., Hay, C.C., Little, C.M. and Mitrovica, J.X., 2015. Geographic variability of sea-level
1325 change. *Current Climate Change Reports*, in press.
- 1326 Kopp, R.E., Horton, R. M. , Little, C.M., Mitrovica, J.X., Oppenheimer, M., Rasmussen, D.J.,
1327 Strauss, B.H. and Tebaldi, C., 2014. Probabilistic 21st and 22nd century sea-level
1328 projections at a global network of tide-gauge sites. *Earth's Future*, 2: 383-406.
- 1329 Kosaka, Y. and Xie, S.-P., 2013. Recent global-warming hiatus tied to equatorial Pacific
1330 surface cooling. *Nature*, 501: 403-407.
- 1331 Kuipers Munneke, P., Ligtenberg, S.R.M., van den Broeke, M.R. and Vaughan, D.G., 2014. Firn
1332 air depletion as a precursor of Antarctic ice-shelf collapse. *Journal of Glaciology*,
1333 60(220): 205-214.
- 1334 Lang, C., Fettweis, X. and Erpicum, M., 2015. Future climate and surface mass balance of
1335 Svalbard glaciers in an RCP8.5 climate scenario: a study with the regional climate
1336 model MAR forced by MIROC5. *Cryosphere*, 9(3): 945-956.

- 1337 Leeson, A.A., Shepherd, A., Briggs, K., Howat, I., Fettweis, X., Morlighem, M. and Rignot, E.,
1338 2015. Supraglacial lakes on the Greenland ice sheet advance inland under warming
1339 climate. *Nature Climate Change*, 5(1): 51-55.
- 1340 Lenaerts, J.T.M., van Meijgaard, E., van den Broeke, M.R., Ligtenberg, S.R.M., Horwath, M.
1341 and Isaksson, E., 2013. Recent snowfall anomalies in Dronning Maud Land, East
1342 Antarctica, in a historical and future climate perspective. *Geophysical Research*
1343 *Letters*, 40(11): 2684-2688.
- 1344 Leuliette, E., 2015. The balancing of the sea-level budget. *Current Climate Change Reports*,
1345 1: 1-7.
- 1346 Levermann, A., Winkelmann, R., Nowicki, S., Fastook, J.L., Frieler, K., Greve, R., Hellmer, H.H.,
1347 Martin, M.A., Meinshausen, M., Mengel, M., Payne, A.J., Pollard, D., Sato, T.,
1348 Timmermann, R., Wang, W.L. and Bindshadler, R.A., 2014. Projecting Antarctic ice
1349 discharge using response functions from SeaRISE ice-sheet models. *Earth System*
1350 *Dynamics*, 5(2): 271-293.
- 1351 Levitus, S., Antonov, J.I., Boyer, T.P., Baranova, O.K., Garcia, H.E., Locarnini, R.A., Mishonov,
1352 A.V., Reagan, J.R., Seidov, D., Yarosh, E.S. and Zweng, M.M., 2012. World ocean heat
1353 content and thermosteric sea level change (0-2000 m), 1955-2010. *Geophys. Res.*
1354 *Lett.*, 39(10): L10603.
- 1355 Lichter, M., Vafeidis, A.T., Nicholls, R.J. and Kaiser, G., 2011. Exploring data-related
1356 uncertainties in analyses of land area and population in the 'low-elevation coastal
1357 zone' (LECZ). *Journal of Coastal Research*, 27: 757-768.
- 1358 Little, C.M., Horton, R.M., Kopp, R.E., Oppenheimer, M. and Yip, S., 2015. Uncertainty in
1359 Twenty-First-Century CMIP5 Sea Level Projections. *Journal of Climate*, 28(2): 838-
1360 852.
- 1361 Loeb, N.G., Lyman, J.M., Johnson, G.C., Allan, R.P., Doelling, D.R., Wong, T., Soden, B.J. and
1362 Stephens, G.L., 2012. Observed changes in top-of-the-atmosphere radiation and
1363 upper-ocean heating consistent within uncertainty. *Nature Geosci*, 5(2): 110-113.
- 1364 Lowe, J. and Gregory, J.M., 2010. A sea of uncertainty. *Nature Reports Climate Change*, 4: 42-
1365 43.
- 1366 Lyman, J.M., Good, S.A., Gouretski, V.V., Ishii, M., Johnson, G.C., Palmer, M.D., Smith, D.M. and
1367 Willis, J.K., 2010. Robust warming of the global upper ocean. *Nature*, 465(7296):
1368 334-337.
- 1369 Lyu, K.W., Zhang, X.B., Church, J.A., Slangen, A.B.A. and Hu, J.Y., 2014. Time of emergence for
1370 regional sea-level change. *Nature Climate Change*, 4(11): 1006-1010.
- 1371 MacAyeal, D.R., Scambos, T.A., Hulbe, C.L. and Fahnestock, M.A., 2003. Catastrophic ice-shelf
1372 break-up by an ice-shelf-fragment-capsize mechanism. *Journal of Glaciology*, 49: 22-
1373 36.
- 1374 Marcos, M. and Amores, A., 2014. Quantifying anthropogenic and natural contributions to
1375 thermosteric sea level rise. *Geophysical Research Letters*, 41(7): 2502-2507.
- 1376 Marzeion, B., Cogley, J.G., Richter, K. and Parkes, D., 2014a. Attribution of global glacier
1377 mass loss to anthropogenic and natural causes. *Science*, 345(6199): 919-921.
- 1378 Marzeion, B., Jarosch, A.H. and Gregory, J.M., 2014b. Feedbacks and mechanisms affecting
1379 the global sensitivity of glaciers to climate change. *Cryosphere*, 8(1): 59-71.
- 1380 Marzeion, B., Jarosch, A.H. and Hofer, M., 2012. Past and future sea-level change from the
1381 surface mass balance of glaciers. *Cryosphere*, 6(6): 1295-1322.

- 1382 Marzeion, B., Leclercq, P.W., Cogley, J.G. and Jarosch, A.H., 2015. Brief communication:
1383 Global glacier mass loss reconstructions during the 20th century are consistent. *The*
1384 *Cryosphere Discussions*, 9: 3807–3820.
- 1385 Mayaud, J.R., Banwell, A.F., Arnold, N.S. and Willis, I.C., 2014. Modeling the response of
1386 subglacial drainage at Paakitsoq, west Greenland, to 21st century climate change.
1387 *Journal of Geophysical Research-Earth Surface*, 119(12): 2619-2634.
- 1388 McGranahan, G., Balk, D. and Anderson, B., 2007. The rising tide: assessing the risks of
1389 climate change and human settlements in low elevation coastal zones. *Environment*
1390 *and Urbanization*, 19: 17-37.
- 1391 McInnes, K.L., Church, J.A., Monselesan, D., Hunter, J.R., O’Grady, J.G., Haigh, I.D. and Zhang,
1392 X., 2015. Sea-level Rise Projections for Australia: Information for Impact and
1393 Adaptation Planning. *Australian Meteorological and Oceanographic Journal*, in press.
- 1394 McMillan, M., Shepherd, A., Sundal, A., Briggs, K., Muir, A., Ridout, A., Hogg, A. and Wingham,
1395 D., 2014. Increased ice losses from Antarctica detected by CryoSat-2. *Geophysical*
1396 *Research Letters*, 41(11): 3899-3905.
- 1397 Meehl, G.A., Arblaster, J.M., Fasullo, J.Y., Hu, A. and Trenberth, K.E., 2011. Model-based
1398 evidence of deep-ocean heat uptake during surface-temperature hiatus periods.
1399 *Nature Climate Change*, 1: 360-364.
- 1400 Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., Gaye, A.T., Gregory, J.M., Kitoh, A.,
1401 Knutti, R., Murphy, J.M., Noda, A., Raper, S.C.B., Watterson, I.G., Weaver, A.J. and Zhao,
1402 Z.-C., 2007. Global climate projections. In: S. Solomon, D. Qin, M. Manning, Z. Chen, M.
1403 Marquis, K.B. Averyt, M. Tignor and H.L. Miller (Editors), *Climate Change 2007: The*
1404 *Physical Science Basis. Contribution of Working Group I to the Fourth Assessment*
1405 *Report of the Intergovernmental Panel on Climate Change Cambridge University*
1406 *Press, Cambridge, UK, pp. 748-845.*
- 1407 Meinshausen, M., Smith, S.J., Calvin, K., Daniel, J.S., Kainuma, M.L.T., Lamarque, J.F.,
1408 Matsumoto, K., Montzka, S.A., Raper, S.C.B., Riahi, K., Thomson, A., Velders, G.J.M. and
1409 van Vuuren, D.P.P., 2011. The RCP greenhouse gas concentrations and their
1410 extensions from 1765 to 2300. *Climatic Change*, 109(1-2): 213-241.
- 1411 Merrifield, M.A., Thompson, P.R. and Lander, M., 2012. Multidecadal sea level anomalies
1412 and trends in the western tropical Pacific. *Geophysical Research Letters*, 39:
1413 L13602.
- 1414 Mooney, C., 2014. The world’s climate change watchdog may be underestimating global
1415 warming, *Washington Post*.
- 1416 Moss, R.H., Edmonds, J.A., Hibbard, K.A., Manning, M.R., Rose, S.K., van Vuuren, D.P., Carter,
1417 T.R., Emori, S., Kainuma, M., Kram, T., Meehl, G.A., Mitchell, J.F.B., Nakicenovic, N.,
1418 Riahi, K., Smith, S.J., Stouffer, R.J., Thomson, A.M., Weyant, J.P. and Wilbanks, T.J.,
1419 2010. The next generation of scenarios for climate change research and assessment.
1420 *Nature*, 463(7282): 747-756.
- 1421 Nick, F.M., Vieli, A., Andersen, M.L., Joughin, I., Payne, A., Edwards, T.L., Pattyn, F. and van de
1422 Wal, R.S.W., 2013. Future sea-level rise from Greenland's main outlet glaciers in a
1423 warming climate. *Nature*, 497(7448): 235-238.
- 1424 Nieves, V., Willis, J.K. and Patzert, W.C., 2015. Recent hiatus caused by decadal shift in Indo-
1425 Pacific heating. *Science*, 349(6247): 532-535.
- 1426 Nowicki, S., Bindschadler, R.A., Abe-Ouchi, A., Aschwanden, A., Bueller, E., Choi, H., Fastook,
1427 J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U.C., Jackson, C., Johnson, J.,

- 1428 Khroulev, C., Larour, E., Levermann, A., Lipscomb, W.H., Martin, M.A., Morlighem, M.,
1429 Parizek, B.R., Pollard, D., Price, S.F., Ren, D., Rignot, E., Saito, F., Sato, T., Seddik, H.,
1430 Seroussi, H., Takahashi, K., Walker, R. and Wang, W.L., 2013. Insights into spatial
1431 sensitivities of ice mass response to environmental change from the SeaRISE ice
1432 sheet modeling project I: Antarctica. *Journal of Geophysical Research: Earth Surface*,
1433 118: 1002-1024.
- 1434 Olivieri, M. and Spada, G., 2013. Intermittent sea-level acceleration. *Global and Planetary*
1435 *Change*, 109: 64-72.
- 1436 Orlic, M. and Pasarić, Z., 2015. Some pitfalls of the semiempirical method used to project
1437 sea level. *Journal of Climate*, 28(9): 3779-3785.
- 1438 Palanisamy, H., Meyssignac, B., Cazenave, A. and Delcroix, T., 2015. Is anthropogenic sea
1439 level fingerprint already detectable in the Pacific Ocean? *Environmental Research*
1440 *Letters*, 10: 084024.
- 1441 Palmer, M.D. and McNeall, D.J., 2014. Internal variability of Earth's energy budget simulated
1442 by CMIP5 climate models. *Environmental Research Letters*, 9(3): 034016.
- 1443 Paolo, F.S., Fricker, H.A. and Padman, L., 2015. Volume loss from Antarctic ice shelves is
1444 accelerating. *Science*, 348(6232): 327-331.
- 1445 Pardaens, A.K., Gregory, J.M. and Lowe, J.A., 2011. A model study of factors influencing
1446 projected changes in regional sea level over the twenty-first century. *Climate*
1447 *Dynamics*, 36(9-10): 2015-2033.
- 1448 Pfeffer, W.T., Harper, J.T. and O'Neel, S., 2008. Kinematic constraints on glacier
1449 contributions to 21st-century sea-level rise. *Science*, 321: 1340-1343.
- 1450 Piecuch, C.G. and Ponte, R.M., 2015. Inverted barometer contributions to recent sea level
1451 changes along the northeast coast of North America. *Geophysical Research Letters*,
1452 42(14): 5918-5925.
- 1453 Pollard, D., DeConto, R.M. and Alley, R.B., 2015. Potential Antarctic Ice Sheet retreat driven
1454 by hydrofracturing and ice cliff failure. *Earth and Planetary Science Letters*, 412:
1455 112-121.
- 1456 Purkey, S.G. and Johnson, G.C., 2010. Warming of Global Abyssal and Deep Southern Ocean
1457 Waters between the 1990s and 2000s: Contributions to Global Heat and Sea Level
1458 Rise Budgets. *Journal of Climate*, 23(23): 6336-6351.
- 1459 Purkey, S.G., Johnson, G.C. and Chambers, D.P., 2014. Relative contributions of ocean mass
1460 and deep steric changes to sea level rise between 1993 and 2013. *Journal of*
1461 *Geophysical Research-Oceans*, 119(11): 7509-7522.
- 1462 Radic, V., Bliss, A., Beedlow, A.C., Hock, R., Miles, E. and Cogley, J.G., 2014. Regional and
1463 global projections of twenty-first century glacier mass changes in response to
1464 climate scenarios from global climate models. *Climate Dynamics*, 42: 37-58.
- 1465 Rahmstorf, S., 2013. Sea level in the 5th IPCC report, RealClimate.org.
- 1466 Rhein, M., Rintoul, S.R., Aoki, S., Campos, E., Chambers, D., Feely, R.A., Gulev, S., Johnson, G.C.,
1467 Josey, S.A., Kostianoy, A., Mauritzen, C., Roemmich, D., Talley, L.D. and Wang, F.,
1468 2013. Observations: Oceans. In: T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K.
1469 Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (Editors), *Climate*
1470 *Change 2013: The Physical Science Basis. Contribution of Working Group I to the*
1471 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*
1472 Cambridge University Press, Cambridge, U.K., pp. 255-315.

- 1473 Richter, K. and Marzeion, B., 2014. Earliest local emergence of forced dynamic and steric
1474 sea-level trends in climate models. *Environmental Research Letters*, 9(11).
- 1475 Rignot, E., Jacobs, S., Mouginot, J. and Scheuchl, B., 2013. Ice-Shelf Melting Around
1476 Antarctica. *Science*, 341(6143): 266-270.
- 1477 Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H. and Scheuchl, B., 2014. Widespread,
1478 rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers,
1479 West Antarctica, from 1992 to 2011. *Geophysical Research Letters*, 41(10): 3502-
1480 3509.
- 1481 Roemmich, D., Church, J., Gilson, J., Monselesan, D., Sutton, P. and Wijffels, S., 2015.
1482 Unabated planetary warming and its ocean structure since 2006. *Nature Clim.*
1483 *Change*, 5(3): 240-245.
- 1484 Schmidtko, S., Heywood, K.J., Thompson, A.F. and Aoki, S., 2014. Multidecadal warming of
1485 Antarctic waters. *Science*, 346(6214): 1227-1231.
- 1486 Seroussi, H., Morlighem, M., Rignot, E., Mouginot, J., Larour, E., Schodlok, M. and Khazendar,
1487 A., 2014. Sensitivity of the dynamics of Pine Island Glacier, West Antarctica, to
1488 climate forcing for the next 50 years. *Cryosphere*, 8(5): 1699-1710.
- 1489 Shannon, S.R., Payne, A.J., Bartholomew, I.D., van den Broeke, M.R., Edwards, T.L., Fettweis,
1490 X., Gagliardini, O., Gillet-Chaulet, F., Goelzer, H., Hoffman, M.J., Huybrechts, P., Mair,
1491 D.W.F., Nienow, P.W., Perego, M., Price, S.F., Smeets, C.J.P.P., Sole, A.J., van de Wal,
1492 R.S.W. and Zwinger, T., 2013. Enhanced basal lubrication and the contribution of the
1493 Greenland ice sheet to future sea-level rise. *Proceedings of the National Academy of*
1494 *Sciences of the United States of America*, 110(35): 14156-14161.
- 1495 Simpson, M.R., Breili, K. and Kierulf, H., 2014. Estimates of twenty-first century sea-level
1496 changes for Norway. *Climate Dynamics*, 42(5-6): 1405-1424.
- 1497 Slangen, A., Church, J.A., Zhang, X. and Monselesan, D., in press. The sea-level response to
1498 external forcings in CMIP5 climate models. *Journal of Climate*.
- 1499 Slangen, A.B.A., Carson, M., Katsman, C.A., van de Wal, R.S.W., Kohl, A., Vermeersen, L.L.A.
1500 and Stammer, D., 2014a. Projecting twenty-first century regional sea-level changes.
1501 *Climatic Change*, 124(1-2): 317-332.
- 1502 Slangen, A.B.A., Church, J.A., Zhang, X.B. and Monselesan, D., 2014b. Detection and
1503 attribution of global mean thermosteric sea level change. *Geophysical Research*
1504 *Letters*, 41(16): 5951-5959.
- 1505 Slangen, A.B.A., Katsman, C.A., van de Wal, R.S.W., Vermeersen, L.L.A. and Riva, R.E.M., 2012.
1506 Towards regional projections of twenty-first century sea-level change based on IPCC
1507 SRES scenarios. *Climate Dynamics*, 38: 1191-1209.
- 1508 Smith, D.M., Allan, R.P., Coward, A.C., Eade, R., Hyder, P., Liu, C., Loeb, N.G., Palmer, M.D.,
1509 Roberts, C.D. and Scaife, A.A., 2015. Earth's energy imbalance since 1960 in
1510 observations and CMIP5 models. *GEOPHYSICAL RESEARCH LETTERS*, 42(4):
1511 2014GL062669.
- 1512 Sole, A., Nienow, P., Bartholomew, I., Mair, D., Cowton, T., Tedstone, A. and King, M.A., 2013.
1513 Winter motion mediates dynamic response of the Greenland Ice Sheet to warmer
1514 summers. *Geophysical Research Letters*, 40: 3940-3944.
- 1515 Spada, G., Olivieri, M. and Galassi, G., 2015. A heuristic evaluation of long-term global sea
1516 level acceleration. *Geophysical Research Letters*, 42: 4166-4172.
- 1517 Sun, S., Cornford, S.L., Liu, Y. and Moore, J.C., 2014. Dynamic response of Antarctic ice
1518 shelves to bedrock uncertainty. *Cryosphere*, 8(4): 1561-1576.

- 1519 Sundal, A.V., Shepherd, A., Nienow, P., Hanna, E., Palmer, S. and Huybrechts, P., 2011. Melt-
1520 induced speed-up of Greenland ice sheet offset by efficient subglacial drainage.
1521 *Nature*, 469: 521-524.
- 1522 Sutterley, T.C., Velicogna, I., Rignot, E., Mouginot, J., Flament, T., van den Broeke, M.R., van
1523 Wessem, J.M. and Reijmer, C.H., 2014. Mass loss of the Amundsen Sea Embayment of
1524 West Antarctica from four independent techniques. *Geophysical Research Letters*,
1525 41(23): 8421-8428.
- 1526 Suzuki, T. and Ishii, M., 2011. Regional distribution of sea level changes resulting from
1527 enhanced greenhouse warming in the Model for Interdisciplinary Research on
1528 Climate version 3.2. *Geophysical Research Letters*, 38: L02601.
- 1529 Swingedouw, D., Rodehacke, C.B., Behrens, E., Menary, M., Olsen, S.M., Gao, Y.Q.,
1530 Mikolajewicz, U., Mignot, J. and Biastoch, A., 2013. Decadal fingerprints of freshwater
1531 discharge around Greenland in a multi-model ensemble. *Climate Dynamics*, 41(3-4):
1532 695-720.
- 1533 Thompson, D.W.J., Solomon, S., Kushner, P.J., England, M.H., Grise, K.M. and Karoly, D.J.,
1534 2011. Signatures of the Antarctic ozone hole in Southern Hemisphere surface
1535 climate change. *Nature Geoscience*, 4(11): 741-749.
- 1536 van Angelen, J.H., van den Broeke, M.R., Wouters, B. and Lenaerts, J.T.M., 2014.
1537 Contemporary (1960-2012) evolution of the climate and surface mass balance of the
1538 Greenland Ice Sheet. *Surveys in Geophysics*, 35(5): 1155-1174.
- 1539 Vaughan, D.G., Comiso, J.C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T.,
1540 Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K. and Zhang, T., 2013. Observations:
1541 Cryosphere. In: T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung,
1542 A. Nauels, Y. Xia, V. Bex and P.M. Midgley (Editors), *Climate Change 2013: The
1543 Physical Science Basis. Contribution of Working Group I to the Fifth Assessment
1544 Report of the Intergovernmental Panel on Climate Change*. Cambridge University
1545 Press, Cambridge, UK, pp. 317-382.
- 1546 Velicogna, I., Sutterley, T.C. and van den Broeke, M.R., 2014. Regional acceleration in ice
1547 mass loss from Greenland and Antarctica using GRACE time-variable gravity data.
1548 *Geophysical Research Letters*, 41: 8130-8137.
- 1549 Visser, H., Dangendorf, S. and Petersen, A.C., 2015. A review of trend models applied to sea
1550 level data with reference to the "acceleration-deceleration debate". *Journal of
1551 Geophysical Research: Oceans*, 120.
- 1552 von Schuckmann, K., Sallée, J.B., Chambers, D., Le Traon, P.Y., Cabanes, C., Gaillard, F.,
1553 Speich, S. and Hamon, M., 2014. Consistency of the current global ocean observing
1554 systems from an Argo perspective. *Ocean Sci.*, 10(3): 547-557.
- 1555 Watson, C.S., White, N.J., Church, J.A., King, M.A., Burgette, R.J. and Legresy, B., 2015.
1556 Unabated global mean sea-level rise over the satellite altimeter era. *Nature Climate
1557 Change*.
- 1558 Wenzel, M. and Schroter, J., 2014. Global and regional sea level change during the 20th
1559 century. *Journal of Geophysical Research-Oceans*, 119(11): 7493-7508.
- 1560 Wong, P.P., Losada, I.J., Gattuso, J.-P., Hinkel, J., Khattabi, A., McInnes, K.L., Saito, Y. and
1561 Sallenger, A., 2014. Coastal systems and low-lying areas. In: C.B. Field, V.R. Barros,
1562 D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O.
1563 Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R.
1564 Mastrandrea and L.L. White (Editors), *Climate Change 2014: Impacts, Adaptation,*

- 1565 and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working
1566 Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate
1567 Change. Cambridge University Press, Cambridge, U.K., pp. 361-409.
- 1568 Wouters, B., Bamber, J.L., van den Broeke, M.R., Lenaerts, J.T.M. and Sasgen, I., 2013. Limits
1569 in detecting acceleration of ice sheet mass loss due to climate variability. *Nature*
1570 *Geoscience*, 6(8): 613-616.
- 1571 Wright, A.P., Le Brocq, A.M., Cornford, S.L., Bingham, R.G., Corr, H.F.J., Ferraccioli, F., Jordan,
1572 T.A., Payne, A.J., Ripplin, D.M., Ross, N. and Siegert, M.J., 2014. Sensitivity of the
1573 Weddell Sea sector ice streams to sub-shelf melting and surface accumulation.
1574 *Cryosphere*, 8(6): 2119-2134.
- 1575 Wu, X. and Heflin, M.B., 2015. A global assessment of accelerations in surface mass
1576 transport. *Geophysical Research Letters*, 42: 6716-6723.
- 1577 Wunsch, C. and Heimbach, P., 2014. Bidecadal thermal changes in the abyssal ocean.
1578 *Journal of Physical Oceanography*, 44(8): 2013-2030.
- 1579 Yi, S., Sun, W., Heki, K. and Qian, A., 2015. An increase in the rate of global mean sea level
1580 rise since 2010. *Geophysical Research Letters*, 42: 3998-4006.
- 1581 Yin, J.J., 2012. Century to multi-century sea level rise projections from CMIP5 models.
1582 *Geophysical Research Letters*, 39.
- 1583 Yin, J.J., Schlesinger, M.E. and Stouffer, R.J., 2009. Model projections of rapid sea-level rise
1584 on the northeast coast of the United States. *Nature Geoscience*, 2(4): 262-266.
- 1585 Zhang, X.B. and Church, J.A., 2012. Sea level trends, interannual and decadal variability in
1586 the Pacific Ocean. *Geophysical Research Letters*, 39.
- 1587 Zwally, H.J., Abdalati, W., Herring, T., Larson, K., Saba, J. and Steffen, K., 2002. Surface Melt-
1588 Induced Acceleration of Greenland Ice-Sheet Flow. *Science*, 297: 218-222.
- 1589




Figure 1. Thermosteric sea level change (m) with respect to 2005. CMIP5 multimodel mean (red) $\pm 2\sigma$ (light grey), realization mean (blue), individual realizations (dark grey). Numbers of models/realizations in brackets. Observations (black lines): Domingues (solid + uncertainties), Levitus (dashes), Ishii (dots). **(a)** Internal variability, **(b)** natural forcing only, **(c)** greenhouse gas forcing only, **(d)** aerosol forcing only, **(e)** anthropogenic forcing only, and **(f)** all forcings combined. (From Slangen et al., 2104)




Figure 2. Net change in volume above flotation in the WAIS over the course of the combined experiment in of both atmospheric and oceanic forcing derived from regional models that were forced, in turn, using climate models for the A1B and E1 (strong mitigation) scenarios. Only the Amundsen Sea Embayment experiences a net loss (ΔV) in all of the combined experiments. Nonetheless, the result is a net loss over West Antarctica as a whole. Note that Thwaites glacier does not retreat in the combined anomaly experiments (which use the synthetic accumulation), and the ASE could contribute an extra $9 \times 10^3 \text{ km}^3$ loss by 2100 and $40 \times 10^3 \text{ km}^3$ by 2200. (From Cornford et al. 2015)




Figure 3. Observed sea level change trends (mm/year) from satellite altimetry between 1993–2012. The hatching indicates trends that are significant (at the 5 % level) with respect to at least 2/3 of CMIP5 pre-industrial control simulations.




Figure 4. Multimodel ensemble median time-of-emergence (ToE) for regional sea-level change under RCP8.5. Different change signals are used: **a**, ocean dynamic topography change (dynamic sea-level); **b**, ocean climate change (dynamic sea-level change plus global mean thermosteric sea-level change); **c**, relative sea-level change (total sea-level change). Warm (cold) colours represent rising (falling) sea level; light grey areas have no emergence before 2080; deep grey colour means no agreement among models; white colour means no data coverage or over land. (From Lyu et al. 2014.)




Figure 5. The cumulative fraction of the total area with the emergence of change signals before the given time from the multimodel ensemble median patterns. Dynamic sea level (DSL, blue), ocean climate change (dynamic sea level (DSL) plus global mean thermosteric sea level (GMTSL), black), relative sea-level (total sea level, red), surface air temperature (green). Dash-dot lines are for RCP4.5 and solid lines for RCP8.5. (From Lyu et al. 2014)




Figure 6. (a) CMIP5 model ensemble mean and (b) standard deviation of the forced patterns of ocean dynamic topography change ($\text{m}/^\circ\text{C}$) calculated using ocean volume mean temperature ($^\circ\text{C}$) as predictor for the historical + RCP4.5 simulations between 1993–2099. (From Bilboa et al. 2015)