

# *The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate*

Article

Published Version

Creative Commons: Attribution 3.0 (CC-BY)

Open access

Turnock, S. T., Butt, E. W., Richardson, T. B., Mann, G. W., Reddington, C. L., Forster, P. M., Haywood, J., Crippa, M., Janssens-Maenhout, G., Johnson, C. E., Bellouin, N. ORCID: <https://orcid.org/0000-0003-2109-9559>, Carslaw, K. S. and Spracklen, D. V. (2016) The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate. *Environmental Research Letters*, 11 (2). 024010. ISSN 1748-9326 doi: <https://doi.org/10.1088/1748-9326/11/2/024010> Available at <https://centaur.reading.ac.uk/62044/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1088/1748-9326/11/2/024010>

To link to this article DOI: <http://dx.doi.org/10.1088/1748-9326/11/2/024010>

Publisher: Institute of Physics

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 024010

(<http://iopscience.iop.org/1748-9326/11/2/024010>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.225.109.63

This content was downloaded on 06/04/2016 at 11:31

Please note that [terms and conditions apply](#).

## Environmental Research Letters



## LETTER

## The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate

## OPEN ACCESS

## RECEIVED

18 September 2015

## REVISED

23 January 2016

## ACCEPTED FOR PUBLICATION

25 January 2016

## PUBLISHED

12 February 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



S T Turnock<sup>1</sup>, E W Butt<sup>1</sup>, T B Richardson<sup>1</sup>, G W Mann<sup>1,2</sup>, C L Reddington<sup>1</sup>, P M Forster<sup>1</sup>, J Haywood<sup>3</sup>, M Crippa<sup>4</sup>, G Janssens-Maenhout<sup>4</sup>, C E Johnson<sup>3</sup>, N Bellouin<sup>5</sup>, K S Carslaw<sup>1</sup> and D V Spracklen<sup>1</sup>

<sup>1</sup> Institute of Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK

<sup>2</sup> National Centre for Atmospheric Science, University of Leeds, Leeds, UK

<sup>3</sup> Met Office, Fitzroy Road, Exeter, Devon, UK

<sup>4</sup> European Commission, Joint Research Centre, Institute for Environment and Sustainability, Air and Climate Unit, Via Fermi, 2749, I-21027 Ispra, Italy

<sup>5</sup> Department of Meteorology, University of Reading, Reading, UK

E-mail: [s.t.turnock12@leeds.ac.uk](mailto:s.t.turnock12@leeds.ac.uk)

**Keywords:** air quality, health, climate, aerosols, emissions, mitigation

Supplementary material for this article is available [online](#)

### Abstract

European air quality legislation has reduced emissions of air pollutants across Europe since the 1970s, affecting air quality, human health and regional climate. We used a coupled composition-climate model to simulate the impacts of European air quality legislation and technology measures implemented between 1970 and 2010. We contrast simulations using two emission scenarios; one with actual emissions in 2010 and the other with emissions that would have occurred in 2010 in the absence of technological improvements and end-of-pipe treatment measures in the energy, industrial and road transport sectors. European emissions of sulphur dioxide, black carbon (BC) and organic carbon in 2010 are 53%, 59% and 32% lower respectively compared to emissions that would have occurred in 2010 in the absence of legislative and technology measures. These emission reductions decreased simulated European annual mean concentrations of fine particulate matter (PM<sub>2.5</sub>) by 35%, sulphate by 44%, BC by 56% and particulate organic matter by 23%. The reduction in PM<sub>2.5</sub> concentrations is calculated to have prevented 80 000 (37 000–116 000, at 95% confidence intervals) premature deaths annually across the European Union, resulting in a perceived financial benefit to society of US\$232 billion annually (1.4% of 2010 EU GDP). The reduction in aerosol concentrations due to legislative and technology measures caused a positive change in the aerosol radiative effect at the top of atmosphere, reduced atmospheric absorption and also increased the amount of solar radiation incident at the surface over Europe. We used an energy budget approximation to estimate that these changes in the radiative balance have increased European annual mean surface temperatures and precipitation by  $0.45 \pm 0.11$  °C and by  $13 \pm 0.8$  mm yr<sup>-1</sup> respectively. Our results show that the implementation of European legislation and technological improvements to reduce the emission of air pollutants has improved air quality and human health over Europe, as well as having an unintended impact on the regional radiative balance and climate.

### 1. Introduction

Air pollutants have a detrimental effect on atmospheric visibility (Vautard *et al* 2009, Stjern *et al* 2011), ecosystems (Fowler *et al* 2009) and human health (Seaton *et al* 1995, Brunekreef and Holgate 2002). The World Health Organization (WHO) attributed 3.7

million annual premature deaths worldwide to the effects of ambient air pollution in 2012 (WHO 2014). Air pollutants can either be gas phase species (e.g. ozone) or particulate matter. This study will focus solely on particulate matter air quality. Exposure to fine particulate matter (PM<sub>2.5</sub>) is responsible for ~3.3 million deaths worldwide each year (Cohen *et al* 2005,

Lelieveld *et al* 2015), with in excess of 400 000 premature deaths annually across Europe (EEA 2014). Particulate matter (also known as aerosols) also influences climate by altering the radiation budget at both the top of atmosphere (TOA) and surface (Boucher *et al* 2013). Aerosols scatter and absorb radiation, termed as aerosol radiation interactions (ari), and can also alter the properties of clouds, termed as aerosol cloud interactions (aci) (Boucher *et al* 2013). Both interactions caused a negative radiative forcing of the Earth's radiation budget over the industrial period resulting in a cooling effect on climate (IPCC 2013).

Europe and North America implemented legislation from the 1970s to reduce emissions of air pollutants (UNECE 2004). Emissions of sulphur dioxide (SO<sub>2</sub>) fell by 73% across Europe over the period 1980–2004 and by 70% in the United States from 1980 to 2010 (Vestreng *et al* 2007, US EPA 2011). Consequently, European surface concentrations of sulphate aerosol mass decreased by between 65% and 78% over the period 1980–2010 (Tørseth *et al* 2012, Turnock *et al* 2015). In addition to the beneficial health effects from improved air quality, such reductions in particulate matter will also have changed the regional radiative balance and climate (Turnock *et al* 2015).

The benefits to human health from reductions in anthropogenic emissions have been highlighted in a number of studies (Anenberg *et al* 2011, 2012, Shindell *et al* 2012). Reducing PM<sub>2.5</sub> concentrations across more than 20 European cities to 15 µg m<sup>-3</sup> was calculated to reduce the number of annual premature deaths by nearly 17 000 (Boldo *et al* 2006), with larger PM<sub>2.5</sub> reductions to the WHO annual mean guideline value of 10 µg m<sup>-3</sup> estimated to avoid 19 000 premature deaths (Pascal *et al* 2013). Considering specific emission reductions from power plants over the United States, Levy *et al* (2007) calculated that 17 000–21 000 annual premature deaths could be avoided. These studies highlight the large potential benefit to human health that can be achieved from reducing the emissions of air pollutants.

Policies that reduce anthropogenic emissions of aerosols may cause inadvertent impacts on climate (Andreae *et al* 2005, Arneth *et al* 2009, Fiore *et al* 2012, 2015, von Schneidmesser *et al* 2015). The air quality and climate impact of reductions in particulate matter depends on the contribution from purely scattering (sulphate) and more absorbing (black carbon (BC)) components (Shindell and Faluvegi 2009), as well as the complex interactions between aerosols and clouds.

Increased surface temperatures over Europe since the 1980s have been attributed to a combination of reductions in aerosols and increases in greenhouse gases (Ruckstuhl *et al* 2008, Philipona *et al* 2009). Increased surface radiation over Europe (solar brightening) has been observed since the 1980s across the continent (Wild 2009 and references within). This

solar brightening was simulated in all of the Fifth Coupled Model Intercomparison Project (CMIP5) models from 1990 to 2005 and was attributed to a reduction in aerosol optical depth caused by decreases in sulphate aerosols (Cherian *et al* 2014). Changes to aerosols over Europe between the 1980s and 2000s are estimated to have led to a positive change in the TOA aerosol radiative effect (RE) of 1–2 W m<sup>-2</sup> and increased surface radiation by 2 W m<sup>-2</sup> (Marriner *et al* 2007, Pozzoli *et al* 2011). The associated changes in cloud albedo are estimated to have caused a positive change in the RE of 1.5 W m<sup>-2</sup> over central Europe from the late 1980s to late 1990s (Krüger and Graßl 2002). In the United States, a reduction in anthropogenic aerosol precursor emissions between 1990 and 2010 coincided with RE<sub>ari</sub> and RE<sub>aci</sub> decreasing by 0.8 W m<sup>-2</sup> and 1.0 W m<sup>-2</sup> respectively (Leibensperger *et al* 2012a). The observed rapid warming over this period of 0.5 °C was potentially attributed in part to a reduction in anthropogenic aerosols (Leibensperger *et al* 2012b). Gan *et al* (2014) analysed the effect of air pollution controls in the US and found that an increase in the surface shortwave radiation occurred over the same time period as negative trends in AOD and PM<sub>2.5</sub>.

Previous studies have assessed the impact of changes in historical (Colette *et al* 2011, Koch *et al* 2011, Turnock *et al* 2015) and future emissions (Kloster *et al* 2008, Unger *et al* 2008, Colette *et al* 2013) on air quality and climate. Changes in historical emissions have resulted from the combined effects of legislation, abatement technology, economic situation and makeup of the national energy supply (Vestreng *et al* 2007). However, it is not known what fraction of the changes in historical emissions and associated impacts on air quality, climate and human health can be attributed to the implementation of legislation and technological developments. Such studies are needed to understand the efficacy of historical policy. In recent work, Crippa *et al* (2015) studied the impact of historical European air pollution mitigation measures on emissions of SO<sub>2</sub>, BC and organic carbon (OC). They found that emissions of SO<sub>2</sub> and PM<sub>2.5</sub> in 2010 would have been substantially higher over Europe if technology improvements and end of pipe abatement measures had not been implemented since 1970.

Here we use emissions scenarios from Crippa *et al* (2015), combined with a global composition climate model to assess the effect of historical legislative and technology measures on particulate matter air quality, climate and human health. The effect of these emission changes on ozone concentrations will be the subject of a future study. The achievements from past policy initiatives are highlighted and will aid in the design of future emission mitigation measures to improve air quality and climate across other regions of the globe.

## 2. Methods

### 2.1. HadGEM3-UKCA composition-climate model

We used the composition-climate model HadGEM3-UKCA, as described in Turnock *et al* (2015), to simulate surface aerosol mass concentrations as well as the impact on aerosol REs. HadGEM3-UKCA is similar in complexity to other models that have participated in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP, (Lamarque *et al* 2013)) and the AeroCom initiative (Mann *et al* 2014). Simulations had a horizontal resolution of  $1.875^\circ$  by  $1.275^\circ$  ( $\sim 140$  km at mid latitudes). Our previous work has demonstrated that HadGEM3-UKCA is able to capture the regional changes in annual mean surface aerosol mass concentrations and surface solar radiation observed over Europe in response to changes in anthropogenic emissions during the period 1980–2010 (Turnock *et al* 2015). Our simulations therefore capture regional changes in  $PM_{2.5}$  (at the scale of  $\sim 100$  km) and are suitable for isolating the effects of policy driven changes in regional aerosol precursor emissions. Our model underestimates  $PM_{2.5}$  concentrations in urban areas; we estimate the impact of this in section 2.3. Further details on the model description and experimental set-up are in supplementary methods section 1.

### 2.2. Anthropogenic emission scenarios

Monthly mean anthropogenic emissions of  $CO$ ,  $SO_2$ ,  $NO_x$ , NMVOC, OC and BC are based on version v4.3 of the Emission Database for Global Atmospheric Research (EDGAR) (EC-JRC/PBL 2015). The emissions used in this study are described in Crippa *et al* (2015) and are summarised below. Emission fields are provided at  $0.5^\circ$  by  $0.5^\circ$  resolution for the energy, industry, transport, agriculture, residential, waste and other sources sectors (<http://edgar.jrc.ec.europa.eu/pegasos>).

We compare two simulations with different anthropogenic emissions. All other factors in the model are the same. The reference simulation (REF) uses actual 2010 emissions. The NO-AQ simulation uses anthropogenic emissions that would have occurred in 2010 if measures to reduce air pollutant emissions between 1970 and 2010 did not occur. The NO-AQ scenario is based on 2010 economic activity data but assumes 1970 emission factors to simulate the stagnation of technological improvements in the energy, industry and road transport sectors. It also assumes that no further end-of-pipe abatement measures (e.g., particle filter traps, fuel desulphurisation) occurred in the energy and road transportation sectors after 1970 (Crippa *et al* 2015). The NO-AQ scenario does however account for changes in the makeup of the national energy supply (e.g. coal to gas) that occurred without specific legislation. These assumptions

alter emissions globally, since stagnation of technology improvements are applied globally and some European emission standards are used outside Europe (Crippa *et al* 2016). However, we focus on the impact of these measures over Europe (defined here as the domain  $10^\circ W$ – $36^\circ E$  and  $36^\circ N$ – $61^\circ N$ ).


Figures 1(a)–(c) shows the spatial distribution of the difference in emissions between the two scenarios (REF minus NO-AQ). This difference represents the emissions that were averted in 2010 due to the implementation of technology improvements and end-of-pipe abatement measures. The largest difference in emissions occurs over central and eastern Europe and the UK. Total European  $SO_2$ , BC and OC emissions over all sectors are reduced, compared to what would have occurred, by  $9.4$  Tg  $SO_2$   $yr^{-1}$  (53%),  $0.4$  Tg BC  $yr^{-1}$  (59%) and  $0.3$  Tg OC  $yr^{-1}$  (32%) (figure 1(d)). Reductions in  $SO_2$  emissions are dominated by changes in the energy sector, whereas reductions in BC and OC are dominated by changes in the transport sector (table S1).

Figure 1(d) compares total European emissions in the NO-AQ scenario against the actual 2010 (REF) and 1970 emissions from the EDGAR and MACCity inventories. The emissions of  $SO_2$  and OC in both the 2010 REF and NO-AQ scenarios are lower than emissions in 1970. In contrast, BC emissions from the 2010 NO-AQ scenario are larger than in 1970 but are lower in the 2010 REF scenario. This suggests that in the absence of technological improvements and end-of-pipe abatement measures, emissions of  $SO_2$  and OC would still have declined substantially due to changes in the makeup of the energy supply and consumption rates whereas emissions of BC would have been larger than in 1970. This highlights the large impact of legislation and technological improvements on reducing BC emissions compared to  $SO_2$  and OC and that historical changes to anthropogenic emissions are determined by the combined effect of legislation, technology and economic factors.

### 2.3. Human health and economic impacts

Concentration response functions (CRFs) for long-term exposure to  $PM_{2.5}$  have been used here to quantify the adverse health effects in terms of excess premature mortality from ischemic heart disease, stroke, chronic obstructive pulmonary disease, lung cancer and acute lower respiratory infection in infants. Here, the link between  $PM_{2.5}$  concentrations and relative risk of the above diseases is based on the integrated exposure-response functions developed by Burnett *et al* (2014), which also uses the baseline rate of disease and spatial distribution of the population.

The resolution of global aerosol models means that they typically underestimate urban  $PM_{2.5}$  concentrations. To estimate the impact of this underestimation on simulated human health impacts, we complete another calculation where we increase the



PM<sub>2.5</sub> concentrations in European urban areas in both the REF and NO-AQ simulations. We then apply the CRFs to these modified concentrations. We estimate that the underprediction of PM<sub>2.5</sub> in urban areas in our global model (Turnock *et al* 2015) causes a bias in our mortality estimates of 13%–16%, substantially less than the uncertainty in the CRF (see supplementary methods). Our method is outlined fully in supplementary methods section 2.

The impact on human health from poor air quality can also have a perceived economic effect on society. The methods and data within WHO Regional Office for Europe (2015) have been used to calculate the perceived economic benefit to society that result from the health benefits associated with the reduction in particulate matter concentrations due to the implementation of air pollution mitigation measures (see supplementary methods section 3).

#### 2.4. Climate response

Based on the climate response to regional radiative forcing (Shindell and Faluvegi 2009), and the atmospheric energy budget (Andrews *et al* 2010, Richardson *et al* 2015), we estimate the temperature and precipitation response due to changes in the radiative balance


from the implementation of air pollutant (particulate matter) mitigation measures (see supplementary methods section 4).

### 3. Results and discussion

#### 3.1. Effects on air quality

Figure 2 shows the simulated change in surface annual mean PM<sub>2.5</sub> concentrations due to the implementation of European air pollutant emission reduction measures (REF versus NO-AQ simulations). European average concentrations are calculated as the 5 year annual mean ( $\pm$  standard deviation) over the continental land mass within the domain 10°W–36°E and 36°N–61°N. Across Europe, annual mean PM<sub>2.5</sub> concentrations reduce by  $1.75 \pm 0.04 \mu\text{g m}^{-3}$  (35%), sulphate by  $1.1 \pm 0.03 \mu\text{g m}^{-3}$  (44%), BC by  $0.27 \pm 0.01 \mu\text{g m}^{-3}$  (56%) and particulate organic matter (POM) by  $0.36 \pm 0.01 \mu\text{g m}^{-3}$  (23%).

Reductions in PM<sub>2.5</sub> mass concentration of 40%–50% occur across much of central, eastern and parts of southern Europe (figure 2(a)). Sulphate mass concentrations decrease by more than 50% across most of central Europe and south eastern Europe (figure 2(b)), mainly due to the substantial decrease in SO<sub>2</sub>




emissions in these regions (figure 1(a)) that mostly occurs in the energy sector (table S1). The largest absolute reduction in sulphate mass concentrations (of  $>5 \mu\text{g m}^{-3}$ ) occurs in south-eastern Europe (figure S1). This is attributed to large reductions in SO<sub>2</sub> emissions from the energy sector combined with atmospheric conditions that enhance the efficiency of sulphate formation (higher oxidant concentrations and temperatures than northern Europe). Reductions in BC mass concentrations of more than 70% occur across all of western Europe and Scandinavia (figure 2(c)), mainly due emissions reductions over Europe from the road transport sector (table S1). Fractional reductions in POM mass concentrations are typically less than 40% (figure 2(d)), despite a similar absolute reduction compared to BC (figure S1). The smaller fractional change in POM concentrations are due to the larger background concentrations of POM from natural emissions, which are the same in both simulations.

Legislative and technology measures have therefore led to reduced aerosol mass concentrations across Europe, with potential impacts for both regional climate and human health. We compared the change in aerosol concentration calculated due to these air

quality mitigation measures against the change in aerosol concentrations that actually occurred over the period 1970–2009 simulated by Turnock *et al* (2015) using the same model. Using this method we estimate that air pollutant mitigation measures are responsible for 29% of the actual reductions in PM<sub>2.5</sub> that occurred over the same period. Air pollutant mitigation contributes the most to changes in BC concentrations (73%) compared to sulphate (33%) or POM (15%). This difference reflects the different causes behind the changes in anthropogenic emissions and highlights the important effect of enacted European legislation on air quality.

Ammonium nitrate aerosol (absent from our simulations) has increased by 15%–30% over Europe since the 1980s due to the reduction in SO<sub>2</sub> and sulphate aerosols (Fagerli and Aas 2008). The increase in ammonium nitrate aerosol concentrations would partially offset some of our simulated improvements to air quality and health but would not alter the overall net decrease in European mean particulate matter concentrations ( $1.75 \mu\text{g m}^{-3}$ ) due to the larger declines in sulphate (44%), BC (56%) and POM (23%) simulated here.





### 3.2. Health and economic effects

Figure 3 shows the impacts of improvements in regional air quality from air pollutant mitigation measures on public health. The simulated reduction in  $PM_{2.5}$  concentrations due to air pollutant mitigation measures reduces the number of premature deaths across Europe, with the largest numbers prevented across central Europe (Germany and The Netherlands) (figure 3(a)). Air pollutant mitigation measures prevent 3 to 4 premature deaths annually per 10 000 people in central and eastern Europe (figure 3(b)) and 5 to 6 premature deaths annually per 10 000 people in south eastern Europe (Romania and Bulgaria), where the largest reductions in  $PM_{2.5}$  occurred (figures 2(a) and S1(a)).

Across all the European Union (EU), reductions in  $PM_{2.5}$  due to air pollutant mitigation measures prevented 80 250 (37 380–115 840, at 95% confidence intervals of the CRFs) premature deaths annually (individual country totals on figure 3(c) and table S2), which is 19% of the estimated premature deaths due to present-day (2011)  $PM_{2.5}$  concentrations across Europe.

We compared the improved public health due to these air quality mitigation measures against an estimate of the improved public health due to the actual change in  $PM_{2.5}$  concentrations that occurred over the same period. Using the change in simulated  $PM_{2.5}$  concentrations between 1970–74 and 2005–2009 from Turnock *et al* (2015), we estimate that 210 000 annual premature deaths were prevented over this period. Therefore European policies to reduce air pollutants and technological improvements can be considered responsible for ~38% of the total number of premature deaths prevented due to the improved air quality across the EU from 1970 to 2009 (figure 3(c)).

We estimate that the improvement in human health from implementing air pollutant mitigation measures results in a perceived economic benefit to society of US\$232 billion across the EU (individual country totals on figure 3(d) and table S3), 1.4% of the EU's GDP in 2010. In particular, the perceived economic benefits to society for some countries in eastern Europe, such as Hungary and Bulgaria, are relatively more important and represent a greater percentage of their national GDP (>5%). The WHO calculated that

**Table 1.** Changes in the European annual mean all-sky aerosol radiative effects due to air pollutant mitigation measures. Clear-sky and cloudy-sky RE at the TOA and surface between the REF and NO-AQ simulations over the European continental land mass within the domain 10°W–36°E and 36°N–61°N. Values in parentheses are changes over the period 1970–2009 from Turnock *et al* (2015).

	All sky ( $\text{W m}^{-2}$ )	Clear sky ( $\text{W m}^{-2}$ )	Cloudy sky ( $\text{W m}^{-2}$ )
TOA	2.26 (3.1)	1.67 (2.0)	0.59 (1.1)
Surface	3.30	2.70	0.60
Atmosphere	−1.04	−1.03	−0.01

in 2010, premature mortality due to ambient concentrations of particulate matter was costing the EU US\$700 billion, or 4% of the EU's GDP (WHO Regional Office for Europe 2015). We therefore suggest that the perceived economic costs of poor air quality would be 33% greater than they are currently if European legislation and technological improvements had not been implemented.

### 3.3. Effects on climate

Table 1 shows the difference in European annual mean aerosol RE at the TOA and at the surface due to the implementation of the air pollutant emission mitigation measures. The reduction of aerosol concentrations in the atmosphere over Europe (figure 2) causes a positive change in the aerosol RE at the TOA, reduces the atmospheric absorption of radiation (mainly due to reductions in BC) and also increases the shortwave radiation incident at the surface. The annual European mean all-sky aerosol RE at the TOA increases by  $2.3 \pm 0.06 \text{ W m}^{-2}$  with clear-sky TOA radiation increasing by  $1.7 \pm 0.05 \text{ W m}^{-2}$ . The shortwave radiation incident at the surface increases under both all-sky ( $3.3 \pm 0.07 \text{ W m}^{-2}$ ) and clear-sky conditions ( $2.7 \pm 0.05 \text{ W m}^{-2}$ ). Air pollutant mitigation measures have reduced atmospheric absorption of radiation over Europe by  $\sim 1 \text{ W m}^{-2}$  in both all-sky and clear-sky conditions (table 1 and figure 4) due to the large reductions in BC aerosols. Our model shows that air pollution mitigation has resulted in a warming effect on climate compared to what would have occurred, dominated by changes in the direct aerosol RE (clear-sky). This occurs from a combination of warming due to reductions in scattering aerosols (sulphate and POM) and the cooling response from the reduction in absorbing BC aerosols.

We can compare the change in aerosol RE calculated here against an estimate of the RE that actually occurred over the period 1970–2009 (table 1). In transient simulations with historical emissions, Turnock *et al* (2015) estimated that the annual European mean all-sky aerosol RE increased by  $3.1 \text{ W m}^{-2}$  between 1970 and 2009. The larger change in RE over the period 1970–2009 reflects the similar changes in BC but larger reductions in sulphate and POM that occurred in reality. Therefore our model simulations show that

European legislation and technological improvements to reduce air pollutants are responsible for approximately 74% of the change in all-sky TOA aerosol RE from the 1970s to present day.


Over the 20th Century nitrate aerosols (not present within our model) were shown by Bellouin *et al* (2011) to have had a relatively small impact on the negative global aerosol forcing, enhancing it by less than 20%. The calculated changes in aerosol REs in this study will have been overestimated due to the absence of the negative aerosol REs from ammonium nitrate.

Figure 4 shows the spatial pattern of changes to the aerosol RE at the TOA and at the surface. The largest increase in TOA clear-sky aerosol RE occurs over south-east and central Europe (figure 4(b)), corresponding with the largest reduction in sulphate mass (figures 2(b) and S1). The largest reductions in atmospheric absorption (TOA minus surface) occur over The Netherlands and Germany where the largest reductions in BC aerosols occurs (figure 2(c)). The TOA warming response over these areas would have been even larger without the reductions of the absorbing BC aerosols (cooling response).

The all-sky changes are also caused by changes to cloud properties (aci). Air pollution mitigation reduces concentrations of cloud condensation nuclei (CCN) (figure S3(a)) leading to larger, less numerous cloud droplets (figure S3(b)), increasing shortwave all-sky radiation at the TOA and surface (figures 4(a) and (c)). Air pollutant mitigation causes the largest increase in all-sky radiation (both TOA and surface) over coastal areas of western Europe (figures 4(a) and (c)), corresponding with the largest reduction in CCN, cloud droplet number concentrations and larger cloud fractions (figures S3 and S4). Over central Europe, CCN and cloud drop concentrations are larger (shown by larger total number concentrations in figure S2), meaning air pollutant mitigation causes smaller fractional changes and a smaller RE here. The difference between all-sky and clear-sky radiation changes (figure S3(d)) gives an indication of the cloudy sky RE, which is greatest over north west Europe.

Overall reductions in aerosols from the implementation of air pollutant mitigation measures resulted in a positive change in the aerosol RE at the TOA and also increased the amount of solar radiation incident at the surface under both all-sky and clear-sky conditions. The simulated change in European all-sky TOA aerosol RE due to air pollutant mitigation is 2.5 times the change in global mean  $\text{CO}_2$  radiative forcing over the period 1970–2009 (Myhre *et al* 2013), indicating the large regional impact of air pollutant mitigation measures. Such changes to the radiative balance over Europe could have non-negligible effects on the climate, impacting both surface temperature and precipitation patterns.

Using the change in TOA aerosol RE between the two simulations, we estimate that the reduction in



particulate matter from the implementation of air pollutant mitigation measures have increased European mean surface temperatures by  $0.45 \pm 0.11 \text{ }^{\circ}\text{C}$  compared to temperatures that would have occurred without mitigation measures (for methods see section 4 of supplementary methods). This change in European surface temperature is anticipated to be slightly smaller than that presented here due to the absence of ammonium nitrate from the simulations. In addition, the reduced atmospheric absorption of radiation caused by mitigation measures will induce a rapid adjustment to increase European precipitation by  $13 \pm 0.8 \text{ mm yr}^{-1}$ . Future work is needed to contrast the human health and economic costs associated with these changes in climate against those quantified here due to air quality improvements.

#### 4. Conclusions

Air quality legislation and technology improvements implemented over Europe have reduced the emission of air pollutants. Here we used a coupled composition-climate model to assess the impact of these reduced emissions on air quality, human health and the Earth's radiative balance.

We used two simulations to quantify the impact of air pollutant mitigation measures on emissions across the energy, industry and road transport sectors over the period 1970–2010 (Crippa *et al* 2015). One simulation used actual 2010 emissions whereas the other simulation used a 2010 emission scenario in which air pollutant mitigation measures were not implemented, represented by the absence of improvements to end-of-pipe abatement measures and technology between 1970 and 2010. The implementation of these air pollutant mitigation measures is estimated to have reduced European anthropogenic emissions of  $\text{SO}_2$  (53%), BC (59%) and OC (32%) compared to what would have occurred in the absence of legislative and technology measures.

We estimate that this reduction in emissions improved regional air quality over Europe by reducing concentrations of fine particulate matter ( $\text{PM}_{2.5}$  by 35%) and their component masses (sulphate by 44%, BC by 56% and POM by 23%). We estimate that these policy measures and technology changes are responsible for  $\sim 29\%$  of the actual reductions in  $\text{PM}_{2.5}$  that occurred from 1970 to 2009.

The improvement in air quality from the implementation of European air pollutant mitigation measures has benefitted human health by preventing

80 000 premature deaths (37 000–116 000, at 95% confidence intervals of the CRF) annually across Europe. These improvements to health have had a perceived economic benefit to society estimated to be US \$232 billion annually, representing 1.4% of the EU's GDP in 2010.

The spatial resolution of our global model (~140 km) means that our results apply to regional changes in PM<sub>2.5</sub>. For this reason our model typically underestimates PM<sub>2.5</sub> concentrations in urban areas of Europe, leading to a ~15% bias in our premature mortality estimates (based on calculations where we adjust simulated PM<sub>2.5</sub> concentrations in urban areas to match observed concentrations). This error is substantially smaller than the factor 3 uncertainty in estimated premature mortality calculated using the 95% confidence intervals of the CRF. Better understanding of the health impacts of PM<sub>2.5</sub>, in addition to studies using higher resolution models would improve future calculations of the impact of air pollutants on human health.

Changes in fine particulate matter concentrations from measures to reduce air pollutants can have an unintended impact on climate by altering the radiative balance. The reduction in aerosol concentrations has caused a positive change to the all-sky aerosol RE at the TOA ( $2 \text{ W m}^{-2}$ ), reduced the absorption of radiation in the atmosphere ( $-1 \text{ W m}^{-2}$ ) and increased the amount of shortwave radiation incident at the surface ( $3 \text{ W m}^{-2}$ ). Changes are largest over central and eastern Europe for clear sky conditions, indicating that aerosol radiation interactions are dominant here, but for all-sky conditions the changes are largest over western Europe, signifying that aerosol cloud interactions are more important here.

We have estimated that these changes in the radiative balance from air pollutant mitigation measures would have caused European mean surface temperatures to increase by  $\sim 0.45 \pm 0.11 \text{ }^\circ\text{C}$  and precipitation by  $\sim 13 \pm 0.8 \text{ mm yr}^{-1}$  compared to those that would have occurred without mitigation measures.

Implementing measures to reduce emissions of air pollutants has led to an improvement in air quality and reduced the impact on human health over Europe. However, the reduction in aerosol concentrations has also had an unintended warming (positive) response on the Earth's radiative balance. The calculated effect on air quality, climate and human health from European legislation and technological improvements implemented over the last 40 years should be used to help inform the design of future multi-pollutant mitigation strategies.

## Acknowledgments

Steven Turnock would like to thank the Natural Environment Research Council (NERC) and Met Office for providing the funding for this PhD

studentship. This work was also made possible by participation in the EU Framework 7 PEGASOS project (no: 265148). We acknowledge the use of the MONSooN system, a collaborative facility supplied under the Joint Weather and Climate Research Programme, a strategic partnership between the Met Office and NERC. We would like to acknowledge the use of the EDGAR anthropogenic emissions inventory and the scenarios used in this study provided by the Joint Research Centre, Italy. Edward Butt is funded by the United Bank of Carbon.

## References

- Andreae M O, Jones C D and Cox P M 2005 Strong present-day aerosol cooling implies a hot future *Nature* **435** 1187–90
- Andrews T, Forster P M, Boucher O, Bellouin N and Jones A 2010 Precipitation, radiative forcing and global temperature change *Geophys. Res. Lett.* **37** L14701
- Anenberg S C, Talgo K, Arunachalam S, Dolwick P, Jang C and West J J 2011 Impacts of global, regional, and sectoral black carbon emission reductions on surface air quality and human mortality *Atmos. Chem. Phys.* **11** 7253–67
- Anenberg S C *et al* 2012 Global air quality and health co-benefits of mitigating near-term climate change through methane and black carbon emission controls *Environ. Health Perspect.* **120** 831–9
- Arnett A, Unger N, Kulmala M and Andreae M O 2009 Clean the air, heat the planet? *Science* **326** 672–3
- Bellouin N, Rae J, Jones A, Johnson C, Haywood J and Boucher O 2011 Aerosol forcing in the climate model intercomparison project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate *J. Geophys. Res.* **116** D20206
- Boldo E, Medina S, LeTertre A, Hurley F, Mücke H-G, Ballester F, Aguilera I and Eilstein D 2006 Apehis: health impact assessment of long-term exposure to PM(2.5) in 23 European cities *Eur. J. Epidemiology* **21** 449–58
- Boucher O *et al* 2013 Clouds and aerosols *Climate Change 2013: The Physical Science Basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press)
- Brunekreef B and Holgate S T 2002 Air pollution and health *Lancet* **360** 1233–42
- Burnett R T *et al* 2014 An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure *Environ. Health Perspect.* **122** 397–403
- Cherian R, Quaas J, Salzmann M and Wild M 2014 Pollution trends over Europe constrain global aerosol forcing as simulated by climate models *Geophys. Res. Lett.* **41** 2176–81
- Cohen A J *et al* 2005 The global burden of disease due to outdoor air pollution *J. Toxicology Environ. Health A* **68** 1301–7
- Colette A *et al* 2011 Air quality trends in Europe over the past decade: a first multi-model assessment *Atmos. Chem. Phys.* **11** 11657–78
- Colette A *et al* 2013 European atmosphere in 2050, a regional air quality and climate perspective under CMIP5 scenarios *Atmos. Chem. Phys.* **13** 7451–71
- Crippa M, Janssens-Maenhout G, Dentener F, Guizzardi D, Sindelarova K, Muntean M, Dingenen R V and Granier C 2015 Forty years of improvements in European air quality: the role of EU policy-industry interplay *Atmos. Chem. Phys. Discuss.* **15** 20245–85
- Crippa M, Janssens-Maenhout G, Guizzardi D and Galmarini S 2016 Understanding the rapid precipitation response to CO<sub>2</sub> and aerosol forcing on a regional scale *J. Climate* **29** 583–94
- EC-JRC/PBL 2015 *EDGAR version 4.3*. (online) 2015 Available from: <http://edgar.jrc.ec.europa.eu/>

- European Environment Agency (EEA) 2014 *Air Quality in Europe—2014 Report* (<http://eea.europa.eu/publications/air-quality-in-europe-2014>)
- Fagerli H and Aas W 2008 Trends of nitrogen in air and precipitation: model results and observations at EMEP sites in Europe, 1980–2003 *Environ. Pollut.* **154** 448–61
- Fiore A M, Naik V and Leibensperger E M 2015 Air quality and climate connections *J. Air Waste Manage. Assoc.* **65** 645–85
- Fiore A M *et al* 2012 Global air quality and climate *Chem. Soc. Rev.* **41** 6663–83
- Fowler D *et al* 2009 Atmospheric composition change: ecosystems–atmosphere interactions *Atmos. Environ.* **43** 5193–267
- Gan C-M, Pleim J, Mathur R, Hogrefe C, Long C N, Xing J, Roselle S and Wei C 2014 Assessment of the effect of air pollution controls on trends in shortwave radiation over the United States from 1995 through 2010 from multiple observation networks *Atmos. Chem. Phys.* **14** 1701–15
- IPCC 2013 Summary for policymakers *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Kloster S, Dentener F, Feichter J, Raes F, van Aardenne J, Roekner E, Lohmann U, Stier P and Swart R 2008 Influence of future air pollution mitigation strategies on total aerosol radiative forcing *Atmos. Chem. Phys.* **8** 6405
- Koch D *et al* 2011 Coupled aerosol–chemistry–climate twentieth-century transient model investigation: trends in short-lived species and climate responses *J. Clim.* **24** 2693–714
- Krüger O and Graßl H 2002 The indirect aerosol effect over Europe *Geophys. Res. Lett.* **29** 3–6
- Lamarque J-F *et al* 2013 The atmospheric chemistry and climate model intercomparison project (ACCMIP): overview and description of models, simulations and climate diagnostics *Geosci. Model Dev.* **6** 179–206
- Leibensperger E M, Mickley L J, Jacob D J, Chen W-T, Seinfeld J H, Nenes A, Adams P J, Streets D G, Kumar N and Rind D 2012a Climatic effects of 1950–2050 changes in US anthropogenic aerosols: I. Aerosol trends and radiative forcing *Atmos. Chem. Phys.* **12** 3333–48
- Leibensperger E M, Mickley L J, Jacob D J, Chen W-T, Seinfeld J H, Nenes A, Adams P J, Streets D G, Kumar N and Rind D 2012b Climatic effects of 1950–2050 changes in US anthropogenic aerosols: II. Climate response *Atmos. Chem. Phys.* **12** 3349–62
- Lelieveld J, Evans J S, Fnais M, Giannadaki D and Pozzer A 2015 The contribution of outdoor air pollution sources to premature mortality on a global scale *Nature* **525** 367–71
- Levy J I, Wilson A M and Zwack L M 2007 Quantifying the efficiency and equity implications of power plant air pollution control strategies in the United States *Environ. Health Perspect.* **115** 743–50
- Mann G W *et al* 2014 Intercomparison and evaluation of global aerosol microphysical properties among AeroCom models of a range of complexity *Atmos. Chem. Phys.* **14** 4679–713
- Marmar E, Langmann B, Fagerli H and Vestreng V 2007 Direct shortwave radiative forcing of sulfate aerosol over Europe from 1900 to 2000 *J. Geophys. Res.* **112** D23S17
- Myhre G *et al* 2013 Anthropogenic and natural radiative forcing *Climate Change 2013: The Physical Science Basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Pascal M *et al* 2013 Assessing the public health impacts of urban air pollution in 25 European cities: results of the Aphekom project *Sci. Total Environ.* **449** 390–400
- Philipona R, Behrens K and Ruckstuhl C 2009 How declining aerosols and rising greenhouse gases forced rapid warming in Europe since the 1980s *Geophys. Res. Lett.* **36** L02806
- Pozzoli L, Janssens-Maenhout G, Diehl T, Bey I, Schultz M G, Feichter J, Vignati E and Dentener F 2011 Re-analysis of tropospheric sulfate aerosol and ozone for the period 1980–2005 using the aerosol–chemistry–climate model ECHAM5–HAMMOZ *Atmos. Chem. Phys.* **11** 9563–94
- Richardson T, Forster P, Andrews T and Parker D 2015 Understanding the rapid precipitation response to CO<sub>2</sub> and Aerosol forcing on a regional scale *J. Clim.* in press
- Ruckstuhl C *et al* 2008 Aerosol and cloud effects on solar brightening and the recent rapid warming *Geophys. Res. Lett.* **35** L12708
- Seaton A, Godden D, MacNee W and Donaldson K 1995 Particulate air pollution and acute health effects *Lancet* **345** 176–8
- Shindell D and Faluvegi G 2009 Climate response to regional radiative forcing during the twentieth century *Nat. Geosci.* **2** 294–300
- Shindell D T *et al* 2012 Simultaneously mitigating near-term climate change and improving human health and food security *Science* **335** 183–9
- Stjern C W, Stohl A and Kristjánsson J E 2011 Have aerosols affected trends in visibility and precipitation in Europe? *J. Geophys. Res.* **116** D02212
- Tørseth K, Aas W, Breivik K, Fjæraa A M, Fiebig M, Hjellbrekke A G, Lund Myhre C, Solberg S and Yttri K E 2012 Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009 *Atmos. Chem. Phys.* **12** 5447–81
- Turnock S T *et al* 2015 Modelled and observed changes in aerosols and surface solar radiation over Europe between 1960 and 2009 *Atmos. Chem. Phys.* **15** 9477–500
- US Environmental Protection Agency 2011 *Technology Transfer Network Clearinghouse for Inventories and Emission Factors, National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data* ([www.epa.gov/air-emissions-inventories](http://www.epa.gov/air-emissions-inventories)) (accessed 23 July 2015)
- Unger N, Shindell D T, Koch D and Streets D G 2008 Air pollution radiative forcing from specific emissions sectors at 2030 *J. Geophys. Res.* **113** D02306
- United Nations Economic Commission for Europe 2004 *Clearing the Air: 25 years of the Convention on Long-Range Transboundary Air Pollution* ed J Sliggers and W Kakebeeke ([www.unece.org/fileadmin/DAM/env/lrtap/ExecutiveBody/BOOKscreen.pdf](http://www.unece.org/fileadmin/DAM/env/lrtap/ExecutiveBody/BOOKscreen.pdf))
- Vautard R, Yiou P and van Oldenborgh G J 2009 Decline of fog, mist and haze in Europe over the past 30 years *Nat. Geosci.* **2** 115–9
- Vestreng V, Myhre G, Fagerli H, Reis S and Tarrasón L 2007 Twenty-five years of continuous sulphur dioxide emission reduction in Europe *Atmos. Chem. Phys.* **7** 3663–81
- von Schneidmesser E *et al* 2015 Chemistry and the linkages between air quality and climate change *Chem. Rev.* **115** 3856–97
- WHO Regional Office for Europe 2015 *Economic Cost of the Health Impact of Air Pollution in Europe: Clean Air, Health and Wealth* (Copenhagen: WHO Regional Office for Europe) ([www.euro.who.int/en/media-centre/events/events/2015/04/ehp-mid-term-review/publications/economic-cost-of-the-health-impact-of-air-pollution-in-europe](http://www.euro.who.int/en/media-centre/events/events/2015/04/ehp-mid-term-review/publications/economic-cost-of-the-health-impact-of-air-pollution-in-europe))
- Wild M 2009 Global dimming and brightening: a review *J. Geophys. Res.* **114** D00D16
- World Health Organization 2014 *Burden of Disease from Ambient Air Pollution for 2012* ([http://who.int/phe/health\\_topics/outdoorair/databases/AAP\\_BoD\\_results\\_March2014.pdf](http://who.int/phe/health_topics/outdoorair/databases/AAP_BoD_results_March2014.pdf)) (accessed 3 May 2015)