

# *Pigs and cattle in Gaul: the role of Gallic societies in the evolution of husbandry practices*

Article

Accepted Version

Fremondeau, D., Nuviala, P. and Duval, C. (2017) Pigs and cattle in Gaul: the role of Gallic societies in the evolution of husbandry practices. *European Journal of Archaeology*, 20 (3). pp. 494-509. ISSN 1741-2722 doi: <https://doi.org/10.1017/eea.2016.10> Available at <https://centaur.reading.ac.uk/69133/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1017/eea.2016.10>

To link to this article DOI: <http://dx.doi.org/10.1017/eea.2016.10>

Publisher: Maney

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# Pigs and Cattle in Gaul: The Role of Gallic Societies in the Evolution of Husbandry Practices

DELPHINE FRÉMONDEAU<sup>1,2</sup>, PAULINE NUVALA<sup>3</sup> AND COLIN DUVAL<sup>1,4</sup>

<sup>1</sup>UMR 7209, *Muséum national d'Histoire naturelle, Paris, France*

<sup>2</sup>*Center for Archaeological Sciences, KU Leuven, Belgium*

<sup>3</sup>UMR 6298 *ArTeHis, Université de Bourgogne, Dijon, France*

<sup>4</sup>UMR 7324 *CITERES, Université François-Rabelais, Tours*

In this article we present a comparative study of pig and cattle morphologies, and stable isotope analysis relating to pig demographic management at Levroux Les Arènes (Indre, France), to evaluate changes in husbandry practices between the Iron Age and the Roman period in Gallic societies. Results indicate the establishment of new production and distribution structures, probably before the second century BC, along with the implementation of a specific size/weight selection for the specialized production of pork. Pig and cattle size evolves progressively from the end of the third century BC. These changes are likely to be the result of an internal evolution within Gallic societies, based on local herds, but possibly they are a response to a broader changing economic climate. Within the Western Roman Empire, each province, and Italy, follows its own evolutionary pattern, which also differs between pig and cattle, suggesting that each region adapted its husbandry strategies according to its agro-pastoral characteristics, capacities, or ambitions.

Keywords: cattle, pig, morphometry, stable isotope analysis, Iron Age and Roman husbandry, Gaul

## INTRODUCTION

The rise of the Roman Empire has long been invoked to explain changes observed in husbandry practices, most notably in the evolution of domestic animal growth and morphology, in Gaul and elsewhere (e.g. Brunaux & Méniel, 1983; Audoin- Rouzeau, 1991, 1995; Méniel, 1996). These changes, believed to coincide with the Roman conquest (first century BC to first century AD), were considered to be a direct manifestation of the process of Rome assuming power over the provinces. However, the hypotheses regarding the chronology of events and their triggers are being reassessed. The economic, and especially the agricultural, changes affecting Gaul actually occurred earlier, most probably from the third century BC onwards, in response to a developing, more dynamic exchange network (e.g. Woolf, 2000, 2001; Keay & Terrenato, 2001; Le Roux, 2004; Rieckhoff, 2006 and subsequent issues of *Bibracte* volume 12<sup>1</sup>; Buchsenschutz, 2007, 2015; Ouzoulias & Tranoy, 2010; Reddé et al., 2011). This new line of enquiry has increasingly led researchers not only to reconsider

---

<sup>1</sup> Rieckhoff, 2006 is the first issue of volume 12 *Celtes et Gaulois, l'Archéologie face à l'Histoire* of the *Bibracte* collection. This volume consists of six publications published from 2006 to 2010 (in ascending order: Vitali, 2006; Szabó, 2006; Haselgrove, 2006; Paunier 2006; Goudineau, et al., 2010), each presenting the proceedings of one of the six international Round Tables organized in 2005–2006 and aimed at synthesising the last twenty years of scientific knowledge obtained about protohistoric Celts.

the chronological and geographical origins and mechanisms of change, but also to reassess the relative level of Roman influence over this change (e.g. Malrain & Blancquaert, 2016). Zooarchaeological research in this domain has likewise advanced, addressing issues such as herd management strategies and the organization of meat supply and their impact on domestic animals (Lepetz, 1995; Forest & Rodet-Belarbi, 1997, 2002; Lepetz & Matterné, 2003; Schlumbaum, et al., 2003, 2006; Forest, 2008). The growing quantity of available data, development of new methods, renewal of archaeological approaches, and comparison of results from different studies now permits a more accurate understanding of the agro-pastoral practices and ties that existed between Italy and the various provinces of the Western Roman Empire between the Iron Age and the Roman period.

This comparative study aims to throw light on the socio-economic changes occurring between the Late Iron Age and Roman period that may have affected husbandry strategies in Gaul and neighbouring territories. To do so, two domestic animals, characterized by different statuses and modes of exploitation—pig and cattle—are considered. First, the evolution of the size of both species is analysed in Gaul and its neighbouring territories (Germania Magna, Italy, and more eastern provinces such as Germania Inferior and Superior, Rhaetia, Noricum, and Pannonia Superior) between the sixth century BC and the seventh century AD. These wide geographical and chronological frameworks allow us to address both animal growth dynamics and size diversity by geographical region well before, during, and after the Roman period, and enable us to answer questions concerning the origin (indigenous or external) of the morphological changes observed in the livestock. Second, the stable isotope analysis of pigs from Levroux Les Arènes (Indre, France) is presented, providing an overview of husbandry and meat production capacities among the Bituriges Cubi (a Gallic people) at the beginning of the second century BC. The Levroux Les Arènes case study gives further insights into the potential socio-economic mechanisms behind the changes in husbandry strategies— notably those affecting livestock morphology—over the course of the Late Iron Age in Gaul.

## ANALYSIS OF PIG AND CATTLE MORPHOLOGY

### Material and methods

Pig and cattle morphometric analysis was carried out over the territory of Gaul (Figure 1) as one assemblage to document broad growth trends. Early Imperial administrative divisions were then used to address geographical diversity in size. For pigs, the analysis was based on osteometric data (9194 measurements) from 307 Gallic, German, and Italian sites, distributed over 139 parishes in modern France, Belgium, Switzerland, Italy, and the Netherlands (Figure 1A). For cattle, the 14,938 measurements came from 341 sites (representing 168 parishes) in Gaul, Roman Italy, and the provinces adjoining the limes, as well as from Germania Magna, outside the boundaries of the Roman Empire (Figure 1B). Only limb bone length measurements were taken following standards established by von den Driesch (1976).




Figure 1. Distribution maps of the parishes (black dots) that yielded (A) pig and (B) cattle bone measurements (1. Germania Magna, 2. Germania Inferior, 3. Germania Superior, 4. Gallia Belgica, 5. Gallia Lugdunensis, 6. Gallia Aquitania, 7. Gallia Narbonensis, 8. Italia, 9. Rhaetia, 10. Noricum, 11. Pannonia Superior).

In order to identify broad morphological trends, to make the best use of the data, and to ensure that samples are reliable, the Log Size Index (LSI) methodology was applied. This method permits the combined use of measurements taken from different bones of the skeleton and therefore makes it possible to study otherwise inaccessible small assemblages. First developed by G.G. Simpson (1941) and adapted by R.H. Meadow (1999), the LSI method consists of calculating the difference between the decimal logarithms for each measurement taken on archaeological material ( $x$ ) and the corresponding dimensions for an individual or reference group ( $y$ ):  $LSI = \log(x) - \log(y)$ . The reference used for pig LSI calculations corresponds to the averaged measurements from three sows from Gaul discovered in the Région Centre (France), specifically in Neuville-aux-Bois (Loiret), Prasville, and Allonnes (Eure-et-Loir) (see Bayle & Josset, 2012; for detailed information: Duval, 2015). These skeletons are dated to the Early and Middle La Tène period (between 400 and 150 BC). For cattle, a third-century AD bull from the site of Fresnes-lès-Montauban (Pas-de-Calais, France) (Lepetz, 1996) was used as reference (metric data published in Duval et al., 2012). Its age was estimated at 42 months and its wither height 1.35 m.

For the morphological analysis undertaken for Gaul, results are displayed in diagrams showing both log size mean values per sites—or groups of sites when data were too scarce—(Figure 2A and 2B, individual circles), and best fit polynomial curves reflecting the general trends in size variation through time. These trends were statistically tested, grouping log ratios by century. As the normality and the homoscedasticity of these newly constituted samples were not always proved (as revealed by Shapiro-Wilk and Levene tests respectively), the adopted statistical approach was to perform a Kruskal-Wallis test, followed by post-hoc comparisons using a Mann-Whitney pairwise test with Bonferroni correction. All results are given in the Supplementary Material.

As far as comparisons between Early Imperial administrative divisions are concerned (Figures 4 and 5), only the best fit polynomial curves are displayed to keep the figures easily readable. For each curve, the number of data points and the strength of the relationship ( $R^2$  values, ranging between 0.204 and 0.782 for pigs, and between 0.392 and 0.869 for cattle) are directly shown on the

diagrams. The detail of site means and standard deviations of log ratios is provided both for Figure 2A and 2B and for Figures 3 and 4 in the Supplementary Material.




Figure 2. Evolution of (A) pig and (B) cattle bone length measurements from the sixth century BC to the seventh century AD in Gaul. The circles on the two plots correspond to site mean values. The best fit polynomial curves summarize the general trend in size variation (the strength of the relationship [R<sup>2</sup>] and the number of samples used [n] are indicated for both curves). Animal silhouettes were drawn by Michel Coutureau, in collaboration with Vianney Forest (INRAP), ©1996 ArcheoZoo.org.

### An early and continuous evolution of pig and cattle morphology

The osteometric analysis reveals that, after a conspicuous drop in pig size from the start of the period considered, pig bone lengths increase from the very end of the third century BC (Figure 2A). This growth appears progressive until the first century AD, when pig size rapidly increases (significant pairwise comparisons between the first century BC and the second century AD, see Supplementary Material), and exhibits an increasing diversity, documented by sites with higher or much smaller length mean values (initially in the first century BC, then in second century AD; Ansari-Bradley unilateral tests, p values = 1). This trend is at its most prominent during the second century AD. That century appears to be a turning point, as a relatively stagnant phase is observed afterwards, which may correspond to reaching a morphological optimum, and is accompanied by a progressive decrease in the range of length variation (Ansari-Bradley unilateral test, p value = 1). From the fourth century onwards, pig bone lengths progressively drop.

Cattle size (Figure 2B) seems to conspicuously decrease between the sixth and the third centuries BC. Thereafter, between the end of the third century and the beginning of the second century BC, bone lengths progressively increase. This growth accelerates during the course of the first century BC (significant pairwise comparisons between the second century BC and the third century AD, see Supplementary Material), which is one century earlier than for pigs. As with pigs, this rapid growth is associated with greater size diversity (initially in the first century BC, then in second and third

centuries AD; Ansari-Bradley unilateral tests,  $p$  values = 1), and is rapidly followed by a drop in values from the start of the fourth century AD; this is slightly earlier than for pigs.




Figure 3. Evolution of pig bone length from the fifth century BC to the seventh century AD in different provinces of the Roman Empire. The best fit polynomial curves summarize the general trend in size variation (the strength of the relationship [ $R^2$ ] and the number of samples used [ $n$ ] for the curve construction are indicated between parenthesis for each administrative region). Animal silhouette was drawn by Michel Coutureau, in collaboration with Vianney Forest (INRAP). See online version for the code colour. ©1996 ArcheoZoo.org.

In light of these results, it appears that domestic animal size starts to increase well before the Roman conquest and the subsequent rise of the Roman model in Gaul. Gallic herders would therefore appear to have been instrumental in the evolution of the size of pigs and cattle within a broader La Tène context of demographic, urban, and economic growth (Buchenschutz, 2007). Nevertheless, the Roman period saw further change for these two species. A slight increase in size growth, coupled with greater size diversity, is noticeable from the first century AD onwards. The combination of these two phenomena may indicate the strengthening of economic frameworks, the intensification of exchanges and influences, and the emergence of more numerous animal types. This latter may result from the complexification of social and administrative hierarchies, the expression of a stronger regionalism in response to the penetration of the Roman economic model, the import of animals of various types, and/or the reshaping of agro-pastoral systems within the territory of Gaul.




Figure 4. Evolution of cattle bone length from the fifth century BC to the fifth century AD in different provinces of the Roman Empire and in unoccupied Germany. The best fit polynomial curves summarize the general trend in size variation (the strength of the relationship [R<sup>2</sup>] and the number of samples used [n] for the curve construction are indicated between parenthesis for each administrative region). Animal silhouette was drawn by Michel Coutureau in collaboration with Vianney Forest (INRAP). See online version for the code colour. ©1996 ArcheoZoo.org.

To refine on our perception of these broad morphological trends, the data have been split between the different provinces of Gaul and compared with those from other territories of the Western Roman Empire.

#### Regional diversity in livestock sizes

Pig size evolution is compared in six different regions (Figure 3). The largest sizes are observed in the provinces of Gallia Belgica and Lugdunensis. It is notable that animals from both provinces are characterized by similar sizes that change in relatively parallel patterns, despite a slight time-lag. In Germania, the size increase follows a similar pattern, but pigs do not become as tall as in Gallia Belgica and Lugdunensis. In Gallia Narbonensis and Aquitania, pigs seem to be taller than in other regions during the La Tène period. However, in these two provinces, pig size increases relatively little through time and, consequently, pigs become comparatively smaller. These first five curves show a geographical progression (from north to south) in diachronic size evolution trends, which is also shown by the Italian curve. In Roman Italy, pigs display the smallest size and are also characterized by a less intense growth than in Gallia Belgica and Lugdunensis. These results strongly challenge the previously held ideas that Italy was the epicentre from which changes in pig husbandry spread. Indeed, Italy cannot have initiated, at least from a zootechnical or genetic point of view, the growth of pigs observed in Gaul.

The data present a different picture for cattle (Figure 4). Among all the cattle from the corpus (spread over nine different regions), the largest are those from Gallia Narbonensis and Italy. In Gallia Belgica, Gallia Lugdunensis, and Aquitania, as well as in the eastern provinces and Germania Superior, cattle size follows a similar increasing pattern and finally catches up with the Mediterranean size during the course of the third century AD. In Germania Inferior, although an increase in size is clearly visible, cattle appear quite small. Germania Magna stands out, with cattle



characterized by a rather stable size through time, as if this region had not been culturally or economically affected by the new economic networks during the time-span considered.

By paring down the geographical subdivisions and taking other territories of the Roman Empire into account, a greater diversity in livestock morphology and change dynamics emerges within the Roman Empire and on its fringes. This diversity exists before the Roman conquest and changes through time according to the region considered. Furthermore, pigs and cattle follow different trends, likely to be the result of different modes of exploitation and economic stakes. The existence of such diversity suggests that each territory, including Roman Italy, was driven by an overarching changing economic context and underwent internal mutations related to its own production capacity, know-how, and the livestock available locally. Moreover such a trend is also visible at more local scales, at the level of civitates, valleys, or even individual sites (MacKinnon, 2010; Duval et al., 2012, 2013; Duval, 2015; Nuviala, 2014, 2015, 2016), reflecting local environmental and cultural characteristics, the agro-pastoral (crop/livestock) balance, or supply strategies.




Figure 5. (A) Pig kill-off pattern from the early second century BC at Levroux Les Arènes, established following Horard-Herbin’s 1997 methodology based on the percentage of the corrected number of teeth (for further explanation, see Frémondeau et al., 2015). (B) The sequentially sampled specimens that participate in the main slaughter peak—the horizontal bar represents the estimated age at death of each specimen (following Horard-Herbin’s 1997 method). In the following section, this local scale is used to highlight the mechanisms at the basis of the diversity in livestock morphology and change dynamics in the case-study of Levroux Les Arènes. A stable isotope analysis carried out on pigs from this Iron Age village provides further insights into husbandry strategies implemented to promote meat production well before the Roman conquest.

#### COMPLEX ORGANIZATION OF MEAT PRODUCTION IN GAUL IN THE SECOND CENTURY BC

The Late Iron Age industrial settlement of Levroux Les Arènes was established at the beginning of the second century BC within the territory of the Bituriges Cubi in central France. It remained occupied until around 80 BC, after which the village was largely abandoned (Buchsenschutz et al., 2000). Levroux Les Arènes played an important economic role as a specialized production centre, and as an intermediary between local rural sites, for metalworking. It was also a trading post

supplying networks at a regional scale and further afield (Buchsenschutz et al., 2000; Berranger & Fluzin, 2009). More than 99 per cent of the identified faunal remains from the excavation are ascribed to domestic species, with the pig predominating. All the main parts of the pig skeleton are present in the faunal assemblage, which indicates that pigs were slaughtered within the village. However, some meat-bearing bones (femora, ribs, and vertebra) are under-represented, suggesting that pork was also exported and consumed outside the village. Interestingly, animals slaughtered at 20–24 months are over-represented among the dental remains (Figure 5A; Horard-Herbin, 1997; Frémondeau et al., 2015). In the particular economic context of Levroux Les Arènes, one may wonder what the reasons for such an emphasis on this particular age group are. In an attempt to document the scale of the Bituriges Cubi pig husbandry, and in order to identify the demographic management strategies of pig herds, a stable isotope study was conducted on the faunal assemblage from this site (Frémondeau, 2012), the results of which have been partly published elsewhere (Frémondeau et al., 2013, 2015; Duval et al., 2016).

The  $\delta^{18}\text{O}$  sequences measured in fifteen lower incisors are used to assess seasonality in pig birth and slaughter (e.g. Balasse et al., 2003, 2012) and to determine whether the standardized demographic management resulted from a strong seasonality of birth and slaughter, or from an intentional selection of an age class and therefore a slaughter weight; the latter would strongly support the hypothesis of specialized meat production at the site. The carbon and nitrogen stable isotope composition of bone collagen from the main domestic species helps determine pig diet and, from this, the environment they were raised in, assuming that omnivorous pigs reflect feeding from human waste activities in the village (e.g. Ervynck et al., 2007; Hammond & O'Connor, 2013; Balasse et al., 2013, 2015). In addition, the sequential oxygen and carbon stable isotope analysis of nine male lower canines better characterizes pig diet on a seasonal level (e.g. Balasse, 2002; Frémondeau et al., 2012).

The C and N stable isotope composition of bone collagen reveals that pigs had a herbivorous diet (Frémondeau et al., 2013). Moreover, the combined interpretation of the  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  sequences measured in the canines suggests that pigs were raised under tree cover at least seasonally: in all nine sampled specimens,  $\delta^{13}\text{C}$  values increase when  $\delta^{18}\text{O}$  values decrease and reach a minimum, probably corresponding to autumn and winter (Figure 6; see Supplementary Material; Frémondeau, 2012). This pattern, previously observed in modern Corsican wild pigs living under tree cover (Frémondeau et al., 2012), may correspond to feeding on forest fruits during autumn and winter. These combined pieces of evidence strongly suggest that the pigs slaughtered in Levroux Les Arènes were not raised within the village, but were mainly herded in surrounding farms with herders likely to be exploiting forest resources to feed their livestock.

Results from the  $\delta^{18}\text{O}$  sequential analysis indicate that the births of the fifteen pigs sampled are spread over roughly nine months, indicating that there was no strong seasonality of pig births within Bituriges Cubi farms (Figure 7; Frémondeau et al., 2015). The age at death of these fifteen pigs was estimated from tooth eruption and wear stages of their mandibles following Horard-Herbin's 1997 method. Seven of them display a tooth development stage compatible with slaughter at around the age of two years, which corresponds to the main slaughter peak (Figure 5B). The  $\delta^{18}\text{O}$  results suggest that their births were spread over roughly nine months—even excluding *Sus d. 13* and *18* for which age-at-death estimations are less precise (Figure 7). Therefore, if these seven pigs were actually killed at the same age, this would mean that their slaughter was also spread over nine months. Provided that these specimens are representative of the age class of 20–24 months, the killing of the pigs at Levroux Les Arènes would therefore not have been targeting a specific season of the year, but rather a specific age group or slaughter weight (Frémondeau et al., 2015). This

deliberate and systematic selection of animals approximately two years in age strongly suggests a standardization of meat production. Furthermore, if pork was exported, as suggested by the under-representation of some of the ham-bearing bones, a specialized meat production in Levroux Les Arènes can be proposed (Flad & Hruby, 2005).




Figure 6. Intra-tooth  $\delta^{18}\text{O}$  (open circles) and  $\delta^{13}\text{C}$  (filled circles) variations (B) measured in a male lower canine (A) dated to the early second century BC from Levroux Les Arènes. The arrows indicate the seasonal increase in  $\delta^{13}\text{C}$  values, a pattern suggesting the seasonal consumption of forest fruits in autumn and winter.

The case-study of Levroux Les Arènes thus provides good evidence for a complex organization of meat production. This efficient and organized production system implies the exploitation of a specific environment (the forest) in order to produce a standardized finished product (two-year old pigs) within either a network of farms or in a few specialized farms. Zooarchaeological data, however suggest a specialization in pig husbandry at the scale of the whole civitas (Bayle et al., 2016). Pigs were selected and brought on foot to the settlement (involving regular animal movements and likely necessitating regulation or management). These pigs were killed and their carcasses processed by a limited number of specialists (Horard-Herbin, 1997) and some of the meaty cuts may have been used to make high value-added products (cured pork) that were subsequently put into circulation in the existing trade networks. Therefore, right from the beginning of the occupation of the settlement, the system was well established, indicating an earlier initial development time. In turn, this complex meat production system may have triggered further specialization in animal husbandry strategies, encouraging herders (or making it more profitable for

them) to raise larger animals that would consequently produce larger carcasses. Indeed, pigs killed at Levroux Les Arènes are, on average, among the tallest in Gaul for the period considered (Horard-Herbin, 1997; Duval et al., 2016).




Figure 7. Birth distribution over an annual cycle of fifteen sequentially sampled pig specimens dated to the early second century BC from Levroux Les Arènes. For each specimen the  $x_0/X$  value is the position (in mm) of the measure  $\delta^{18}O$  maximum value over crown height divided by the distance (in mm) over which an annual cycle is recorded (Frémondeau et al., 2015). The specimens likely to be 20–24 months old when killed (i.e. the main slaughter peak) are identified by grey symbols.

## CONCLUSION

In relation to Gaul, changes in domestic animal morphology and management are visible from the second century BC and result most probably from an internal and voluntary impetus, in response to a changing economic context—the Celtic world playing an active role in the then developing market economy (Buchsenschutz, 2004, 2015). The Levroux Les Arènes case-study gives us a glimpse of this new economic context and its impact on husbandry strategies. The stable isotope analysis revealed the existence of a well-developed and complex meat production system—perhaps organized at the wider civitas level (Bailey et al., 2016)—already by the beginning of the second century BC, which implies a development at a much earlier age. This new economic organization may, in turn, have fostered the size increase observed in pigs in the upper north of Gaul from the third century BC onwards. It is therefore from Gallic initiatives and available livestock that changes would have arisen.

Comparing results from different territories in the Western Roman Empire has shown that each province possessed herds with their own characteristics, which followed their own distinctive pattern of evolution. Like neighbouring regions, livestock in Roman Italy underwent size changes, potentially as a result of a more dynamic and open developing market. This heterogeneity of patterns, based on a variety of existing situations from the start of the time-span under study, appears also between species, with pigs and cattle following different evolutionary trends. This could reflect the existence of different species' status depending on the territory considered and the adaptation to the characteristics, strategies, and/or (quality or aesthetic) selection criteria existing in the various regions. In a context of generalized economic growth, each region seems to adapt according to its agro-pastoral characteristics, capacities, or ambitions.

In conclusion, it appears that communities in Gaul (and others in the Roman Empire) may have benefited from the blooming of a favourable economic context. Yet, well before the Roman conquest, they adapted according to their own agricultural characteristics and their own level of engagement in the economic market. The practical organization of husbandry production in western Europe, however, remains unclear over the period considered in this study (sixth century BC to seventh century AD). The pre-eminence of cattle from Roman Italy and pigs from northern Gaul in terms of size, and the great diversity in trends in the evolution of size within the various provinces, tend to cloud our view of the interactions between external influences, local initiatives, and the reflection of more indirect factors such as environmental conditions, or the farming, cultural, and broader economic frameworks within the Roman Empire. New light on these issues may come from increasing the database available for the geographical and chronological range considered in this article and enlarging these frameworks, comparing pig and cattle evolution and taking into account other domestic taxa (sheep, goat, horse, or dog). Finally, further case studies like that undertaken for Levroux Les Arènes, integrating traditional zooarchaeological methods and isotope analysis, are needed to deepen our understanding of inter-site variation in husbandry strategies.

#### ACKNOWLEDGEMENTS

The osteometric analysis of pig and cattle stature evolution would not have been feasible without the help of the numerous zooarchaeologists who agreed to share their data. We heartily thank: Rose-Marie Arbogast, Thierry Argant, Ginette Auxiette, Alessio Bandelli, Anna Baudry, Grégory Bayle, Francesco Boschin, Céline Bemilli, Estelle Bidault, Aurélie Borvon, Cécile Callou, David Cambou, Nadia Cavanhié, Cécile Callou, Paola Catalano, Benoît Clavel, Aurélien Creuzieux, Sabine Deschler-Erb, Jessie Duval, Alexandre Fontaine, Vianney Forest, Sylvain Foucras, Stéphane Frère, Armelle Gardeisen, Chloé Genies, David Germinet, Alexandra Griffon, Maaïke Groot, Jacopo de Grossi Mazzorin, Marie-Pierre Horard-Herbin, Gaëtan Jouanin, Virginie Labonne, Caroline Lachiche, Dominique Lalaï, Jean-Claude Le Blay, Martine Leguilloux, Sébastien Lepetz, Julie Massendari, Patrice Méniel, Elsa Paillot, Fabienne Pigière, Frédéric Poupon, Eric Pucher, Olivier Putelat, Emeline Quilez, Audrey Renaud, Julie Rivière, Opale Robin, Maryline Salin, Barbara Stopp, Antonio Tagliacozzo, Olivier Talluault and Christian Vallet. Stable isotope analyses were carried out at the Service de Spectrométrie de Masse Isotopique du Muséum national d'Histoire naturelle (SSMIM) in Paris, thanks to the financial support of the ERC Starting Grant GA-202881 (SIANHE, directed by M. Balasse), of UMR 7324 CITERES and the Académie d'Agriculture de France, within a PhD funded by the SIANHE Starting Grant. Delphine Frémondeau is currently supported by a post-doctoral fellowship from the Center for Archaeological Sciences (CAS), KU Leuven, Belgium. We thank O. Buchsenschutz for giving us access to the Levroux Les Arènes faunal assemblages. We are also grateful to G. Fahy and two anonymous reviewers whose helpful comments significantly improved the manuscript.

## REFERENCES

- Audoin-Rouzeau, F. 1991. La taille du bœuf domestique en Europe de l'Antiquité aux temps modernes. *Fiches d'ostéologie animale pour l'archéologie*, 2 (40): 2–40.
- Audoin-Rouzeau, F. 1995. La taille des animaux d'élevage à l'époque romaine et leur exportation. In: E. Baratay, ed. *Homme et Animal dans l'Antiquité*. Tours: Presses Universitaires de Tours, pp. 79–101.
- Balasse, M. 2002. Reconstructing Dietary and Environmental History from Enamel Isotopic Analysis: Time Resolution of Intra-tooth Sequential Sampling. *International Journal of Osteoarchaeology*, 12: 155–65.
- Balasse, M., Bałașescu, A., Janzen, A., Ughetto-Monfrin, J., Mirea, P. & Andreescu, R. 2013. Early Herding at Mağura-Boldul lui Moșlvañuș (Early Sixth Millennium BC, Romania): Environments and Seasonality from Stable Isotope Analysis. *European Journal of Archaeology*, 16: 221–46.
- Balasse, M., Bałașescu, A., Tornero, C.,
- Frémonteau, D., Hovsepyan, R., Gillis, R., Ughetto-Monfrin, J. & Popovici, D. 2015. Investigating the Scale of Herding in Chalcolithic Pastoral Communities Settled along the Danube River in the 5th Millennium BC: A Case Study Bordsușani-Popinăand Hârșova-tell (Romania). *Quaternary International*. Available online at: <http://dx.doi.org/10.1016/j.quaint.2015.07.030>.
- Balasse, M., Obein, G., Ughetto-Monfrin, J. & Mainland, I. 2012. Investigating Seasonality and Season of Birth in Past Herds: A Reference Set of Sheep Enamel Stable Oxygen Isotope Ratios. *Archaeometry*, 54: 349–68.
- Balasse, M., Smith, A.B., Ambrose, S.H. & Leigh, S.R. 2003. Determining Sheep Birth Seasonality by Analysis of Tooth Enamel Oxygen Isotope Ratios: The Late Stone Age Site of Kasteelberg (South Africa). *Journal of Archaeological Science*, 30: 205–15.
- Bayle, G. & Josset, D. 2012. Dépôt de truies en silos et fossés dans des sites gaulois de Beauce. *Archéopages*, 35: 32–33.
- Bayle, G., Auxiette, G., Frère, S., Germinet, D., Horard-Herbin, M.-P., Poupon, F., Rivière, J. & Salin, M. 2016. Première synthèse sur l'alimentation carnée dans le sud du Bassin parisien à l'âge du Fer: traditions, particularismes et influences externes. In: F. Malrain & G. Blancquaert, eds. *Évolution des sociétés gauloises du Second âge du Fer, entre mutations internes et influences externes (Revue Archéologique de Picardie, n° spécial 30)*. Senlis: Société des antiquités historiques de Picardie, pp. 565–82.
- Berranger, M. & Fluzin, P. 2009. The Celtic Village of Levroux (Indre, France): The Nature and Organization of the Ironworking Activities (2nd–1st centuries BC). In: O. Buchsenschutz, M.-B. Chardenoux, S. Krausz & M. Vaginay, eds. *L'âge du Fer dans la boucle de la Loire. Les Gaulois sont dans la ville. (Supplément à la Revue archéologique du Centre de la France 35)*. Tours: Fédération pour l'édition de la Revue archéologique du Centre de la France, pp. 19–38.
- Brunaux, J.-L. & Méniel, P. 1983. L'importation du bœuf à la période romaine: premières données. Les fouilles de Gournay-sur-Aronde. *Revue Archéologique de Picardie*, 4: 15–20.

Buchsenschutz, O. 2004. Les Celtes et la formation de l'empire romain. *Annales. Histoire, Sciences sociales*, 59: 337–61.

Buchsenschutz, O. 2007. *Les Celtes*. Paris: Armand Colin.

Buchsenschutz, O. ed. 2015. *L'Europe celtique à l'âge du Fer VIIIe: Ier siècles*. Paris: Presses Universitaires de France.

Buchsenschutz, O., Colin, A., Firmin, G., Fischer, B., Guillaumet, J.-P., Krausz, S., Levéry, M., Marinval, P., Orellana, L. & Pierret, A. 2000. Le village celtique des Arènes à Levroux: synthèses (Supplément à la *Revue Archéologique du Centre de la France* 19). Tours: Fédération pour l'édition de la *Revue archéologique du Centre de la France*.

Duval, C. 2015. Evolution et diversité de la forme du cochon entre la fin de l'âge du Fer et la période moderne en Gaule et en France. Régionalisme, acquisitions zoo-techniques et implications historiques (unpublished PhD dissertation, Université François Rabelais de Tours).

Duval, C., Frémondeau, D., Lepetz, S. & Horard-Herbin, M.-P. 2016. L'élevage du porc: un savoir-faire gaulois? Apport croisé des études isotopique et ostéométrique des os de cochon. In: F. Malrain & G. Blancquaert, eds. *Évolution des sociétés gauloises du Second âge du Fer, entre mutations internes et influences externes* (*Revue Archéologique de Picardie*, n° spécial 30). Senlis: Société des antiquités historiques de Picardie, pp. 583–96.

Duval, C., Horard-Herbin, M.-P. & Lepetz, S. 2013. Morphological Changes in Domestic Cattle in Gaul, from the Second Century BC to the Fifth Century AD: Diversity of Herds in the Seine Valley (France) and Northern Gaul. *Journal of Archaeological Science*, 40: 3977–90.

Duval, C., Lepetz, S. & Horard-Herbin, M.-P. 2012. Diversité des cheptels et diversification des morphotypes bovins dans le tiers nord-ouest des Gaules entre la fin de l'âge du Fer et la période romaine. *Gallia*, 69: 79–114.

Ervynck, A., Lentacker, A., Müldner, G., Richards, M.P. & Dobney, K. 2007. An Investigation into the Transition from Forest Dwelling Pigs to Farm Animals in Medieval Flanders, Belgium. In: U. Albarella, K. Dobney, A. Ervynck & P. Rowley-Conwy, eds. *Pigs and Humans: 10,000 Years of Interaction*. Oxford: Oxford University Press, pp. 170–93.

Flad, R.K. & Hruby, Z.X. 2005. 'Specialized' Production in Archaeological Contexts: Rethinking Specialization, the Social Value of Products, and the Practice of Production. *Archaeological Papers of the American Anthropological Association*, 17: 1–19.

Forest, V. 2008. Équidés de La Tène finale et de la période romaine en Gaule: approche ostéométrique. In: J.-L. Blanchard & L. Izac-Imberd, eds. *L'exploitation agricole dans son environnement à la fin de l'âge du Fer*. Toulouse: Archives d'Écologie Préhistorique, pp. 61–71.

Forest, V. & Rodet-Belarbi, I. 1997. Augmentation du format des bovins en Gaule romaine: problèmes méthodologiques et innovation technique. In: D. Meeks & D. Garcia, eds. *Techniques et économie antiques et médiévales. Le temps de l'innovation*. Paris: Errance, pp. 166–71.

Forest, V. & Rodet-Belarbi, I. 2002. A propos de la corpulence des bovins en France durant les périodes historiques. *Gallia*, 59: 273–306.

Frémondeau, D. 2012. Rythmes saisonniers des élevages porcins archéologiques: propositions pour un protocole d'analyse isotopique (unpublished PhD dissertation, Muséum national d'Histoire naturelle, Paris).

Frémondeau, D., Cucchi, T., Casabianca, F., Ughetto-Monfrin, J., Horard-Herbin, M.-

P. & Balasse, M. 2012. Seasonality of Birth and Diet of Pigs from Stable Isotope Analyses of Tooth Enamel ( $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$ ): A Modern Reference Data Set from Corsica, France. *Journal of Archaeological Science*, 39: 2023–35.

Frémondeau, D., Horard-Herbin, M.-P., Ughetto-Monfrin, J. & Balasse, M. 2013. L'alimentation des troupeaux porcins et la production de viande à Levroux Les Arènes (Indre): une analyse isotopique. In: A. Colin & F. Verdin, eds. *L'âge du Fer en Aquitaine et sur ses marges. Mobilité des hommes, diffusion des idées, circulation des biens dans l'espace européen à l'âge du Fer*. Bordeaux: Fédération Aquitania, pp. 747–52.

Frémondeau, D., Horard-Herbin, M.-P., Buchsenschutz, O., Ughetto-Monfrin, J. & Balasse, M. 2015. Standardized Pork Production at Levroux Les Arènes (France, 2nd c. BC): Evidence from Kill-off Patterns and Birth Seasonality Inferred from Enamel  $\delta^{18}\text{O}$  Analysis. *Journal of Archaeological Sciences: Reports*, 2: 215–26.

Goudineau, C., Guichard, V. & Kaenel, G. 2010. Celtes et Gaulois, l'archéologie face à l'histoire, 6. Colloque de synthèse (Paris, Collège de France, 3–7 juillet 2006) (Bibracte 12/6). Glux-en-Glenne: Centre archéologique européen.

Hammond, C. & O'Connor, T. 2013. Pig Diet in Medieval York: Carbon and Nitrogen Stable Isotopes. *Archaeological and Anthropological Sciences*, 5: 123–27.

Haselgrove, C. 2006. Celtes et Gaulois, l'archéologie face à l'histoire, 4 : les mutations de la fin de l'âge du Fer (Actes de la table ronde de Cambridge, juillet 2005) (Bibracte 12/4). Glux-en-Glenne: Centre archéologique européen.

Horard-Herbin, M.-P. 1997. Le village celtique des Arènes à Levroux. L'élevage et les productions animales dans l'économie de la fin du second âge du Fer (Levroux 4, Supplément à la Revue archéologique du Centre de la France, 12). Tours: Fédération pour l'édition de la Revue archéologique du Centre de la France.

Keay, S. & Terrenato, N. 2001. *Italy and the West: Comparative Issues in Romanization*. Oxford: Oxbow Books.

Lepetz, S. 1995. L'amélioration des races à l'époque gallo-romaine: l'exemple du bœuf. In: E. Baratay, ed. *Homme et Animal dans l'Antiquité*. Tours: Presses Universitaires de Tours, pp. 67–79.

Lepetz, S. 1996. L'animal dans la société gallo-romaine de la France du Nord (Supplément à la Revue Archéologique de Picardie 12). Senlis: Société des antiquités historiques de Picardie.

Lepetz, S. & Matterné, V. 2003. Élevage et agriculture dans le Nord de la Gaule durant l'époque gallo-romaine: une confrontation des données archéologiques et carpologiques. *Revue Archéologique de Picardie*, 1(2): 23–35.

Le Roux, P. 2004, La romanisation en question. *Annales. Histoire, Sciences Sociales*, 59: 287–311.

Mackinnon, M. 2010. Cattle 'Breed' Variation and Improvement in Roman Italy: Connecting the Zooarchaeological and Ancient Textual Evidence. *World Archaeology*, 42: 55–73.

Malrain, F. & Blancquaert, G. eds. 2016. Évolution des sociétés gauloises du Second âge du Fer, entre mutations internes et influences externes (Revue Archéologique de Picardie, n° spécial 30). Senlis: Société des antiquités historiques de Picardie.



- Meadow, R.H. 1999. The Use of Size Index Scaling Techniques for Research on Archaeozoological Collections from the Middle East. In: C. Becker, H. Manhart, J. Peters & J. Schibler, eds. *Historia Animalium ex Ossibus. Beiträge zur Paläoanatomie, Archäologie, Ägyptologie, Ethnologie und Geschichte der Tiermedizin*. Rahden/ Westfalen: Marie Leidorf, pp. 285–300.
- Méniel, P. 1996. Importation de grands animaux romains et amélioration du cheptel à la fin de l'âge du Fer en Gaule Belgique. *Revue Archéologique de Picardie*, 3: 113–22.
- Nuviala, P. 2014. La méthode des contours sur les dents de bœufs: perspectives de recherche pour l'étude des variations morphologiques. In: A. Gardeisen & C. Chandezon, eds. *Équidés et bovidés de la Méditerranée antique, Actes du colloque « Équidés et bovidés de la Méditerranée antique » 26–28 avril 2012*. Arles: Publication de l'UMR 5140 du CNRS, pp. 285–94.
- Nuviala, P. 2015. La révolution zootechnique romaine et la diffusion des grands bœufs et des grands chevaux dans l'est de la Gaule (unpublished PhD dissertation, Université de Bourgogne, Dijon).
- Nuviala, P. 2016. La production des « grands bœufs » dans l'est de la Gaule: entre évolutions gauloises et influences romaines. In: F. Malrain & G. Blancquaert, eds. *Évolution des sociétés gauloises du Second âge du Fer, entre mutations internes et influences externes (Revue Archéologique de Picardie, n° spécial 30)*. Senlis: Société des antiquités historiques de Picardie, pp. 597–609.
- Ouzoulias, P. & Tranoy, L. eds. 2010. *Comment les Gaules devinrent romaines*. Paris: La Découverte.
- Paunier, D. 2006. Celtes et Gaulois, l'archéologie face à l'histoire, 5 : la romanisation et la question de l'héritage celtique (Actes de la table ronde de Lausanne, juin 2005) (Bibracte 12/5). Glux-en-Glenne: Centre archéologique européen.
- Reddé, M., Barral, P. & Favory, F. eds. 2011. *Aspects de la romanisation dans l'Est de la Gaule*. Glux-en-Glenne: Bibracte, Centre archéologique européen.
- Rieckhoff, S. 2006. Celtes et Gaulois, l'archéologie face à l'histoire, 1 : Celtes et Gaulois dans l'Histoire, l'historiographie et l'idéologie moderne (Actes de la table ronde de Leipzig, juin 2005) (Bibracte 12/1). Glux-en-Glenne: Centre archéologique européen.
- Schlumbaum, A., Stopp, B., Breuer, G., Rehazek, A., Blatter, R., Turgay, M. & Schibler, J. 2003. Combining Archaeo-zoology and Molecular Genetics: The Reason behind the Changes in Cattle Size between 150 BC and 700 AD in Northern Switzerland. *Antiquity Project Gallery Issue 298*, December 2003. Available online at: <http://antiquity.ac.uk/projgall/schlumbaum298/> [accessed 15 August 2016].
- Schlumbaum, A., Turgay, M. & Schibler, J. 2006. Near East mtDNA Haplotype Variants in Roman Cattle from Augusta Raurica, Switzerland, and in the Swiss Evolène breed. *Animal Genetics*, 37: 373–75.
- Simpson, G.G. 1941. Explanation of Ratio Diagrams: Large Pleistocene Felines of North America. *American Museum Novitates*, 1136: 23–25.
- Szabó, M. 2006. Celtes et Gaulois, l'archéologie face à l'histoire, 3 : les Civilisés et les Barbares (du Ve au IIe siècle avant J.-C.) (Actes de la table ronde de Budapest, juin 2005) (Bibracte 12/3). Glux-en-Glenne: Centre archéologique européen.

Vitali, D. 2006. Celtes et Gaulois, l'archéologie face à l'histoire, 2 : la préhistoire des Celtes (Actes de la table ronde de Bologne- Monterezenzio, mai 2005) (Bibracte 12/2). Glux-en-Glenne: Centre archéologique européen.

von den Driesch, A. 1976. A Guide to Measurement of Animal Bones from Archaeological Sites (Peabody Museum Bulletin 1). Cambridge (MA): Peabody Museum of Archaeology and Ethnology.

Woolf, G. 2000. *Becoming Roman: The Origins of Provincial Civilization in Gaul*. Cambridge: Cambridge University Press.

Woolf, G. 2001. The Roman Cultural Revolution in Gaul. In: S.J. Keay & N. Terrenato, eds. *Italy and the West: Comparative Issues in Romanization*. Oxford: Oxbow Books, pp. 173–86.

## SUPPLEMENTARY MATERIAL

List of sites included in the study.

Results from the statistical tests carried out on pig LSI data grouped by century.

Results from the statistical tests carried out on cattle LSI data grouped by century. Isotopic results from the sequential sampling of LEV Sus 15 canine.

To view supplementary material for this article, please visit <https://doi.org/10.1017/ eaa.2016.10>.

## BIOGRAPHICAL NOTES

Delphine Frémondeau is a post-doctoral researcher at the Center for Archaeological Sciences (CAS, KU Leuven, Belgium), specialist in the stable isotope analysis of faunal remains. Her PhD, completed in 2012 at the Muséum national d'Histoire naturelle in Paris, aimed at establishing a protocol of stable isotope analysis for the study of the seasonal rhythms of pig husbandry in the past. Her main research interests are husbandry practices, land use, and food supply in urbanized societies.

Address: CAS, Geo Instituut, KU Leuven, Celestijnenlaan 200E, bus 2408, 3001 Leuven, Belgium. [email: [delphine.fremondeau@kuleuven.be](mailto:delphine.fremondeau@kuleuven.be)]

Pauline Nuviala is a zooarchaeologist, affiliated to the ArTeHis laboratory (UMR 6298, Dijon). Her main area of interest is the husbandry techniques from the Gallic period to the Roman period onwards, a research topic that she has been working on since her PhD, completed in 2015 at the University of Bourgogne. Favouring an interdisciplinary approach, she is notably using morphometrics and isotopic analysis.

Address: CNRS, UMR 6298 ArTeHiS, Université de Bourgogne, 6 bld Gabriel, 21000 Dijon, France.  
[email: p.nuviala@ gmail.com]

Colin Duval is a French zooarchaeologist, working at the Université François Rabelais (Tours) and the Muséum national d'Histoire naturelle (Paris). His PhD, completed in 2015, examines the economic and agro-pastoral changes that occurred in Gaul between the Iron Age and the Roman period. This research mainly focuses on the evolution of pig and cattle morphology in western Europe (using both osteometry and geometric morphometrics).

Address: UMR 7324, Laboratoire Archéologie et Territoires, Université François-Rabelais, MSH Villes et Territoires, USR 3501 - BP 60449, 33 allée Ferdinand de Lesseps, 37204, Tours, Cedex 03, France.  
[email: duval.colin@ gmail.com]

Le porc et le bœuf en Gaule : le rôle des sociétés gauloises dans l'évolution des pratiques d'élevage

Pour mieux comprendre les changements dans les pratiques d'élevage survenus entre l'âge du Fer et la période romaine en Gaule, cet article présente l'étude combinée de l'évolution morphologique des porcs et des bœufs et de la gestion démographique des porcs du site de Levroux Les Arènes (Indre, France). Les résultats indiquent la mise en place, sans doute avant le deuxième siècle av. J.-C., de nouveaux systèmes de production et de distribution, et l'existence d'une sélection des porcs abattus à Levroux Les Arènes en fonction d'un format - ou d'un poids – pour une production spécialisée de viande. En Gaule, la taille des porcs et des bœufs évolue progressivement à partir de la fin du troisième siècle av. J.-C. Ces changements résultent vraisemblablement d'une évolution interne aux sociétés gauloises, basée sur des cheptels locaux, mais potentiellement en réponse à des changements économiques plus globaux. Au sein de l'Empire Romain d'Occident, chaque province, ainsi que l'Italie elle-même, semble suivre son propre schéma évolutif, qui en outre diffère entre porcs et bovins, ce qui suggère que chaque région a pu adapter ses stratégies d'élevage en accord avec ses propres caractéristiques agro-pastorales, ses capacités ou ses ambitions.

Mots-clés: bœuf, porc, morphométrie, analyse des isotopes stables, âge du Fer, époque romaine, élevage, Gaule

Schwein und Rind in Gallien: die Rolle der gallischen Gemeinschaften in der Entwicklung der Tierhaltung

In diesem Artikel wird eine vergleichende Untersuchung der Morphologie von Schweinen und Rinder vorgelegt sowie eine Analyse stabiler Isotopen, die die Führung des Schweinebestands in Levroux Les

Arènes (Indre, Frankreich) betrifft, um die Veränderungen in den Tierhaltungspraktiken zwischen der Eisenzeit und der Römerzeit in der gallischen Gesellschaft auszuwerten. Unsere Ergebnisse zeigen, dass neue Produktionsstrukturen und Vertriebssysteme wahrscheinlich vor dem 2. Jahrhundert v. Chr. eingeführt wurden, und dass bestimmte Größen/Gewichte in der spezialisierten Erzeugung von Schweinefleisch ausgesucht wurden. Die Statur der Schweine und Rinder nimmt zunehmend vom späten 3. Jahrhundert an. Diese Veränderungen sind wahrscheinlich auf eine Entwicklung innerhalb der gallischen Gesellschaft zurückzuführen, die sich auf einheimische Herden stützte. Vielleicht waren sie aber auch eine Reaktion zu Umwandlungen im weiteren wirtschaftlichen Umfeld. Im westlichen Römerreich folgt jede Provinz, und auch Italien, ihren eigenen Entwicklungsprozess, der sich für Schweine und Rinder unterschiedlich entfaltet; dies deutet darauf hin, dass jedes Gebiet seine Tierhaltungsstrategie je nach seinen agropastoralen Eigenschaften, Leistungsfähigkeiten und Ziele angepasst hatte. Translation by Madeleine Hummler

Stichworte: Rind, Schwein, Morphometrie, Analyse stabiler Isotopen, Eisenzeit, Römerzeit, Tierhaltung, Gallien