

Resilience and food security: rethinking an ecological concept

Article

Accepted Version

Bullock, J. M., Dhanjal-Adams, K. L., Milne, A., Oliver, T. H. ORCID: https://orcid.org/0000-0002-4169-7313, Todman, L. C. ORCID: https://orcid.org/0000-0003-1232-294X, Whitmore, A. P. and Pywell, R. F. (2017) Resilience and food security: rethinking an ecological concept. Journal of Ecology, 105 (4). pp. 880-884. ISSN 0022-0477 doi: https://doi.org/10.1111/1365-2745.12791 Available at https://centaur.reading.ac.uk/69863/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

To link to this article DOI: http://dx.doi.org/10.1111/1365-2745.12791

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the <u>End User Agreement</u>.

www.reading.ac.uk/centaur



CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Resilience and food security: rethinking an ecological concept

James M. Bullock¹, Kiran L. Dhanjal-Adams¹, Alice Milne², Tom H. Oliver³, Lindsay C. Todman², Andrew P. Whitmore² and Richard F. Pywell¹

¹NERC Centre for Ecology and Hydrology, Benson Lane, Wallingford, OX10 8BB, UK

²Rothamsted Research, Harpenden AL5 2JQ, UK

³School of Biological Sciences, Harborne Building, University of Reading, Whiteknights, Reading, RG6 6AS, UK Summary

- 1. Focusing on food production, in this paper we define resilience in the food security context as *maintaining production of sufficient and nutritious food in the face of chronic and acute environmental perturbations*. In agri-food systems, resilience is manifest over multiple spatial scales: field, farm, regional and global. Metrics comprise production and nutritional diversity as well as socio-economic stability of food supply.
- Approaches to enhancing resilience show a progression from more ecologically-based methods at small scales to more socially-based interventions at larger scales. At the field scale, approaches include the use of mixtures of crop varieties, livestock breeds and of forage species, polycultures, and boosting ecosystem functions. Stress-tolerant crops, or with greater plasticity, provide technological solutions.
- 3. At the farm scale, resilience may be conferred by diversifying crops and livestock and by farmers implementing adaptive approaches in response to perturbations. Biodiverse landscapes may enhance resilience, but the evidence is weak. At regional to global scales, resilient food systems will be achieved by coordination and implementation of resilience approaches among farms, advice to farmers and targeted research.
- Synthesis. Threats to food production are predicted to increase under climate change and land degradation.
 Holistic responses are needed that integrate across spatial scales. Ecological knowledge is critical, but should be implemented alongside agronomic solutions and socio-economic transformations.

Resilience is a multi-faceted concept for food security

The resilience concept has been developed in many disciplines, including ecology, engineering, agriculture and economics (Altieri *et al.* 2015; Martin & Sunley 2015; Oliver *et al.* 2015), with much cross-fertilization of ideas. There is ongoing debate about precise definitions (Newton 2016). We do not aim to add to this debate, but to consider a pragmatic application of the resilience concept to food security. In ecology, resilience generally refers to the ability of an ecological system to resist a change in state or to recover in response to perturbations; but numerous state variables are used, including population size, community composition and ecological functions (Oliver *et al.* 2015). With regard to food security, resilience has been conceptualised in multiple ways, e.g.: a) in terms of international development, by considering social structures and capacity building (Bene *et al.* 2016; Pelletier *et al.* 2016); b) by analysing sensitivity in models linking human population growth to food supply (Suweis *et al.* 2015); and c) maintaining agricultural production under climate change (Altieri *et al.* 2015).

In this mini-review we explore this subject in terms of approaches for resilient food production in agricultural systems. We focus on multi-scale and multi-disciplinary issues to provide fresh view of how food production might become more resilient. Taking an agro-ecological perspective, we implement a working definition of resilience in terms of food security as: *maintaining production of sufficient and nutritious food in the face of chronic and acute environmental perturbations*. Following Oliver *et al.* (2015), a quantification of resilience would therefore represent the amount of time that food production is below a "sufficient and nutritious" threshold, reflecting Target 2 of the Sustainable Development Goals. Our definition overlaps with that of *food system* resilience by Tendall *et al.* (2015): the capacity over time of a food system and its units at multiple levels, to provide sufficient, appropriate and accessible food to all, in the face of various and even unforeseen disturbances. While the definition of Tendall *et al.* (2015) includes aspects of demand – distribution, access and utilisation – we focus on the production side of food security. While related concepts such as stability, robustness and vulnerability can be studied individually (Urruty, Tailliez-Lefebvre & Huyghe 2016), they are subtly different metrics of dynamical systems, and resilience provides a pragmatic and overarching structure (Fig. 1). For example, instability of food supply can indicate low resilience (Suweis *et al.* 2015). Indeed, to apply generally to

food security, resilience concepts must combine considerations of production responses to ongoing fluctuations in drivers, large perturbations or 'shocks' and long-term changes in these drivers (Fig. 1).

Food quantity, in terms of absolute amount of production or food calories, is the usual metric of food security and thus resilience. The need for varied and nutritious diets, to avoid both malnutrition and obesity, means nutritional diversity should also be factored into resilience considerations (Khoury *et al.* 2014; De Keersmaecker *et al.* 2016). Nutritional diversity and sufficient micronutrients are generally supplied through a varied diet, especially by including foodstuffs other than the cereal staples. Furthermore, relevant measures of resilience may also relate to economic and social aspects of food systems at multiple scales (Bene *et al.* 2016).

Resilience is manifest over multiple spatial scales in agri-food systems

In agri-food systems, resilience is manifest over multiple spatial scales. Most agro-ecologists focus on crop and livestock production at small, e.g. field, scales, but resilience of food production should also consider the farm, regional and global scales (Fig. 2).

To inform approaches for enhancing resilience, we summarise the major factors affecting food production. Infield crop yields vary primarily in response to climatic fluctuations, with temperature and water availability being the most common drivers (Ray *et al.* 2015). Other environmental drivers of both the quantity and quality of crop yields include pests and pathogens, and increasingly important chronic factors include rising salinity, deteriorating soils, new pests and decreasing pollinators (Lal 2009; Jaggard, Qi & Ober 2010; Potts *et al.* 2016). Similarly, climate, disease, and poor forage quality are among the environmental factors that affect production in livestock systems (Nardone *et al.* 2010).

At larger spatial scales, these environmental drivers remain important, with their overall impact depending on their spatial and temporal patterns; for example el Niño has synchronous impacts on crop yields over large areas (lizumi *et al.* 2014). But, from the farm to global scales, socio-economic perturbations become important. Considering the supply-side of food security, these include changes in food needs, price volatilities for inputs, policy decisions, and scarcities in land, labour, finances and machinery (Rodriguez *et al.* 2011; Golub *et al.* 2013). For example, globalisation and urbanisation are suggested to have increased homogeneity of food supplies globally, with consequences for nutrition (Khoury *et al.* 2014).

Resilience at these different scales is not necessarily nested, meaning food security may not always require resilience at small scales; for example, poor production in a field, farm or region may be balanced by increased production elsewhere (Fig 1), so that there is resilience at larger scales (Reidsma *et al.* 2010). Furthermore, because food production has multiple drivers, some degree of resilience may arise through asynchrony or conflicts among these drivers (Fig. 1). For example, adverse weather which decreases crop growth may also limit pests (Caffarra *et al.* 2012). These multiple scales and the interacting forms of perturbation mean that approaches to improve resilience of food production should be multi-faceted. These approaches need to take account of both acute and chronic perturbations; for example, while production in one year may be greatly decreased due to an extreme weather event but recover the following year, chronic soil degradation may cause ongoing declines in production (Fig. 1). Complementary approaches to enhancing resilience should be considered at different scales, and we describe a progression from more ecologically-based interventions at small-scales to more socially-based approaches at larger scales (Fig. 2).

Approaches to enhancing resilience

At the field scale, ecologically-inspired approaches invoke the role of diversity at some taxonomic level, which is hypothesised to impart resilience through different taxa having complementary responses to perturbations. Thus, varietal mixtures of a crop can achieve greater yield resilience than mono-varietal plantings (Creissen, Jorgensen & Brown 2016). Yield shows greater resilience to climatic perturbations in more species-rich or more genetically diverse forage systems (Prieto *et al.* 2015), and it is suggested that stocking multiple livestock species might also enhance resilience (Duru & Therond 2015). While there is little evidence for such benefits from planting multiple crop species in polycultures (Thiessen Martens, Entz & Wonneck 2015), increased diversity of crop types in a rotation can enhance yield resilience (Gaudin *et al.* 2015). Biotech approaches involve breeding crop and forage varieties that are less affected by perturbations, through being more resistant to a specific driver,

4

such as drought (Birthal *et al.* 2015; Kissoudis *et al.* 2016), or having greater plasticity (Bloomfield, Rose & King 2014). Some crops, such as maize, exhibit local adaptation to more extreme climates (Butler & Huybers 2013), which provides a similar resource. A final set of approaches considers the agroecosystem more broadly, enhancing resilience of production by bolstering the resilience of key functions. Examples are: improving soil condition, such as by agroforestry or adding organic matter (Rivest *et al.* 2013; Zhang *et al.* 2016); or by maintaining wild bee diversity to provide a resilient pollination service (Potts et al. 2016).

At the farm scale, more diverse crop rotations can aid yield resilience (Davis *et al.* 2012). More generally, adaptive responses by the farmer to perturbations provide an integrated way to manage for resilience (Reidsma *et al.* 2010; Rodriguez *et al.* 2011). This involves flexibility in crop, forage or livestock selection and husbandry and implementing new technologies in response to changing conditions. For example, Birthal *et al.* (2015) suggest rice yields in India are becoming resilient to increased drought frequency through farmers selecting appropriate varieties, expanding irrigation and utilising technologies such as micro-irrigation. Some aspects of traditional farming may confer resilience; such as the raised bed cultivation systems used in Latin American seasonally-flooded savannahs (Altieri *et al.* 2015). But reinstating traditional farming should not be seen as a panacea as such systems do not necessarily have higher resilience (De Keersmaecker *et al.* 2016), and it is likely they are poorly equipped to cope with climate and other enviornmental changes. Similarly, while it is hypothesised that biodiverse landscapes in and around farms enhance agroecosystem resilience (Tscharntke *et al.* 2005), there is little evidence.

At regional to global scales, approaches to resilient food production should ensure the approaches detailed for field to farm scales are implemented widely in a coordinated fashion. While policies across the world have tended to erode farm system resilience (Anderies, Ryan & Walker 2006), new policies such as the EU's crop diversification measure could be beneficial (Mahy *et al.* 2015). Knowledge transfer to and among farmers, building capacity and enhancing social networks may be more effective approaches, allowing farmers to self-organise to address resilience issues in an adaptive fashion (Carlisle 2014; Pelletier *et al.* 2016), and this can be addressed by policies

5

(Anderies *et al.* 2013). Many national and international institutional structures have complex and interacting consequences for food production resilience, through impacts on resources, costs and collective actions; and there is a need for these institutions to adapt and transform to new types of perturbation (Anderies *et al.* 2013). Ecological and agronomic research institutions can help, for example, by developing new crop types (Kissoudis *et al.* 2016) and elucidating links between biodiversity and agroecosystem resilience (Oliver *et al.* 2015).

Enhancing resilience in terms of food security requires holistic, multi-scale actions

Threats to food security and the resilience of agri-food systems are predicted to increase under climate change and ongoing land degradation. Ecological principles inform responses to enhance the resilience of agricultural production at field and farm scales, generally through diversity-resilience relationships. One challenge for ecologists is to provide better evidence for such relationships; for example, whether landscape-scale biodiversity confers agroecosystem resilience. Ecologists might also work with plant breeders in developing crops and varieties showing complementary response to perturbations and in designing resilient mixtures and rotations. The approaches to resilience that we detail mostly concern agricultural production as there is little relevant research on nutritional diversity. This is a clear knowledge gap and, for example, Haddad *et al.* (2016) call for research into yield resilience of neglected nutritious fruits and vegetables. Ultimately, there is a need to merge ecological approaches into a holistic interpretation of resilience that considers both production and nutritional diversity at multiple scales, with socially-based approaches from the farm up to the global food system. In this way, ecologists can be active in the design of adaptive agri-food systems based on socio-ecological evidence that can help in enhancing resilience of food security.

References

Altieri, M.A., Nicholls, C.I., Henao, A. & Lana, M.A. (2015) Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*, **35**, 869-890.

Anderies, J.M., Folke, C., Walker, B. & Ostrom, E. (2013) Aligning key concepts for global change policy: robustness, resilience, and sustainability. *Ecology and Society*, **18**, 16.

6

- Anderies, J.M., Ryan, P. & Walker, B.H. (2006) Loss of resilience, crisis, and institutional change: Lessons from an intensive agricultural system in southeastern Australia. *Ecosystems*, **9**, 865-878.
- Bene, C., Headey, D., Haddad, L. & von Grebmer, K. (2016) Is resilience a useful concept in the context of food security and nutrition programmes? Some conceptual and practical considerations. *Food Security*, 8, 123-138.
- Birthal, P.S., Negi, D.S., Khan, M.T. & Agarwal, S. (2015) Is Indian agriculture becoming resilient to droughts? Evidence from rice production systems. *Food Policy*, **56**, 1-12.
- Bloomfield, J.A., Rose, T.J. & King, G.J. (2014) Sustainable harvest: managing plasticity for resilient crops. *Plant Biotechnology Journal*, **12**, 517-533.
- Butler, E.E. & Huybers, P. (2013) Adaptation of US maize to temperature variations. *Nature Climate Change*, **3**, 68-72.
- Caffarra, A., Rinaldi, M., Eccel, E., Rossi, V. & Pertot, I. (2012) Modelling the impact of climate change on the interaction between grapevine and its pests and pathogens: European grapevine moth and powdery mildew. *Agriculture, Ecosystems & Environment,* **148**, 89-101.
- Carlisle, L. (2014) Diversity, flexibility, and the resilience effect: lessons from a social-ecological case study of diversified farming in the northern Great Plains, USA. *Ecology and Society*, **19**.
- Creissen, H.E., Jorgensen, T.H. & Brown, J.K.M. (2016) Increased yield stability of field-grown winter barley (Hordeum vulgare L.) varietal mixtures through ecological processes. *Crop Protection*, **85**, 1-8.
- Davis, A.S., Hill, J.D., Chase, C.A., Johanns, A.M. & Liebman, M. (2012) Increasing cropping system diversity balances productivity, profitability and environmental health. *PLoS ONE*, **7**.
- De Keersmaecker, W., van Rooijen, N., Lhermitte, S., Tits, L., Schaminee, J., Coppin, P., Honnay, O. & Somers, B.
 (2016) Species-rich semi-natural grasslands have a higher resistance but a lower resilience than intensively managed agricultural grasslands in response to climate anomalies. *Journal of Applied Ecology*, 53, 430-439.

- Duru, M. & Therond, O. (2015) Livestock system sustainability and resilience in intensive production zones: which form of ecological modernization? Regional Environmental Change, 15, 1651-1665.
- Gaudin, A.C.M., Tolhurst, T.N., Ker, A.P., Janovicek, K., Tortora, C., Martin, R.C. & Deen, W. (2015) Increasing crop diversity mitigates weather variations and improves yield stability. *PLoS ONE*, **10**.
- Golub, A.A., Henderson, B.B., Hertel, T.W., Gerber, P.J., Rose, S.K. & Sohngen, B. (2013) Global climate policy impacts on livestock, land use, livelihoods, and food security. *Proceedings of the National Academy of Sciences*, **110**, 20894-20899.
- Haddad, L., Hawkes, C., Webb, P., Thomas, S., Beddington, J., Waage, J. & Flynn, D. (2016) A new global research agenda for food. *Nature*, 540, 30-32.
- Iizumi, T., Luo, J.-J., Challinor, A.J., Sakurai, G., Yokozawa, M., Sakuma, H., Brown, M.E. & Yamagata, T. (2014)
 Impacts of El Niño Southern Oscillation on the global yields of major crops. *Nature Communications*, 5, 3712.
- Jaggard, K.W., Qi, A. & Ober, E.S. (2010) Possible changes to arable crop yields by 2050. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 2835-2851.
- Khoury, C.K., Bjorkman, A.D., Dempewolf, H., Ramirez-Villegas, J., Guarino, L., Jarvis, A., Rieseberg, L.H. & Struik, P.C. (2014) Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences*, **111**, 4001-4006.
- Kissoudis, C., van de Wiel, C., Visser, R.G.F. & van der Linden, G. (2016) Future-proof crops: challenges and strategies for climate resilience improvement. *Current Opinion in Plant Biology*, **30**, 47-56.
- Lal, R. (2009) Soils and food sufficiency. A review. Agronomy for Sustainable Development, 29, 113-133.
- Mahy, L., Dupeux, B., Van Huylenbroeck, G. & Buysse, J. (2015) Simulating farm level response to crop diversification policy. *Land Use Policy*, **45**, 36-42.
- Martin, R. & Sunley, P. (2015) On the notion of regional economic resilience: conceptualization and explanation. *Journal of Economic Geography*, **15**, 1-42.

Nardone, A., Ronchi, B., Lacetera, N., Ranieri, M.S. & Bernabucci, U. (2010) Effects of climate changes on animal production and sustainability of livestock systems. *Livestock Science*, 130, 57-69.

Newton, A.C. (2016) Biodiversity risks of adopting resilience as a policy goal. Conservation Letters, 9, 369-376.

- Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, D.L., Petchey, O.L., Proenca, V., Raffaelli, D., Suttle, K.B., Mace, G.M., Martin-Lopez, B., Woodcock, B.A. & Bullock, J.M. (2015) Biodiversity and resilience of ecosystem functions. *Trends in Ecology & Evolution*, **30**, 673-684.
- Pelletier, B., Hickey, G.M., Bothi, K.L. & Mude, A. (2016) Linking rural livelihood resilience and food security: an international challenge. *Food Security*, **8**, 469-476.
- Potts, S.G., Imperatriz-Fonseca, V., Ngo, H.T., Aizen, M.A., Biesmeijer, J.C., Breeze, T.D., Dicks, L.V., Garibaldi, L.A., Hill, R., Settele, J. & Vanbergen, A.J. (2016) Safeguarding pollinators and their values to human wellbeing. *Nature*, 540, 220-229.
- Prieto, I., Violle, C., Barre, P., Durand, J.-L., Ghesquiere, M. & Litrico, I. (2015) Complementary effects of species and genetic diversity on productivity and stability of sown grasslands. *Nature Plants*, 1, 15033.
- Ray, D.K., Gerber, J.S., MacDonald, G.K. & West, P.C. (2015) Climate variation explains a third of global crop yield variability. *Nature Communications*, **6**.
- Reidsma, P., Ewert, F., Lansink, A.O. & Leemans, R. (2010) Adaptation to climate change and climate variability in European agriculture: The importance of farm level responses. *European Journal of Agronomy*, **32**, 91-102.
- Rivest, D., Lorente, M., Olivier, A. & Messier, C. (2013) Soil biochemical properties and microbial resilience in agroforestry systems: Effects on wheat growth under controlled drought and flooding conditions. *Science of the Total Environment*, **463**, 51-60.
- Rodriguez, D., deVoil, P., Power, B., Cox, H., Crimp, S. & Meinke, H. (2011) The intrinsic plasticity of farm businesses and their resilience to change. An Australian example. *Field Crops Research*, **124**, 157-170.

- Suweis, S., Carr, J.A., Maritan, A., Rinaldo, A. & D'Odorico, P. (2015) Resilience and reactivity of global food security. *Proceedings of the National Academy of Sciences of the United States of America*, **112**, 6902-6907.
- Tendall, D.M., Joerin, J., Kopainsky, B., Edwards, P., Shreck, A., Le, Q.B., Kruetli, P., Grant, M. & Six, J. (2015) Food system resilience: Defining the concept. *Global Food Security*, **6**, 17-23.
- Thiessen Martens, J.R., Entz, M.H. & Wonneck, M.D. (2015) Review: Redesigning Canadian prairie cropping systems for profitability, sustainability, and resilience. *Canadian journal of plant science*, **95**, 1049-1072.
- Tscharntke, T., Klein, A.M., Kruess, A., Steffan-Dewenter, I. & Thies, C. (2005) Landscape perspectives on agricultural intensification and biodiversity ecosystem service management. *Ecology Letters*, **8**, 857-874.
- Urruty, N., Tailliez-Lefebvre, D. & Huyghe, C. (2016) Stability, robustness, vulnerability and resilience of agricultural systems. A review. *Agronomy for Sustainable Development*, **36**.
- Zhang, X.B., Sun, N., Wu, L.H., Xu, M.G., Bingham, I.J. & Li, Z.F. (2016) Effects of enhancing soil organic carbon sequestration in the topsoil by fertilization on crop productivity and stability: Evidence from long-term experiments with wheat-maize cropping systems in China. *Science of the Total Environment*, **562**, 247-259.

Fig 1. Types of perturbation and forms of resilience in agri-food systems. Relevant perturbations comprise: ongoing fluctuations such as annual climatic variations; chronic perturbations, which are long-term and potentially permanent changes in drivers, such as rising average temperature; and acute, i.e. short-term, perturbations such as extreme weather events. Resilience (for example, in yield) can be achieved through resisting or recovering from perturbations, in contrast to large, and sometimes enduring, losses in production. In addition, because food production is affected by multiple drivers, resilience may be delivered by opposing patterns in these drivers. Finally, while perturbations may affect production at a local scale, losses in some locations may be countered by enhanced yield at other locations, leading to resilience at larger scales.

Types of perturbations	Resilient responses		
Fluctuating	Direct		
≩ ^{high}	high Resistance high Recovery high Poor resilience		
which which have been a series of the series			
Time	Time Time Time		
Chronic	Opposing drivers		
Environmental duality why why why why why why why why why wh	high low Time Driver A Jow Time Driver B Jow Driver B Jow Time Driver B Jow Driver		
Acute	Scaling effects		
Euvironmental quality	High Location 1 Image Image		

Fig 2. Resilience in the production of sufficient and nutritious food is manifest at multiple scales. This conceptual diagram illustrates three broad scales and the multi-faceted approaches to conferring resilience discussed in the paper, which progress from more ecologically-based methods at smaller scales to more socially-based interventions at larger scales. A holistic approach to resilience involves working across these scales and combining ecological and social interventions.

Holistic integration	Global/regional scale Supportive policies Knowledge transfer Farmer-led initiatives Capacity building Institutional transformation Targeted research Diverse crop rotations Variety of livestock Adaptive management New technology Traditional farming Landscape biodiversity <u>Field scale</u> Varietal mixtures Multiple plant/animal species Stress-tolerant varieties Plastic varieties Pollinator diversity Soil management	
-------------------------	---	--