

# *Intraseasonal variability of air-sea fluxes over the Bay of Bengal during the southwest monsoon*

Article

Supplemental Material

Sanchez-Franks, A., Kent, E. C., Matthews, A. J., Webber, B. G. M., Peatman, S. C. and Vinayachandran, P. N. (2018) Intraseasonal variability of air-sea fluxes over the Bay of Bengal during the southwest monsoon. *Journal of Climate*, 31 (17). pp. 7087-7109. ISSN 1520-0442 doi: <https://doi.org/10.1175/JCLI-D-17-0652.1> Available at <https://centaur.reading.ac.uk/77492/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/JCLI-D-17-0652.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

Intraseasonal Variability of Air-Sea Fluxes over the Bay of Bengal during the  
Southwest Monsoon

**Supplementary materials**

**Table Captions:**

**Table. S1.** Summary statistics, mean, standard deviation (std), maximum and minimum for the meteorological and flux variables from the RAMA buoy, b28, and the reanalysis products, ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR. The values were computed for the summer (JJAS) from 2007 to 2015.

**Figure Captions:**

**Fig. S1.** Difference (a), correlation (b), variance ratio (c), and skill score (d) for reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) against data from the RAMA buoy at b27 (12°N and 90°E). The variables evaluated are the meteorological, SST (°C),  $V$  ( $\text{m s}^{-1}$ ),  $T_a$  (°C),  $q_a$  ( $\text{g kg}^{-1}$ ),  $\Delta T$  (°C),  $\Delta q$  ( $\text{g kg}^{-1}$ ), and fluxes,  $Q_{\text{SW}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{LW}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{SH}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{LH}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{net}}$  ( $\text{W m}^{-2}$ ), for summer (JJAS) from 2007 to 2015. Panel (a) shows a split scale to differentiate between meteorological and flux parameters.

**Fig. S2.** Difference (a), correlation (b), variance ratio (c), and skill score (d) for reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) against data from the RAMA buoy at b26 (8°N and 90°E). The variables evaluated are the meteorological, SST (°C),  $V$  ( $\text{m s}^{-1}$ ),  $T_a$  (°C),  $q_a$  ( $\text{g kg}^{-1}$ ),  $\Delta T$  (°C),  $\Delta q$  ( $\text{g kg}^{-1}$ ), and fluxes,  $Q_{\text{SW}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{LW}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{SH}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{LH}}$  ( $\text{W m}^{-2}$ ),  $Q_{\text{net}}$  ( $\text{W m}^{-2}$ ), for summer (JJAS) from 2007 to 2015. Panel (a) shows a split scale to differentiate between meteorological and flux parameters.

**Fig. S3.** The mean and standard deviation of the fluxes,  $Q_{\text{SH}}$  (a, f),  $Q_{\text{LH}}$  (b, g),  $Q_{\text{SW}}$  (c, h),  $Q_{\text{LW}}$  (d, i),  $Q_{\text{net}}$  (e, j) across an ensemble of the reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) for summer (JJAS) from 2007 to 2015.

**Fig. S4.** Mean  $Q_{\text{net}}$  ( $\text{W m}^{-2}$ ) for ERA-I (a), TropFlux (b), JRA-55 (c), MERRA-2 (d), and CFSR (e). All fields are averaged for the SW monsoon season (JJAS) from 2007 to 2015. The black square indicates the location of the RAMA buoy, b28, in the Bay of Bengal.

**Fig. S5.** RAMA b28 time series of  $V$  (a),  $Q_{\text{SW}}$  (b),  $Q_{\text{LH}}$  (c), and  $Q_{\text{NET}}$  (d) for SW monsoon season (JJAS) of 2015. Phases 2 to 8 of a single BSISO1 event are marked above panel (a), and phases 2 and 5 are highlighted in grey boxes.

| <b>Mean</b>      | <b>B28</b> | <b>ERA-I</b> | <b>TropFlux</b> | <b>JRA-55</b> | <b>MERRA2</b> | <b>CFSR</b> |
|------------------|------------|--------------|-----------------|---------------|---------------|-------------|
| SST              | 29.03      | 28.85        | 28.88           | 29.02         | 28.85         | 29.08       |
| V                | 7.92       | 7.82         | 8.21            | 7.55          | 8.02          | 8.74        |
| T <sub>a</sub>   | 28.56      | 28.13        | 28.55           | 28.38         | 28.41         | 28.70       |
| q <sub>a</sub>   | 20.99      | 20.20        | 21.30           | 20.49         | 20.50         | 20.16       |
| Q <sub>SW</sub>  | 176.23     | 172.25       | 162.25          | 172.02        | 147.81        | 247.67      |
| Q <sub>LW</sub>  | -34.02     | -36.74       | -27.79          | -43.85        | -40.33        | -42.83      |
| Q <sub>SH</sub>  | -5.10      | -10.75       | -4.74           | -24.30        | -4.78         | -3.66       |
| Q <sub>LH</sub>  | -104.94    | -146.36      | -97.94          | -154.90       | -147.29       | -167.02     |
| Q <sub>net</sub> | 31.61      | -21.60       | 31.77           | -51.03        | -44.6         | 34.15       |
| <b>STD</b>       | <b>B28</b> | <b>ERA-I</b> | <b>TropFlux</b> | <b>JRA-55</b> | <b>MERRA2</b> | <b>CFSR</b> |
| SST              | 0.52       | 0.58         | 0.60            | 0.49          | 0.55          | 0.52        |
| V                | 2.40       | 2.44         | 2.70            | 2.31          | 2.41          | 2.86        |
| T <sub>a</sub>   | 0.64       | 0.69         | 0.73            | 0.63          | 0.52          | 0.54        |
| q <sub>a</sub>   | 0.91       | 0.70         | 0.73            | 0.58          | 0.72          | 0.71        |
| Q <sub>SW</sub>  | 73.86      | 47.86        | 54.29           | 41.53         | 42.26         | 48.53       |
| Q <sub>LW</sub>  | 7.62       | 5.66         | 6.62            | 3.34          | 5.69          | 6.50        |
| Q <sub>SH</sub>  | 6.42       | 8.07         | 8.59            | 7.55          | 4.74          | 6.17        |
| Q <sub>LH</sub>  | 30.48      | 36.76        | 30.18           | 36.24         | 41.97         | 64.20       |
| Q <sub>net</sub> | 89.51      | 72.62        | 70.37           | 65.36         | 60.43         | 88.00       |
| <b>Max</b>       | <b>B28</b> | <b>ERA-I</b> | <b>TropFlux</b> | <b>JRA-55</b> | <b>MERRA2</b> | <b>CFSR</b> |
| SST              | 30.88      | 30.80        | 30.83           | 30.44         | 30.77         | 30.75       |
| V                | 13.40      | 17.96        | 18.95           | 14.40         | 15.18         | 18.27       |
| T <sub>a</sub>   | 30.26      | 30.42        | 30.93           | 30.24         | 30.06         | 30.68       |
| q <sub>a</sub>   | 24.10      | 22.93        | 24.30           | 22.74         | 23.35         | 23.27       |
| Q <sub>SW</sub>  | 295.56     | 283.95       | 280.99          | 280.61        | 272.36        | 298.68      |
| Q <sub>LW</sub>  | -18.81     | -20.22       | -11.12          | -34.79        | -26.5         | -21.93      |
| Q <sub>SH</sub>  | 10.04      | 6.55         | 8.09            | -4.08         | 7.68          | 11.75       |
| Q <sub>LH</sub>  | -26.99     | -57.74       | -33.71          | -70.85        | -42.41        | -31.25      |
| Q <sub>net</sub> | 175.07     | 153.34       | 193.50          | 109.22        | 149.04        | 190.50      |
| <b>Min</b>       | <b>B28</b> | <b>ERA-I</b> | <b>TropFlux</b> | <b>JRA-55</b> | <b>MERRA2</b> | <b>CFSR</b> |
| SST              | 27.49      | 26.83        | 26.89           | 28.06         | 27.41         | 27.48       |
| V                | 1.07       | 0.76         | 0.07            | 0.89          | 1.05          | 1.39        |
| T <sub>a</sub>   | 26.58      | 26.30        | 26.71           | 26.19         | 26.92         | 27.27       |
| q <sub>a</sub>   | 17.72      | 17.96        | 19.01           | 17.11         | 17.77         | 17.66       |
| Q <sub>SW</sub>  | 11.16      | 6.68         | 21.29           | 69.35         | 56.25         | 34.75       |
| Q <sub>LW</sub>  | -67.16     | -65.95       | -61.73          | -64.78        | -69.56        | -77.75      |
| Q <sub>SH</sub>  | -26.33     | -70.01       | -73.12          | -58.75        | -26.18        | -22.75      |
| Q <sub>LH</sub>  | -230.27    | -400.42      | -293.36         | -335.40       | -332.67       | -454.75     |
| Q <sub>net</sub> | -225.25    | -497.19      | -311.69         | -308.41       | -270.05       | -348.63     |

Table S1. Summary statistics, mean, standard deviation (std), maximum and minimum for the meteorological and flux variables from the RAMA buoy, b28, and the reanalysis products, ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR. The values were computed for the summer (JJAS) from 2007 to 2015.




Fig. S1. Difference (a), correlation (b), variance ratio (c), and skill score (d) for reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) against data from the RAMA buoy at b27 (12°N and 90°E). The variables evaluated are the meteorological, SST (°C), V (m s<sup>-1</sup>),  $T_a$  (°C),  $q_a$  (g kg<sup>-1</sup>),  $\Delta T$  (°C),  $\Delta q$  (g kg<sup>-1</sup>), and fluxes,  $Q_{SW}$  (W m<sup>-2</sup>),  $Q_{LW}$  (W m<sup>-2</sup>),  $Q_{SH}$  (W m<sup>-2</sup>),  $Q_{LH}$  (W m<sup>-2</sup>),  $Q_{net}$  (W m<sup>-2</sup>), for summer (JJAS) from 2007 to 2015. Panel (a) shows a split scale to differentiate between meteorological and flux parameters.




Fig. S2. Difference (a), correlation (b), variance ratio (c), and skill score (d) for reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) against data from the RAMA buoy at b26 (8°N and 90°E). The variables evaluated are the meteorological, SST (°C), V (m s<sup>-1</sup>),  $T_a$  (°C),  $q_a$  (g kg<sup>-1</sup>),  $\Delta T$  (°C),  $\Delta q$  (g kg<sup>-1</sup>), and fluxes,  $Q_{SW}$  (W m<sup>-2</sup>),  $Q_{LW}$  (W m<sup>-2</sup>),  $Q_{SH}$  (W m<sup>-2</sup>),  $Q_{LH}$  (W m<sup>-2</sup>),  $Q_{net}$  (W m<sup>-2</sup>), for summer (JJAS) from 2007 to 2015. Panel (a) shows a split scale to differentiate between meteorological and flux parameters.




Fig. S3. The mean and standard deviation of the fluxes,  $Q_{SH}$  (a, f),  $Q_{LH}$  (b, g),  $Q_{SW}$  (c, h),  $Q_{LW}$  (d, i),  $Q_{net}$  (e, j) across an ensemble of the reanalysis products (ERA-I, TropFlux, JRA-55, MERRA-2 and CFSR) for summer (JJAS) from 2007 to 2015.




Fig. S4. Mean  $Q_{net}$  ( $W m^{-2}$ ) for ERA-I (a), TropFlux (b), JRA-55 (c), MERRA-2 (d), and CFSR (e). All fields are averaged for the SW monsoon season (JJAS) from 2007 to 2015. The black square indicates the location of the RAMA buoy, b28, in the Bay of Bengal.






Fig. S5. RAMA b28 time series of  $V$  (a),  $Q_{SW}$  (b),  $Q_{LH}$  (c), and  $Q_{NET}$  (d) for SW monsoon season (JJAS) of 2015. Phases 2 to 8 of a single BSISO1 event are marked above panel (a), and phases 2 and 5 are highlighted in grey boxes.